




Contents lists available at ScienceDirect

Life Sciences

journal homepage: www.elsevier.com/locate/lifescie

Reactive oxygen species mediate adipocyte differentiation in mesenchymal stem cells

Yasunari Kanda^{a,c,*}, Takashi Hinata^a, Sang Won Kang^b, Yasuhiro Watanabe^a

^a Department of Pharmacology, National Defense Medical College, Saitama 359-8513, Japan

^b Center for Cell Signaling Research and Division of Molecular Life Science, Ewha Womans University, Seoul 120-750, Republic of Korea

^c Division of Pharmacology, National Institute of Health Sciences, Setagaya 158-8501, Japan

ARTICLE INFO

Article history:

Received 30 September 2010

Accepted 6 June 2011

Keywords:

Adipogenesis

CREB

Mesenchymal stem cells

NAD(P)H oxidase

Reactive oxygen species

ABSTRACT

Aims: Mesenchymal stem cells (MSC) have the potential to differentiate into various cell lineages, including adipocytes and osteoblasts. The formation of adipose tissue involves the commitment of MSC to the preadipocyte lineage and the differentiation of preadipocytes into mature adipocytes. In the present study, we investigated the involvement of reactive oxygen species (ROS) in adipocyte differentiation from MSC.

Main methods: ROS signaling was evaluated by the effects of antioxidant N-acetyl-L-cysteine (NAC) or shRNA against NAD(P)H oxidase in the multipotent mesenchymal stem cell line 10T1/2 cells. Intracellular ROS was measured using an H₂DCF dye.

Key findings: We found that NAC blocked adipocyte differentiation in MSC. An H₂DCF assay revealed that differentiation-inducing agents induced ROS generation. These data suggest that ROS is involved in adipocyte differentiation in MSC. Next, we examined the source of ROS. Knockdown of NAD(P)H oxidase 4 (Nox4) by RNA interference inhibited ROS production and adipocyte differentiation by differentiation-inducing agents. Furthermore, treatment with NAC blocked the transcriptional activation of CREB, and the expression of dominant-negative mutants of CREB inhibited adipocyte differentiation.

Significance: The findings suggest that the increase in the intracellular ROS level via Nox4 mediates adipocyte differentiation through CREB in MSC. This data will provide new insight into the drug development for obesity.

© 2011 Elsevier Inc. All rights reserved.

Introduction

Obesity is the major risk factor for the development of type 2 diabetes, heart disease, hypertension, and stroke (Rexrode et al., 1996). It is caused due to a positive energy balance in the body that results in an increase in adipose tissue by an increase in either the number or the size of adipocytes (Spiegelman and Flier, 2001). Adipocytes could arise from mesenchymal stem cells (MSC) that have the potential to differentiate into various cell lineages, including adipocytes and osteoblasts (Pittenger et al., 1999). Thus, further understanding of the mechanism underlying adipocyte differentiation from MSC might facilitate the development of novel targets for new drugs in order to modify the in vivo function of adipocytes.

As determined using the murine preadipocyte cell line 3T3-L1, adipocyte differentiation in both preadipocytes and mature adipocytes is a well-defined process that occurs via a series of transcriptional events that involve several key transcription factors. Differentiation is initiated by the induction of the CCAAT/enhancer-binding protein (C/EBP) β that in turn induces C/EBP α and nuclear

receptor peroxisome proliferator-activated receptor γ (PPAR γ) (Rosen et al., 2000). Once expressed, C/EBP α and PPAR γ enhance each other's production and are involved in terminal differentiation. In addition to the C/EBPs and PPAR γ , cAMP response element-binding protein (CREB) has also been implicated as an early regulator of the adipocyte differentiation program. A previous study showed that CREB overexpression in 3T3-L1 preadipocytes promotes the expression of adipocyte markers and the accumulation of triglycerides (Reusch et al., 2000). Adipogenic agents induced the phosphorylation of Ser133 CREB, thereby increasing its capacity to activate adipocyte differentiation-related transcription.

In contrast to differentiation of preadipocytes to mature adipocytes, adipocyte differentiation of MSC has not been entirely elucidated. Although 3T3-L1 preadipocytes are an excellent model cell system with which to elucidate the terminal adipocyte differentiation program, a more primitive cell line should be useful in investigating earlier mechanisms in adipocyte development. In particular, 10T1/2 cells, which are derived from C3H mouse embryos (Reznikoff et al., 1973), have been widely used as a mesenchymal stem cell line for studies on stem cell differentiation (Pinney and Emerson, 1989; Taylor and Jones, 1979).

Reactive oxygen species (ROS), such as superoxide anions (O₂⁻) and hydrogen peroxide (H₂O₂), are produced by the NAD(P)H oxidase

* Corresponding author at: 1-18-1, Kamiyoga, Setagaya 158-8501, Japan. Tel./fax: +81 3 3700 9704.

E-mail address: kanda@nihs.go.jp (Y. Kanda).

enzyme complex in phagocytic cells (e.g., macrophages and neutrophils) (Babior et al., 2002). Growing evidence suggests that ROS production is not restricted to phagocytic cells, and these species function as important secondary messengers in intracellular signal transduction pathways for several functions such as proliferation and differentiation (Sundaresan et al., 1995; Nishio and Watanabe, 1997; Sauer et al., 2000). Nonphagocytic ROS are produced by an enzyme that shows a high degree of homology with the phagocytic NAD(P)H oxidase and is now believed to belong to the Nox family, which comprises Nox1, Nox2 (gp91), Nox3, Nox4, and Nox5. ROS generation during adipogenesis in 3T3-L1 preadipocytes has already been reported (Lee et al., 2009). However, the mechanisms by which ROS affect MSC are largely unknown.

In the present study, we demonstrated that ROS mediate adipocyte differentiation in MSC. Furthermore, we show that the ROS generated via Nox4 play a key role in adipocyte differentiation.

Materials and methods

Materials

We obtained N-acetyl-L-cysteine (NAC) from Kanto Chemical Co. (Tokyo, Japan). 4'-hydroxy-3'-methoxyacetophenone (apocynin) was from Calbiochem (La Jolla, CA, USA). Anti-PPAR γ polyclonal antibody (H-100) was from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Anti-CREB polyclonal antibody was from Upstate Biotechnology (Charlottesville, VA, USA). Anti-Prx II polyclonal antibody was from LabFrontier (Seoul, Korea). All other reagents were of analytical grade and obtained from commercial sources.

MSC culture

Rat bone marrow-derived mesenchymal stem cells were isolated by the previous method with slight modifications (Azizi et al., 1998). Eight-week-old Sprague–Dawley rats (Tokyo Laboratory Animal Science, Tokyo, Japan) were sacrificed, and their femurs and tibiae were carefully cleaned of the adherent soft tissue. The marrow was harvested by inserting a syringe needle (23 gauge) into one end of the bone and flushing with Dulbecco's modified Eagle's medium (DMEM; Nissui Pharmaceutical Co., Ltd., Tokyo, Japan) and were seeded in DMEM containing 10% fetal bovine serum (FBS; JRH Biosciences, Lenexa, KS, USA), 100 U/ml penicillin (Gibco BRL), and 100 μ g/ml streptomycin (Gibco BRL). After the cells were allowed to adhere to culture dishes for 3 days, the nonadherent cells (such as hematopoietic stem cells) were removed by changing the medium and the adherent cells were maintained in culture. The medium was changed every 3 days. All the procedures involving animals were in accordance with the guidelines of National Defense Medical College. The 10T1/2 cell line (clone 8; Japanese Cancer Research Resources Bank) was cultured in basal medium Eagle (BME; Gibco BRL, Invitrogen Corp., Carlsbad, CA, USA) supplemented with 10% FBS, 2 mM L-glutamine, 100 U/ml penicillin, and 100 μ g/ml streptomycin.

Adipocyte differentiation

Adipocytes were differentiated as described previously with a slight modification (Cheng et al., 2003). Briefly, the cells were cultured to confluence in a normal growth medium. At 2 days post-confluence, the medium was changed to differentiation medium supplemented with 10% FBS, 5 μ g/ml insulin, 0.1 μ M dexamethasone, and 50 μ M indomethacin. After 3 days, the medium was changed back to the normal growth medium, and the cells were further cultured for 4 days.

Oil red-O staining

Oil red O-staining was carried out as described previously (Green and Kehinde, 1975). The cells were washed with PBS and fixed in 4% paraformaldehyde for 20 min. After they were washed twice with PBS, the cells were stained with 0.3% oil red-O solution (Sigma-Aldrich, St. Louis, MO, USA) at room temperature for 30 min and visualized by standard light microscopy (Olympus IX70; Olympus, Okaya, Japan). Staining was quantified by extracting oil red-O from the stained cells with isopropanol and determining the absorbance of the solution at 518 nm using a Hitachi U-3010 spectrophotometer (Tokyo, Japan).

Measurement of intracellular ROS

In a 96-well plate (Perkin-Elmer Life Sciences, Wellesley, MA, USA), cells were seeded at a density of 1×10^4 cells per well, and they were allowed to adhere overnight. The cells were then incubated at 37 °C in serum-free medium containing 10 μ M 2',7'-dichlorofluorescein (H₂DCF; Molecular Probes, Eugene, OR, USA). After they were washed with PBS, the cells were incubated in the differentiation medium. The change in fluorescence was measured using a Fusion™ Multilabel Reader (Perkin-Elmer) at excitation and emission wavelengths of 485 and 530 nm, respectively.

Cell lysis and immunoblotting

Cell lysates were prepared as described previously (Kanda and Watanabe, 2005). Briefly, the cells were lysed in a buffer [50 mM Tris (pH 7.4), 150 mM NaCl, 10 mM Na₄P₂O₇, 20 mM NaF, 2 mM EDTA, 1% Triton X-100, 1 mM phenylmethylsulfonyl fluoride (PMSF), 1 mM Na₃VO₄, and 10 μ g/ml aprotinin] on ice. After incubation on ice for another 30 min, the lysed cells were centrifuged at 4 °C for 20 min at 15000 \times g, and the supernatant was collected. The protein concentration was determined using the BCA Protein Assay Reagent Kit (Pierce, Rockford, IL, USA). Samples were analyzed by 10% sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS-PAGE), and they were electrophoretically transferred (15 V, 90 min) onto polyvinylidene difluoride (PVDF) membranes (Millipore, Bedford, MA, USA) by a semi-dry blotting apparatus (ATTO, Tokyo, Japan). The membranes were blocked in 5% skimmed milk in PBS containing 0.2% Tween 20 (PBS-T) for 1 h at room temperature and were incubated with specific antibodies overnight at 4 °C. The blots were then washed with PBS and incubated with peroxidase-conjugated secondary antibodies (1:1000; Cell Signaling Technology) for 1 h at room temperature. They were washed with PBS-T, and peroxidase activity was detected by measuring enhanced chemiluminescence (ECL Detection Kit; Amersham Pharmacia Biotech, Buckinghamshire, UK).

Plasmids and retrovirus

Plasmids encoding dominant-negative mutant of CREB were kindly provided by Dr. Goodman (Oregon Health and Science University, Portland, OR, USA). The plasmids were transfected into 10T1/2 cells by using FuGENE6 (Roche Diagnostics, Indianapolis, IN, USA), according to the manufacturer's protocol. Prx II retrovirus were previously reported (Choi et al., 2005).

Short hairpin RNA

pSuper.retro.neo (OligoEngine, Seattle, WA, USA) was used as the mammalian RNA interference expression vector. It was linearized with *Hind*III and *Bgl*II, and the annealed 64-nucleotide oligos (synthesized by Genosys) including the Nox4 sequence (GCTGTCCCTAAACGTCTACT) was ligated into the linearized vector. The resulting positive clones of the recombinant expression vector were identified by *Eco*RI and *Hind*III digestion. Cells transfected with

Lipofectamine 2000 were selected with G418 (1 mg/ml), and polyclonal cells were used in the experiments. Empty vector was used as the control.

RNA isolation and real-time PCR

Total cellular RNA was extracted using the ISOGEN reagent (Nippon Gene, Tokyo, Japan). Reverse transcription was performed in a 20- μ l reaction volume comprising 1 μ g total RNA, 1 \times reaction buffer, 0.5 mM dNTP mixture, 2.5 μ M oligo(dT)₂₀ primer, 40 U RNase OUT, and 200 U Superscript III reverse transcriptase (RT; Gibco). Real-

time Polymerase chain reaction (PCR) was performed with the cDNA samples and SYBR PCR mixture (Takara) using an ABI PRISM 7900 sequence detector (Applied Biosystems). The results are expressed in 2^{- Δ Ct} using β -actin as a reference. The sequences of the primers used in real-time PCR are as follows: rC/EBP α (forward, 5'-GAATCTCC-TAGTCTGGCTC-3'; reverse, 5'-GATGAGAACAGAACGAGTAC-3'); rC/EBP β (forward, 5'-GCCACGGACACCTTCGAGG-3'; reverse, 5'-CGGCTCCGCTTGAGCTGC-3'); rPPAR γ (forward, 5'-GCGGAGATCC-CAGTGATATC-3'; reverse, 5'-TCAGCGACTGGGACTTTTCT-3') r β -actin (forward, 5'-AGGGAATCGTCGCTGAC-3'; reverse, 5'-CGCTCATTGCC-GATAGTG-3'); mNox4 (forward, 5'-TGTTGGCCTAGGATTGTGT-3';




Fig. 1. ROS-mediated adipocyte differentiation in MSC. (A) Rat bone marrow-derived MSC were pretreated with or without NAC (5 mM), and then differentiation was induced for 14 days. The cells were fixed and stained with oil red-O. The oil red-O was extracted with isopropanol, and absorbance was measured at 518 nm. *P<0.05 vs diff. (B) Expression of adipogenesis markers were measured by real-time PCR using rat bone marrow-derived MSC with or without NAC. Open column indicates mRNA level at day 0. Closed column indicates mRNA level at day 2 (C/EBP β) or day 4 (C/EBP α and PPAR γ) in the presence of differentiation-inducing agents. *P<0.05 vs control day 2 or day 4. (C) 10T1/2 cells were pretreated with or without NAC (5 mM), and then differentiation was induced for 7 days. The cells were fixed and stained with oil red-O. The oil red-O was extracted with isopropanol, and absorbance was measured at 518 nm. *P<0.05 vs diff. (D) Expression of adipogenesis markers were measured by real-time PCR using 10T1/2 cells with or without NAC. Open column indicates mRNA level at day 0. Closed column indicates mRNA level at day 2 (C/EBP β) or day 4 (C/EBP α and PPAR γ) in the presence of differentiation-inducing agents. *P<0.05 vs control day 2 or day 4. (E) 10T1/2 cells were loaded with H₂DCF (10 μ M) and incubated with differentiation-inducing agents for the indicated times. *P<0.05 vs time 0. (F) 10T1/2 cells were pretreated with NAC (5 mM) for 30 min, loaded with H₂DCF, and incubated with differentiation-inducing agents for 30 min. Intracellular ROS production was detected by measuring the fluorescence of H₂DCF as described in the Materials and methods section. *P<0.05 vs diff.




Fig. 1 (continued).

reverse, 5'-AGGGACCTTCTGTGATCCTCG-3'); mNox2 (forward, 5'-ACTCCTGGGTCAGCACTGG-3'; reverse, 5'-GTTCTGTCCAGTTGTCTTCG-3'); mC/EBPα (forward, 5'-GAACAGCAACGAGTACCGGTA-3'; reverse, 5'-GCCATGGCCTTGACCAAGGAG-3'); mC/EBPβ (forward, 5'-CCAGAA-GAAGGTGGAGCAACTG-3'; reverse, 5'-TCGGGCAGCGTCTTGAAC-3'); mPPARγ (forward, 5'-AGTCTGCTGATCTGCGAGCC-3'; reverse, 5'-CTTTCTGTCAAGATCGCCC-3') mβ-actin (forward, 5'-ACCACACCTTCTA-CAATGAG-3'; reverse, 5'-AGCACAGCCTGGATGG-3').

PCR analysis

PCR was performed using the 2720 Thermal Cycler (Applied Biosystems, Foster City, CA) in a 100-μl reaction volume comprising 1 μl RT product, 1× PCR buffer, 0.5 μM each of the sense and antisense primers, and 2.5 U long and accurate Taq (Sigma). The following primers were used: Nox4 (forward, 5'-GAAGCCATTGAGGAGTCA-3'; reverse, 5'-GGGTCCACAGCAGAAAATC-3'). The cycling conditions were as

follows: 1 min at 94 °C, 30 cycles of 1 min at 94 °C, 1 min at 52 °C, and 1 min at 72 °C, followed by 7 min at 72 °C.

Luciferase assay

pCRE-Luc (Stratagene, La Jolla, CA, USA) was transfected into 10T1/2 cells by FuGENE6. The cells were lysed 2 days after transfection, and luciferase activity was determined using the luciferase assay system (Promega, Madison, WI, USA), according to the manufacturer's protocol. To normalize the transfection efficiency, the pRL-CMV plasmid was co-transfected in all the experiments.

Statistical analysis

Data represent the mean ± S.D. of at least three separate experiments. The mean values were compared with control by using an unpaired Student's *t* test using Microsoft Excel-based application. A value of *P* less than 0.05 was considered statistically significant.




Fig. 1 (continued).

Results

Effects of antioxidants on adipocyte differentiation

To investigate the role of ROS on adipocyte differentiation from MSC, we first tested the effects of an antioxidant on adipocyte differentiation. Since MSC in mice has been shown to be more difficult than that in rat, due to the unwanted growth of non-MSC during the expansion in culture (Lindolfo Da Silva and Nance, 2003), we used rat bone marrow-derived MSC. As shown in Fig. 1A, treatment with differentiation-inducing agents induced formation of fat droplets, which triglycerol is stained by oil red-O. The antioxidant NAC inhibited the lipid accumulation in the cells. To confirm the effect of NAC, we performed real-time PCR for C/EBP α , C/EBP β and PPAR γ , which are well-known adipocyte markers. As shown in Fig. 1B, NAC inhibited the induction of C/EBP β (at day2), C/EBP α and PPAR γ (at day4) in rat bone marrow-derived MSC with the differentiation-inducing agents. Treatment with NAC for 30 min had little effects on the expression of the markers. Since primary MSC are heterogeneous, we then used murine mesenchymal stem cell line 10T1/2, which is widely used as a model of adipogenesis from MSC and do not require post-confluence mitosis (Cho and Jefcoate, 2004), for further analysis of the differentiation mechanism from MSC. Consistent with the data using primary cells, similar inhibitory effects of NAC on differentiation were observed in 10T1/2 cells (Fig. 1C and D). Thus, the observed inhibitory effects of antioxidants on adipocyte differentiation in both primary and 10T1/2 cells suggest that ROS might function during the early stage of adipocyte differentiation in MSC.

ROS generation by differentiation-inducing agents

Next, we examined the level of ROS in 10T1/2 cells by using the ROS-reactive dye H₂DCF. As shown in Fig. 1E, treatment with differentiation-inducing agents for 30 min induced an increase in the fluorescence

intensity of H₂DCF. When the cells were treated with differentiation-inducing agents for 1 h, some cells underwent cell death (data not shown). Since this might be due to the cytotoxicity, ROS production was measured for 30 min. Moreover, this increase was abolished by NAC (Fig. 1F), suggesting that the fluorescence intensity was derived from ROS. NAC alone did not inhibit the basal intensity. These data suggest that ROS are generated after the induction of differentiation in 10T1/2 cells.

Effect of peroxiredoxin on adipocyte differentiation

To further confirm the involvement of ROS in adipocyte differentiation, we studied the effect of the overexpression of H₂O₂-scavenging enzyme peroxiredoxin II (Prx II) (Choi et al., 2005). As shown in Fig. 2A, retroviral transduction of Prx II inhibited lipid accumulation. Retrovirus against an empty vector did not affect lipid accumulation. Expression level of endogenous Prx II was not affected during differentiation. In addition, retroviral transduction of Prx II decreased the fluorescence intensity of H₂DCF (Fig. 2B), confirming the activity of Prx II. When 10T1/2 cells were treated with 100 μ M H₂O₂, the cells did not induce lipid accumulation (data not shown), suggesting that ROS is necessary, but not sufficient in 10T1/2 cells. In contrast to differentiation, cell growth was not inhibited by Prx II expression in the presence or absence of differentiation-inducing agents (Fig. 2C), suggesting that ROS are involved in differentiation, not proliferation.

Inhibition of NAD(P)H oxidase impairs adipocyte differentiation

Subsequently, we investigated the source of ROS in adipocyte differentiation. Because NAD(P)H oxidase is considered to generate low levels of ROS, we suspected that it plays a role in ROS production and adipocyte differentiation in 10T1/2 cells. To examine the involvement of NAD(P)H oxidase in adipocyte differentiation, we tested the effect of apocynin, a selective inhibitor of NAD(P)H oxidase, on adipocyte differentiation in 10T1/2 cells. As shown in Fig. 3A, apocynin abolished lipid accumulation. In contrast, L-NAME (an NO inhibitor) or allopurinol (a xanthine oxidase inhibitor) did not inhibit lipid accumulation. To further elucidate the role of NAD(P)H oxidase in adipocyte differentiation, we studied the expression level of the NAD(P)H oxidase family. Since NAD(P)H oxidase isoforms Nox2 and Nox4 are detected by real-time PCR in 10T1/2 cells, we checked their expression levels during differentiation. As shown in Fig. 3B, Nox4 expression was significantly decreased after differentiation. In contrast, Nox2 expression was not changed during differentiation. To assess the involvement of Nox4 in adipocyte differentiation, we silenced its expression by using the RNA interference. Nox4 knockdown was confirmed by RT-PCR and an H₂DCF assay (Fig. 3C, D). A significant decrease in lipid accumulation was observed in the Nox4-silenced cells, whereas normal lipid accumulation was observed in the cells that were transfected with a control vector (Fig. 3E). These data suggest that the ROS produced by Nox4 mediate adipocyte differentiation in 10T1/2 cells.

Effect of CREB on adipocyte differentiation

We examined the molecular mechanisms by which ROS regulate adipocyte differentiation. Since CREB is known to regulate adipocyte differentiation in 3T3-L1 preadipocytes (Reusch et al., 2000), we studied its effects on 10T1/2 cells. As shown in Fig. 4A, the expression of dominant-negative mutants of CREB (DN-CREB) inhibited lipid accumulation. Differentiation-inducing agents stimulated the transcriptional activation of CREB, while antioxidants (NAC or apocynin) suppressed the activation (Fig. 4B). Furthermore, treatment with H₂O₂ induced CREB transcriptional activation, confirming the presence of a signaling




Fig. 2. Effect of peroxiredoxin II on adipocyte differentiation in 10T1/2 cells. (A) 10T1/2 cells were transduced with control or peroxiredoxin II (Prx II) retrovirus and were then incubated with differentiation-inducing agents for 7 days. Expression of Prx II was analyzed by immunoblotting with anti-Prx II antibody (upper panel). After the cells were stained with oil red-O, the oil red-O was extracted with isopropanol, and absorbance at 518 nm was measured (lower panel). * $P < 0.05$ vs control diff. (B) 10T1/2 cells transduced with Prx II were loaded with H_2DCF and incubated with differentiation-inducing agents for 30 min. Intracellular ROS production was detected by measuring the fluorescence of H_2DCF as described in the Materials and methods section. * $P < 0.05$ vs control diff. (C) 10T1/2 cells transduced with control or Prx II retrovirus were incubated with or without differentiation-inducing agents. After 3 days, cell number was determined using a hemocytometer.

pathway from ROS to CREB. These data suggest that adipocyte differentiation via ROS is mediated by CREB in 10T1/2 cells.

Effect of MAPK on adipogenesis

Previous studies using 3T3-L1 preadipocytes have reported that MAPK plays a role in adipocyte differentiation (Sale et al., 1995). Therefore, we examined the contribution of the MAPK pathway to adipocyte differentiation in 10T1/2 cells. We tested the effect of U0126, which is a potent and selective inhibitor of MAPK/ERK kinase (MEK; an upstream ERK kinase). As shown in Fig. 5A, U0126 decreased lipid accumulation by approximately 35%. PD98059, another MEK inhibitor, yielded similar results. Differentiation-inducing agents induced ERK phosphorylation, while the MEK inhibitors completely inhibited the ERK phosphorylation (Fig. 5B). Since ROS have been shown to induce heparin-binding epidermal growth factor (EGF)-like growth factor (HB-EGF) production (Zhang et al., 2004), we studied the effects of AG1478 (a selective EGF receptor kinase inhibitor) and CRM197 (a toxin that inhibits HB-EGF production) on differentiation. Both inhibitors did not inhibit lipid accumulation. As shown in Fig. 5C, NAC inhibited the phosphorylation of ERK by differentiation-inducing agents, suggesting the presence of

a signaling pathway from ROS to ERK. These results suggest that the ROS-ERK pathway partially mediates adipocyte differentiation in 10T1/2 cells.

Discussion

In the present study, we demonstrated that ROS play an important role during the early stage of adipocyte differentiation in MSC. We showed that adipocyte differentiation is impaired in the presence of ROS scavengers or RNA interference against Nox4. Furthermore, we showed that CREB act downstream of ROS in adipocyte differentiation.

We showed that ROS are related to differentiation, not proliferation, by using the antioxidant NAC (Fig. 1) and Prx II retrovirus (Fig. 2). NAC inhibited the expression of adipocyte markers including C/EBP α , C/EBP β and PPAR γ . This is consistent with a previous report showing that ROS enhance C/EBP β activation in 3T3-L1 cells (Lee et al., 2009). Since H_2O_2 induced CREB transcriptional activation (Fig. 4B) and CREB has been shown to regulate C/EBP β (Reusch et al., 2000), it is possible that ROS act at an upstream of CREB, and subsequently induce C/EBP β expression in MSC. Inconsistent with our observations, recent study using the 3T3-L1 preadipocytes has shown that ROS stimulate mitotic clonal expansion during adipogenesis in 3T3-L1 preadipocytes (Lee et al., 2009). The




Fig. 3. Role of Nox4 in adipocyte differentiation in 10T1/2 cells. (A) 10T1/2 cells were pretreated with apocynin (1 μ M), allupurinol (50 μ M), L-NAME (50 μ M) or vehicle and incubated with differentiation-inducing agents for 7 days. After fixation, the cells were stained with oil red-O. The oil red-O was extracted with isopropanol, and absorbance at 518 nm was measured. * P <0.05 vs vehicle. (B) After 10T1/2 cells were incubated with differentiation-inducing agents for the indicated times, total RNA was isolated and Nox2 and Nox4 were amplified using specific primers. * P <0.05 vs day 0. (C) Forty-eight hours after the cells were transfected with shRNA against Nox4 or control shRNA, total RNA was isolated, and Nox4 was amplified using specific primers. (D) The shRNA-transfected cells were loaded with H₂DCF and incubated with differentiation-inducing agents for 30 min. Intracellular ROS production was detected by measuring the fluorescence of H₂DCF, as described in the Materials and methods section. (E) The shRNA-transfected cells were incubated with differentiation-inducing agents for 7 days. After the cells were fixed, they were stained with oil red-O. The oil red-O was extracted with isopropanol, and absorbance was measured at 518 nm. * P <0.05 vs control diff.

discrepancy might be due to the differences of cell type or the stage of differentiation. 10T1/2 cells that we used here do not require post-confluence mitosis, whereas 3T3-L1 cells require one or two cell divisions before differentiation (Cho and Jefcoate, 2004; Tang et al., 2003). In addition, since 10T1/2 cells are considered to be multi-potent and more primitive than 3T3-L1 preadipocytes. Taken together, ROS could act at not only the mitotic clonal expansion but also the early differentiation stage.

We demonstrated NAD(P)H oxidase as the source of ROS in MSC by a pharmacological approach. Among the NAD(P)H oxidase family, our data showed that Nox4 was down-regulated after induction of differentiation in MSC and Nox4 shRNA inhibited lipid accumulation (Fig. 3). Consistent with our results, several reports have shown that Nox4 is down-regulated in mature adipocytes (Mouche et al., 2007;

Schröder et al., 2009). Taken together, Nox4 is considered to play an important role in adipogenesis from MSC. Since Nox4 contains multiple transmembrane segments and has been reported to localize in the endoplasmic reticulum and nucleus (Chen et al., 2008; Kuroda et al., 2005), ROS generated via Nox4 could diffuse in the cells and function as secondary messengers. The intracellular localization of Nox4 in MSC remains to be determined.

We further demonstrated that the generated ROS induce differentiation via CREB in MSC. CREB has been identified as a transcription factor of adipocyte differentiation in 3T3-L1 preadipocytes (Reusch et al., 2000). Thus, it may also functional in MSC, which are more primitive than preadipocytes. Many CREB targets have already been identified (Shaywitz and Greenberg, 1999). MAPK is one of the candidates for differentiation. MAPK has been shown to induce




Fig. 4. Effects of antioxidants on transcriptional activation of CREB in 10T1/2 cells. (A) 10T1/2 cells were transfected with dominant-negative mutants of CREB (DN-CREB) or control vector. The cells were incubated with differentiation-inducing agents for 7 days. The cells were fixed and stained with oil red-O. The oil red-O was extracted with isopropanol, and absorbance was measured at 518 nm. *P<0.05 vs control. (B) 10T1/2 cells were transiently transfected with a CRE-luciferase construct. After pretreatment with or without ROS scavengers (NAC, 5 mM or Apocynin, 1 μM) for 30 min, the cells were incubated with differentiation-inducing agents or H₂O₂ (1 mM) for 1 h. Cell lysates were then analyzed for luciferase activity. pRL-CMV vector was co-transfected to normalize transfection efficiency. *P<0.05 vs diff.

adipocyte differentiation in 3T3-L1 cells and acts downstream of ROS (Sale et al., 1995). However, its effect is only limited in 10T1/2 cells (Fig. 5). These data suggest the presence of other CREB targets in MSC. PGC-1, a regulator of mitochondrial biogenesis, is another good candidate which acts at a downstream of CREB. Oxidative stress has been shown to activate the PGC-1 promoter via CREB in 10T1/2 cells (St-Pierre et al., 2006). We are currently conducting studies to examine the involvement of PGC-1 in adipocyte differentiation.

In addition to MSC, several reports demonstrate the role of ROS in other kinds of stem cells. For example, Nox1 stimulates RANKL-induced osteoclast differentiation in bone marrow monocyte-macrophage lineage cells (Lee et al., 2005). Ito et al. revealed that ROS determine the lifespan of hematopoietic stem cells (Ito et al., 2006). Thus, stem cell differentiation utilizes the common route of ROS generation, and the NAD(P)H oxidase family appears to be a key regulator in this differentiation. Future studies will determine the common mechanisms by which ROS initiate stem cell differentiation.

Conclusions

We demonstrated that ROS mediate adipocyte differentiation using rat bone marrow-derived mesenchymal stem cells and murine multipotent mesenchymal stem cell line 10T1/2 cells. In addition, we identified that NAD(P)H oxidase family Nox4 is responsible for production of ROS at the early stage of differentiation in 10T1/2 cells. Furthermore, we showed that CREB act downstream of ROS in adipocyte differentiation. Thus, Nox4–ROS–CREB signaling provides a




Fig. 5. Effects of kinase inhibitors on adipocyte differentiation in 10T1/2 cells. (A) 10T1/2 cells were pretreated with PD98059 (20 μM), U0126 (1 μM), AG1478 (10 μM), CRM197 (10 μM), or vehicle for 30 min and then incubated with differentiation-inducing agents for 7 days. After the cells were fixed, they were stained with oil red-O. The oil red-O was extracted with isopropanol, and absorbance was measured at 518 nm. (B) 10T1/2 cells were pretreated with MEK inhibitors (PD98059, 20 μM or U0126, 1 μM) or vehicle for 30 min and then treated with differentiation-inducing agents for 10 min. ERK phosphorylation was detected by immunoblotting with the anti-phospho-ERK or anti-ERK antibody. (C) 10T1/2 cells were pretreated with ROS scavengers (NAC, 5 mM or Apocynin, 1 μM) for 30 min or vehicle and then treated with differentiation-inducing agents for 10 min. ERK phosphorylation was analyzed by immunoblotting with the anti-phospho-ERK or anti-ERK antibody.

unique pathway to understand the early stage of adipocyte differentiation. The activity of Nox4 is a potential therapeutic target for anti-obesity drug discovery.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

Acknowledgments

We would like to thank Dr. Goodman, for providing the plasmids. This work was supported in part by a grant from the Smoking Research Foundation (Y.K. and Y.W.), a Grant-in-Aid for Scientific Research (No. 2179025, 23590322) from the Ministry of Education, Culture, Sports, Science, and Technology, Japan (Y.K.) and a Health and Labour Sciences Research Grant from the Ministry of Health, Labour and Welfare, Japan (Y.K.).

References

- Azizi SA, Stokes D, Augelli BJ, DiGirolamo C, Prockop DJ. Engraftment and migration of human bone marrow stromal cells implanted in the brains of albino rats—similarities to astrocyte grafts. *Proc Natl Acad Sci* 1998;95(7):3908–13.
- Babior BM, Lambeth JD, Nauseef W. The neutrophil NADPH oxidase. *Arch Biochem Biophys* 2002;397(2):342–4.
- Chen K, Kirber MT, Xiao H, Yang Y, Keaney Jr JF. Regulation of ROS signal transduction by NADPH oxidase 4 localization. *J Cell Biol* 2008;181(7):1129–39.
- Cheng SL, Shao JS, Charlton-Kachigian N, Loewy AP, Towler DA. MSX2 promotes osteogenesis and suppresses adipogenic differentiation of multipotent mesenchymal progenitors. *J Biol Chem* 2003;278(46):45969–77.
- Cho YC, Jefcoate CR. PPAR γ 1 synthesis and adipogenesis in C3H10T1/2 cells depends on S-phase progression but does not require mitotic clonal expansion. *J Cell Biochem* 2004;91(2):336–53.
- Choi MH, Lee IK, Kim GW, Kim BU, Han YH, Yu DY, et al. Regulation of PDGF signalling and vascular remodelling by peroxiredoxin II. *Nature* 2005;435(7040):347–53.
- Green H, Kehinde O. An established preadipose cell line and its differentiation in culture. II. Factors affecting the adipose conversion. *Cell* 1975;5(1):19–27.
- Ito K, Hirao A, Arai F, Takubo K, Matsuoka S, Miyamoto K, et al. Reactive oxygen species act through p38 MAPK to limit the lifespan of hematopoietic stem cells. *Nat Med* 2006;12(4):446–51.
- Kanda Y, Watanabe Y. Thrombin-induced glucose transport via Src-p38MAPK pathway in vascular smooth muscle cells. *Br J Pharmacol* 2005;146(1):60–7.
- Kuroda J, Nakagawa K, Yamasaki T, Nakamura K, Takeya R, Kuribayashi F, et al. The superoxide-producing NAD(P)H oxidase Nox4 in the nucleus of human vascular endothelial cells. *Genes Cells* 2005;10(12):1139–51.
- Lee NK, Choi YG, Baik JY, Han SY, Jeong D, Bae YS, et al. A crucial role for reactive oxygen species in RANKL-induced osteoclast differentiation. *Blood* 2005;106(3):852–9.
- Lee H, Lee YJ, Choi H, Ko EH, Kim JW. Reactive oxygen species facilitate adipocyte differentiation by accelerating mitotic clonal expansion. *J Biol Chem* 2009;284(16):10601–1–6–9.
- Lindolfo Da Silva M, Nance BN. Murine marrow-derived mesenchymal stem cell: isolation, *in vitro* expansion, and characterization. *Br J Haematol* 2003;123(4):702–11.
- Mouche S, Mkaddem SB, Wang W, Katic M, Tseng YH, Carnesecchi S, et al. Reduced expression of the NADPH oxidase NOX4 is a hallmark of adipocyte differentiation. *Biochim Biophys Acta* 2007;1773(7):1015–27.
- Nishio E, Watanabe Y. The involvement of reactive oxygen species and arachidonic acid in α 1-adrenoceptor-induced SMC proliferation and migration. *Br J Pharmacol* 1997;121(4):665–70.
- Pinney DF, Emerson Jr CP. 10T1/2 cells: an *in vitro* model for molecular genetic analysis of mesodermal determination and differentiation. *Environ Health Perspect* 1989;80:221–7.
- Pittenger MF, Mackay AM, Beck SC, Jaiswal RK, Douglas R, Mosca JD, et al. Multilineage potential of adult human mesenchymal stem cells. *Science* 1999;284(5411):143–7.
- Reusch JE, Colton LA, Klemm DJ. CREB activation induces adipogenesis in 3T3-L1 cells. *Mol Cell Biol* 2000;20(3):1008–20.
- Rexrode KM, Manson JE, Hennekens CH. Obesity and cardiovascular disease. *Curr Opin Cardiol* 1996;11(5):490–5.
- Reznikoff CA, Brankow DW, Heidelberger C. Establishment and characterization of a cloned line of C3H mouse embryo cells sensitive to postconfluence inhibition of division. *Cancer Res* 1973;33(12):3231–8.
- Rosen ED, Walkey CJ, Puigserver P, Spiegelman BM. Transcriptional regulation of adipogenesis. *Genes Dev* 2000;14(11):1293–307.
- Sale EM, Kinson PG, Sale GJ. Requirement of MAP kinase for differentiation of fibroblasts to adipocytes, for insulin activation of p90 S6 kinase and for insulin or serum stimulation of DNA synthesis. *EMBO J* 1995;14(4):674–84.
- Sauer H, Rahimi G, Hescheler J, Wartenberg M. Role of reactive oxygen species and phosphatidylinositol 3-kinase in cardiomyocyte differentiation of embryonic stem cells. *FEBS Lett* 2000;476(3):218–23.
- Schröder K, Wandzioch K, Helmcke I, Brandes RP. Nox4 acts as a switch between differentiation and proliferation in preadipocytes. *Arterioscler Thromb Vasc Biol* 2009;29(2):239–45.
- Shaywitz AJ, Greenberg ME. CREB: a stimulus-induced transcription factor activated by a diverse array of extracellular signals. *Annu Rev Biochem* 1999;68:821–61.
- Spiegelman BM, Flier JS. Obesity and the regulation of energy balance. *Cell* 2001;104(4):531–43.
- St-Pierre J, Drori S, Uldry M, Silvaggi JM, Rhee J, Jager S, et al. Suppression of reactive oxygen species and neurodegeneration by the PGC-1 transcriptional coactivators. *Cell* 2006;127(2):397–408.
- Sundaresan M, Yu ZX, Ferrans VJ, Irani K, Finkel T. Requirement for generation of H₂O₂ for platelet-derived growth factor signal transduction. *Science* 1995;270(5234):296–9.
- Tang QQ, Otto TC, Lane MD. Mitotic clonal expansion: a synchronous process required for adipogenesis. *Proc Natl Acad Sci* 2003;100(1):44–9.
- Taylor SM, Jones PA. Multiple new phenotypes induced in 10T1/2 and 3T3 cells treated with 5-azacytidine. *Cell* 1979;17(4):771–9.
- Zhang H, Chalothorn D, Jackson LF, Lee DC, Faber JE. Transactivation of epidermal growth factor receptor mediates catecholamine-induced growth of vascular smooth muscle. *Circ Res* 2004;95(10):989–97.

Role of $\alpha 7$ -Nicotinic Acetylcholine Receptor in Normal and Cancer Stem Cells

Waka Lin, Naoya Hirata, Yuko Sekino and Yasunari Kanda*

Division of Pharmacology, National Institute of Health Sciences, Setagaya 158-8501, Japan

Abstract: The $\alpha 7$ -nicotinic acetylcholine receptor ($\alpha 7$ -nAChR) is widely known as a neurotransmitter receptor in nervous systems. $\alpha 7$ -nAChR is also present in a variety of non-neuronal tissues, where it has been implicated in the regulation of essential cellular functions including proliferation, survival, differentiation and communication. We have recently found in breast cancer that $\alpha 7$ -nAChR is involved in the proliferation of cancer stem cells, which constitute a minor subpopulation responsible for tumor development and metastasis. Since growing evidence suggests that $\alpha 7$ -nAChR is present not only in mature tissues and organs but also in undifferentiated stem cells and progenitor cells, $\alpha 7$ -nAChR emerges as a key mediator in the regulation of self-renewal and differentiation. We provide here an overview of the recent works on the expression and function of $\alpha 7$ -nAChR in normal and cancer stem cells, and their relevance to disease-related cellular dysfunction. Understanding the role of $\alpha 7$ -nAChR in stem cells would be of great interest for its application potential in drug discovery and in regenerative medicine.

Keywords: $\alpha 7$ -nicotinic acetylcholine receptor, breast cancer, cancer stem cell, nicotine, stem cells.

INTRODUCTION

Acetylcholine (ACh) is known as one of the most important neurotransmitters in nervous systems. The effect of ACh is mediated through two types of membrane ACh receptors (AChR): the G protein-coupled muscarinic AChR and the nicotinic AChR (nAChR). nAChR are ligand-gated ion channels and can be activated by nicotine as agonist. They are plasma-membrane receptors composed of a hetero- or homo-pentamer of five subunits enclosing a central ion channel. The nAChR that contains $\alpha 7$ subunits ($\alpha 7$ -nAChR) presents the distinctive features to be homomeric and to be selectively permeable to Ca^{2+} [1]. The specific expression and function of $\alpha 7$ -nAChR can also be revealed by selective antagonists such as α -Bungarotoxin (α -BTX). $\alpha 7$ -nAChR is one of the dominant nAChR in brain and has been implicated in several nervous and cognitive functions of significance in neurodegenerative disorders.

Recent studies have also accumulated evidence that AChR are expressed and functional in a wide diversity of non-neuronal mammalian cells and tissues [2]. $\alpha 7$ -nAChR has been demonstrated to be involved in a number of basic cellular processes including proliferation, differentiation, survival and intercellular communication. There is growing interest in modulating the function of AChR, especially of $\alpha 7$ -nAChR, for the study and treatment of both neuronal and non-neuronal diseases. It is also especially noteworthy that $\alpha 7$ -nAChR has been suggested to be involved in the stimulation of cancer development, which provides a direct link between cancer risk and exposure to nicotine [3, 4]. nAChR were found to be expressed in human lung cancer and to regulate proliferation and apoptosis [5, 6]. Since then, $\alpha 7$ -nAChR has been implicated in various cellular processes regulating tumor growth and survival, as reviewed in other chapters of this issue of Current Drug Targets.

Stem cells are undifferentiated cells defined by two fundamental characteristics. First, they are able to self-renew, that is to sustain themselves for long periods without losing their properties through mitotic cell division. Second, they can be induced under certain conditions to proliferate and differentiate into a variety of mature cells, allowing the maintenance and repair of various tissues throughout life. Increasing evidence also pointed to a relationship between the mechanisms regulating the development of stem cells and those of tumor cells. Cancers are composed of heterogeneous cell populations proliferating uncontrollably with the risk of invading adjacent tissues and of spreading to other tissues by metastasis. The "cancer stem cell hypothesis" stipulates that a minor subpopulation of cells, characterized by their stem cell-like abilities, is able to initiate and maintain new tumors by reconstituting the whole cancer cellular heterogeneity, whereas the remaining bulk of the tumor lacks this potential [7, 8]. Cancer stem cells were first defined in acute myeloid leukemia by the demonstration that only a small subset of hematopoietic stem cell-like population is necessary and sufficient to allow serial leukemic transplantations in immunodeficient mice [9, 10]. Cancer stem cells were identified some years later in human breast cancer [11], followed rapidly by other types of solid tumors [8]. Although the clinical relevance of cancer stem cells still remains an issue and controversies have been raised in some types of tumors such as colon and brain cancers due to the difficulties in accurately defining and targeting the stem cell fraction [12, 13], accumulating evidence in various tissues including breast and lung cancers has demonstrated correlations between enrichment of cancer stem cells and tumor progression stage, enhanced tumorigenic and metastatic potential, and resistance to chemotherapies [8, 13-15].

As $\alpha 7$ -nAChR appears implicated in major cellular processes in diverse tissues, considering its relevance to stem cells would be of particular interest to appreciate its role in tissue homeostasis and the consequences of nicotine exposure. Indeed, as summarized in Table 1, several works have

*Address correspondence to this author at the 1-18-1, Kamiyoga, Setagaya-ku 158-8501, Japan; Tel: +81-3-3700-9704; Fax: +81-3-3700-9704; E-mail: kanda@nihs.go.jp

Table 1. Expression of $\alpha 7$ -nAChR in Stem Cells

Stem/Progenitor cell types	Evidence for the presence of $\alpha 7$ -nAChR	Suggested function of $\alpha 7$ -nAChR	Refs.
<i>Embryonic stem cells</i>			
Mouse ESC	RT-PCR, intracellular Ca^{2+} response	Inhibition of proliferation, survival	[24]
Human ESC	RT-PCR, microarray, immunostaining	Proliferation, differentiation	[21, 22]
P19 embryonic carcinoma	RT-PCR, immunostaining, intracellular Ca^{2+} response	Inhibition of proliferation	[25, 26]
<i>Hematopoietic stem/progenitor cells</i>			
Mouse bone marrow stem cells	Staining with labeled α -BTX, immunostaining	Proliferation, mobilization, communication with niche cells	[36]
Mouse myeloid, erythroid, lymphoid progenitor cells	Staining with labeled α -BTX, effects of $\alpha 7$ -nAChR knock-out	Proliferation, survival	[34, 35]
<i>Mesenchymal stem cells</i>			
Human bone marrow stem cells	RT-PCR, immunostaining, intracellular Ca^{2+} response, inhibition by α -BTX	Proliferation, apoptosis, inhibition of chemotaxis	[41, 42]
<i>Neural stem/progenitor cells</i>			
Early embryonic mouse cerebral cortex	Immunostaining, intracellular Ca^{2+} response, membrane potential response	Proliferation, differentiation	[47]
P19-derived progenitor cells	RT-PCR, immunostaining, intracellular Ca^{2+} response	Proliferation, differentiation	[25, 26]
Adult rat hippocampal progenitor cells	RT-PCR, staining with labeled α -BTX	Apoptosis	[51]
Adult mouse hippocampal progenitor cells	Staining with labeled α -BTX, effects of $\alpha 7$ -nAChR knock-out	Proliferation, survival	[52,53]
<i>Cancer stem cells</i>			
MCF-7 breast cancer cell line	RT-PCR, inhibition by α -BTX	Proliferation, survival	[76]

reported the expression of $\alpha 7$ -nAChR in different categories of stem cells and have suggested its involvement in their regulation. Most interestingly, recent works in our laboratory demonstrated that $\alpha 7$ -nAChR is involved in cancer stem cell proliferation in breast cancer, shading a new light on the mechanisms underlying tumor development and progression in presence of nicotinic agonists. Here we review the multiple facets of the expression and function of $\alpha 7$ -nAChR in stem cells, including embryonic, hematopoietic, mesenchymal and neural stem cells, as schematically illustrated in Fig. (1). Then we focus on the most recent findings in breast cancer stem cells and on the insights these data may provide for understanding disease mechanisms and for developing therapeutic strategies.

EMBRYONIC STEM CELLS

Embryonic stem cells (ESC) are undifferentiated cells originally derived from the inner cell mass of blastocysts. They can be maintained *in vitro* and they are characterized by their pluripotency, meaning that they can be induced to differentiate into any cell lineage of all three embryonic germ layers, as opposed to more restricted somatic stem cells. Since the first successful establishment of human ESC [16, 17], research in this field has been the focus of considerable interest for developmental studies as well as for its huge therapeutic application potential [18].

ESC have been shown to be able to reconstitute a fully functional cholinergic system upon differentiation into ESC-

derived neurons, hematopoietic and endothelial cells [19-22]. Interestingly, several members of the cholinergic signaling system, including AChR and enzymes for ACh synthesis (choline acetyltransferase) and degradation (acetyl- and butyryl-cholinesterases), have been demonstrated to be already expressed in undifferentiated ESC [23]. Expression analyses have confirmed the presence of nAChR in mouse ESC [24], undifferentiated human ESC [22] and human embryoid bodies [21]. The type of the nAChR subunits that were detected in ESC depended on the experimental conditions, but $\alpha 7$ -nAChR was reliably found to be expressed in all of these studies [21, 22, 24]. Remarkably, $\alpha 5$, $\alpha 7$ and $\alpha 9$ -nAChR in human ESC were shown to be significantly higher expressed in undifferentiated ESC than after initiation of differentiation, which suggests the possible importance of these receptors in regulating ESC in its undifferentiated state [22].

The function of nAChR has been demonstrated in mouse ESC by monitoring the increase of intracellular Ca^{2+} concentration in the presence of ACh or nicotine [24]. In contrast, muscarine did not have any effect on intracellular Ca^{2+} [24], suggesting that muscarinic AChR might not be fully functional in mouse ESC despite being transcriptionally expressed [23]. ESC proliferation was found to be decreased by exposure to nicotine [24] and by treatment with a butyryl-cholinesterase inhibitor, which increases the cholinergic signal by inhibiting ACh degradation [23]. In addition, ESC viability was increased by exposure to nicotine [24]. These results suggest that nAChR are involved in modulating ESC




Fig. (1). Overview of the involvement of $\alpha 7$ -nAChR in the regulation of essential biological processes in stem cells.

proliferation and promoting survival. In parallel, investigations in mouse P19 embryonic carcinoma cells showed that the nAChR-dependent increase of intracellular Ca^{2+} inhibits proliferation [25, 26]. Indeed, P19 cell proliferation was decreased in a dose-dependent fashion by nicotine exposure but not by muscarine, and the effect was blocked by the nAChR antagonist mecamylamine and the extracellular Ca^{2+} chelant EGTA [25]. Since $\alpha 7$ -nAChR was reported to be the most expressed nAChR in P19 cells [26] and that muscarinic receptors do not appear functional in mouse ESC and P19 cells, it would be tempting to suggest that $\alpha 7$ -nAChR is the main AChR involved in the regulation of ESC proliferation and survival.

Since ESC express functional receptors as well as enzymes for the synthesis or degradation of their ligand, these might be required in some local intercellular signal pathway controlling the earliest stages of embryonic development. The concentration of nicotine in the plasma of chronic smokers has been reported to range from 10 nM to 10 μM [27]. It would be of interest to determine if exposure to nicotine in this range of concentrations could perturb the nAChR-mediated signaling system and consequently affect embryonic development *in vivo*. Initiation of embryonic differentiation can be studied to some extent in embryoid bodies, which are formed by aggregation of ESC in spheroid clusters when cultured in non-adherent conditions. Human embryoid bodies have been shown to express nAChR, especially $\alpha 3$ and $\alpha 7$. Their stimulation by nicotine at 10 nM induced a rapid phosphorylation of mitogen-activated protein kinases (MAPK) [21], which activate a highly conserved pathway known to stimulate growth and differentiation. Furthermore, when embryoid bodies were induced to differentiate into hematopoietic lineages, nicotine-treated emb-

ryoid bodies contained a higher number of hemangioblasts than non-treated ones. Thus nAChR appear involved in controlling proliferation and cell fate determination at the onset of differentiation. ESC has been also used as a model system to study the adverse effects of prenatal nicotine absorption [28], which might explain some of the pregnancy complications and fetal defects at early stages associated with maternal smoking [29]. Exposure of human ESC to nicotine at doses above 1.8 μM was shown to inhibit cell adhesion, alter cell morphology and induce apoptosis, an effect blocked by addition of the nAChR antagonist d-tubocurarine [28]. Exposure to nicotine was also found to decrease the expression of several known markers of pluripotency including OCT-4 [28], the amount of which should be tightly controlled for proper maintenance of ESC self-renewal and lineage commitment [30].

Thus, the function of $\alpha 7$ -nAChR in ESC appears to be important in regulating proliferation, survival, and differentiation, most likely using ACh as one of the signaling molecules orchestrating embryonic development. Exposure to nicotine might directly affect the embryo *in vivo*, since many of these effects were observed at concentrations of nicotine comparable to those in the plasma of cigarette smokers.

HEMATOPOIETIC STEM CELLS

Hematopoietic stem cells (HSC) are multipotent cells that are present throughout the adult life to allow the regeneration of blood cells. They are the best characterized somatic stem cells so far, as they can be identified by immunophenotyping and by assaying their ability to reconstitute all blood-cell types, including myeloid and lymphoid lineages, which

allows for instance to entirely repopulate the bone marrow of lethally irradiated animals [31]. Somatic stem cells are known to be maintained in specific cellular microenvironments referred to as niches. The cross-talk of signals between stem cells and niche constituents is believed to be essential for the proper control of their function. HSC are mainly located in the bone-marrow niche [32], where they are maintained in a relatively quiescent state. HSC can also exit the bone marrow by a process of mobilization, migrate through the systemic circulation and reintegrate again the bone marrow niche or facultative niches in other tissues such as the spleen or liver. This system of HSC trafficking and homing to the appropriate niche allows a fast proliferation and distribution for maintenance of blood cell homeostasis and immune surveillance [33].

The role of nAChR in hematopoiesis has been of particular interest because of the concerns about the effects of smoking on cardiovascular diseases and deficits in immune and inflammatory responses. Knockout and chimera mice experiments showed that α 4 β 2 and α 7-nAChR are required for survival and proliferation of B lymphocytes precursors [34]. Myeloid and erythroid progenitor cells also express α 7-nAChR and the number of progenitor cells was found to be reduced in α 7-nAChR knockout mice [35]. Thus, α 7-nAChR seems to play a role in all blood-cell lineages at different stages of the hematopoietic development. Regarding undifferentiated HSC, a recent report has demonstrated that HSC purified from mouse bone marrow also express α 7-nAChR as shown by binding of labeled α -BTX [36]. Exposure of mice to nicotine has been shown to increase the number of circulating white blood cells, in correlation with an increase of both the frequency and number of HSC in the bone-marrow. This effect might be mediated by a direct action of nicotine via α 7-nAChR. On the other hand, nicotine also induced an increase of the plasma levels of soluble Kit ligand (sKitL or stem cell factor), a marker of stem cell activation [36]. This suggests that HSC mobilization and proliferation can be also stimulated by an indirect effect via the production of sKitL by the stroma cells of the HSC supportive niches. Another series of studies has also shown that nicotine induces concerted responses in both HSC and niche stroma cells. Nicotine interfered with HSC homing into the bone marrow by arresting HSC on the endothelium via nAChR [37] and by decreasing the expression of the CD44 adhesion molecule in bone stroma cells [38] and of β 2-integrin in HSC [39]. Exposure of mice to nicotine during gestation was also found to affect hematopoiesis and HSC homing during fetal and newborn development, an effect accompanied by a perturbation of the profile of cytokines and chemokines released in the niche, as well as the expression of corresponding receptors in HSC [21].

These results strongly suggest that α 7-nAChR and the cholinergic system plays a role in regulating HSC either by acting directly on HSC by nAChR-mediated effects, or by affecting the surrounding niche cells, or most likely by a synergy between both effects. Interfering on the signaling balance by excessive exposure to nicotine might induce disease-related consequences such as fetal and postpartum developmental defects [21], increased number of white blood cells, which has been shown to be a risk factor for heart diseases [36], and pathological extramedullary hematopoiesis [39].

MESENCHYMAL STEM/STROMAL CELLS

Mesenchymal stem cells (MSC) are multipotent non-hematopoietic stromal cells found especially in the bone marrow and originating from the mesodermal germ layer. They can differentiate into various cell lineages, including osteoblasts, chondrocytes and adipocytes depending on the appropriate stimuli, and they modulate immune and inflammatory responses. Their great potential for clinical applications, by virtue of their relative ease of culture and their capacity to migrate to sites of injury for damage repair, has been the object of intense investigation in recent years [40].

The expression of α 7-nAChR in human MSC has been confirmed by flow-cytometry [41] and by confocal microscopy [42]. Other nAChR subunits such as α 3, α 5, β 2 and β 4 were also reported to be expressed in MSC [41, 42] but they were not always detected, probably depending on the experimental conditions and the intrinsic heterogeneity of MSC [42]. Human MSC have been shown to respond to nicotine (1 μ M) by an increased influx of Ca²⁺ and by activation of the MAPK pathway [41]. Further confirmation of the involvement of α 7-nAChR has been provided by the observation of a similar increase in MAPK phosphorylation in mouse C3H10T1/2 mesenchymal cells overexpressing α 7-nAChR [41]. The activation of the MAPK pathway suggests a role in the regulation of proliferation and differentiation of MSC. However, the exposure of MSC to high concentrations of nicotine over the threshold of 1 μ M has also been shown to induce cell death by apoptosis [42], so that any putative effect of nicotine on proliferation might be possibly masked by its toxic effect in these experimental conditions. A moderate dose of nicotine (less than 1 μ M) has been shown to inhibit chemotaxis *in vitro*, which has been confirmed by *in vivo* observations of a defect in the migration of MSC to the bone marrow and spleen of mice injected with nicotine [42]. The induction of apoptosis and the inhibition of chemotaxis were blocked by the specific antagonist α -BTX, which suggests the involvement of α 7-nAChR in mediating these effects. These findings could be of importance for the regulation of survival and homing of MSC systemically administered in therapeutics applications.

NEURAL STEM/PROGENITOR CELLS

Neural stem cells (NSC) are defined by their ability for self-renewal and differentiation into all cell types of the nervous system, including neurons, astrocytes and oligodendrocytes. In the adult mammalian brain, neurogenesis has been demonstrated to occur in the subventricular zone (SVZ) and in the dentate gyrus (DG) of the hippocampus [43]. NSC and their derived progenitor cells can be enriched and propagated *in vitro* by their ability to form floating aggregates called neurospheres [44]. NSC can also be isolated from the developing brain of the embryo, where they proliferate to generate the complete nervous system of the organism. Understanding the control of NSC differentiation by neurotransmitters such as ACh is the focus of growing interest with the prospects for developing cell therapies for neurodevelopmental and neurodegenerative diseases [45]. nAChR are among the major players of neurotransmission and recent advances have shown their involvement in diverse other functions such as neuroprotection and regulation of

neurogenesis, which makes them a target of choice in therapeutic applications against nervous-system disorders [46]. The role of $\alpha 7$ -nAChR in the human central nervous system being extensively reviewed in other chapters of this issue of *Current Drug Targets*, we provide here a summary restricted to the recent findings on its expression and functions in neural stem/progenitor cells.

The involvement of $\alpha 7$ -nAChR in the early development of the nervous system has been analyzed by characterizing NSC in embryonic/fetal organs. The early embryonic mouse cerebral cortex at embryonic day 10 (E10) consists of a large majority of neural stem and progenitor cells as identified by stem cell-specific markers [47]. Dissociated cells from the early embryonic mouse cerebral cortex at E10 have been shown to express $\alpha 3$, $\alpha 4$, and $\alpha 7$ -nAChR by immunofluorescence staining. The presence of functional receptors that respond to ACh and nicotine has been demonstrated by patch-clamp recordings. The response to nicotine was blocked by the $\alpha 7$ -nAChR antagonist methyllycaconitine [47]. Interestingly, higher levels of expression were observed at early embryogenesis (E10) than at later stages (E18). The kinetics of ACh-induced cytosolic Ca^{2+} response showed a faster decay at later stages, suggesting that $\alpha 7$ -nAChR is involved in early development and that its function might change with embryonic age [47]. The function of nAChR was also examined in P19 murine embryonic carcinoma cells induced to differentiate into neurons by retinoic acid treatment, which is commonly used as a model of neuronal development. As we have previously described for ESC cells, P19 cells express functional nAChR, which mediate an inhibition of proliferation by nicotine [25, 26]. Remarkably, the expression of several nAChR, including $\alpha 7$ -nAChR, decreased in P19-derived neuronal progenitor cells and then increased again in mature neurons [25]. At the same time, muscarinic receptors, which were not functional in undifferentiated embryonic cells, were shown to become functional in progenitor cells. Both nicotinic and muscarinic receptors appeared to stimulate proliferation and differentiation by some mechanism involving Ca^{2+} mobilization from intracellular stores, which was suggested to override the inhibitory effects mediated by nAChR in undifferentiated cells [25]. This is consistent with the idea that the function of nAChR might change depending on the cellular state and on the presence of other factors acting in concert, notably of muscarinic AChR, which have also been reported as regulators of neuronal stem cell proliferation [48, 49].

In the adult central nervous system, several studies reported various effects of nicotine exposure on neural stem and progenitor cells, with some discrepancies regarding the suggested role of $\alpha 7$ -nAChR in inducing apoptosis or on the contrary in promoting survival and proliferation. Exposure to nicotine induced a decrease of neurogenesis and cell death in the rat hippocampal DG *in vivo* [50], as well as in experiments using isolated primary progenitor cells or the rat HC2S2 immortalized hippocampal progenitor cells [51]. The expression of $\alpha 7$ -nAChR was confirmed in hippocampal cells and the induction of apoptosis in undifferentiated progenitor cells was blocked by α -BTX and dependent on the increase of intracellular Ca^{2+} concentrations [51]. In contrast, apoptosis was not induced in differentiated HC2S2 cells, despite their expression of $\alpha 7$ -nAChR. The greater sensitivity of undifferentiated progenitor cells to calcium

toxicity was suggested to be due to the lack of calbindin D28K, which is expressed at much higher levels in differentiated HC2S2 cells and may provide a protective effect by calcium buffering [51].

As opposed to the induction of apoptosis, a protective effect of nicotine has been described in the mouse DG and the rat SVZ. $\alpha 7$ -nAChR was found to be expressed in the neuronal progenitor cells of the mouse DG and in a subpopulation of progenitor cells derived from the SVZ [52, 53]. In the DG, neural progenitor cell survival was shown to be promoted when intracellular ACh was increased by a long-term treatment of mice with the cholinesterase inhibitor donepezil [53]. Mice heterozygous for $\alpha 7$ -nAChR also showed a reduced cell proliferation in the DG, suggesting that the expression levels of $\alpha 7$ -nAChR might be important in regulating proliferation [52]. Nicotine-induced proliferation of progenitor cells was also reported in the rat SVZ. [54]. The positive effect on proliferation was accompanied by an increase in the local expression of fibroblast growth factor (FGF-2) and of the corresponding FGF receptor (FGFR-1). A proposed model is that nicotine promotes the creation of a micro-environment favorable to proliferation in the SVZ by stimulating the release of growth factors in the stem cell niche and the coordinated expression of the appropriate receptors on neural progenitor cells [54].

Thus, the response of neural stem and progenitor cells to nicotine appears to differ depending on the cell type, the tissue localization, the animal species and the experimental conditions including the concentrations and the mode of administration of nicotine. Further investigation would be necessary to determine the extent of the contribution of $\alpha 7$ -nAChR in stem cell proliferation and survival in the DG and SVZ.

CANCER STEM CELLS

Cancer stem cells (CSC) have been proposed to constitute a specific cellular subpopulation that is responsible for tumor progression and metastasis. Like somatic stem cells, CSC can be selected on the basis of a combination of cell surface markers such as $\text{CD}44^+/\text{CD}24^-$ in breast cancer [11] and $\text{CD}133^+$ initially used in brain [55]. Aldehyde dehydrogenase (ALDH), a detoxifying enzyme responsible for the oxidation of intracellular aldehydes, is another marker with increased activity in stem cells, which has been successfully used in recent years to identify a subpopulation enriched in CSC and correlated to a poor prognosis in breast and lung cancer patients [56, 57], increased metastasis [58, 59] and chemotherapy resistance [60]. The demonstration of CSC properties can be performed by *in vivo* observation of tumor growth after serial transplantation of a small amount of selected cells. CSC population can also be characterized by the selective formation of spheres [61], reminiscent of neurospheres formed by NSC, when cultured *in vitro* in non-adherent conditions. The maintenance of CSC has been shown to depend on their stem cell-like capacity for self-renewal, and major developmental signaling pathways such as Notch, Hedgehog and Wnt have been implicated in their viability and proliferation [62-64]. Notably, regarding breast cancer, overexpression of Notch in transgenic mice was shown to promote mammary tumor development [65],

aberrant Notch activation was reported in various subtypes of breast carcinoma [66] and the coexpression of Notch-1 and one of its ligand Jagged-1 was correlated with poor prognosis [67].

A number of studies have provided evidence that nicotine, although not carcinogenic in itself, can stimulate the development and progression of cancer. Research on the effects of nicotine has been especially active in the field of lung cancers, which could be initiated by other carcinogens of tobacco smoke, then maintained and stimulated by exposure to nicotine [68]. Breast cancer risk has been also associated to tobacco smoking by epidemiologic studies [69, 70] and clinical reports indicating that smoking enhance breast cancer mortality and metastasis from breast to lung [71, 72]. However, it is only recently that experimental data have suggested direct links between breast cancer and nicotine exposure. Nicotine has thus been shown in human

MCF-7 breast cancer cells to induce a resistance to the anti-cancer drug doxorubicin [73] and to promote their migration by activation of protein kinase C (PKC) [74]. Nicotine also promotes proliferation, invasion and epithelial-to-mesenchymal transition (EMT), which is a process associated to the enhancement of tumor cell migration and metastasis, by calcium and Src dependent pathways in various cell lines expressing $\alpha 7$ -nAChR, including the human MCF-7 and MDA-MB-468 breast cancer cells [75].

In a recent report from our laboratory, we have demonstrated for the first time that nicotine increases breast CSC population via $\alpha 7$ -nAChR [76]. A possible model for the process inferred from our results is illustrated in Fig. (2). We showed that exposure to nicotine, at concentrations comparable to those reported in the plasma of cigarette smokers, increased in a dose-dependent manner both the frequency and the absolute number of CSC identified as the




Fig. (2). Proposed model for cancer stem cell proliferation via $\alpha 7$ -nAChR. The binding of nicotine to $\alpha 7$ -nAChR induces an increase of the intracellular Ca^{2+} influx and the consequent activation of calcium-dependent signaling molecules such as protein kinase C (PKC) and activate the Notch signaling pathway. In the presence of notch ligands, Notch is cleaved by ADAM and by γ -secretase to release the Notch intracellular domain (NICD). NICD translocates to the nucleus and activates the transcription of downstream regulatory factors such as HES1, thereby promoting stem-cell proliferation and self-renewal. The inhibitors α -BTX, DAPT and GFX allow to target specific molecules of this pathway as indicated.

ALDH-positive fraction in MCF-7 cells. The increase of a population with stem-cell like properties was confirmed by observing a corresponding increase in mammosphere formation. The proliferation of ALDH-positive cells was blocked by α -BTX, suggesting that the effect was dependent on $\alpha 7$ -nAChR. The increase of the ALDH-positive population was also reproduced when the MCF-7 cells were treated with the $\alpha 7$ -nAChR-specific agonist PHA543613 (PHA). We confirmed the expression of $\alpha 7$ -nAChR by RT-PCR in non-treated MCF-7 cells [76] and in both ALDH-positive and negative populations (unpublished data). Furthermore, we investigated the involvement of stem-cell related pathways, and we showed that nicotine induced an expression of the Notch target gene HES1. The increase in HES1 expression was blocked by DAPT, which prevents Notch signaling by inhibiting the cleavage of activated Notch receptor by γ -secretase. The nicotine-induced increase of ALDH-positive population was blocked by DAPT and by the PKC inhibitor GF109203X (GFX). This suggests that the Notch-Hes1 pathway is involved in the increase of CSC population and that it acts downstream of $\alpha 7$ -nAChR and PKC. The mechanisms underlying the $\alpha 7$ -nAChR-mediated activation of Notch have not been fully investigated yet, but PKC activity has been shown to induce the membrane translocation in glioblastoma of the ADAM protease [77], which is required for the cleavage of Notch receptors. CSC proliferation might be therefore induced by the upregulation of Notch signaling via the $\alpha 7$ -nAChR-mediated PKC activation.

CSC maintenance and proliferation have been correlated to EMT [78], activation of anti-apoptotic pathways [79, 80] and increase of chemoresistance [60]. Our results indicating the involvement of CSC in nicotine-dependent breast carcinogenesis appear therefore in concordance with the previously mentioned other studies showing various effects of nicotine in breast cancer [73-75]. Our results also suggest that $\alpha 7$ -nAChR is the main receptor controlling CSC self-renewal, although we do not exclude the contribution of other nAChR subunits, since it was reported that in some breast cancer cell lines, nicotine-induced tumor growth could be mediated by $\alpha 9$ -nAChR [81]. It is noteworthy that $\alpha 7$ -nAChR has been shown to be expressed in many different types of cancer cells in various tissues, including in particular non-small cell lung carcinoma (NSCLC) [3]. Tumorigenic ALDH-positive population has been recently identified and related to poor prognosis in lung cancers [57] and its maintenance shown to be dependent on Notch signaling in NSCLC [82]. $\alpha 7$ -nAChR is already being considered as a target for the treatment of NSCLC and mesothelioma [83, 84]. Therefore, it would be also of great interest to consider the effects of $\alpha 7$ -nAChR on CSC populations in other types of tumor such as NSCLC. The finding that $\alpha 7$ -nAChR might be involved in CSC maintenance and proliferation sheds a new light on the effects of nicotine on tumor progression and opens the way to the development of new therapies to eradicate CSC by targeting $\alpha 7$ -nAChR or the downstream signaling pathways.

CONCLUSIONS

A growing body of evidence suggests that $\alpha 7$ -nAChR plays a key role in the regulation of stem cells. As

summarized in Fig. (1), current data point to the prominent involvement of $\alpha 7$ -nAChR in the control of maintenance, proliferation, and stem cell fate determination. Cellular signals mediated by $\alpha 7$ -nAChR have been shown to promote survival and to limit proliferation of undifferentiated ESC, whereas they stimulate proliferation and differentiation of more developed somatic stem cells including hematopoietic and neuronal progenitors. Several works reviewed in this issue demonstrated that in the presence of an appropriate agonist, namely Ach in physiological conditions, $\alpha 7$ -nAChR activates developmental signaling pathways such as MAPK and Notch. $\alpha 7$ -nAChR-mediated effects also appeared to depend on the state of the cell itself, including the presence or absence of other receptors and factors modulating the cytosolic Ca^{2+} response. Thus, $\alpha 7$ -nAChR possibly plays a central role in integrating intra- and extra-cellular information through the cholinergic signaling to coordinate stem cell maintenance and differentiation. To this effect, $\alpha 7$ -nAChR has also been shown to enhance the cross-talk between stem cells and their niche to finely tune cellular development, as best demonstrated in experiments on hematopoietic or neural progenitor cells treated with nicotine *in vivo*. This also implies that stem cells are sensitive to environmental change through $\alpha 7$ -nAChR, and in several experimental systems, the disruption of the signal balance was shown to induce detrimental effects such as apoptosis or anomalous differentiation and migration. Finally, the involvement of $\alpha 7$ -nAChR in the promotion of stem cell maintenance and proliferation is of particular significance with regard to CSC in breast cancer, as this might explain nicotine-induced tumor progression and metastasis. The finding that $\alpha 7$ -nAChR plays a major role in the regulation of various types of stem cells provides important insights for the prospects of developing new therapeutic strategies. The wide distribution of $\alpha 7$ -nAChR in the organism and their involvement in a variety of functions imply that the risks for unexpected side effects must be carefully assessed. A precise evaluation of risk factors and of drug delivery systems would be required to validate $\alpha 7$ -nAChR as target in therapeutic applications.

ABBREVIATIONS

ACh	=	Acetylcholine
AChR	=	Acetylcholine receptor
ALDH	=	Aldehyde dehydrogenase
α -BTX	=	α -bungarotoxin
CSC	=	Cancer stem cells
DG	=	Dentate gyrus
EMT	=	Epithelial-to-mesenchymal transitions
ESC	=	Embryonic stem cells
GFX	=	GF109203X
HSC	=	Hematopoietic stem cells
MAPK	=	Mitogen-activated protein kinases
MSC	=	Mesenchymal stem cells

nAChR	=	Nicotinic acetylcholine receptor
NSC	=	Neural stem cells
NSCLC	=	Non-small cell lung carcinoma
PHA	=	PHA543613
PKC	=	Protein kinase C
RT-PCR	=	Reverse transcription polymerase chain reaction
sKitL	=	Soluble Kit ligand
SVZ	=	Subventricular zone.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest.

ACKNOWLEDGEMENTS

This work was supported in part by a grant from the Program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation (NIBIO) (No.09-02 to Y.K.), a Grant-in-Aid for Scientific Research (No. 2179025, 23590322 to Y.K.) from the Ministry of Education, Culture, Sports, Science, and Technology, Japan (Y.K.), a Health and Labour Sciences Research Grant from the Ministry of Health, Labour and Welfare, Japan (Y.K.), and a grant from the Smoking Research Foundation (Y.K.).

REFERENCES

[1] Lindstrom J, Anand R, Gerzanich V, *et al.* Structure and function of neuronal nicotinic acetylcholine receptors. *Prog Brain Res* 1996; 109: 125-37.

[2] Wessler I, Kirkpatrick CJ. Acetylcholine beyond neurons: the non-neuronal cholinergic system in humans. *Br J Pharmacol* 2008; 154: 1558-71.

[3] Egleton RD, Brown KC, Dasgupta P. Nicotinic acetylcholine receptors in cancer: multiple roles in proliferation and inhibition of apoptosis. *Trends Pharmacol Sci* 2008; 29: 151-8.

[4] Schuller HM. Is cancer triggered by altered signalling of nicotinic acetylcholine receptors? *Nat Rev Cancer* 2009; 9: 195-205.

[5] Maneckjee R, Minna JD. Opioid and nicotine receptors affect growth regulation of human lung cancer cell lines. *Proc Natl Acad Sci USA* 1990; 87: 3294-8.

[6] Schuller HM. Cell type specific, receptor-mediated modulation of growth kinetics in human lung cancer cell lines by nicotine and tobacco-related nitrosamines. *Biochem Pharmacol* 1989; 38: 3439-42.

[7] Reya T, Morrison SJ, Clarke MF, Weissman IL. Stem cells, cancer, and cancer stem cells. *Nature* 2001; 414: 105-11.

[8] Visvader JE, Lindeman GJ. Cancer stem cells in solid tumours: accumulating evidence and unresolved questions. *Nat Rev Cancer* 2008; 8: 755-68.

[9] Lapidot T, Sirard C, Vormoor J, *et al.* A cell initiating human acute myeloid leukaemia after transplantation into SCID mice. *Nature* 1994; 367: 645-8.

[10] Bonnet D, Dick JE. Human acute myeloid leukemia is organized as a hierarchy that originates from a primitive hematopoietic cell. *Nat Med* 1997; 3: 730-7.

[11] Al-Hajj M, Wicha MS, Benito-Hernandez A, Morrison SJ, Clarke MF. Prospective identification of tumorigenic breast cancer cells. *Proc Natl Acad Sci USA* 2003; 100: 3983-8.

[12] Prestegarden L, Enger PO. Cancer stem cells in the central nervous system--a critical review. *Cancer Res* 2010; 70: 8255-8.

[13] Papailiou J, Bramis KJ, Gazouli M, Theodoropoulos G. Stem cells in colon cancer. A new era in cancer theory begins. *Int J Colorectal Dis* 2011; 26: 1-11.

[14] McDermott SP, Wicha MS. Targeting breast cancer stem cells. *Mol Oncol* 2010; 4: 404-19.

[15] Perona R, Lopez-Ayllon BD, de Castro Carpeno J, Belda-Iniesta C. A role for cancer stem cells in drug resistance and metastasis in non-small-cell lung cancer. *Clin Transl Oncol* 2011; 13: 289-93.

[16] Thomson JA, Itskovitz-Eldor J, Shapiro SS, *et al.* Embryonic stem cell lines derived from human blastocysts. *Science* 1998; 282: 1145-7.

[17] Shambloot MJ, Axelman J, Wang S, *et al.* Derivation of pluripotent stem cells from cultured human primordial germ cells. *Proc Natl Acad Sci USA* 1998; 95: 13726-31.

[18] Klimanskaya I, Rosenthal N, Lanza R. Derive and conquer: sourcing and differentiating stem cells for therapeutic applications. *Nat Rev Drug Discov* 2008; 7: 131-42.

[19] Nilbratt M, Porras O, Marutle A, Hovatta O, Nordberg A. Neurotrophic factors promote cholinergic differentiation in human embryonic stem cell-derived neurons. *J Cell Mol Med* 2010; 14: 1476-84.

[20] Li XJ, Du ZW, Zarnowska ED, *et al.* Specification of motoneurons from human embryonic stem cells. *Nat Biotechnol* 2005; 23: 215-21.

[21] Seroby N, Jagannathan S, Orlovskaya I, *et al.* The cholinergic system is involved in regulation of the development of the hematopoietic system. *Life Sci* 2007; 80: 2352-60.

[22] Yu J, Huang NF, Wilson KD, *et al.* nAChRs mediate human embryonic stem cell-derived endothelial cells: proliferation, apoptosis, and angiogenesis. *PLoS One* 2009; 4: e7040.

[23] Paraoanu LE, Steinert G, Koehler A, Wessler I, Layer PG. Expression and possible functions of the cholinergic system in a murine embryonic stem cell line. *Life Sci* 2007; 80: 2375-9.

[24] Landgraf D, Barth M, Layer PG, Sperling LE. Acetylcholine as a possible signaling molecule in embryonic stem cells: studies on survival, proliferation and death. *Chem Biol Interact* 2010; 187: 115-9.

[25] Resende RR, Alves AS, Britto LR, Ulrich H. Role of acetylcholine receptors in proliferation and differentiation of P19 embryonal carcinoma cells. *Exp Cell Res* 2008; 314: 1429-43.

[26] Resende RR, Gomes KN, Adhikari A, Britto LR, Ulrich H. Mechanism of acetylcholine-induced calcium signaling during neuronal differentiation of P19 embryonal carcinoma cells in vitro. *Cell Calcium* 2008; 43: 107-21.

[27] Gritz ER, Baer-Weiss V, Benowitz NL, Van Vunakis H, Jarvik ME. Plasma nicotine and cotinine concentrations in habitual smokeless tobacco users. *Clin Pharmacol Ther* 1981; 30: 201-9.

[28] Zdravkovic T, Genbacev O, LaRocque N, McMaster M, Fisher S. Human embryonic stem cells as a model system for studying the effects of smoke exposure on the embryo. *Reprod Toxicol* 2008; 26: 86-93.

[29] Rogers JM. Tobacco and pregnancy. *Reprod Toxicol* 2009; 28: 152-60.

[30] Niwa H, Miyazaki J, Smith AG. Quantitative expression of Oct-3/4 defines differentiation or self-renewal of ES cells. *Nat Genet* 2000; 24: 372-6.

[31] Orkin SH, Zon LI. Hematopoiesis: an evolving paradigm for stem cell biology. *Cell* 2008; 132: 631-44.

[32] Wilson A, Trumpp A. Bone-marrow haematopoietic-stem-cell niches. *Nat Rev Immunol* 2006; 6: 93-106.

[33] Laird DJ, von Andrian UH, Wagers AJ. Stem cell trafficking in tissue development, growth, and disease. *Cell* 2008; 132: 612-30.

[34] Skok MV, Grailhe R, Agenes F, Changeux JP. The role of nicotinic receptors in B-lymphocyte development and activation. *Life Sci* 2007; 80: 2334-6.

[35] Koval LM, Zverkova AS, Grailhe R, *et al.* Nicotinic acetylcholine receptors alpha4beta2 and alpha7 regulate myelo- and erythropoiesis within the bone marrow. *Int J Biochem Cell Biol* 2008; 40: 980-90.

[36] Chang E, Forsberg EC, Wu J, *et al.* Cholinergic activation of hematopoietic stem cells: role in tobacco-related disease? *Vasc Med* 2010; 15: 375-85.

- [37] Serobyann N, Schraufstatter IU, Strongin A, Khaldoyanidi SK. Nicotinic acetylcholine receptor-mediated stimulation of endothelial cells results in the arrest of haematopoietic progenitor cells on endothelium. *Br J Haematol* 2005; 129: 257-65.
- [38] Khaldoyanidi S, Sikora L, Orlovskaya I, *et al.* Correlation between nicotine-induced inhibition of hematopoiesis and decreased CD44 expression on bone marrow stromal cells. *Blood* 2001; 98: 303-12.
- [39] Pandit TS, Sikora L, Muralidhar G, Rao SP, Sriramarao P. Sustained exposure to nicotine leads to extramedullary hematopoiesis in the spleen. *Stem Cells* 2006; 24:2373-81.
- [40] Salem HK, Thiemermann C. Mesenchymal stromal cells: current understanding and clinical status. *Stem Cells* 2010; 28: 585-96.
- [41] Hoogduijn MJ, Cheng A, Genever PG. Functional nicotinic and muscarinic receptors on mesenchymal stem cells. *Stem Cells Dev* 2009; 18: 103-12.
- [42] Schraufstatter IU, DiScipio RG, Khaldoyanidi SK. Alpha 7 subunit of nAChR regulates migration of human mesenchymal stem cells. *J Stem Cells* 2009; 4: 203-15.
- [43] Zhao C, Deng W, Gage FH. Mechanisms and functional implications of adult neurogenesis. *Cell* 2008; 132: 645-60.
- [44] Reynolds BA, Weiss S. Generation of neurons and astrocytes from isolated cells of the adult mammalian central nervous system. *Science* 1992; 255: 1707-10.
- [45] Trujillo CA, Schwandt TT, Martins AH, *et al.* Novel perspectives of neural stem cell differentiation: from neurotransmitters to therapeutics. *Cytometry A* 2009; 75: 38-53.
- [46] Taly A, Corringer PJ, Guedin D, Lestage P, Changeux JP. Nicotinic receptors: allosteric transitions and therapeutic targets in the nervous system. *Nat Rev Drug Discov* 2009; 8: 733-50.
- [47] Atluri P, Fleck MW, Shen Q, *et al.* Functional nicotinic acetylcholine receptor expression in stem and progenitor cells of the early embryonic mouse cerebral cortex. *Dev Biol* 2001; 240: 143-56.
- [48] Diamandis P, Wildenhain J, Clarke ID, *et al.* Chemical genetics reveals a complex functional ground state of neural stem cells. *Nat Chem Biol* 2007; 3: 268-73.
- [49] Zhou C, Wen ZX, Shi DM, Xie ZP. Muscarinic acetylcholine receptors involved in the regulation of neural stem cell proliferation and differentiation in vitro. *Cell Biol Int* 2004; 28: 63-7.
- [50] Abrous DN, Adriani W, Montaron MF, *et al.* Nicotine self-administration impairs hippocampal plasticity. *J Neurosci* 2002; 22: 3656-62.
- [51] Berger F, Gage FH, Vijayaraghavan S. Nicotinic receptor-induced apoptotic cell death of hippocampal progenitor cells. *J Neurosci* 1998; 18: 6871-81.
- [52] Koike K, Hashimoto K, Okamura N, *et al.* Decreased cell proliferation in the dentate gyrus of alpha 7 nicotinic acetylcholine receptor heterozygous mice. *Prog Neuropsychopharmacol Biol Psychiatry* 2004; 28: 517-20.
- [53] Kaneko N, Okano H, Sawamoto K. Role of the cholinergic system in regulating survival of newborn neurons in the adult mouse dentate gyrus and olfactory bulb. *Genes Cells* 2006; 11: 1145-59.
- [54] Mudo G, Belluardo N, Mauro A, Fuxe K. Acute intermittent nicotine treatment induces fibroblast growth factor-2 in the subventricular zone of the adult rat brain and enhances neuronal precursor cell proliferation. *Neuroscience* 2007; 145: 470-83.
- [55] Singh SK, Hawkins C, Clarke ID, *et al.* Identification of human brain tumour initiating cells. *Nature* 2004; 432: 396-401.
- [56] Ginestier C, Hur MH, Charafe-Jauffret E, *et al.* ALDH1 is a marker of normal and malignant human mammary stem cells and a predictor of poor clinical outcome. *Cell Stem Cell* 2007; 1: 555-67.
- [57] Jiang F, Qiu Q, Khanna A, *et al.* Aldehyde dehydrogenase 1 is a tumor stem cell-associated marker in lung cancer. *Mol Cancer Res* 2009; 7: 330-8.
- [58] Marcato P, Dean CA, Pan D, *et al.* Aldehyde dehydrogenase activity of breast cancer stem cells is primarily due to isoform ALDH1A3 and its expression is predictive of metastasis. *Stem Cells* 2011; 29: 32-45.
- [59] Charafe-Jauffret E, Ginestier C, Iovino F, *et al.* Breast cancer cell lines contain functional cancer stem cells with metastatic capacity and a distinct molecular signature. *Cancer Res* 2009; 69: 1302-13.
- [60] Tanei T, Morimoto K, Shimazu K, *et al.* Association of breast cancer stem cells identified by aldehyde dehydrogenase 1 expression with resistance to sequential Paclitaxel and epirubicin-based chemotherapy for breast cancers. *Clin Cancer Res* 2009; 15: 4234-41.
- [61] Dontu G, Abdallah WM, Foley JM, *et al.* In vitro propagation and transcriptional profiling of human mammary stem/progenitor cells. *Genes Dev* 2003; 17: 1253-70.
- [62] Merchant AA, Matsui W. Targeting Hedgehog--a cancer stem cell pathway. *Clin Cancer Res* 2010; 16: 3130-40.
- [63] Pannuti A, Foreman K, Rizzo P, *et al.* Targeting Notch to target cancer stem cells. *Clin Cancer Res* 2010; 16: 3141-52.
- [64] Takahashi-Yanaga F, Kahn M. Targeting Wnt signaling: can we safely eradicate cancer stem cells? *Clin Cancer Res* 2010; 16: 3153-62.
- [65] Hu C, Dievert A, Lupien M, *et al.* Overexpression of activated murine Notch1 and Notch3 in transgenic mice blocks mammary gland development and induces mammary tumors. *Am J Pathol* 2006; 168: 973-90.
- [66] Stylianou S, Clarke RB, Brennan K. Aberrant activation of notch signaling in human breast cancer. *Cancer Res* 2006; 66: 1517-25.
- [67] Reedijk M, Odorcic S, Chang L, *et al.* High-level coexpression of JAG1 and NOTCH1 is observed in human breast cancer and is associated with poor overall survival. *Cancer Res* 2005; 65: 8530-7.
- [68] Russo P, Catassi A, Cesario A, Servent D. Development of novel therapeutic strategies for lung cancer: targeting the cholinergic system. *Curr Med Chem* 2006; 13: 3493-512.
- [69] Johnson KC. Accumulating evidence on passive and active smoking and breast cancer risk. *Int J Cancer* 2005; 117: 619-28.
- [70] Nagata C, Mizoue T, Tanaka K, *et al.* Tobacco smoking and breast cancer risk: an evaluation based on a systematic review of epidemiological evidence among the Japanese population. *Jpn J Clin Oncol* 2006; 36: 387-94.
- [71] Murin S, Inciardi J. Cigarette smoking and the risk of pulmonary metastasis from breast cancer. *Chest* 2001; 119: 1635-40.
- [72] Murin S, Pinkerton KE, Hubbard NE, Erickson K. The effect of cigarette smoke exposure on pulmonary metastatic disease in a murine model of metastatic breast cancer. *Chest* 2004; 125: 1467-71.
- [73] Zhou Y, Gu X, Ashayeri E, Zhang R, Sridhar R. Nicotine decreases the cytotoxicity of doxorubicin towards MCF-7 and KB-3.1 human cancer cells in culture. *J Natl Med Assoc* 2007; 99: 319-27.
- [74] Guo J, Ibaragi S, Zhu T, *et al.* Nicotine promotes mammary tumor migration via a signaling cascade involving protein kinase C and CDC42. *Cancer Res* 2008; 68: 8473-81.
- [75] Dasgupta P, Rizwani W, Pillai S, *et al.* Nicotine induces cell proliferation, invasion and epithelial-mesenchymal transition in a variety of human cancer cell lines. *Int J Cancer* 2009; 124: 36-45.
- [76] Hirata N, Sekino Y, Kanda Y. Nicotine increases cancer stem cell population in MCF-7 cells. *Biochem Biophys Res Commun* 2010; 403: 138-43.
- [77] Kohutek ZA, diPierro CG, Redpath GT, Hussaini IM. ADAM-10-mediated N-cadherin cleavage is protein kinase C-alpha dependent and promotes glioblastoma cell migration. *J Neurosci* 2009; 29: 4605-15.
- [78] Mani SA, Guo W, Liao MJ, *et al.* The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell* 2008; 133: 704-15.
- [79] Korkaya H, Paulson A, Charafe-Jauffret E, *et al.* Regulation of mammary stem/progenitor cells by PTEN/Akt/beta-catenin signaling. *PLoS Biol* 2009; 7: e1000121.
- [80] Zhou J, Wulfschuh J, Zhang H, *et al.* Activation of the PTEN/mTOR/STAT3 pathway in breast cancer stem-like cells is required for viability and maintenance. *Proc Natl Acad Sci USA* 2007; 104: 16158-63.
- [81] Lee CH, Huang CS, Chen CS, *et al.* Overexpression and activation of the alpha9-nicotinic receptor during tumorigenesis in human breast epithelial cells. *J Natl Cancer Inst* 2010; 102: 1322-35.
- [82] Sullivan JP, Spinola M, Dodge M, *et al.* Aldehyde dehydrogenase activity selects for lung adenocarcinoma stem cells dependent on notch signaling. *Cancer Res* 2010; 70: 9937-48.

[83] Grozio A, Paleari L, Catassi A, *et al.* Natural agents targeting the alpha7-nicotinic-receptor in NSCLC: a promising prospective in anti-cancer drug development. *Int J Cancer* 2008; 122: 1911-5.

[84] Catassi A, Paleari L, Servent D, *et al.* Targeting alpha7-nicotinic receptor for the treatment of pleural mesothelioma. *Eur J Cancer* 2008; 44: 2296-311.

Received: June 01, 2011

Revised: July 25, 2011

Accepted: August 01, 2011

Discovery of a Tamoxifen-Related Compound that Suppresses Glial L-Glutamate Transport Activity without Interaction with Estrogen Receptors

Kaoru Sato,^{*,‡,†} Jun-ichi Kuriwaki,^{‡,†} Kanako Takahashi,[‡] Yoshihiko Saito,[§] Jun-ichiro Oka,[§] Yuko Otani,^{||} Yu Sha,^{||} Ken Nakazawa,[‡] Yuko Sekino,[‡] and Tomohiko Ohwada^{*,||}


[‡]Laboratory of Neuropharmacology, Division of Pharmacology, National Institute of Health Sciences, 1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan

[§]Laboratory of Pharmacology, Faculty of Pharmaceutical Sciences, Tokyo University of Science, 2541 Yamazaki, Noda-city, Chiba 278-8510, Japan

^{||}Laboratory of Organic and Medicinal Chemistry, Graduate School of Pharmaceutical Sciences, University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

ABSTRACT: We recently found that tamoxifen suppresses L-glutamate transport activity of cultured astrocytes. Here, in an attempt to separate the L-glutamate transporter-inhibitory activity from the estrogen receptor-mediated genomic effects, we synthesized several compounds structurally related to tamoxifen. Among them, we identified two compounds, **1** (YAK01) and **3** (YAK037), which potently inhibited L-glutamate transporter activity. The inhibitory effect of **1** was found to be mediated through estrogen receptors and the mitogen-activated protein kinase (MAPK)/phosphatidylinositol 3-kinase (PI3K) pathway, though **1** showed greatly reduced transactivation activity compared with that of 17 β -estradiol. On the other hand, compound **3** exerted its inhibitory effect through an estrogen receptor-independent and MAPK-independent, but PI3K-dependent pathway, and showed no transactivation activity. Compound **3** may represent a new platform for developing novel L-glutamate transporter inhibitors with higher brain transfer rates and reduced adverse effects.

KEYWORDS: Tamoxifen, astrocyte, L-glutamate transporter, ER α , tetrasubstituted ethylene, nongenomic pathway



L-Glutamate (L-Glu) is one of the major excitatory neurotransmitters in the central nervous system (CNS), but high concentrations of extracellular L-Glu cause excessive stimulation of L-Glu receptors in the CNS, leading to neurotoxicity.^{1,2} Astrocyte L-Glu transporters are the only machinery available to remove L-Glu from extracellular fluid and to maintain a low and nontoxic concentration of L-Glu.³ Consequently, dysfunction of astrocyte L-Glu transporters is considered to be implicated in the pathology of neurodegenerative conditions.⁴ Therefore, exogenous compounds that can regulate the function of L-Glu transporters may provide chemical tools to investigate the regulatory mechanisms of these transporters at the molecular level, and would also be candidate therapeutic agents.

There is growing evidence that estrogen receptor (ER) α , which is a nuclear ER (nER) that mediates genomic effects, can also be translocated to plasma membranes and mediate acute nongenomic effects in some cases. We have clarified that 17 β -estradiol (E2) inhibits L-Glu transporters via a nongenomic pathway involving membrane-associated ER α (mER α).⁵ Tamoxifen (Tam), a synthetic estrogen analogue that is clinically used in the treatment of breast cancer to block the proliferative action of estrogens,⁶ also inhibited astrocyte L-Glu transporters at picomolar concentration, probably through the same nongenomic pathway as

E2.⁷ Because overexpression of astrocyte L-Glu transporters is often associated with neuropsychiatric disorders,⁴ inhibitors of L-Glu transporters may be clinically useful to ameliorate these disorders.⁸ However, Tam also acts on genomic pathways involving nuclear estrogen receptors (nERs) α and β , depending on the cell type and promoter context,⁹ and so may cause adverse effects including endometrial changes, depression and weight gain.^{10,11} Therefore, Tam-inspired compounds that retain the inhibitory effect on L-Glu transporters, but lack the nER-mediated genomic effects, would be useful tools for biological research, as well as candidate therapeutic agents.

Tam is a tetrasubstituted triphenylethylene derivative, in which the four substituents on the olefinic carbon atoms are different. This structural complexity makes the stereospecific synthesis of Tam-related derivatives difficult. We thus focused on Tam-inspired compounds bearing identical substituents on at least one of the olefinic carbon atoms.¹² It is well-known that the *N,N*-dimethylaminoethyl substituent on the phenolic oxygen atom and the regiochemistry of the tetrasubstituted

Received: September 29, 2011

Accepted: November 14, 2011

Published: November 14, 2011