

Table 1. How to determine the total activity of individual newborn

A	B	C	D	E	F	
No.	Real time	Time after start (s)	Weighting value of pup ^a (g)	Difference between the weighing value and that 0.1 s beforehand ($D_n = C_n - C_{n-1}$)	Absolute value ($E_n = D_n $)	Total activity ^b ($F_n = E_n + F_{n-1}$)
1	12:01:02			Before start of determination		
2	12:01:02			Before start of determination		
3	12:01:03	0	C3 = 2.0227	$D3 = +0$ ($C3 - C2 = 2.0227 - 2.0227$)	$E3 = 0.0000$	$F3 = 0.0000$ ($E3 + F2 = 0.0000 + 0.0000$)
4	12:01:03	0.1	C4 = 2.0231	$D4 = +0.0004$ ($C4 - C3 = 2.0231 - 2.0227$)	$E4 = 0.0004$	$F4 = 0.0004$ ($E4 + F3 = 0.0000 + 0.0004$)
5	12:01:03	0.2	C5 = 2.0233	$D5 = +0.0002$ ($C5 - C4 = 2.0233 - 2.0231$)	$E5 = 0.0002$	$F5 = 0.0006$ ($E5 + F4 = 0.0004 + 0.0002$)
6	12:01:03	0.3	C6 = 2.0233	$D6 = +0$ ($C6 - C5 = 2.0233 - 2.0233$)	$E6 = 0.0000$	$F6 = 0.0006$ ($E6 + F5 = 0.0006 + 0.0000$)
7	12:01:03	0.4	C7 = 2.0231	$D7 = -0.0002$ ($C7 - C6 = 2.0231 - 2.0233$)	$E7 = 0.0002$	$F7 = 0.0008$ ($E7 + F6 = 0.0006 + 0.0002$)
8	12:01:03	0.5	C8 = 2.0228	$D8 = -0.0003$ ($C8 - C7 = 2.0228 - 2.0231$)	$E8 = 0.0003$	$F8 = 0.0011$ ($E8 + F7 = 0.0008 + 0.0003$)
9	12:01:03	0.6	C9 = 2.0225	$D9 = -0.0003$ ($C9 - C8 = 2.0225 - 2.0228$)	$E9 = 0.0003$	$F9 = 0.0014$ ($E9 + F8 = 0.0011 + 0.0003$)
10	12:01:03	0.7	C10 = 2.0222	$D10 = -0.0003$ ($C10 - C9 = 2.0222 - 2.0225$)	$E10 = 0.0003$	$F10 = 0.0017$ ($E10 + F9 = 0.0014 + 0.0003$)
11	12:01:03	0.8	C11 = 2.0219	$D11 = -0.0003$ ($C11 - C10 = 2.0219 - 2.0222$)	$E11 = 0.0003$	$F11 = 0.0020$ ($E11 + F10 = 0.0017 + 0.0003$)
12	12:01:03	0.9	C12 = 2.0217	$D12 = -0.0002$ ($C12 - C11 = 2.0217 - 2.0219$)	$E12 = 0.0002$	$F12 = 0.0022$ ($E12 + F11 = 0.0020 + 0.0002$)
13	12:01:04	1.0	C13 = 2.0216	$D13 = -0.0001$ ($C13 - C12 = 2.0216 - 2.0217$)	$E13 = 0.0001$	$F13 = 0.0023$ ($E13 + F12 = 0.0022 + 0.0001$)
14	12:01:04	1.1	C14 = 2.0217	$D14 = +0.0001$ ($C14 - C13 = 2.0217 - 2.0216$)	$E14 = 0.0001$	$F14 = 0.0024$ ($E14 + F13 = 0.0023 + 0.0001$)
15	12:01:04	1.2	C15 = 2.0218	$D15 = +0.0001$ ($C15 - C14 = 2.0218 - 2.0217$)	$E15 = 0.0001$	$F15 = 0.0025$ ($E15 + F14 = 0.0024 + 0.0001$)

^aWeighing value of pup is a mouse newborn weight sent from the balance every 0.1 s.

^bIn order to determine the total activity (F) of a newborn from the start time of weighing to an individual time (B), each absolute value (E) was added up ($E_n + F_{n-1}$).




Figure 1. Total activities of intact newborn for 6 min (A) and the magnification (B, 0.4 s; see Table 1) of the area indicated by a broken line in (A).

Table 2. Integration of absolute values showing large movements of newborn

A	E	F
No.	Absolute value ($E_n = D_n $)	Total activity ($F_n = E_n + F_{n-1}$)
50	E50 = 0.0001	
51	E51 = 0.0001	
52	E52 = 0.0000	
53	E53 = 0.0003	F53 = 0.0003
54	E54 = 0.0005	F54 = 0.0008
55	E55 = 0.0007	F55 = 0.0015
56	E56 = 0.0011	F56 = 0.0026
57	E57 = 0.0008	F57 = 0.0034
58	E58 = 0.0004	F58 = 0.0038
59	E59 = 0.0002	F59 = 0.0040
60	E60 = 0.0001	
61	E61 = 0.0001	
62	E62 = 0.0000	
63	E63 = 0.0001	
64	E64 = 0.0001	

The large movements of the newborns were defined as activities showing an absolute value (E) of 0.0002 or more. Bold and nonbold entries show the large movements and small movements, respectively, of newborns.




Figure 2. An integration of only absolute values showing large movements of newborn.

group and the control group. In addition, viability on PND 5 in the MNU-treated group was comparable to that in the controls. On PND 5, 20 male offspring from five litters in each group were weighed, and we measured the length of the interhemispheric fissure and the length of rostral-caudal telencephalon to determine (the length of the interhemispheric fissure/the length of rostral-caudal telencephalon) ratio. The ratio in the group treated with MNU was significantly smaller than that in the controls. The body weight of male and female offspring treated with MNU was comparable to that of the control offspring when evaluated on PNDs 1 and 5. In addition, male and female pups had comparable body weights in both groups (data not shown).

In our previous studies, we demonstrated that embryos treated with MNU at 10 mg kg^{-1} on day 13 of gestation show thinning of the cortical plate and subventricular zone/ventricular zone in the dorsal telencephalon, and that embryonic day 13 is a critical period for microcephaly in mice (Fujiyama *et al.*, 2007; Komada *et al.*, 2010). Thus, we investigated the activity of PND 1 newborns exposed to MNU on day 13 of gestation in order to elucidate the scientific utilization of the developed behavioral testing method for evaluating the risk of neurotoxicity of MNU.

Activity of Newborns Exposed Prenatally to MNU

Figure 3 shows representative patterns concerning the activities for 6 min of male newborns exposed prenatally to MNU at 10 mg kg^{-1} and control newborns on PND 1. The shape, slope and area under the curve in each graph were clearly different between the MNU-treated group and the control group. Subsequently, the mean total activities of male newborns from 0 to 2 min, from 2 to 4 min and from 4 to 6 min at 12 p.m. on PND 1 were determined in the MNU-treated group and the control group according to the methods shown in Table 1 and Fig. 1(B). The weight data collected from 0 to 2, from 2 to 4 and from 4 to 6 min were analyzed with an approach that take into account the multiple measures from each animal using a repeated measures ANOVA. There were significant differences in the activity and measurement time in both sexes [males, $F(2, 27) = 6.36$, $P = 0.0093$; females, $F(2, 27) = 4.24$, $P = 0.0334$]. Activities for the first two time blocks in the MNU-treated group showed no significant difference from those in the controls, and those from 4 to 6 min in the group treated with MNU were significantly higher than those in the controls. The ANOVA for males and females revealed main effects of treatment [$F(1, 8) = 17.47$, $P = 0.0031$, and $F(1, 8) = 22.94$, $P = 0.0014$, respectively]. No significant differences were detected between those from 0 to 2 and from 2 to 4 min in the group treated with MNU and those in the controls in both sexes. The total activities for 6 min in the MNU-treated group were significantly higher than those in the control group in both sexes (Fig. 4). The ANOVA for males and females revealed main effects of treatment [$F(1, 8) = 7.75$, $P = 0.0238$, and $F(1, 8) = 24.32$, $P = 0.0011$, respectively]. In addition, the ANOVA revealed no effect of MNU on body weight in either males or females [$F(1, 8) = 0.15$, $P = 0.7112$, and $F(1, 8) = 0.091$, $P = 0.7704$, respectively].

There are numerous behavioral tests to evaluate learning and memory, and the activity of rats and mice with microcephaly induced by neurotoxicants using Biel maze and radial arm maze, shuttlebox and wheel cage (Akaike *et al.*, 1994). Ohta *et al.* (1997) demonstrated that rat offspring with microcephaly induced by prenatal MNU exposure exhibited increased wheel cage activity when evaluated at 6 and 9 weeks of age, suggesting hyperactivity in adult rats exposed *in utero* to a neurotoxicant. However, few neurobehavioral tests have been developed in




Figure 3. Representative pattern of activities for 6 min of PND 1 male newborns from three dams each in the control group and the MNU-treated group.




Figure 4. Mean total activities of PND 1 newborns from five dams each in the control group and the MNU-treated group. Vertical lines represent standard deviations. * Significantly different from the control, $P < 0.05$; ** significantly different from the control, $P < 0.01$.

order to evaluate the newborn behavior and activity. Hashimoto and Mizutani (1991) showed early neurobehavioral disorders in microcephalic rat offspring induced by prenatal exposure to MNU at 5 mg kg^{-1} . The microcephalic pups during the suckling period, PND 0–22, were retarded in terms of neurobehavioral ontogeny. They showed impaired performance, such as paired limb movement, clumsy locomotion or hyperreflexive reaction, without growth retardation. In their study, some behavioral disorders such as strengthened hyperactivity, increased susceptibility to seizure and deficit in rotarod performance were detected in the neonatal rats after PND 3. Neurobehavioral changes of

newborns on PND 0 or 1 have not been detected in any developed behavioral testing methods. In the present study, we clearly showed the quantitative changes of neurobehavior of mouse newborns on PND 1.

Of particular importance is consideration of the limitations and potential pitfalls of the proposed technique. The measurement proposed in this study uses changes in weight on the balance as the measure of activity. The same degree of movement in mouse newborns of different weights (i.e. ages and gender) would produce different changes in weight detected on the balance. Thus, on the basis of changes in weight, this would suggest that newborns of different weights have different levels of activity when they actually have comparable levels of activity. The same amount of movement in PND 1 and PND 5 newborns would translate into different weight changes, suggesting different levels of activity. The consequence of this is that using changes in weight as proposed cannot be used directly to compare movement between animals of different sizes (e.g. males vs females, different ages).

In conclusion, we have developed a rodent newborn behavioral testing method using an analytical balance. This developed technique is a practical approach to solve one of the challenges in assessing early effects of neurotoxicants and it may provide a useful addition to neurobehavioral assessment in very young rodents. As for the advantages of our method, it is considered that there is a possibility of early detection of neurobehavioral abnormalities and early medical treatment by means of the

analyses of newborn activity by this test method, as well as the measurement of newborn activity without stress such as pain. As the next step of our neurobehavioral studies using this technique, the dose–response relationship between newborn activities and the doses of a neurotoxicant should be clarified. The new technique proposed is simple. If this approach is validated, it may have the potential to become widely used.

Acknowledgments

We thank all our colleagues, including M. Shirazaki, T. Inada and H. Irie, who contributed to the present study. This study was partly supported by Grants-in-Aid for Scientific Research from the Japanese Ministry of Health, Labour and Welfare.

Conflict of Interest

The authors declare that there are no conflicts of interest

References

- Akaike M, Ohno H, Tsutsumi S, Omosu M. 1994. Comparison of four spatial maze learning tests with methylnitrosourea-induced microcephaly rats. *Teratology* **49**: 83–89.
- Bearer CF. 2000. The special and unique vulnerability of children to environmental hazards. *Neurotoxicology* **6**: 925–934.
- Burbacher TM, Rodier PM, Weiss B. 1990. Methylmercury developmental neurotoxicity: a comparison of effects in humans and animals. *Neurotoxicol. Teratol.* **12**: 191–202.
- Francis EZ, Kimmel CA, Rees DC. 1990. Workshop on the qualitative and quantitative comparability of human and animal developmental neurotoxicity: summary and implications. *Neurotoxicol. Teratol.* **12**: 285–292.
- Fujiyama F, Saito Y, Nagao T. 2007. Effects of prenatal exposure to methylnitrosourea on the developing brains of mouse offspring. *Congen. Anom.* **47**: A29.
- Hashimoto Y, Mizutani M. 1991. Early neurobehavioral disorders in the micrencephalic offspring induced by prenatal treatment with *N*-methyl-*N*-nitrosourea or methylazoxymethanol in the rat. *J. Vet. Med. Sci.* **53**: 643–649.
- Hass U. 2006. The need for developmental neurotoxicity studies in risk assessment for developmental toxicity. *Reprod. Toxicol.* **22**: 148–156.
- Kimmel CA. 1988. Current status of behavioral teratology: science and regulation. *Crit. Rev. Toxicol.* **19**: 1–10.
- Komada M, Fujiyama F, Yamada S, Shiota K, Nagao T. 2010. Methylnitrosourea induces neural progenitor cell apoptosis and microcephaly in mouse embryos. *Birth Defects Res. (Part B)* **89**: 213–222.
- Ohta R, Matsumoto A, Hashimoto Y, Nagao T, Mizutani M. 1997. Behavioral characteristics of micrencephalic rats in high and low shuttlebox avoidance lines. *Neurotoxicol. Teratol.* **19**: 157–162.

Developmental effects of oral exposure to diethylstilbestrol on mouse placenta

Tetsuji Nagao,^{a*} Nao Kagawa,^b Yoshiaki Saito^c and Munekazu Komada^d

ABSTRACT: Placental growth and function are of biological significance in that placental tissue promotes prenatal life and the maintenance of pregnancy. Exposure to synthetic estrogens causes embryonic mortality and placental growth restriction in mice. The aim of the present study was to examine the effects of diethylstilbestrol (DES) on placenta in mice. DES at 1, 5, 10 or 15 $\mu\text{g kg}^{-1} \text{day}^{-1}$, or 17 β -estradiol (E_2) at 50 $\mu\text{g kg}^{-1} \text{day}^{-1}$, was administered orally to ICR mice on days 4 through to 8 of gestation. Expression of ER α , ER β , ERR β or ERR γ mRNA in the junctional or labyrinth zone of the placentas on day 13 was assessed using RT-PCR, as well as the embryonic mortality, embryonic and placental weight, histological changes of labyrinth and ultrastructural changes of the trophoblast giant cells (TGCs). Embryo mortalities in the DES 10 and 15 $\mu\text{g kg}^{-1} \text{day}^{-1}$ groups were markedly increased. No significant changes in embryonic and placental weight were observed in any DES- or E_2 -exposed groups. Expression of ER α mRNA in the junctional zone with male embryos in the 5 $\mu\text{g kg}^{-1} \text{day}^{-1}$ group was significantly higher than that in the control, whereas expression was not determined in the 15 $\mu\text{g kg}^{-1} \text{day}^{-1}$ group. Histological observation revealed that the placentas exposed to DES at 10 $\mu\text{g kg}^{-1} \text{day}^{-1}$ lacked the developing labyrinth. Ultrastructural observation of the TGCs showed poor rough-surfaced endoplasmic reticulum in the DES 10 $\mu\text{g kg}^{-1} \text{day}^{-1}$ group. The present data suggest that developmental changes induced by DES may be related to interference with the nutrition and oxygen exchange between mother and embryo or decreased protein synthesis, resulting in a high frequency of embryo mortality. Copyright © 2012 John Wiley & Sons, Ltd.

Keywords: diethylstilbestrol; placental development; trophoblast giant cells; labyrinth; rough-surfaced endoplasmic reticulum

Introduction

The placenta is the first organ to form during mammalian embryogenesis. Problems in its formation and function underlie many aspects of early pregnancy loss and pregnancy complications in humans. Trophoblast cells are the first lineage to form in the mammalian conceptus and contribute exclusively to the extraembryonic structures that form the placenta (Cross *et al.*, 1994), which is critical for the survival of mammalian embryos. After implantation, the mural trophoblast differentiates into the primary trophoblast giant cells (TGCs). The development of TGCs is essential to embryo implantation and to the maintenance of pregnancy. The placenta undertakes many functions beyond simple exchange between the maternal and fetal environment. The mouse placenta is a highly tuned exchange bed with supporting cells involved in both structural and functional aspects of fetal–maternal exchange. The mouse and the human both have discoid hemochorial placentas, but the mouse placenta is divided into two morphologically and functionally distinct regions known as junctional (basal) and labyrinth zones. The junctional zone, which includes TGCs and spongiotrophoblast cells, is involved in the formation of hormones, angiogenic factors, vascular endothelial growth factor, tissue remodeling factors and urokinase-type plasminogen activator. The labyrinth zone, on the other hand, is the major site of feto–maternal exchange, the point of nutrient transfer between the maternal and fetal blood spaces (Rossant and Cross, 2001; Georgiades *et al.*, 2002; Malassiné *et al.*, 2003).

Diethylstilbestrol (DES) is a synthetic (stilbene) estrogen with an *in vivo* estrogenic potency similar to that of 17 β -estradiol (E_2), and is structurally related to E_2 . DES (daily dose of 5–125 mg during

pregnancy, Coscrove *et al.*, 1977; Oradell, 1961) was widely used to prevent a threatened miscarriage from the 1940s to 1971. DES can be metabolized to conjugates (glucuronide and sulfate) and to numerous oxidative metabolites (reviewed by Metzler, 1981). The teratogenicity, reproductive toxicity and carcinogenicity of DES in humans are believed to be transduced via the classic estrogen receptors ER α and ER β (Couse and Korach, 1999; Greco *et al.*, 1993). DES has also been shown to induce placental changes that include a reduction of the labyrinth zone and accumulation of TGCs in mid- and late gestation in mice (Scott and Adejokun, 1980). These effects are unlikely to involve the classic ERs as genetic ablation of both receptors, singly or in combination, did not reveal a role for these receptors in placentation (Couse and Korach, 1999). In our previous study, we demonstrated that decidual hypoplasia and subsequent placental hemorrhage cause fetal death as a result of the oral administration of DES during the early stage of pregnancy in mice (Nagao and Yoshimura, 2009).

*Correspondence to: T. Nagao, Department of Life Science, Faculty of Science and Technology, Kinki University, 3-4-1 Kowakae, Higashi-Osaka 577-8502, Japan. E-mail: tnagao@life.kindai.ac.jp

^aLaboratory of Developmental Biology, Department of Life Science, Kinki University, Osaka 577-8502, Japan

^bLaboratory of Animal Genetics, Department of Life Science, Kinki University, Osaka 577-8502, Japan

^cLaboratory of Pathology, Food and Drug Safety Center, Kanagawa 257-8523, Japan

^dDivision of Cell Biology and Neuroscience, Faculty of Medical Sciences, University of Fukui, Fukui 910-1193, Japan

Pharmacological doses of estrogens cause high levels of embryonic mortality (Haddad and Ketchel, 1969), although administration later in pregnancy causes retarded fetal growth with less of an effect on survival (Kuhn *et al.*, 1982). In order for estrogens to exert their biological effects, they need to bind to the ER, which then undergoes a conformational change allowing it to interact with chromatin and to modulate transcription of target genes (Tsai and O'Malley, 1994). ER α and ER β proteins are physiologically relevant; they bind estradiol with high affinity and activate transcription of estrogen-responsive reporter gene constructs expressed in mammalian cell lines (Kuiper *et al.*, 1996, 1997). Little information on the expression profile of ER mRNA in developing rodent placentas is available. The expressions of ER isoforms and variants have been studied in rat placenta on days 16, 19 and 21 of gestation, and it has been demonstrated that ER α and ER β mRNA were detected from as early as day 16 of gestation (Al-Bader, 2006). In mouse placenta, the highest expressions of ER α and ER β mRNA were determined on day 14 of gestation (Nagao *et al.*, unpublished data).

Classic endocrine studies have shown that steroid hormones are required for the maintenance of pregnancy and placental viability. The estrogen receptor-related receptor β (ERR β) is an orphan member of the superfamily of nuclear hormone receptors (Giguère *et al.*, 1988; Knöfler *et al.*, 2001). The protein is homologous to the ER and binds the estrogen response element. Luo *et al.* (1997) reported that ERR β has an important role in early placentation. Tremblay *et al.* (2001) demonstrated that DES promotes coactivator release from ERR β and inhibits its transcriptional activity, and the treatment of trophoblast stem cells with DES led to their differentiation towards the polyploidy giant-cell lineage, and that DES-treated pregnant mice exhibited abnormal early placenta development associated with an overabundance of TGCs and an absence of diploid trophoblasts.

To address the adverse effects of oral DES administration on placentogenesis resulting in embryonic death, the present study was designed to examine the ultrastructural changes of the TGCs in the trophoblast layer between the spongiotrophoblast layer and the decidua basalis, and the expression of ER or ERR mRNA. In addition, the development of the labyrinth zone where most gaseous and nutrient exchange occurs was investigated.

Materials and Methods

Animals

Eight-week-old ICR mice purchased from CLEA (Osaka, Japan) were used in the present study after acclimation for 1 to 2 weeks. The experimental protocols were approved by the Animal Care and Use Committee of Kinki University. Mice were kept under specific pathogen free conditions and housed in polycarbonate cages in a room in which the temperature (23 ± 1 °C), humidity ($50 \pm 5\%$) and light cycle (12: 12 h light/dark cycle; lights on at 07.00 hours) were maintained. To avoid the possibility of stressing the animals, noise levels were kept to a minimum both within the room and in the adjacent areas. Mice were allowed food (Certified Rodent Chow CE-2; CLEA, Osaka, Japan) and drinking water *ad libitum*. Certification analysis of each lot of diet was performed by the manufacturer. The same lots of diet were provided to animals from control and compound-exposed groups at the same times, in order to control across groups for possible variation in the content of the diet. Ten-week-old virgin female mice weighing 29–34 g were cohabited

for 2 h (07.00 to 09.00 hours) on a 1: 1 basis with males aged 11 weeks or older. Females were checked for the presence of a vaginal plug immediately thereafter, and 08.00 hours was defined as the time of conception if a plug was found (Nagao *et al.*, 2000).

Exposure of Pregnant Dams to DES and Isolation of Placental Tissues

Diethylstilbestrol (DES, (E)-3,4-Bis(4-hydroxyphenyl)-3-hexene) and 17 β -estradiol (E₂) were purchased from Sigma Chemical (St. Louis, MO, USA), suspended in corn oil and administered orally on days 4 through to 8 of gestation, a time in the early development of the placenta when ERR β expression is essential (Tremblay *et al.*, 2001). The dose solution was prepared once per 5 days and analyzed prior to dosing. The DES or E₂ concentration was confirmed to be within $\pm 10\%$ of the targeted concentration. Mice administered corn oil (5 ml kg^{-1} body weight) were used as controls. Ten mice each were administered DES at 1, 5, 10 or $15 \mu\text{g kg}^{-1} \text{ day}^{-1}$, or E₂ at $50 \mu\text{g kg}^{-1} \text{ day}^{-1}$. Administration occurred at a defined time (12.00 to 12.15 hours). On day 13 of gestation, pregnant mice were euthanized using carbon dioxide anesthesia followed by cervical dislocation, and the uteri from dams in each group were removed quickly to count the numbers of live and dead embryos as well as to measure embryonic and placental weights, and the determination of sex was carried out by observation of the gonadal morphology and position under a dissecting microscope to examine the embryo-sex-dependent differences. Subsequently, the placentas were quickly washed in sterile Tyrode's solution.

Determination of ER α , ER β , ERR β or ERR γ mRNA in the Developing Placentas Using RT-PCR

In two placentas with female embryos and two placentas with male embryos of each dam, the junctional zone (decidua basalis and spongiotrophoblasts) and the labyrinth zone were separated using a scalpel under a dissecting microscope (SZX12, Olympus Co., Tokyo, Japan). In the separation of zones, the boundary layer between the spongiotrophoblasts and the labyrinth was excluded from the sample for analysis in order to avoid the mixture of the 'peg' of spongiotrophoblast cells into the labyrinthine part of the placenta. Tissues from each zone in the placentas of the groups exposed to DES at 0, 5 or $10 \mu\text{g kg}^{-1} \text{ day}^{-1}$, and E₂ at $50 \mu\text{g kg}^{-1} \text{ day}^{-1}$ were provided for RT-PCR to determine the mRNA levels of ER α , ER β , ERR β and ERR γ .

Total RNA was isolated from placental tissues from each zone using Isogen (Nippon Gene Co., Ltd., Tokyo, Japan). Complementary DNA (cDNA) was synthesized from $1 \mu\text{g}$ of purified total RNA using AMV Reverse Transcriptase XL from the RT-PCR Kit (Takara Bio Inc., Shiga, Japan) with a random primer at 42 °C for 60 min. PCR amplification of ER α , ER β , ERR β or ERR γ was carried out on equal quantities of cDNA product ($20 \mu\text{l}$) using a TaKaRa PCR Thermal Cycler Dice mini (Takara Bio Inc., Shiga, Japan). The sequences of ER α , ER β , ERR β and ERR γ primers and conditions of PCR amplification of cDNA are shown in Table 1. PCR products were run on a 1.5% Agarose S gel (Nippon Gene Co., Ltd., Tokyo, Japan) and visualized by ethidium bromide (Nippon Gene Co., Ltd., Tokyo, Japan) staining under ultraviolet (UV) light. Gene expression levels were normalized by the expression levels of β -actin as an internal control.

Table 1. Sequences of PCR primers

Primer name		Sequences	Products (bps)	Annealing temperature
β -actin	Forward	tgttaccaactgggacgaca	392	58 °C
	Reverse	tctcagctgtggtggaag		
ER α	Forward	atgaaaggcggcatcacgaaag	94	60 °C
	Reverse	caccatttcatttcgccttc		
ER β	Forward	ccagactgcaagccaaatgt	81	60 °C
	Reverse	agaagcgatgattggcagtg		
ERR β	Forward	cgctcgctcccatcacgac	78	60 °C
	Reverse	ccctaccaggcgagagtgttc		
ERR γ	Forward	actccatgcccaagagactg	137	60 °C
	Reverse	ggctggcgagctgtactcta		

Observation of the Placental Morphology

Three to four placentas of each dam in the groups exposed to DES at 0, 1, 5 or 10 $\mu\text{g kg}^{-1} \text{day}^{-1}$, and E₂ at 50 $\mu\text{g kg}^{-1} \text{day}^{-1}$ were fixed for 24 h in 0.1 M phosphate-buffered 10% formalin embedded in paraffin, and sectioned for staining with hematoxylin and eosin. The thicknesses of the whole placenta and labyrinth zone close to the central portion were measured to calculate the proportions of labyrinth zone per whole placenta (Fig. 1). Subsequently, the number of TGCs in the trophoblast layer between the spongiotrophoblast layer and decidua basalis was counted using a calibrated ocular grid in all visual fields with $\times 200$ objective magnification. Three separate fields were randomly selected in each placenta and the number of TGCs was counted according to the methods previously reported (Suzuki *et al.*, 1997).

For ultrastructural observation, one to two placentas in each dam in the groups exposed to DES at 0, 1, 5 or 10 $\mu\text{g kg}^{-1} \text{day}^{-1}$, and E₂ at 50 $\mu\text{g kg}^{-1} \text{day}^{-1}$ were selected at random. The placentas from the uteri were fixed in 0.1 M phosphate-buffered 2% paraformaldehyde and 1.25% glutaraldehyde (pH 7.4). After fixation, placentas were rinsed three times in phosphate buffer, post-fixed for 2 h at 4 °C in 0.1 M phosphate-buffered 2% osmium tetroxide and dehydrated in alcohol. Placentas

were embedded in epoxy resin. Semithin sections (about 1 μm) were stained with toluidine blue, and histopathological morphology was observed. Ultrathin sections stained with uranyl acetate and lead citrate were prepared from a representative area of the basal zone of the placenta and TGCs were observed in the trophoblast layer between the spongiotrophoblast layer and decidua with an electron microscope (HT7700, Hitachi, Japan).

Statistical Analysis

Data are given as the mean \pm standard deviation (SD). Comparisons between the treated groups and controls were performed using $P \leq 0.05$ and $P \leq 0.01$ as levels of significance. Mean fetal weight and placental weight, and the number of TGCs were compared using Student's *t*-test. The proportions of resorbed embryos and labyrinth zone/whole placenta were compared using the Mann-Whitney *U*-test as described by Siegel (1956) to determine the significance of differences. As for RT-PCR, statistical analysis was performed using SPSS (ANOVA followed by post-hoc analysis [LSD]) when the test for homogeneity of variance was fulfilled and using Games-Howell post-hoc analysis when the homogeneity of variance was not attained.

Day 13 murine placenta




Figure 1. Diagrammatic transverse section of placenta showing the measurement of thickness. The measurement was performed with three sections close to the central portion from each placenta, and five areas (No. 1–5) close to the highest portion in each section. Fifteen values of A, B or B/A in each placenta were obtained, and the mean value of A, B or B/A in each placenta was determined. Two to three placentas per dam were used. (A) thickness of whole placenta, (B) thickness of labyrinth zone.

Results

Embryonic Mortality, and the Weight of Embryos and Placentas

No adverse effects on the general conditions: decreased body weight gain, death, moribund, loss of spontaneous activity, crouching, tremor, piloerection, loose stool and so on, in maternal animals were observed in any DES- or E₂-exposed groups. The developmental effects of DES and E₂ on the mortality of embryos, and embryonic and placental weights by sex, when evaluated on days 13 of gestation, are presented in Table 2. After oral exposure to DES on days 4 through to 8 of gestation, the embryonic mortality in the groups treated with DES at 10 or 15 $\mu\text{g kg}^{-1} \text{day}^{-1}$ was significantly increased compared with that of the controls. Most of the dead embryos in the group exposed to DES at 10 $\mu\text{g kg}^{-1} \text{day}^{-1}$ were resorbed at the early to middle developmental stage. Dead embryos in the group exposed to DES at 15 $\mu\text{g kg}^{-1} \text{day}^{-1}$ almost all involved early resorptions, or implantation sites, suggesting death immediately after early administration. Embryonic mortality in the group exposed to

Table 2. Embryonic mortality, placental weight and embryonic weight in dams exposed to diethylstilbestrol (DES) or 17 β -estradiol (E₂) on days 4 through to 8 of gestation

Dose ($\mu\text{g kg}^{-1} \text{ day}^{-1}$)	No. of dams	Embryonic mortality (%)	No. of dams with dead embryos	Placental weight (g, A)		Embryonic weight (g, B)		B/A	
				Male	Female	Male	Female	Male	Female
DES 0	10	2.8 \pm 4.1 ^a (4/117) ^b	2 (0) ^c	0.096 \pm 0.013 (75) ^d	0.091 \pm 0.014 (68) ^d	0.183 \pm 0.034 (75) ^e	0.178 \pm 0.031 (68) ^e	1.919 \pm 0.339	1.937 \pm 0.333
1	10	5.4 \pm 6.3 (8/150)	1 (0)	0.087 \pm 0.009 (76) ^f	0.084 \pm 0.005 (64)	0.220 \pm 0.036 (78)	0.208 \pm 0.035 (64)	2.290 \pm 0.491	2.276 \pm 0.535
5	10	5.5 \pm 5.2 (8/145)	2 (0)	0.094 \pm 0.010 (64)	0.088 \pm 0.012 (73)	0.173 \pm 0.014 (64)	0.167 \pm 0.014 (73)	1.839 \pm 0.307	1.894 \pm 0.281
10	10	68.5 \pm 44.7** (104/151)	9 (2)	0.100 \pm 0.006 (27)	0.095 \pm 0.007 (20)	0.177 \pm 0.013 (27)	0.169 \pm 0.016 (20)	1.773 \pm 0.115	1.781 \pm 0.203
15	10	88.2 \pm 26.5** (121/137)	10 (6)	0.105 \pm 0.008 (9)	0.096 \pm 0.008 (7)	0.163 \pm 0.018 (9)	0.165 \pm 0.007 (7)	1.550 \pm 0.118**	1.628 \pm 0.136*
E ₂ 50	10	4.91 \pm 2.86 (7/152)	1 (0)	0.093 \pm 0.013 (68)	0.090 \pm 0.009 (77)	0.193 \pm 0.018 (68)	0.188 \pm 0.017 (77)	2.008 \pm 0.303	1.998 \pm 0.258

^aMean \pm SD
^bNo. of dead embryos/no. of implants
^cNo. of dams with no survivors.
^dNo. of placentas with male or female embryos.
^eNo. of male or female embryos.
^fTwo placentas were not used for further analysis because they were fused *in utero*.
* Significantly different from the control, $P < 0.05$.
** Significantly different from the control, $P < 0.01$.

DES at 1 or 5 $\mu\text{g kg}^{-1} \text{day}^{-1}$ was comparable to that of the controls. No significant increase in the mortality was observed in the group exposed to E_2 at 50 $\mu\text{g kg}^{-1} \text{day}^{-1}$.

There were no significant differences between the DES-treated groups and the controls in terms of weights of placentas and embryos in both sexes, whereas the placental weight showed a tendency to increase and the embryonic weight showed a tendency to decrease in the group exposed to DES at 10 or 15 $\mu\text{g kg}^{-1} \text{day}^{-1}$ in comparison with those in the controls. The placental or embryonic weight in both sexes of the E_2 -treated group was comparable to that in the control.

ER or ERR mRNA Expression in the Placentas

Expression of $\text{ER}\alpha$ and $\text{ER}\beta$ mRNA, and $\text{ERR}\beta$ and $\text{ERR}\gamma$ mRNA in the placentas of dams exposed to DES or E_2 is shown in Fig. 2.




Figure 2. The expression of $\text{ER}\alpha$, $\text{ER}\beta$, $\text{ERR}\beta$ and $\text{ERR}\gamma$ mRNA in the junctional zone (\square) and labyrinth zone (\blacksquare) of placentas with male or female embryos of dams exposed to diethylstilbestrol (DES) or 17 β -estradiol (E_2) on days 4 through to 8 of gestation. *Significantly different from the control, $P < 0.05$.

Expression of $\text{ER}\alpha$ mRNA in the junctional zone (decidua basalis and spongiotrophoblasts) of placentas with male embryos in the group exposed to DES at 5 $\mu\text{g kg}^{-1} \text{day}^{-1}$ was significantly higher than that in the control ($P = 0.003$). No significant differences in any $\text{ER}\alpha$, $\text{ER}\beta$, $\text{ERR}\beta$ and $\text{ERR}\gamma$ mRNA expression were observed between the group exposed to DES at 10 $\mu\text{g kg}^{-1} \text{day}^{-1}$ or the group exposed to E_2 at 50 $\mu\text{g kg}^{-1} \text{day}^{-1}$ and the controls in both decidua basalis with spongiotrophoblasts and the labyrinth zone of the placentas with male embryos. In the placentas with female embryos, there were no significant differences in the mRNA expression of $\text{ER}\alpha$, $\text{ER}\beta$, $\text{ERR}\beta$ and $\text{ERR}\gamma$ between the groups exposed to DES or E_2 and the controls.

Morphology of Placenta

Light microscopy of the placenta clearly showed that the group exposed to DES at 10 $\mu\text{g kg}^{-1} \text{day}^{-1}$ lacked the developing labyrinth (Fig. 3). The decidua was much less affected than the labyrinth in the DES-exposed group, but decidual cells were smaller than in the controls. Ultrastructural changes in the TGCs of the placentas sampled on day 13 of gestation are shown in Fig. 4. There were rich rough-surfaced endoplasmic reticulum and Golgi apparatus in the TGCs of the group exposed to DES at 1 or 5 $\mu\text{g kg}^{-1} \text{day}^{-1}$ or E_2 at 50 $\mu\text{g kg}^{-1} \text{day}^{-1}$, and in the control group (Fig. 4A). In the group exposed to DES at 10 $\mu\text{g kg}^{-1} \text{day}^{-1}$, poor rough-surfaced endoplasmic reticulum or its atrophy was found (Fig. 4B).

The decidua basalis, spongiotrophoblasts and labyrinth zone were easily discriminated for both control and DES-treated groups. The developmental parameters of placenta are shown in Table 3. The thickness of the whole placenta and that of the labyrinth zone in the group treated with DES at 10 $\mu\text{g kg}^{-1} \text{day}^{-1}$ were significantly decreased compared with those of the controls. Those in the group exposed to DES at 1 or 5 $\mu\text{g kg}^{-1} \text{day}^{-1}$ or E_2 at 50 $\mu\text{g kg}^{-1} \text{day}^{-1}$ were comparable to those of the controls. No significant differences were found in the proportions of the labyrinth zone per whole placenta between the DES- or E_2 -treated groups and the controls. Numbers of TGCs per unit area of decidua basalis and spongiotrophoblasts are also shown in Table 3. The number of TGCs in the group exposed to DES at 10 $\mu\text{g kg}^{-1} \text{day}^{-1}$ was significantly increased compared with that in the controls. There were no significant differences in the number between the group exposed to DES at 1 or 5 $\mu\text{g kg}^{-1} \text{day}^{-1}$ or E_2 at 50 $\mu\text{g kg}^{-1} \text{day}^{-1}$ and the controls.

Discussion

It is well known that estrogen administration elicits hemorrhage around gestational sacs and decreases placental blood flow in the mouse and rabbit (Abdul-Karim and Bruce, 1972; Mahendroo *et al.*, 1997; Scott and Adejokum, 1980). In our previous study, we demonstrated that decidual hypoplasia and subsequent placental hemorrhage cause fetal death as a result of the oral administration of DES during the early stage of pregnancy in mice (Nagao and Yoshimura, 2009).

The present study clearly showed that placental development was affected after oral exposure to DES, resulting in prenatal lethality. It has been reported that DES acts in rats to depress the preterm levels of steroid hormones, which leads to a failure of uterine contraction accompanied by placental detachment and fetal death (Clevenger *et al.*, 1991; Zimmerman *et al.*, 1991). Bartholomeusz *et al.* (1999) demonstrated a decrease in embryo




Figure 3. Light microscopic profiles of transverse sections of the control placenta (A, C) of a dam on day 13 of gestation showing a rich villous network and of the placenta (B, D) of a dam exposed to diethylstilbestrol (DES) at $10 \mu\text{g kg}^{-1} \text{day}^{-1}$ on days 4 through to 8 of gestation showing impaired villous network. Upper panels (A, B) show the peripheral part of placentas and the labyrinth zone (L) was magnified (C, D). Toluidine blue staining.

survival and disturbance of fetal and placental growth after elevation of maternal estradiol blood concentration in rats.

Embryonic Mortality and Placental Growth

The incidences of embryonic mortality after treatment with DES at 0, 1, 5, 10 and $15 \mu\text{g kg}^{-1} \text{day}^{-1}$ were 2.8%, 5.4%, 5.5%, 68.5% and 88.2%, respectively, suggesting that the no observed adverse effect dose level (NOAEL) for embryonic lethality was $5 \mu\text{g kg}^{-1} \text{day}^{-1}$ upon oral exposure on days 4 through to 8 of gestation and evaluation on day 13 in ICR mice. In the 1940s, DES was used to prevent adverse pregnancy outcomes in

women with a history of miscarriage. Evaluating the DES applied dose as $5\text{--}125 \text{ mg day}^{-1}$ (Cosgrove *et al.*, 1977; Oradell, 1961), a 150-pound (assuming an average body weight of 68 kg) pregnant woman could intake $74\text{--}1800 \mu\text{g kg}^{-1} \text{day}^{-1}$, and the dose is 15–360 times higher than the NOAEL for embryonic lethality evaluated in the present study. In a previous study, it was reported that the embryonic mortality after oral exposure to DES at $10 \mu\text{g kg}^{-1} \text{day}^{-1}$ on days 4 through to 8 of gestation was approximately 25% when evaluated on day 9 of gestation in mice (Nagao and Yoshimura, 2009). Thus, it is suggested that many embryos of dams exposed to DES at this dose on days 4 through to 8 proceeded to death during days 9 to 13 of gestation.




Figure 4. Electron micrographs of trophoblast giant cells in the control placenta (A) of a dam on day 13 of gestation showing rich rough-surfaced endoplasmic reticulum (*) and Golgi apparatus (l), and in the placenta (B) of a dam exposed to diethylstilbestrol (DES) at $10 \mu\text{g kg}^{-1} \text{day}^{-1}$ on days 4 through to 8 of gestation showing depletion of rough-surfaced endoplasmic reticulum.

Table 3. Developmental effects of diethylstilbestrol (DES) or 17 β -estradiol (E₂) on labyrinth and trophoblast giant cells (TGCs)

Dose ($\mu\text{g kg}^{-1} \text{ day}^{-1}$)	Thickness of whole placenta (A)	Thickness of labyrinth zone (B)	B/A \times 100	Number of TGCs ^b
DES 0	1382.9 \pm 156.4 ^a	708.3 \pm 142.0 ^a	51.3 \pm 8.4	14.8 \pm 4.4
1	1395.0 \pm 147.1	727.3 \pm 120.9	52.7 \pm 7.8	13.6 \pm 3.7
5	1405.6 \pm 155.0	765.3 \pm 173.9	54.2 \pm 9.8	12.3 \pm 4.4
10	1284.6 \pm 156.2 ^{**}	634.0 \pm 126.5 ^{**}	49.4 \pm 8.1	19.6 \pm 3.6*
E ₂ 50	1409.1 \pm 94.7	688.6 \pm 80.3	49.0 \pm 6.0	16.3 \pm 3.9

^aMean \pm SD, μm .
^bThe number of TGCs was counted under a light microscope at \times 200 magnification in three areas (decidua basalis and spongiotrophoblasts) randomly selected from three to four placentas of each dam.
* Significantly different from the control, $P < 0.05$.
** Significantly different from the control, $P < 0.01$.

Oral exposure to DES at 10 or 15 $\mu\text{g kg}^{-1} \text{ day}^{-1}$ resulted in a high frequency of resorptions of embryos. Interestingly, a significant increase in the resorption did not occur when E₂ at a fairly high dose of 50 $\mu\text{g kg}^{-1} \text{ day}^{-1}$ was applied (4.9%). Orally administered DES is rapidly absorbed, distributed and cleared in maternal organs, and its major site of accumulation is the liver of mice (Shah and McLachlan, 1976). Maternally administered DES crosses the placenta and is rapidly distributed to the fetus in rats, mice, hamsters and monkeys (Fischer *et al.*, 1976; Hill *et al.*, 1980; Maydl and Metzler, 1984; McLachlan, 1977); the fetal level is 2–3 times higher than in the maternal blood (Miller *et al.*, 1982). Estradiol given orally in its native crystalline state is poorly absorbed via the gastro-intestinal tract because of extensive first-pass metabolism (Lobo, 1987). Estrogen given orally in mice does not produce continuously elevated levels of estrogen and is rapidly metabolized to control levels (Gordon *et al.*, 1986).

The biological activity of estrogens is influenced by the degree to which they bind to serum proteins. DES binds poorly to sex hormone binding globulin (SHBG), and DES showed enhanced access in serum (serum modified access, SMA = 6.2). SMA was calculated by dividing the relative binding affinity (RBA, relative to estradiol) measured in 100% serum by the RBA measured in serum-free medium. An SMA $>$ 1 indicated that the xenoestrogen had greater access to estrogen receptors than estradiol from serum. Additional calculations through K_i (inhibition constant) indicated that this corresponded to an effective free fraction of 26.9% for DES in serum (Nagel *et al.*, 1998). Therefore, at equivalent concentrations of estradiol and DES, the concentration of DES at the target cell will be significantly greater than that of estradiol. Arnold *et al.* (1996) demonstrated that DES induced greater β -galactosidase activity than estradiol in the presence of albumin or SHBG, suggesting that the availability of DES is greater than estradiol in the presence of albumin or SHBG (the IC₅₀ for SHBG was 0.035 mg ml⁻¹ for estradiol and 0.15 mg ml⁻¹ for DES).

As for the placental and embryonic weight on day 13 of gestation, the weight of male embryos on day 13 of gestation was greater than that of females in control and DES-treated groups. These results are compatible with those in CD-1 mice reported by Ishikawa *et al.* (2006). Although there were no significant differences in the placental and fetal weights of the DES-exposed groups compared with those in the controls, the trends in fetal and placental weights were opposite. The ratios of fetal: placental weights were increased in a dose-dependent manner, and

significant increases were detected between the 15 $\mu\text{g kg}^{-1} \text{ day}^{-1}$ group and the controls in both sexes, suggesting that oral DES dosing affects the development of embryos (see Table 2).

Effects of DES on the Expression of ER and ERR mRNA

Little information on the expression profile of ER mRNA including ERR mRNA in developing rodent placentas is available. Immunocytochemistry revealed that ER was localized within nuclei of cells of the mesometrial decidua basalis but not in trophoblastic cells (day 10 of gestation) or in cells of the junctional zone or labyrinth zone in the rat (Ogle and George, 1995). In murine placentas, ER β mRNA expression was demonstrated in decidual cells (Kurita *et al.*, 2001). RT-PCR analysis revealed that DES exposure caused an increase in ER α mRNA in the cells of the mesometrial decidua basalis with spongiotrophoblasts but not in cells of the labyrinth zone (see Fig. 2). As for ER β mRNA, there was no change in the expression pattern in the placenta exposed to DES.

The present findings on the disturbance of placental development by DES and earlier studies (Luo *et al.*, 1997; Tremblay *et al.*, 2001) led us to hypothesize that ERR β has an important role in early placentation. To test this hypothesis, we investigated the expression of ERR β mRNA by RT-PCR in the placentas from dams on day 13 of gestation. The present study showed that exposure to DES at 10 $\mu\text{g kg}^{-1} \text{ day}^{-1}$ did not change the expression profile of ERR β mRNA in the junctional and labyrinth zone (see Fig. 2), although DES at this dose induced embryonic mortality at a high frequency. Thus, it is reasonable to suggest that the high rate of embryonic mortality induced by DES is not associated with the ERR β mRNA expression in mouse placenta.

Morphology of Placenta and Ultrastructural Changes of TGCs

The treatment of mice with DES (12.5 $\mu\text{g kg}^{-1} \text{ day}^{-1}$) during mid-gestation (days 9 through to 12 of gestation) induces placental changes that include thinning of the labyrinth zone and accumulation of TGCs (Scott and Adejokun, 1980). In addition, the treatment of rats with estradiol benzoate from days 12 through to 19 of gestation induces trophoblast degeneration and destruction of the placental labyrinth (Matsuura *et al.*, 2004). In the present study, light microscopy of the placenta exposed to DES from days 4 through to 8 of gestation showed the impaired development of the labyrinth where most gaseous and nutrient

exchange occurs (see Fig. 3). Thus, it is suggested that disturbance of the villous network resulting from the reduced villigenesis might be associated with the decreases in the physiological function of the placenta.

TGCs are the first terminally differentiated cell type to form during embryogenesis in rodents and are of vital importance for embryo implantation and the promotion of maternal adaptations to pregnancy. TGCs examined in the present study line the implantation site and are in direct contact with decidual and immune cells in the uterus (Simmons *et al.*, 2007), and have several functions to facilitate implantation and initial maternal vascular connections, as well as to regulate decidual cell differentiation and maternal physiology (Hu and Cross, 2010). TGC differentiation can be induced by DES (Tremblay *et al.*, 2001) and retinoic acid (Yan *et al.*, 2001). When injected into pregnant mice, both of these compounds cause an overabundance of giant cells at the expense of other trophoblast subtypes such as stem cells and spongiotrophoblasts, suggesting the differentiation-promoting activity of retinoic acid and DES. This differentiation-promoting activity of retinoic acid and DES is also observed in trophoblast stem cells *in vitro* (Tremblay *et al.*, 2001; Yan *et al.*, 2001). In addition, in the present study, the number of parietal TGCs of the placenta exposed to DES was significantly increased compared with that in the controls, whereas E₂ treatment did not increase the number of TGCs. A few transcription factors have been suggested to regulate giant-cell formation, such as two members of the basic helix-loop-helix (bHLH) family, *Mash2* and *Hand1*. The targeted deletion of this gene results in an increase in the number of giant cells at the expense of the spongiotrophoblasts (Guillemot *et al.*, 1994; Tanaka *et al.*, 1997). In a previous study (Nagao and Yoshimura, 2009), effects on mouse developing placentas were evaluated on day 9 of gestation after exposure to DES at 10 µg kg⁻¹ day⁻¹ on days 4 through to 8, and the number of implantation sites per litter in the DES group (14.8 ± 1.9) was similar to that in the control (14.6 ± 2.3). This suggests that the increase in TGCs in the DES group seen in the present study may be related to the facilitation of implantation.

In the ultrastructural observation of TGCs, poor rough-surfaced endoplasmic reticulum and its atrophy were found in the TGCs of the placentas of mice exposed to DES at 10 µg kg⁻¹ day⁻¹, suggesting the association with the disruption of synthesis of protein. E₂ at 50 µg kg⁻¹ day⁻¹ or DES at 1 or 5 µg kg⁻¹ day⁻¹ did not induce changes of ultrastructure of TGCs or a high frequency of embryonic mortality.

The placenta is a unique organ; placentation starts at the time of implantation and its functional role terminates at the time of parturition. Recently, a few studies have examined the role of endocrine-disrupting chemicals (EDCs) in fetal growth restriction and pregnancy loss, but evidence suggests that exposure to some EDCs during pregnancy can contribute to incomplete placentation. In laboratory animal studies, early exposure to estrogens has been shown to induce trophoblast degeneration (both apoptosis and placental labyrinth destruction) in pregnant rats treated with a physiological dose of estradiol benzoate during days 12 through to 19 of gestation (Matsuura *et al.*, 2004). On day 20 of gestation, the exposed embryos had reduced weight compared with the control ones, indicating fetal growth restriction associated with trophoblast degeneration. Poor placentation, miscarriage and increased neonatal mortality were also observed in mice exposed during early gestation to bisphenol A (Tachibana *et al.*, 2007). These *in vivo* studies suggest that early estrogen or xenobiotic exposure could limit

trophoblast invasion of the endometrium, placing the fetus at risk of intrauterine growth restriction or prenatal and neonatal mortality. However, more studies are needed to understand the developmental effects of inappropriate hormones or xenobiotic exposure on first-trimester placentation.

In conclusion, oral exposure to DES during early to middle gestation affects the rough-surfaced endoplasmic reticulum of TGCs and the developing labyrinth, and it is suggested that these developmental changes may be related to decreased protein synthesis or disruption of nutrition and oxygen exchange between mother and embryo, resulting in a high frequency of embryonic mortality.

Acknowledgments


We thank all our colleagues, including E. Toyoda, Y. Maehara, S. Asano, Y. Nakamura, T. Yoshida and M. Ono, who contributed to the present study.

This study was partly supported by Grants-in-Aid for Scientific Research from the Japanese Ministry of Health, Labour and Welfare.

References

- Abdul-Karim RW, Bruce NW. 1972. The regulatory effect of estrogen on fetal growth: II. Uterine and placental blood flow in rabbits. *Reprod. Fertil.* **30**: 477–480.
- Al-Bader MD. 2006. Estrogen receptors alpha and beta in rat placenta: detection by RT-PCR, real time PCR and Western blotting. *Reprod. Biol. Endocrinol.* **4**: 13.
- Arnold SF, Robinson MK, Notides AC, Guillette LJ, McLachlan JA. 1996. A yeast estrogen screen for examining the relative exposure of cells to natural and xenoestrogens. *Environ. Health Perspect.* **104**: 544–548.
- Bartholomeusz RK, Bruce NW, Lynch AM. 1999. Embryo survival, and fetal and placental growth following elevation of maternal estradiol blood concentration in the rats. *Biol. Reprod.* **61**: 46–50.
- Clevenger WR, Cornwall GA, Carter MW, Bradshaw WS. 1991. Diethylstilbestrol-induced perinatal lethality in the rat. I. Relationship to reduced maternal weight gain. *Biol. Reprod.* **44**: 575–582.
- Couse JF, Korach KS. 1999. Estrogen receptor null mice: What have we learned and where will they lead us? *Endocrinol. Rev.* **20**: 358–417.
- Coscrove MD, Benton B, Henderson BE. 1977. Male genitourinary abnormalities and maternal diethylstilbestrol. *J. Urol.* **117**: 220–222.
- Cross JC, Werb Z, Fisher SJ. 1994. Implantation and the placenta: Key pieces of the development puzzle. *Science* **266**: 1508–1518.
- Fischer LJ, Weissinger JL, Rickert DE, Hintze KL. 1976. Studies on the biological disposition of diethylstilbestrol in rats and humans. *J. Toxicol. Environ. Health* **1**: 587–605.
- Georgiades P, Ferguson-Smith AC, Burton GJ. 2002. Comparative developmental anatomy of the murine and human definitive placenta. *Placenta* **23**: 3–19.
- Giguère V, Yang N, Segui P, Evans RM. 1988. Identification of a new class of steroid hormone receptors. *Nature* **331**: 91–94.
- Gordon MN, Osterburg HH, May PC, Finch CE. 1986. Effective oral administration of 17 beta-estradiol to female C57BL/6J mice through the drinking water. *Biol. Reprod.* **35**: 1088–1095.
- Greco TL, Duello TM, Gorski J. 1993. Estrogen receptors, estradiol, and diethylstilbestrol in early development: The mouse as a model for the study of estrogen receptors and estrogen sensitivity in embryonic development of male and female reproductive tracts. *Endocrinol. Rev.* **14**: 59–71.
- Guillemot F, Nagy A, Auerbach A, Rossant J, Joyner AL. 1994. Essential role of *Mash-2* in extraembryonic development. *Nature* **371**: 333–336.
- Haddad V, Ketchel MM. 1969. Termination of pregnancy and occurrence of abnormalities following estrone administration during early pregnancy. *Int. J. Fertil.* **14**: 56–63.
- Hill DE, Slikker W Jr, Helton ED, Lipe GW, Newport GD, Szszak TJ, Bailey JR. 1980. Transplacental pharmacokinetics and metabolism of diethylstilbestrol and 17 beta-estradiol in the pregnant rhesus monkey. *J. Clin. Endocrinol. Metab.* **50**: 811–818.

- Hu D, Cross JC. 2010. Development and function of trophoblast giant cells in the rodent placenta. *Int. J. Dev. Biol.* **54**: 341–354.
- Ishikawa H, Seki R, Yokonishi S, Yamauchi T, Yokoyama K. 2006. Relationship between fetal weight, placental growth and litter size in mice from mid- to late-gestation. *Reprod. Toxicol.* **21**: 267–270.
- Knöfler M, Vasicek R, Schreiber M. 2001. Key regulatory transcription factors involved in placental trophoblast development: a review. *Placenta* **22**(Suppl A): S83–92.
- Kuhn ER, Bollen M, Darras V. 1982. Fetal growth inhibition and decreased thyroid activity after injection of oestradiol benzoate into pregnant rats. *J. Endocrinol.* **93**: 55–63.
- Kuiper GG, Enmark E, Pelto-Huikko M, Nilsson S, Gustafsson JA. 1996. Cloning of a novel estrogen receptor expressed in rat prostate and ovary. *Proc. Natl. Acad. Sci. U.S.A.* **93**: 5925–5930.
- Kuiper GG, Carlsson B, Grandien K, Enmark E, Haggblad J, Nilsson S, Gustafsson JA. 1997. Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors α and β . *Endocrinology* **138**: 863–870.
- Kurita T, Lee KJ, Saunders PTK, Cooke PS, Taylor JA, Lubahn DB, Zhao C, Mäkelä S, Gustafsson JA, Dahiya R, Cunha GR. 2001. Regulation of progesterone receptors and decidualization in uterine stroma of the estrogen receptor- α knockout mouse. *Biol. Reprod.* **64**: 272–283.
- Lobo RA. 1987. Absorption and metabolic effects of different types of estrogens and progestogens. *Obstet. Gynecol. Clin. North Am.* **14**: 143–167.
- Luo J, Sladek R, Bader J-A, Matthyssen A, Rossant J, Giguère V. 1997. Placental abnormalities in mouse embryos lacking the orphan nuclear receptor ERR- β . *Nature* **388**: 778–782.
- Mahendroo MS, Cala KM, Landrum DP, Russell DW. 1997. Fetal death in mice lacking 5 α -reductase type I causes estrogen excess. *Mol. Endocrinol.* **11**: 917–927.
- Malassiné A, Frendo J-L, Evain-Brion D. 2003. A comparison of placental development and endocrine functions between the human and mouse model. *Hum. Reprod. Update* **9**: 531–539.
- Matsuura S, Itakura A, Ohno Y, Nakashima Y, Murata Y, Takeuchi M, Kobayashi M, Mizutani S. 2004. Effects of estradiol administration on fetoplacental growth in rats. *Early Hum. Dev.* **77**: 47–56.
- Maydl R, Metzler M. 1984. Oxidative metabolites of diethylstilbestrol in the fetal Syrian golden hamster. *Teratology* **30**: 351–357.
- McLachlan JA. 1977. Prenatal exposure to diethylstilbestrol in mice: toxicological studies. *J. Toxicol. Environ. Health* **2**: 527–537.
- Metzler M. 1981. The metabolism of diethylstilbestrol. *CRC Crit. Rev. Biochem.* **10**: 171–213.
- Miller RK, Heckmann ME, McKenzie RC. 1982. Diethylstilbestrol: placental transfer, metabolism, covalent binding and fetal distribution in the Wistar rat. *J. Pharmacol. Exp. Ther.* **220**: 358–365.
- Nagao T, Saitoh Y, Yoshimura S. 2000. Possible mechanism of congenital malformations induced by exposure of mouse preimplantation embryos to mitomycin C. *Teratology* **61**: 248–261.
- Nagao T, Yoshimura S. 2009. Early embryonic losses in mice induced by diethylstilbestrol. *Congenit. Anom. (Kyoto)* **49**: 269–273.
- Nagel SC, Vom Saal FS, Welshons WV. 1998. The effective free fraction of estradiol and xenoestrogens in human serum measured by whole cell uptake assays: physiology of delivery modifies estrogenic activity. *Proc. Soc. Exp. Biol. Med.* **217**: 300–309.
- Ogle TF, George P. 1995. Regulation of the estrogen receptor in the deciduas basalis of the pregnant rat. *Biol. Reprod.* **53**: 65–77.
- Oradell NJ. 1961. *Physicians' desk reference to pharmaceutical specialties and biologicals (15th edn)*. Medical Economics: New Jersey.
- Rossant J, Cross JC. 2001. Placental development: lessons from mouse mutant. *Nat. Rev. Genet.* **2**: 538–548.
- Scott JN, Adejokun F. 1980. Placental changes due to administration of diethylstilbestrol (DES). *Virchows Arch. B Cell Pathol.* **34**: 261–267.
- Shah HC, McLachlan JA. 1976. The fate of diethylstilbestrol in the pregnant mouse. *J. Pharmacol. Exp. Ther.* **197**: 687–696.
- Siegel S. 1956. *Nonparametric Statistics for the Behavioral Sciences*. McGraw-Hill: New York.
- Simmons DG, Fortier AL, Cross JC. 2007. Diverse subtypes and developmental origins of trophoblast giant cells in the mouse placenta. *Dev. Biol.* **304**: 567–578.
- Suzuki K, Kobayashi M, Kobayashi K, Shiraishi Y, Goto S, Hoshino T. 1997. Structural and functional change of blood vessel labyrinth in maturing placenta of mice. *Trophobl. Res.* **9**: 155–164.
- Tachibana T, Wakimoto Y, Nakamura N, Phichitraslip T, Wakitani S, Kusakabe K. 2007. Effects of bisphenol A (BPA) on placentation and survival of the neonates in mice. *J. Reprod. Dev.* **53**: 509–514.
- Tanaka M, Gertsenstein M, Rossant J, Nagy A. 1997. Mash2 acts cell autonomously in mouse spongiotrophoblast development. *Dev. Biol.* **190**: 55–65.
- Tremblay GB, Kunath T, Bergeron D, Lapointe L, Champigny C, Bader J, Rossant J, Giguère V. 2001. Diethylstilbestrol regulates trophoblast stem cell differentiation as a ligand of orphan nuclear receptor ERR β . *Genes Dev.* **15**: 833–838.
- Tsai MJ, O'Malley BW. 1994. Molecular mechanisms of action of steroid/thyroid receptor superfamily members. *Annu. Rev. Biochem.* **63**: 451–486.
- Yan J, Tanaka S, Oda M, Makino T, Ohgane J, Shiota K. 2001. Retinoic acid promotes differentiation of trophoblast stem cells to a giant cell fate. *Dev. Biol.* **235**: 422–432.
- Zimmerman SA, Clevenger WR, Brimhall BB, Bradshaw WS. 1991. Diethylstilbestrol-induced perinatal lethality in the rat. II. Perturbation of parturition. *Biol. Reprod.* **44**: 583–589.



Maternal bisphenol A oral dosing relates to the acceleration of neurogenesis in the developing neocortex of mouse fetuses

Munekazu Komada^{a,b,*}, Yasuko Asai^c, Mina Morii^c, Michie Matsuki^c, Makoto Sato^{a,b,d}, Tetsuji Nagao^c

^a Division of Cell Biology and Neuroscience, Department of Morphological and Physiological Sciences, Faculty of Medical Sciences, University of Fukui, Eiheiji-cho, Fukui 910-0337, Japan

^b Research and Education Program for Life Science, University of Fukui, Fukui, Fukui 910-8507, Japan

^c Laboratory of Developmental Biology, Department of Life Science, Kinki University, Higashi-Osaka, Osaka 577-8502, Japan

^d Research Center for Child Mental Development, University of Fukui, Eiheiji-cho, Fukui 910-0337, Japan

ARTICLE INFO

Article history:

Received 13 December 2011

Received in revised form 25 February 2012

Accepted 27 February 2012

Available online 7 March 2012

Keywords:

Endocrine disruptors

Development

Bisphenol A

Neocortex

Neurogenesis

ABSTRACT

Bisphenol A (BPA), an endocrine-disruptor, is widely used in the production of plastics and resins. Human perinatal exposure to this chemical has been proposed to be a potential risk to public health. Animal studies indicate that postnatal exposure to BPA may affect neocortex development in embryos by accelerated neurogenesis and causing neuronal migration defects. The detailed phenotypes and pathogenetic mechanisms, especially with regard to the proliferation and differentiation of neural stem/progenitor cells, however, have not been clarified. C57BL/6J pregnant mice were orally administered BPA at 200 µg/kg from embryonic day (E) 8.5 to 13.5, and the fetuses were observed histologically at E14.5. To clarify the histological changes, especially in terms of neurogenesis, proliferation and cell cycle, we performed histological analysis using specific markers of neurons/neural stem cells and cell cycle-specific labeling experiments using thymidine-analog substances. Cortical plate was hyperplastic and the number of neural stem/progenitor cells was decreased after the exposure to BPA. In particular, the maternal BPA oral dosing related to the effects on intermediate progenitor cells (IPCs, neural progenitor cells) in the subventricular zone (SVZ) of dorsal telencephalon. Exposure to BPA associated the promotion of the cell cycle exit in radial glial cells (RGCs, neural stem cells) and IPCs, and decreased the proliferation resulting from the prolong cell cycle length of IPCs in the SVZ. Our data show that maternal oral exposure to BPA related to the disruption of the cell cycle in IPCs and the effects of neurogenesis in the developing neocortex.

© 2012 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

Endocrine-disrupting industrial chemicals are released into the environment and interfere with normal endocrine function. Bisphenol A (BPA; 2,2-bis(4-hydroxy-phenyl) propane) is known to be one of the endocrine-disrupting chemicals because of its weak estrogenic, androgenic and thyroid hormone-like activity (Hiroi et al., 2006; Krishnan et al., 1993; Takayanagi et al., 2006; Xu et al., 2005). BPA is used in polycarbonate plastics, epoxy resins, and dental resin-based composites (Howe and Borodinsky, 1998; Pulgar et al., 2000; Sasaki et al., 2005). BPA has been detected in the serum of pregnant women (1–2 ng/ml), fetus serum (0.2–9.2 ng/ml), amniotic fluid (8.3–8.7 ng/ml), placental tissue (1.0–104.9 ng/ml), and

breast milk (1.1 ng/ml) (Ikezuki et al., 2002; Schonfelder et al., 2002; Ye et al., 2005), suggesting that the human fetus is exposed to this compound during pre- and post-natal development. Several studies reported some adverse effects of high- and low-dose BPA on various organs during the prenatal period (Golub et al., 2010; Kundakovic and Champagne, 2011). During fetal development, there is a sensitive period during which environmental exposure may cause persistent damage to the developing brain. Maternal exposure to low-dose BPA (20 µg/kg/day, intraperitoneally) has been shown to affect cortical development in embryos by accelerated neurogenesis and neuronal migration during the mid-gestational period in mice (Nakamura et al., 2006). In addition, BPA exposure results in abnormal neuronal positioning and aberrant neuronal network formation between the thalamus and the cortex in the mature brain exposed prenatally (Nakamura et al., 2007). However, no findings have been added to these reports about the morphological changes of the neocortex induced by low-dose BPA exposure.

During neocortical development, neural stem/progenitor cells sequentially pass through phases of expansion, neurogenesis and gliogenesis. Radial glial cells (RGCs) in the ventricular zone (VZ), neural stem cells, expand their population by symmetric

* Corresponding author at: Division of Cell Biology and Neuroscience, Department of Morphological and Physiological Sciences, Faculty of Medical Sciences, University of Fukui, Japan. Tel.: +81 776 61 8305; fax: +81 776 61 8155.

E-mail addresses: komada@u-fukui.ac.jp (M. Komada), yasuko@fbs.osaka-u.ac.jp (Y. Asai), morii@life.kindai.ac.jp (M. Morii), s1121376@u.tsukuba.ac.jp (M. Matsuki), makosato@u-fukui.ac.jp (M. Sato), tnagao@life.kindai.ac.jp (T. Nagao).

division that produces two RGCs and simultaneously self-renew and generate more differentiated cells through asymmetric cell division (Gotz and Huttnner, 2005). The daughter cells of ventricular RGCs are often neuronal progenitor cells and can migrate superficially into the subventricular zone (SVZ) to divide (Haubensack et al., 2004; Noctor et al., 2004). RGC divisions are asymmetric and associated with self-renewal, while the daughter progenitor cells (intermediate progenitor cells: IPCs) usually undergo one terminal symmetric division that produces two neurons and depletes the progenitor cells (Noctor et al., 2004). The majority of layer II/III neuron-producing cell divisions are by IPCs during the late neurogenesis stages (Kowalczyk et al., 2009). The number of symmetric IPC divisions could also affect neuron number and be a determinant of neocortical size. A human genetic study pinpointed the disease-causing gene in a family exhibiting congenital microcephaly to the homozygous silencing of *Tbr2* (Baala et al., 2007), a transcriptional factor shown in rodent studies to be a selective marker for IPCs (Englund et al., 2005). These reports suggest the overall importance of IPCs of SVZ in the control of neocortical size.

The purposes of this study are to clarify the mechanisms underlying the abnormal proliferation and differentiation of neural stem cells in fetuses exposed to BPA. Observation of the BPA-treated fetuses revealed the accelerated neurogenesis and a reduced number of neural stem/progenitor cells, especially IPCs, in the dorsal telencephalon. Our data suggested that these aberrations of developing neocortex result in disruption of the cell cycle in IPCs.

2. Materials and methods

2.1. Animals and housing

Eight-week-old male and female C57BL/6J mice were purchased from Japan SLC Inc. to be used as breeder animals in this study. These mice were quarantined and habituated for 2 weeks. These animals were kept under SPF conditions and a constant light–dark cycle (dark period from 7:00 pm to 7:00 am) at $24 \pm 1^\circ\text{C}$ and $55 \pm 5\%$ relative humidity. To avoid the possibility of stressing the animals, noise levels were kept to a minimum both within the room and in the adjacent areas. Diet food (Certified Rodent Chow CE-2, CLEA Japan) and drinking water were available *ad libitum*. Certification analysis of each lot of diet was performed by the manufacturer. The same lots of diet were provided to animals from control and BPA-treated groups and at the same times, in order to control across groups for possible variation in the content of the diet. Water was available via glass bottles with Teflon seals during the exposure period. Pregnant females were housed individually throughout the study in polypropylene plastic tubs with stainless steel lids and corncob bedding. Mice in all experiments were humanely treated according to the guidelines of the Animal Research Committee of Kinki University. Ten-week-old mice were allowed to copulate overnight at a 2:1 or 1:1 female to male ratio. Females were checked at 12 h intervals for the presence of vaginal plugs, indicating copulation, and were separated from the male if a plug was present. The presence of a plug represented embryonic day (E) 0.

2.2. Test substance and treatment regimen

Bisphenol A (BPA, 2,2-bis(4-hydroxyphenyl)propane 4,4'-isopropylidenediphenol, CAS no. 80-05-7, Sigma Aldrich) was suspended in corn oil and administered by oral gavage from E8.5 to E13.5. The dose solution was prepared once per 5 days and analyzed prior to dosing. The BPA concentration was confirmed to be within $\pm 10\%$ of targeted concentration. Administration occurred at a defined time (12:00 pm). In a preliminary study, three dams per group were exposed orally to BPA at 20 or 200 $\mu\text{g}/\text{kg}$ from E8.5 through E13.5, and killed on E14.5 to collect the fetuses. Nine female fetuses in each group were examined histologically in terms of the developing brain (Sup. Table 1). No histological abnormality, such as hyperplasia of cortical plate and the acceleration of cell cycle exit (Sup. Fig. 1, Tables 1 and 2), were detected in the fetuses exposed to BPA at 20 $\mu\text{g}/\text{kg}$. In the fetuses exposed to BPA at 200 $\mu\text{g}/\text{kg}$, hyperplasia of cortical plate and the promotion of neurogenesis were identified. On the basis of these results, BPA dose used in the present study was chosen as 200 $\mu\text{g}/\text{kg}/\text{day}$, which revealed clear effects on the development of neocortex. Control animals received an equal volume of corn oil. At this dose of BPA (200 $\mu\text{g}/\text{kg}/\text{day}$) we calculate intake to be approximately 3 μg of BPA daily; body weight of mice used was approximately 30–40 g during the administration period. In the previous study, when rat dam was orally administered BPA at 6 $\mu\text{g}/\text{kg}/\text{day}$, 4–5 $\mu\text{g}/\text{L}$ BPA was detected in embryonic serum (Yoshida et al., 2004). In addition, prenatal exposure to BPA at 200 $\mu\text{g}/\text{kg}/\text{day}$ BPA induced changes of embryonic body weight and total number of embryos born

per litter (Cagen et al., 1999). These data indicate that the embryos were exposed to BPA in utero in the experimental conditions of the present study.

2.3. Body weight measurement and tissue preparation

Pregnant mice were humanely killed and underwent cesarean section on E14.5, and the fetuses were observed histologically. Fetuses were sampled out of the ostium of the uterus, because there were shown to be no differences in the postnatal growth of the reproductive and endocrine systems, sexual maturation and estrous cycle, or behavior depending on the embryo position in utero (Nagao et al., 2004). For body weight measurement, the body weight before fixation was measured for E14.5 mouse fetuses. Fourteen males and 17 female fetuses from 6 litters in the BPA-treated group and 9 males and 12 females from 6 litters in the control group were used for head size and body weight measurement (Sup. Table 1). Fetuses were fixed for 3 h in periodate lysin paraformaldehyde (PLP) at 4°C , and washed in phosphate-buffered saline (PBS). For paraffin sections, fetuses were embedded in paraffin and sectioned at 5 μm for histological and immunohistochemical observation. For frozen sections, fetuses were embedded in 30% sucrose/PBS for cryoprotection, and sectioned at 10 μm for immunohistochemical observation. Nine female fetuses from 3 litters in both BPA-treated and control groups were sampled on E14.5 for HE staining and immunostaining (Sup. Table 1). Fetal body weight and brain measurement data were analyzed employing Student's *t*-test.

2.4. Immunohistochemistry

The following antibodies were used: mouse monoclonal anti-neural class III β -tubulin (Tuj1, 1:500; Covance); pan-neuronal marker, rabbit monoclonal anti-Ki67 (1:200; Lab Vision, SP6); a proliferative cell marker, rat monoclonal anti-BrdU (CldU) (1:50; Oxford Biotechnology, BU1/75), mouse monoclonal anti-BrdU (IdU) (1:50; Becton Dickinson, B44), rabbit polyclonal anti-Tbr2 (1:200, Abcam); transcriptional factor and IPC marker, rabbit polyclonal anti-Pax6 (1:200, Covance); transcriptional factor and RGC marker, mouse monoclonal anti-Nestin (1:200, BD Pharmingen); marker of radial fiber in RGC. Secondary antibodies were conjugated with Alexa 568 and 488 (1:200; Invitrogen). The nuclei were counterstained with DAPI in mounting medium (Vector Labs). Immunohistochemistry was performed as described previously (Komada et al., 2008), and standard immunostaining procedures were used in E14.5 fetuses. Nine female fetuses from 3 litters in the BPA-treated group and 9 female fetuses from 3 litters in the control group were sampled on E14.5 for immunostaining (Sup. Table 1).

2.5. CldU and IdU incorporation

For *in vivo* labeling of S-phase cells (thymidine analog incorporation), one injection of CldU (105478, MP Biomedicals Inc.) and IdU (17125, Sigma) was made 24 h and 1 h, respectively, prior to cesarean sectioning on E14.5. Fetuses were allowed to develop to E14.5 and then sacrificed and processed for CldU and IdU immunohistochemistry. The quantification of positive cells and their distribution within the cortical layer were analyzed according to the methods of Komada et al. (2008), with 2 anatomically matched sections from each fetus (9 BPA-treated and 9 control female embryos).

2.6. Cell cycle kinetics (analysis of cell cycle length and cell cycle exit)

For estimation of cell cycle length, we counted the number of IPCs labeled by 1 h pulse of IdU in Tbr2-positive cells. The population of IdU/Tbr2 double-positive cells among all Tbr2-positive cells enables estimation of the cell cycle length of IPCs. A smaller population of IdU-positive cells among Tbr2-positive cells indicates a greater cell cycle length (Chenn and Walsh, 2002).

Cell cycle exit was estimated from the ratio of CldU-positive cells/Ki67, Pax6, or Tbr2-negative (postmitotic and/or differentiation) cells to all CldU-positive cells at E14.5 in the dorsal telencephalon (Ki67) or upper SVZ (Pax6 and Tbr2) (Chenn and Walsh, 2002). One injection of CldU was made 24 h prior (at E13.5) to sampling (at E14.5).

2.7. Quantification of cell number for cell cycle exit, proliferation, RGC, IPC, and cell cycle length

For quantification of data obtained using immunofluorescent staining, the counterpart areas in the dorsal telencephalic region of brain of the control and BPA-treated fetuses were selected, and the total number of cells (DAPI-stained) and cells stained with each antibody were manually counted in two 100 μm -wide sampling box (indicated by open boxes in Figs. 2–6) with 2 anatomically matched sections from each fetus (9 BPA-treated and 9 control female embryos) using Adobe photoshop CS4 (Adobe). We performed the counting and quantitative procedures by the method reported previously (Komada et al., 2008).




Fig. 1. Maternal BPA oral dosing induces hyperplasia of cortical plate and promotes neurogenesis. Parasagittal sections of the telencephalons of the control fetuses (A) and BPA-treated fetuses (B) at E14.5 were stained with HE. These panels illustrated that the size of the cortical plate was reduced in BPA-treated fetuses (black arrow). Parasagittal sections of E14.5 dorsal telencephalon were immunostained with anti-neuron-specific class III β -tubulin (Tuj1) antibody (C, D) and expressing regions were measured on the sections (white arrow). (E) Tuj1-stained regions were significantly increased in the dorsal telencephalon of BPA-treated fetuses ($57.6 \pm 2.11\%$, $n=9$, $*P < 0.05$) compared with those of control fetuses ($47.5 \pm 2.09\%$, $n=9$). Scale bar: 200 μm .

3. Results

3.1. Hyperplasia of cortical plate in BPA-treated fetuses

In order to clarify the effects of BPA exposure on the development of neocortex, pregnant mice were treated continuously with BPA at a dose of 200 $\mu\text{g}/\text{kg}$ from E8.5 (this day is the start point of mouse neocortex organogenesis). Fourteen male and 17 female fetuses from 6 litters in the BPA-treated group and 9 male and 12 female fetuses from 6 litters in the control group were sampled at E14.5 (Sup. Table 1). These fetuses were weighed and we measured the rostral-caudal length and width of the head region. The body weight and head length and width of male and female fetuses treated with BPA were comparable to those of control fetuses at E14.5 (date not shown). In addition, there was no fetal lethal effect of BPA when evaluated at E14.5 (date not shown).

As for the histopathological observation of fetuses treated with BPA, hyperplasia of the CP in the dorsal telencephalon was found in the HE staining of the parasagittal sections (Fig. 1A and B). To determine the difference of neurogenesis in fetuses treated with BPA, the Tuj1 (neuron-specific class III β -tubulin; pan-neuronal marker) expression regions were quantified by measuring the total thickness of the dorsal telencephalon and the thickness of CP (Tuj1-expression region) (Fig. 1C and D). There was a significant increase in the thickness of CP in BPA-treated fetuses ($57.6 \pm 2.11\%$, $P < 0.05$) compared with that of control fetuses ($47.5 \pm 2.09\%$) (Fig. 1E). However, hyperplasia of the CP and the expansion of Tuj1-expression area in the dorsal telencephalon were not found in the HE staining (Sup. Fig. 1A–C) and immunostaining of Tuj1 (Sup. Fig. 1D–F, Sup. Table 2) in the 20 $\mu\text{g}/\text{kg}/\text{day}$ BPA-treated fetuses. Nakamura et al. reported that low-dose BPA might disrupt normal neocortical development by accelerated neuronal differentiation/migration

(Nakamura et al., 2006). Our results coincided with these previous data and indicated that the maternal BPA oral dosing associated with the accelerated neurogenesis and hyperplasia of CP during the development of telencephalon.

3.2. Acceleration of cell cycle exit in the neural stem/progenitor cells of the dorsal telencephalon

An adequately controlled cell cycle is important for proper neurogenesis during corticogenesis in the neocortex. To investigate the cause of neurogenesis promotion, we examined cell cycle exit of neural stem/progenitor cells. We performed double immunostaining using anti-Ki67 (nucleoprotein, expressed in proliferative cells) and anti-CldU (thymidine analog) antibodies after 24 h CldU labeling (Fig. 2A and B) (Chenn and Walsh, 2002). Cell cycle exit was determined as the ratio of cells that exited the cell cycle (red, CldU+/Ki67–, indicating cells no longer dividing) to all cells labeled with CldU (red and yellow) after 24 h labeling. Quantification of the experiment showed significantly increased cell cycle exit in BPA-treated fetuses ($44.1 \pm 3.8\%$, $P < 0.01$) compared with that in the control fetuses ($34.3 \pm 4.7\%$) (Fig. 2E). However, in the 20 $\mu\text{g}/\text{kg}/\text{day}$ BPA-treated fetuses, the significant increase in rate of cell cycle exit was not observed in the dorsal telencephalon (Sup. Fig. 1G–I, Sup. Table 2). These data indicated that the hyperplasia of CP was caused by the accelerated neurogenesis of neural stem/progenitor cells in the dorsal telencephalon. Ki67-expression region was decreased and the distribution of double positive cells (Ki67+ and CldU+, yellow cells) was abnormal, remaining in a deep position of Ki67-positive SVZ/VZ, compared with that in control fetuses (Fig. 2A and B). We hypothesized that these phenotypes resulted from the aberration of interkinetic nuclear migration in the SVZ/VZ, in which their nuclei migrate using radial fibers between the apical surface




Fig. 2. The maternal oral exposure to BPA leads to acceleration of cell cycle exit and decreases the number of neural stem/progenitor cells. To estimate cell cycle exit of neural stem/progenitor cells, sections were stained with anti-Ki67 and anti-CldU antibodies 24 h after CldU pulse labeling at E14.5 in the control (A) and BPA-treated (B) fetuses. Parasagittal sections of E14.5 dorsal telencephalon were immunostained with anti-Nestin antibody. In BPA-treated fetuses, Nestin-stained radial fibers were shorter than those of the controls (C, D). (E) Cell cycle exit was determined as ratio of cells that exited the cell cycle (red, CldU+/Ki67–, no longer dividing) to all cells labeled with CldU (red + yellow) after 24 h labeling. Cells were counted in 100- μ m-wide sampling boxes (white boxes in A and B). In the BPA-treated fetuses, the ratio was significantly increased at E14.5 (control fetuses: $34.3 \pm 4.7\%$, $n = 9$, BPA-treated fetuses: $44.1 \pm 3.8\%$, $n = 9$, $*P < 0.01$). Scale bar: 200 μ m. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of the article.)

and the basal part of the SVZ/VZ in synchrony with the cell cycle. Thus, we performed immunostaining using anti-Nestin (the marker of radial fibers of RGCs) antibody in the parasagittal sections. In the BPA-treated fetuses, the Nestin-positive radial fibers were shorter than control ones at E14.5 (Fig. 2C and D). These data suggested that the short radial fibers resulting from the abnormal interkinetic nuclear migration in the SVZ/VZ and abnormal positioning of Ki67+ cells and double-positive (Ki67+ and CldU+) cells.

3.3. Reduction of proliferative neural stem/progenitor cells in the dorsal telencephalon

Accelerated neurogenesis might impact on the proliferation of neural stem/progenitor cells in the dorsal telencephalon. To examine whether cell proliferation was affected in the BPA-treated fetuses, we performed immunostaining using anti-Ki67 antibodies (Fig. 3A and B). At E14.5, the Ki67-immunopositive cell index (Ki67-positive cells/DAPI-stained cells) was significantly decreased in the dorsal telencephalon of BPA-treated fetuses ($41.6 \pm 1.0\%$, $P < 0.01$) compared with that in the control fetuses ($47.2 \pm 1.0\%$) (Fig. 3C). These data suggested that the maternal BPA oral dosing related to the reduction of neural stem/progenitor cells as a result of the promotion of neurogenesis in the dorsal telencephalon.

3.4. BPA specifically affected the intermediate progenitor cells

During the corticogenesis of neocortex, three different types of neural stem/progenitor cells (RGCs, IPCs, and outer radial glial cells) exist in the SVZ/VZ of the dorsal telencephalon (Molnar et al., 2011). These neural stem/progenitor cells are distinguished by their distribution, self-renewal ability, capacity for neurogenesis and expression of transcriptional factors (Molnar et al., 2011). To determine which cell population (RGCs or IPCs) was affected by BPA exposure, we performed immunostaining using anti-Pax6 (transcriptional factor, the marker of RGCs) and anti-Tbr2 (transcriptional factor, the marker of IPCs) (Fig. 4A–D). Pax6-positive cells represent RGCs in VZ and Tbr2-positive cells are IPCs in

SVZ. There was no significant difference in the number of Pax6-positive cells between the BPA-treated ($54.1 \pm 3.1\%$) and control fetuses ($55.1 \pm 3.8\%$) (Fig. 4E). However, the number of Tbr2-positive cells was significantly decreased in the BPA-treated fetuses




Fig. 3. Proliferation of stem/progenitor cells in the dorsal telencephalon is reduced by maternal BPA exposure. Immunostaining with Ki67 (red) of sagittal sections of E14.5 (A, B) fetal brains treated with BPA. Ki67 was expressed in the proliferative cells in the SVZ/VZ of dorsal telencephalon. Cells were counted in 100- μ m-wide sampling boxes (white box). Examples of anti-Ki67 labeling of the dorsal telencephalon of the control (A) or BPA-treated fetuses (B). C: BPA-treated fetuses showed significant decrease in the ratio of Ki67+ cells at E14.5 (control fetuses: $47.2 \pm 1.0\%$, $n = 9$, BPA-treated fetuses: $41.6 \pm 1.0\%$, $n = 9$, $*P < 0.01$). Scale bar: 200 μ m. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of the article.)




Fig. 4. Maternal BPA oral dosing induces the reduction of intermediate progenitor cells and does not affect radial glial cells. Immunostaining on coronal sections of E14.5 dorsal telencephalon with anti-Pax6 (A, B) and anti-Tbr2 (C, D). The Pax6-positive cells were considered as RGCs in VZ of the control (A) and BPA-treated fetuses (B) at E14.5. Cells were counted in 100- μ m-wide sampling boxes (white box). There was no significant difference in Pax6-positive RGC number between the control and BPA-treated groups (E: control fetuses: $55.1 \pm 3.8\%$, $n=9$, BPA-treated fetuses: $54.1 \pm 3.1\%$, $n=9$). The Tbr2-positive cells were identified as intermediate progenitor cells (IPCs) in SVZ of the control (C) and the BPA-treated fetuses (D). (E) Tbr2-positive IPCs of the BPA-treated fetuses ($5.46 \pm 0.32\%$, $n=9$, $*P < 0.05$) showed significant reduction compared with those in the control fetuses ($6.12 \pm 0.24\%$, $n=9$). Scale bar, 100 μ m.

($5.46 \pm 0.32\%$, $P < 0.05$) compared with that in the control fetuses ($6.12 \pm 0.24\%$) (Fig. 4E). These data suggested that the maternal BPA oral dosing specifically associated with the maintenance of IPCs in the SVZ of dorsal telencephalon.

3.5. The acceleration of RGCs and IPCs differentiation by BPA exposure

During the corticogenesis of neocortex, RGCs differentiate to IPCs or neurons and IPCs differentiate to neurons. IPCs mainly produce the projection neurons in layer II/III of neocortex. In evolution, layer II/III is among the most highly evolved regions and IPCs might have important roles in the progression of neocortex. Thus, to determine which cell (RGCs or IPCs) differentiation was affected by maternal BPA exposure, we performed cell cycle exit analysis using thymidine analog labeling with each cell. RGCs and IPCs are proliferative cells labeled by CldU in the SVZ/VZ. In the CP and upper SVZ after 24 h of CldU injection, CldU positive and Pax6/Tbr2 negative cells are defined as postmitotic (differentiated) cells from RGCs/IPCs, respectively. Differentiated RGCs and IPCs were identified by double-immunostaining using anti-Pax6/anti-CldU and anti-Tbr2/anti-CldU antibodies, respectively (Fig. 5A–D). The quantification of these experiments showed that the cell cycle exits of RGCs ($37.4 \pm 2.4\%$, $P < 0.01$) and IPCs ($35.7 \pm 1.2\%$, $P < 0.01$) in the BPA-exposed fetuses were significantly promoted compared with those in the control fetuses (RGCs: $32.8 \pm 2.8\%$, IPC: $30.7 \pm 3.7\%$) at E14.5 (Fig. 5E). These results indicated that the maternal BPA treatment related to the acceleration of neurogenesis of IPCs in the SVZ and differentiation of RGCs in the VZ of dorsal telencephalon.

3.6. BPA leads to prolongation of cell cycle of IPC

To investigate the cause of reduction of IPCs (Fig. 4), we examined the cell cycle length of IPCs. We hypothesized that prolongation of the cell cycle results in decreased proliferation of

IPCs in the SVZ of dorsal telencephalon. We performed double immunostaining using anti-Tbr2 and anti-IdU antibodies (Fig. 6A and B). Cell cycle length of IPC was calculated as the rate of the total number of IPCs (Tbr2-positive cell) and the number of S-phase IPCs for 1 h (Tbr2/IdU double-positive cell). A smaller population of IdU-labeled cells among Tbr2-positive cells indicates a greater cell cycle length (Chenn and Walsh, 2002). In the BPA-treated fetuses, the cell cycle was significantly prolonged ($8.4 \pm 0.6\%$, $P < 0.01$) compared with that in the control fetuses ($16.5 \pm 4.3\%$) at E14.5 (Fig. 6C). These data indicated that the extension of the cell cycle length of IPCs induced the reduction of self-renewal, resulting in the decreasing number of IPCs.

4. Discussion

In previous studies, exposure to BPA (20 μ g/kg/day) induced the accelerated neuronal differentiation/migration in ICR mice (Nakamura et al., 2006). Our data indicated that treatment with BPA at the dose of 20 μ g/kg/day showed no differences in the morphogenesis of CP (Sup. Fig. 1A–C') and neurogenesis of neural stem/precursor cells (Sup. Fig. 1D–I, Sup. Table 2) at E14.5 between the BPA-treated and control fetuses in C57BL/6J mice (see the materials and methods). We hypothesized that the difference of phenotype resulted from the different administration routes (intraperitoneally or orally) or mouse strains used, with variation in sensitivity to BPA. In a previous study involving the maternal BPA oral dosing, number of fetuses born per litter and embryonic body weight were significantly lower in the group exposed to BPA at 200 μ g/kg/day than those in the control group. However, no difference was detected between the group exposed to BPA at 20 μ g/kg/day and the control group (Cagen et al., 1999). Our data also showed that histological changes (cell cycle, proliferation, and differentiation changes) were related to BPA at 200 μ g/kg/day. However, these were not found after oral exposure to BPA at 20 μ g/kg/day (Sup. Fig. 1, Sup. Table 2).




Fig. 5. Differentiation of radial glial cells and intermediate progenitor cells is promoted by maternal BPA exposure. To estimate the differentiation of RGCs and IPCs, sagittal sections were stained with anti-Pax6 or -Tbr2 and anti-CldU antibodies 24 h after CldU pulse labeling at E14.5 in the control (A, C) and the BPA-treated (B, D) fetuses. Differentiation was determined as the ratio of cells that differentiated from RGCs (green, CldU+/Pax6-, A, B) and IPCs (green, CldU+/Tbr2-, C, D) to all cells (blue: DAPI) after 24 h labeling. Cells were counted in 100- μ m-wide sampling boxes (white box) in the SVZ (RGC, A, B) or upper SVZ (IPC, C, D). (E) In the BPA-treated fetuses, the ratio of differentiated cells from RGCs was significantly increased at E14.5 (control fetuses: $32.8 \pm 2.8\%$, $n=9$, BPA-treated fetuses: $37.4 \pm 2.4\%$, $n=9$, $*P < 0.01$) and the ratio of differentiated cells from IPCs was also significantly increased at E14.5 (control fetuses: $30.3 \pm 3.7\%$, $n=9$, BPA-treated fetuses: $35.7 \pm 1.2\%$, $n=9$, $*P < 0.01$). Scale bar: 100 μ m. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of the article.)




Fig. 6. The maternal exposure to BPA leads to prolonged cell cycle of intermediate progenitor cells. After 1 h pulse labeling of IdU, immunostaining of coronal sections was performed with anti-Tbr2 (red) and anti-IdU (green) antibodies at E14.5 (A, B). Cell cycle length was estimated as percentage of Tbr2 and IdU double-positive cells among all Tbr2-positive cells. Smaller percentage represents longer cell cycle. Cells were counted in 100- μ m-wide sampling boxes (white box). (C) The BPA-treated fetuses showed significantly prolonged cycle length of IPC ($8.4 \pm 0.6\%$, $n=9$, $*P < 0.01$) compared with control fetuses ($16.5 \pm 4.3\%$, $n=9$) at E14.5. Scale bar: 100 μ m. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of the article.)

BPA, an endocrine disruptor, induces the promotion of neurogenesis and regulation of proliferation in neural stem/precursor cells. BPA is known as an estrogenic chemical that interacts with human estrogen receptor (ER) (Krishnan et al., 1993), acts as an antagonist for human androgen receptor (Xu et al., 2005) and strongly binds to human estrogen-related receptor gamma (Takayanagi et al., 2006). ERs and AR are expressed in many areas of the developing brain in rodents (Matsuda et al., 2008; Zsarnovszky and Belcher, 2001). In the development of CNS, estrogens have neurotrophic and differentiation-promoting effects. ER α and ER β are crucial for the functions during a critical period of brain development (Beyer, 1999). In analysis using ER β knockout mice, ER β was shown to play important roles in the neuronal migration, differentiation and survival in the developing brain (Wang et al., 2003). In addition, BPA increases ER α expression in the brain (Bromer et al., 2010). BPA may also enhance estrogenic activity through the up-regulation of ERs and up- or down-regulation of estrogenic target gene expression in the target tissues. Presently, we are examining whether the estrogen and/or ER are related to these phenotypes and attempting to identify the impaired expression of estrogenic target genes after BPA exposure using our BPA exposure model fetuses.

In addition, BPA binds to protein disulfide isomerase (PDI), the same binding site as thyroid hormone 3,3',5-triiodo-L-thyronine (T3) (Hiroi et al., 2006). Thyroid hormone is essential for proper brain development. In the neocortex, thyroid hormone deficiency during development results in abnormal cytoarchitecture and neuronal network formation (Cuevas et al., 2005). Thyroid hormone receptor alpha and beta were expressed in the cerebral cortex and hippocampus of rat brain (Di Liegro, 2008). The results of the present study suggest that BPA has possibilities of interaction with each factor (receptors and/or enzymes) during the development of neocortex and can induce excessive signaling of these factors.

Our data suggested that BPA exposure inhibited the proliferation and promoted the differentiation of RGC (Pax6-positive cells) and IPC (Tbr2-positive cells) in the SVZ/VZ of dorsal telencephalon. ER β is detected in the deep layer of the dorsal telencephalon at E16.5 (Fan et al., 2006), indicating that RGC and/or IPC in the SVZ/VZ expressed the ER β . In addition, the estrogen signaling might maintain the stem cell niche of RGC and IPC and regulate neurogenesis (Wang et al., 2003). We hypothesized that BPA acts as an agonist of ER and induces excessive estrogen signaling in the dorsal telencephalon, resulting in the promotion of neurogenesis.

A recent study based on cellular behavior, morphology and gene expression pattern identified three similar progenitors during the development of neocortex: outer radial glial cells, RGCs and IPCs. RGCs span the width from ventricular to pial surface, performing self-renewal from symmetric divisions and producing neurons from asymmetric divisions (Molnar et al., 2011). IPCs are derived from asymmetric divisions of RGCs. IPCs express the transcriptional factor Tbr2 and divide within the SVZ (Noctor et al., 2004). IPCs mainly produce projection neurons in layer II/III of the neocortex during the later neurogenesis and are very important for the evolutionary process of the human brain, upgrading and becoming enormous in the neocortex. In the BPA-treated model, our data indicated that the abnormality of IPC resulted in neurogenesis defect and aberrant corticogenesis. Accordingly, it is reasonable to suggest that the detection of IPC defects is essential for the risk assessment of chemical exposure in the development of human brain.

BPA exposure induced the alteration of gene expression associated with neurogenesis. Basic helix-loop-helix (bHLH) genes have two types, repressor type and activator type. Repressor types, such as Hes1 and Hes5, maintain the neural stem cells and promote gliogenesis; activator types, Mash1, Mash2 and Ngn2, accelerate neurogenesis. Nakamura et al. reported that Mash1 and Ngn2 were significantly up-regulated in BPA-treated embryos at E14.5 (Nakamura et al., 2006). In addition, we also indicated that Mash1 was up-regulated at E14.5 in BPA-treated embryos (data not shown). These data suggested that BPA affected the expression of activator-type bHLH gene, Mash1 or Ngn2, causing accelerated neurogenesis.

5. Conclusion

In our studies, the maternal BPA oral dosing related to hyperplasia of CP during the development of telencephalon (Fig. 1) and shortened the radial fibers of RGCs in the SVZ/VZ (Fig. 2). The phenotypes were induced by the accelerated neurogenesis of neural stem/progenitor cells in the dorsal telencephalon (Fig. 2). In addition, BPA associated with the reduction of neural stem/progenitor cells in the SVZ/VZ as a result of the promotion of neurogenesis in the dorsal telencephalon (Fig. 3). In particular, BPA related to the maintenance and neurogenesis of IPCs in the SVZ of dorsal telencephalon (Figs. 4 and 5). These phenotypes (accelerated neurogenesis and the reduction of IPC number) were induced by the extension of the cell cycle length of IPCs in the SVZ (Fig. 6). In the near future, we will clarify whether these morphological defects in the embryonic brain persists, and result in the functional aberrations after birth. The maternal BPA dosing associated with the disruption of cell cycle in IPCs and related to the effects on neurogenesis in the development of neocortex of fetuses.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

Acknowledgements

We are grateful to Dr. H. Yagi (Hyogo College of Medicine), Dr. M. Xie, Dr. T. Iguchi, Dr. K. Kuroda and Dr. Y. Oka (University of Fukui) for helpful discussions, Ms. S. Kanae (Fukui University) for technical assistance, and Ms. T. Taniguchi (University of Fukui), Ms. N. Hirata and Ms. Y. Yoneda (Kinki University) for secretarial assistance. This work was supported by The Ministry of Education, Culture, Sports, Science and Technology of Japan (Grant-in-Aid for Young Scientists, Grant number 22790186) and in part by Grant-in-Aid from the Ministry of Health, Labour and Welfare of Japan, Multidisciplinary program for elucidating the brain development from molecules to social behavior (Fukui Brain Project), The competitive allocation Fund of University of Fukui.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.tox.2012.02.013.

References

- Baala, L., Briault, S., Etchevers, H.C., Laumonnier, F., Natiq, A., Amiel, J., Bodaert, N., Picard, C., Sbiti, A., Asermouh, A., Attie-Bitach, T., Encha-Razavi, F., Munnich, A., Sefiani, A., Lyonnet, S., 2007. Homozygous silencing of T-box transcription factor EOMES leads to microcephaly with polymicrogyria and corpus callosum agenesis. *Nat. Genet.* 39, 454–456.
- Beyer, C., 1999. Estrogen and the developing mammalian brain. *Anat. Embryol. (Berl.)* 199, 379–390.
- Bromer, J.G., Zhou, Y., Taylor, M.B., Doherty, L., Taylor, H.S., 2010. Bisphenol-A exposure in utero leads to epigenetic alterations in the developmental programming of uterine estrogen response. *FASEB J.* 24, 2273–2280.
- Cagen, S.Z., Waechter Jr., J.M., Dimond, S.S., Breslin, W.J., Butala, J.H., Jekat, F.W., Joiner, R.L., Shiotsuka, R.N., Veenstra, G.E., Harris, L.R., 1999. Normal reproductive organ development in CF-1 mice following prenatal exposure to bisphenol A. *Toxicol. Sci.* 50, 36–44.
- Chenn, A., Walsh, C.A., 2002. Regulation of cerebral cortical size by control of cell cycle exit in neural precursors. *Science* 297, 365–369.
- Cuevas, E., Auso, E., Telefont, M., Morreale de Escobar, G., Sotelo, C., Berbel, P., 2005. Transient maternal hypothyroxinemia at onset of corticogenesis alters tangential migration of medial ganglionic eminence-derived neurons. *Eur. J. Neurosci.* 22, 541–551.
- Di Liegro, I., 2008. Thyroid hormones and the central nervous system of mammals. *Mol. Med. Report* 1, 279–295 (Review).
- Englund, C., Fink, A., Lau, C., Pham, D., Daza, R.A., Bulfone, A., Kowalczyk, T., Hevner, R.F., 2005. Pax6, Tbr2, and Tbr1 are expressed sequentially by radial glia, intermediate progenitor cells, and postmitotic neurons in developing neocortex. *J. Neurosci.* 25, 247–251.
- Fan, X., Warner, M., Gustafsson, J.A., 2006. Estrogen receptor beta expression in the embryonic brain regulates development of calretinin-immunoreactive GABAergic interneurons. *Proc. Natl. Acad. Sci. U. S. A.* 103, 19338–19343.
- Golub, M.S., Wu, K.L., Kaufman, F.L., Li, L.H., Moran-Messen, F., Zeise, L., Alexeeff, G.V., Donald, J.M., 2010. Bisphenol A: developmental toxicity from early prenatal exposure. *Birth Defects Res. B Dev. Reprod. Toxicol.* 89, 441–466.
- Gotz, M., Huttner, W.B., 2005. The cell biology of neurogenesis. *Nat. Rev. Mol. Cell Biol.* 6, 777–788.
- Haubensak, W., Attardo, A., Denk, W., Huttner, W.B., 2004. Neurons arise in the basal neuroepithelium of the early mammalian telencephalon: a major site of neurogenesis. *Proc. Natl. Acad. Sci. U. S. A.* 101, 3196–3201.
- Hiroi, T., Okada, K., Imaoka, S., Osada, M., Funae, Y., 2006. Bisphenol A binds to protein disulfide isomerase and inhibits its enzymatic and hormone-binding activities. *Endocrinology* 147, 2773–2780.
- Howe, S.R., Borodinsky, L., 1998. Potential exposure to bisphenol A from food-contact use of polycarbonate resins. *Food Addit. Contam.* 15, 370–375.
- Ikezuki, Y., Tsutsumi, O., Takai, Y., Kamei, Y., Taketani, Y., 2002. Determination of bisphenol A concentrations in human biological fluids reveals significant early prenatal exposure. *Hum. Reprod.* 17, 2839–2841.
- Komada, M., Saitou, H., Kinboshi, M., Miura, T., Shiota, K., Ishibashi, M., 2008. Hedgehog signaling is involved in development of the neocortex. *Development* 135, 2717–2727.
- Kowalczyk, T., Pontious, A., Englund, C., Daza, R.A., Bedogni, F., Hodge, R., Attardo, A., Bell, C., Huttner, W.B., Hevner, R.F., 2009. Intermediate neuronal progenitors (basal progenitors) produce pyramidal-projection neurons for all layers of cerebral cortex. *Cereb. Cortex* 19, 2439–2450.
- Krishnan, A.V., Stathis, P., Permeth, S.F., Tokes, L., Feldman, D., 1993. Bisphenol-A: an estrogenic substance is released from polycarbonate flasks during autoclaving. *Endocrinology* 132, 2279–2286.
- Kundakovic, M., Champagne, F.A., 2011. Epigenetic perspective on the developmental effects of bisphenol A. *Brain Behav. Immun.* 25, 1084–1093.