

effect on activated T cells, arginase inhibitors (nor-NOHA) and anti-IL-10 antibodies were added to the patient's serum. Figure 5C shows that the arginase inhibitor almost completely blocked the inhibitory effect of acute hepatitis serum, whereas the effect of anti-IL-10 was only partial. Furthermore, the inhibitory effect of acute hepatitis sera was reversed by the addition of L-arginine in cell culture (Figure 5D).

Discussion

In this work, we analyzed how the HBV-specific CD8⁺ T cell response is modulated in AHB. We showed that arginase could inhibit the T-cell function during the acute phase of liver inflammation, preferentially acting on activated CD8⁺ T cells. This mechanism is likely to play a crucial regulatory role in limiting excessive liver pathology of different etiologies caused by host adaptive immunity.

By measuring the quantity of activated/proliferating T cells and their function, we observed that HBV infection triggers a robust expansion of activated and proliferating T cells,¹⁴ most of which are, however, impaired in their ability to produce cytokines and degranulate. The observation, by itself, is not novel. A "stunned functional phenotype" was previously described in acute HCV infection²⁸ and was also partially described in a previous analysis of acute HBV infection.^{29,30} In addition, the inability to produce cytokines was observed in murine HBV-specific CD8⁺ T cells adoptively transferred and activated in HBV-transgenic mice.³¹ However, this altered functional state of virus-specific CD8⁺ T cells was interpreted as an antigen-specific tolerogenic phenomenon, caused by excessive antigenic stimulation,²⁹ lack of cytokines,²⁴ and overexpression of inhibitory molecules such as PD-1 typically up-regulated in exhausted T cells.²¹

Here, we argued that a further critical mechanism able to suppress CD8⁺ T cell function during acute hepatitis is non-antigen specific and mediated by arginase, a class of enzymes that metabolize L-arginine and have been shown to suppress T-cell function in different pathological conditions.¹⁶

The importance of such a mechanism *in vivo* during hepatitis was clearly indicated by the important decrease in the ability of CD8⁺ T cells specific for heterologous

viruses to produce antiviral cytokines at the peak of liver damage compared with the recovery phase. The relative major impact of the non-antigen-specific mechanisms of CD8⁺ T cell regulation in comparison to antigen-specific mechanisms was also further supported by the fact that hyperactivation of negative costimulatory or apoptotic pathways was not primarily involved because PD-1 and CTLA-4 blockade as well as the use of antiapoptotic compounds had no effect on *ex vivo* HBV-specific CD8⁺ T cell responses. Moreover, *ex vivo* T-cell responsiveness could not be restored by the simple addition of T-cell stimulatory cytokines (IL-2, IL-7, IL-15) and by increasing doses of antigen.

In contrast, we could observe that either HBV-specific or heterologous virus-specific CD8⁺ T cells could recover their functionality by cell culture in complete medium. More importantly, the ability of acute viral hepatitis serum to inhibit CD8⁺ T cell function revealed that the suppressive effect detected at the peak of acute hepatitis was mediated by soluble factors. Because this effect was not only observed by incubating CD8⁺ T cells with the serum of patients with AHB but also with sera of patients with virus-unrelated hepatitis, such an effect could not be mediated by HBV antigens, which have been reported to inhibit innate immune responses,³² but by soluble factors associated with liver damage. We believed that IL-10 and arginase could be the 2 potential candidates. IL-10 is increasing during acute and chronic hepatic flares^{10,33} and has been implicated in delaying the expansion of virus-specific CD8⁺ T cells in AHB.¹⁰ However, because it acts directly on antigen-presenting cells, IL-10 preferentially exerts its effect on the induction and not on the effector phase of T-cell responses.²⁷ Arginase instead suppresses the function of terminally differentiated effector CD8⁺ T cells.¹⁶ In line with this prediction, data presented here showed that levels of both arginase and IL-10 were elevated during the peak of acute hepatitis and correlated temporally with the inhibition of HBV-specific and heterologous virus-specific CD8⁺ T cell functions. However, only arginase, and not IL-10, at the concentrations detected in the circulation of patients can inhibit the virus-specific CD8⁺ T cell function, and only arginase inhibitor or addition of L-arginine, but not anti-IL-10, prevents the

Figure 5. Arginase, but not IL-10, inhibits CD8⁺ T cell function. (A) HBV core18-27 specific CD8⁺ T cell clone was incubated with arginase or IL-10 at indicated concentration for 2 days, and killing assay and intracellular cytokine staining for IFN- γ was performed with GFP-expressing HepG2 cells as targets. The plots show percent inhibition of cytotoxicity or IFN- γ production (measured as mean fluorescent intensity) compared with that of untreated T cells. (B) Healthy PBMCs were incubated either with IL-10 (10 pg/mL) or arginase (250 pg/mL) with or without IL-15 for 48 hours and specific IFN- γ production was evaluated. The cells were gated on an HCMV pentamer-positive population (B, left plot). The level of activation of HCMV-specific T cells is shown based on HLA-DR/CD38⁺, the histogram plots show the levels of IFN- γ production by HCMV-specific T cells, and the bar charts indicate the percent inhibition of IFN- γ production. Experiments were performed in 2 individuals 3 times. (C) Arginase inhibitor reverted the suppressive effect of the sera from patients with AHB. HBV pentamer-positive CD8⁺ T cells present in patients with AHB were stimulated with HBV peptide after 2 days of culture with IL-2, IL-7, and IL-15 with or without patients' serum (ALT 3100 U/L, arginase 210 pg/mL, IL-10 7 pg/mL), arginase inhibitor (nor-NOHA), or anti-IL-10 antibody. The percent inhibition of IFN- γ production by HBV-specific CD8⁺ T cells compared with culture without serum is shown. (D) Addition of L-arginine reverted the suppressive effect of the sera from patients with AHB. A total of 0.3 μ mol/L arginine was added to PBMCs cultured in the presence of sera from patients with AHB (ALT 3100 U/L, arginase 210 pg/mL). Here, the arginine level was adjusted according to the arginine concentration in culture medium (AIM-V), used to restore T-cell responses. HCMV-specific IFN- γ response was evaluated after 5 hours of activation. Percent inhibition is shown. Experiments were performed in 2 individuals 3 times.

T-cell inhibitory effect of the sera of patients with acute hepatitis.

The inhibition of T-cell function caused by arginase through depletion of L-arginine was suggested to act through TCR ζ chain down-regulation.¹⁶ We analyzed the expression of the TCR ζ chain on total and activated T cells present at the peak of acute hepatitis, but we were unable to detect any reproducible evidence of TCR ζ chain down-regulation (data not shown). This could be explained by the fact that inhibition of T-cell function by arginase can also occur in the absence of TCR ζ chain down-regulation, for example, in T-cell infiltrating human prostate cancers.³⁴ Alternatively, our measurement of TCR ζ chain expression was likely performed also on cells with minimal functional inhibition. Indeed, because of sample availability, TCR ζ chain expression was measured not only on the highly unresponsive HBV-specific CD8⁺ T cells, but on total activated CD8⁺ T cells that also comprise activated heterologous-specific CD8⁺ T cells that, even though inhibited, still conserve a degree of antiviral functionality.

Thus, despite acting in a non-antigen-specific way, arginase maintains a level of selectivity because it acts on activated CD8⁺ T cells with highly active metabolism and therefore preferentially inhibits HBV-specific CD8⁺ T cells that are stimulated by a high quantity of viral antigen and are in a higher activation state than heterologous virus-specific CD8⁺ T cells. This relative selectivity is the likely explanation of the absence of clinical signs of generalized immunosuppression in patients with acute hepatitis. Note that arginase reduces the overall quantity of cytokine produced by the activated T cells but does not seem to interfere with the initial CD8⁺ T cell triggering. This differential effect can explain why HBV-specific CD8⁺ T cells are activated at the peak of acute hepatitis despite the presence of arginase. This interpretation is also consistent with our previous observation that activated heterologous virus-specific CD8⁺ T cells during acute viral hepatitis have a low activation threshold.¹⁴

Overall, our data contribute to describe the existence of distinct mechanisms of T-cell functional regulation, which have the potential to modulate different phases of the antiviral T-cell immunity in the liver.

A representation of such distinct mechanisms is displayed in Figure 6. We propose that early after HBV infection, T-cell expansion can be modulated by IL-10 and by the balance between different costimulatory signals, such as those mediated by PD-1.⁹ However, despite the effect of these negative regulatory mechanisms, HBV-specific T cells expand vigorously and are likely the principal effectors of liver cell lysis. Release of arginase ensuing from hepatocyte damage¹³ or from neutrophils¹⁵ can in turn suppress efficiently effector T-cell function by virtue of a preferential effect on activated T cells (Figure 6).

Thus, arginase can act as a simple direct suppressive mechanism helping to preserve liver viability in the course of the exuberant T-cell activation triggered by acute HBV infection. Interestingly, such an effect is not only re-




Figure 6. Regulation of immune T-cell response during AHB infection. Schematic representation of different mechanisms of T-cell functional modulation potentially active during AHB.

stricted to acute hepatitis, because we have recently observed inhibition of heterologous (HCMV and influenza) virus-specific CD8⁺ T cell IFN- γ production in hepatic flares of chronic HBV infection (unpublished observation, August 2010). In conclusion, arginase-mediated inhibition is a robust antigen-independent mechanism of T-cell functional suppression that is active during hepatitis and may play an important role in preserving the viability of infected parenchymal organs when they are targeted by large quantities of activated virus-specific CD8⁺ T cells.

Supplementary Material

Note: To access the supplementary material accompanying this article, visit the online version of *Gastroenterology* at www.gastrojournal.org, and at <http://dx.doi.org/10.1053/j.gastro.2012.03.041>.

References

1. Ganem D, Prince AM. Hepatitis B virus infection—natural history and clinical consequences. *N Engl J Med* 2004;350:1118–1129.
2. Jilbert A, Wu T, England J, et al. Rapid resolution of duck hepatitis B virus infections occurs after massive hepatocellular involvement. *J Virol* 1992;66:1377–1388.
3. Kajino K, Jilbert AR, Saputelli J, et al. Woodchuck hepatitis virus infections: very rapid recovery after prolonged viremia and infection of virtually every hepatocyte. *J Virol* 1994;68:5792–5803.
4. Guidotti L, Chisari F. To kill or to cure: options in host defense against viral infection. *Curr Opin Immunol* 1996;8:478–483.
5. Guidotti LG, Ishikawa T, Hobbs MV, et al. Intracellular inactivation of the hepatitis B virus by cytotoxic T lymphocytes. *Immunity* 1996;4:25–36.
6. Biermer M, Puro R, Schneider RJ. Tumor necrosis factor alpha inhibition of hepatitis B virus replication involves disruption of capsid integrity through activation of NF-kappaB. *J Virol* 2003;77:4033–4042.
7. Kafrouni MI, Brown GR, Thiele DL. Virally infected hepatocytes are resistant to perforin-dependent CTL effector mechanisms. *J Immunol* 2001;167:1566–1574.
8. Barrie MB, Stout HW, Abougergi MS, et al. Antiviral cytokines induce hepatic expression of the granzyme B inhibitors, proteinase inhibitor 9 and serine proteinase inhibitor 6. *J Immunol* 2004;172:6453–6459.

9. Zhang Z, Jin B, Zhang JY, et al. Dynamic decrease in PD-1 expression correlates with HBV-specific memory CD8T-cell development in acute self-limited hepatitis B patients. *J Hepatol* 2009;50:1163–1173.
10. Dunn C, Peppas D, Khanna P, et al. Temporal analysis of early immune responses in patients with acute hepatitis B virus infection. *Gastroenterology* 2009;137:1289–1300.
11. Wands KR, Isselbacher KJ. Lymphocyte cytotoxicity to autologous liver cells in chronic active hepatitis. *Proc Natl Acad Sci U S A* 1975;72:1301–1303.
12. Chisari FV, Routenberg JA, Fiala M, et al. Extrinsic modulation of Human T-lymphocyte E rosette function associated with prolonged hepatocellular injury after viral hepatitis. *J Clin Invest* 1977;59:134–142.
13. Chisari FV, Nakamura M, Milich DR, et al. Production of two distinct and independent hepatic immunoregulatory molecules by the perfused rat liver. *Hepatology* 1985;5:735–743.
14. Sandalova E, Laccabue D, Boni C, et al. Contribution of herpesvirus specific CD8T cells to anti-viral T cell response in humans. *PLoS Pathog* 2010;6.
15. Munder M, Schneider H, Luckner C, et al. Suppression of T-cell functions by human granulocyte arginase. *Blood* 2006;108:1627–1634.
16. Bronte V, Zanovello P. Regulation of immune responses by L-arginine metabolism. *Nat Rev Immunol* 2005;5:641–654.
17. Tan AT, Loggi E, Boni C, et al. Host ethnicity and virus genotype shape the hepatitis B virus-specific T-cell repertoire. *J Virol* 2008;82:10986–10997.
18. Gehring AJ, Xue SA, Ho ZZ, et al. Engineering virus-specific T cell immunity to target chronic hepatitis B virus infection and HBV-related hepatocellular carcinoma cell lines. *J Hepatol* 2011;55:103–110.
19. Chisari F. Cytotoxic T cells and viral hepatitis. *J Clin Invest* 1997;99:1472–1477.
20. Maier H, Isogawa M, Freeman GJ, et al. PD-1:PD-L1 interactions contribute to the functional suppression of virus-specific CD8+ T lymphocytes in the liver. *J Immunol* 2007;178:2714–2720.
21. Fisicaro P, Valdatta C, Massari M, et al. Antiviral intrahepatic T-cell responses can be restored by blocking programmed death-1 pathway in chronic hepatitis B. *Gastroenterology* 2010;138:682–693.
22. Schurich A, Khanna P, Lopes AR, et al. Role of the coinhibitory receptor cytotoxic T lymphocyte antigen-4 on apoptosis-prone CD8T cells in persistent hepatitis B virus infection. *Hepatology* 2011;53:1494–1503.
23. Lopes AR, Kellam P, Das A, et al. Bim-mediated deletion of antigen-specific CD8T cells in patients unable to control HBV infection. *J Clin Invest* 2008;118:1835–1845.
24. Radziewicz H, Ibegbu CC, Fernandez ML, et al. Liver-infiltrating lymphocytes in chronic human hepatitis C virus infection display an exhausted phenotype with high levels of PD-1 and low levels of CD127 expression. *J Virol* 2007;81:2545–2553.
25. Das A, Hoare M, Davies N, et al. Functional skewing of the global CD8T cell population in chronic hepatitis B virus infection. *J Exp Med* 2008;205:2111–2124.
26. Sitia G, Isogawa M, Kakimi K, et al. Depletion of neutrophils blocks the recruitment of antigen-nonspecific cells into the liver without affecting the antiviral activity of hepatitis B virus-specific cytotoxic T lymphocytes. *Proc Natl Acad Sci U S A* 2002;99:13717–13722.
27. Groux H, Bigler M, de Vries JE, et al. Inhibitory and stimulatory effects of IL-10 on human CD8+ T cells. *J Immunol* 1998;160:3188–3193.
28. Lechner F, Wong D, Dunbar P, et al. Analysis of successful immune responses in person infected with hepatitis C virus. *J Exp Med* 2000;191:1499–1512.
29. Maini MK, Boni C, Ogg GS, et al. Direct ex vivo analysis of hepatitis B virus-specific CD8+ T cells associated with the control of infection. *Gastroenterology* 1999;117:1386–1396.
30. Boettler T, Panther E, Bengsch B, et al. Expression of the interleukin-7 receptor alpha chain (CD127) on virus-specific CD8+ T cells identifies functionally and phenotypically defined memory T cells during acute resolving hepatitis B virus infection. *J Virol* 2006;80:3532–3540.
31. Isogawa M, Furuichi Y, Chisari FV. Oscillating CD8(+) T cell effector functions after antigen recognition in the liver. *Immunity* 2005;23:53–63.
32. Wu J, Meng Z, Jiang M, et al. Hepatitis B virus suppresses toll-like receptor-mediated innate immune responses in murine parenchymal and nonparenchymal liver cells. *Hepatology* 2009;49:1132–1140.
33. Tan AT, Koh S, Goh W, et al. A longitudinal analysis of innate and adaptive immune profile during hepatic flares in chronic hepatitis B. *J Hepatol* 2010;52:330–339.
34. Bronte V, Kasic T, Gri G, et al. Boosting antitumor responses of T lymphocytes infiltrating human prostate cancers. *J Exp Med* 2005;201:1257–1268.

Received October 11, 2011. Accepted March 26, 2012.

Reprint requests


Address requests for reprints to: Antonio Bertoletti, MD, Singapore Institute for Clinical Sciences, 30 Medical Drive, Singapore 117609. e-mail: antonio@sics.a-star.edu.sg; fax: (65) 67766837.

Acknowledgments


The authors thank Shruthi Kumar for help with text/figure editing.

Conflicts of interest


The authors disclose no conflicts.



Supplementary Figure 1. (A) Dysfunction of HBV-specific CD8⁺ T cells in AHB infection. PBMCs of patients with AHB were stimulated with pools of peptides covering the whole HBV proteome, or with staphylococcal enterotoxin B, and the cells were incubated for 12 hours for enzyme-linked immunospot assay. Results of an enzyme-linked immunospot assay for 3 representative acute patients are displayed. (B) The frequencies of IFN- γ pentamer-positive CD8⁺ T cells are plotted together with total HBV core 18-27 pentamer-positive cells in 2 representative patients over time. Corresponding ALT values are shown below each graph.



Supplementary Figure 2. Healthy PBMCs were incubated with either IL-10 (10 pg/mL) or arginase (250 ng/mL) with or without IL-15 for 48 hours, and specific CD107a up-regulation was evaluated. The cells were gated on an EBV or influenza pentamer-positive population. The bar charts indicate the percent inhibition of degranulation (CD107a up-regulation). Experiments were performed in 2 individuals 3 times.



Supplementary Figure 3. Arginase, but not IL-10, inhibits EBV- and influenza-specific responses. Healthy PBMCs were incubated with either IL-10 (10 pg/mL) or arginase (250 ng/mL) with or without IL-15 for 48 hours, and specific IFN- γ production was evaluated. The cells were gated on an EBV or influenza pentamer-positive population. The bar charts indicate the percent inhibition of IFN- γ production. Experiments were performed in 2 individuals 3 times.



OPEN ACCESS

ORIGINAL ARTICLE

Hepatitis C virus kinetics by administration of pegylated interferon- α in human and chimeric mice carrying human hepatocytes with variants of the *IL28B* gene

Tsunamasa Watanabe,¹ Fuminaka Sugauchi,² Yasuhito Tanaka,¹ Kentaro Matsuura,³ Hiroshi Yatsuhashi,⁴ Shuko Murakami,¹ Sayuki Iijima,¹ Etsuko Iio,³ Masaya Sugiyama,⁵ Takashi Shimada,⁶ Masakazu Kakuni,⁶ Michinori Kohara,⁷ Masashi Mizokami⁵

► Additional supplementary files are published online only. To view these files please visit the journal online (<http://dx.doi.org/10.1136/gutjnl-2012-302553>).

¹Department of Virology and Liver Unit, Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan

²Department of Gastroenterology, Nagoya City Kosein Medical Welfare Center, Nagoya, Japan

³Department of Gastroenterology and Metabolism, Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan

⁴Department of Therapeutic Research, National Hospital Organization (NHO) Nagasaki Medical Center, Nagasaki, Japan

⁵The Research Center for Hepatitis and Immunology, National Center for Global Health and Medicine, Ichikawa, Japan

⁶PhoenixBio Co. Ltd., Higashi-Hiroshima, Japan

⁷Tokyo Metropolitan Institute of Medical Science, Tokyo, Japan

Correspondence to

Dr Masashi Mizokami, The Research Center for Hepatitis and Immunology, National Center for Global Health and Medicine 1-7-1, Kohnodai, Ichikawa 272-8516, Japan; mmizokami@hospk.ncgm.go.jp

Revised 4 October 2012
Accepted 9 October 2012

ABSTRACT

Objective Recent studies have demonstrated that genetic polymorphisms near the *IL28B* gene are associated with the clinical outcome of pegylated interferon α (peg-IFN- α) plus ribavirin therapy for patients with chronic hepatitis C virus (HCV). However, it is unclear whether genetic variations near the *IL28B* gene influence hepatic interferon (IFN)-stimulated gene (ISG) induction or cellular immune responses, lead to the viral reduction during IFN treatment.

Design Changes in HCV-RNA levels before therapy, at day 1 and weeks 1, 2, 4, 8 and 12 after administering peg-IFN- α plus ribavirin were measured in 54 patients infected with HCV genotype 1. Furthermore, we prepared four lines of chimeric mice having four different lots of human hepatocytes containing various single nucleotide polymorphisms (SNP) around the *IL28B* gene. HCV infecting chimeric mice were subcutaneously administered with peg-IFN- α for 2 weeks.

Results There were significant differences in the reduction of HCV-RNA levels after peg-IFN- α plus ribavirin therapy based on the *IL28B* SNP rs8099917 between TT (favourable) and TG/GG (unfavourable) genotypes in patients; the first-phase viral decline slope per day and second-phase slope per week in TT genotype were significantly higher than in TG/GG genotype. On peg-IFN- α administration to chimeric mice, however, no significant difference in the median reduction of HCV-RNA levels and the induction of antiviral ISG was observed between favourable and unfavourable human hepatocyte genotypes.

Conclusions As chimeric mice have the characteristic of immunodeficiency, the response to peg-IFN- α associated with the variation in *IL28B* alleles in chronic HCV patients would be composed of the intact immune system.

INTRODUCTION

Hepatitis C is a global health problem that affects a significant portion of the world's population. The WHO estimated that, in 1999, 170 million hepatitis C virus (HCV)-infected patients were present worldwide, with 3–4 million new cases appearing per year.¹

The standard therapy for hepatitis C still consists of pegylated interferon- α (peg-IFN- α), administered once weekly, plus daily oral ribavirin for 24–48 weeks

Significance of this study

What is already known on this subject?

- Genetic polymorphisms near the *IL28B* gene are associated with a chronic HCV treatment response.
- HCV-infected patients with the *IL28B* homozygous favourable allele had a more rapid decline in HCV kinetics in the first and second phases by peg-IFN- α -based therapy.
- During the acute phase of HCV infection, a strong immune response among patients with the *IL28B* favourable genotype could induce more frequent spontaneous clearance of HCV.

What are the new findings?

- In chronically HCV genotype 1b-infected chimeric mice that have the characteristic of immunodeficiency, no significant difference in the reduction in serum HCV-RNA levels and the induction of antiviral hepatic ISG by the administration of peg-IFN- α was observed between favourable and unfavourable human hepatocyte *IL28B* genotypes.
- By comparison of serum HCV kinetics between human and chimeric mice, the viral decline in both the first and second phases by peg-IFN- α treatment was affected by the variation in *IL28B* genotypes only in chronic hepatitis C patients.

How might it impact on clinical practice in the foreseeable future?

- The immune response according to *IL28B* genetic variants could contribute to the first and second phases of HCV-RNA decline and might be critical for HCV clearance by peg-IFN- α -based therapy.

in countries where protease inhibitors are not available.² This combination therapy is quite successful in patients with HCV genotype 2 or 3 infection, leading to a sustained virological response (SVR) in approximately 80–90% of patients treated; however, in patients infected with HCV genotype 1 or 4, only approximately half of all treated individuals achieved a SVR.^{3 4}

Viral hepatitis

Table 1 Characteristics of 54 patients infected HCV genotype 1

	<i>IL28B</i> SNP rs8099917		p Value
	TT (n=34)	TG (n=19) + GG (n=1)	
Age (years)	55.6±10.1	54.7±11.3	0.746
Gender (male %)	70	50	0.199
Body mass index (kg/m ²)	24.6±3.1	24.7±3.3	0.870
Viral load at therapy (log IU/ml)	6.0±0.7	5.8±0.8	0.357
SVR rate (%)	50	11	0.012
Serum ALT level (IU/l)	100.3±80.8	79.3±45.0	0.226
Platelet count (×10 ⁴ /μl)	17.1±9.0	16.5±5.8	0.771
Fibrosis (F3+4 %)	42	40	0.877

HCV, hepatitis C virus; SNP, single nucleotide polymorphism; SVR, sustained virological response.

Host factors were shown to be associated with the outcome of the therapy, including age, sex, race, liver fibrosis and obesity.⁵ Genome-wide association studies have demonstrated that genetic variations in the region near the interleukin-28B (*IL28B*) gene, which encodes interferon (IFN)-λ3, are associated with a chronic HCV treatment response.^{6–10} Furthermore, it was demonstrated that genetic variations in the *IL28B* gene region are also associated with spontaneous HCV clearance.^{11–12}

Interestingly, a recent report showed the effect of genetic polymorphisms near the *IL28B* gene on the dynamics of HCV during peg-IFN-α plus ribavirin therapy in Caucasian, African American and Hispanic individuals;¹³ HCV-infected patients with the *IL28B* homozygous favourable allele had a more rapid decline of HCV in the first phase, which is associated with the inhibition of viral replication as well as the second phase associated with immuno-destruction of viral-infected hepatocytes.¹⁴ However, it is unknown how a direct effect by the *IL28B* genetic variation, such as the induction of IFN-stimulated genes (ISG) or cellular immune responses, would influence the viral kinetics during IFN treatment. Over recent periods, engineered severe combined immunodeficient (SCID) mice transgenic for urokinase-type plasminogen activator (uPA) received human hepatocyte transplants (hereafter referred to as chimeric mice)^{15–17} and are suitable for experiments with hepatitis viruses in vivo.^{18–19} We have also reported that these chimeric mice carrying human hepatocytes are a robust animal model to evaluate the efficacy of IFN and other anti-HCV agents.^{20–21}

The purpose of this study was to reveal the association between genetic variations in the *IL28B* gene region and viral decline during peg-IFN-α treatment in patients with HCV, and to clarify the association between different *IL28B* alleles of human hepatocytes in chimeric mice and the response to peg-IFN-α without immune response. These studies will elucidate whether the immune response by the *IL28B* genetic variation affects the viral kinetics during peg-IFN-α treatment.

MATERIALS AND METHODS

Patients

Fifty-four Japanese patients with chronic HCV genotype 1 infection at Nagasaki Medical Center and Nagoya City

University were enrolled in this study (table 1). Patients received peg-IFN-α2a (180 μg) or 2b (1.5 μg/kg) subcutaneously every week and were administered a weight-adjusted dose of ribavirin (600 mg for <60 kg, 800 mg for 60–80 kg, and 1000 mg for >80 kg daily), which is the recommended dosage in Japan. Patients with other hepatitis virus infection or HIV coinfection were not included in the study. The study protocol conformed to the ethics guidelines of the 1975 Declaration of Helsinki as reflected by earlier approval by the institutions' human research committees.

Laboratory tests

Blood samples were obtained before therapy, as well as on day 1 and at weeks 1, 2, 4, 8 and 12 after the start of therapy and were analysed for the HCV-RNA level by the commercial Abbott Real-Time HCV test with a lower limit of detection of 12 IU/ml (Abbott Molecular Inc., Des Plaines, Illinois, USA). Genetic polymorphism in the *IL28B* gene (rs8099917), a single nucleotide polymorphism (SNP) recently identified to be associated with treatment response,^{6–8} was tested by the TaqMan SNP genotyping assay (Applied Biosystems, Foster City, California, USA).

HCV infection of chimeric mice with the liver repopulated for human hepatocytes

SCID mice carrying the uPA transgene controlled by an albumin promoter were injected with 5.0–7.5×10⁵ viable hepatocytes through a small left-flank incision into the inferior splenic pole, thereafter chimeric mice were generated. The chimeric mice were purchased from PhoenixBio Co, Ltd (Hiroshima, Japan).¹⁷ Human hepatocytes with the *IL28B* homozygous favourable allele, heterozygous allele or homozygous unfavourable allele were imported from BD Biosciences (San Jose, California, USA) (table 2). Murine serum levels of human albumin and the body weight were not significantly different among four chimeric mice groups, providing a reliable comparison for anti-HCV agents.²² Three different serum samples were obtained from three chronic HCV patients (genotype 1b).^{21–22} Each mouse was intravenously infected with serum sample containing 10⁵ copies of HCV genotype 1b. Administration of peg-IFN-α2a (Pegasys; Chugai Pharmaceutical Co., Ltd., Tokyo, Japan) at the dose formulation (30 μg/kg) was consecutively applied to each mouse on days 0, 3, 7 and 10 (table 3).

HCV-RNA quantification

HCV-RNA in mice sera (days 0, 1, 3, 7 and 14) was quantified by an in-house real-time detection PCR assay with a lower quantitative limit of detection of 10 copies/assay, as previously reported.²¹

Quantification of IFN-stimulated gene-expression levels

For analysis of endogenous ISG levels, total RNA was isolated from the liver using the RNeasy RNA extraction kit (Qiagen, Valencia, California, USA) and complementary DNA synthesis

Table 2 Four lines of uPA/SCID mice from four different lots of human hepatocytes (donor) containing various SNP around the *IL28B* gene

uPA/SCID mice	Donor	Race	Age	Gender	rs8103142	rs12979860	rs8099917
PXB mice	A	African American	5 Years	Male	CC	TT	TG
	B	Caucasian	10 Years	Female	CC	TT	TG
	C	Hispanic	2 Years	Female	TT	CC	TT
	D	Caucasian	2 Years	Male	TT	CC	TT

PXB mice; urokinase-type plasminogen activator/severe combined immunodeficiency (uPA/SCID) mice repopulated with approximately 80% human hepatocytes. SCID, severe combined immunodeficient; SNP, single nucleotide polymorphism.

Table 3 Dosage and time schedule of pegIFN- α 2a* treatment for HCV genotype 1b infected chimeric mice

Donor hepatocytes†	No of chimeric mice	Inoculum	Test compound	Dose			
				Level (μ g/kg)	Concentration (μ g/ml)	Volume (ml/kg)	Frequency
A	3	Serum A	Peg-IFN- α 2a	30	3	10	Day 0, 3, 7, 10
B	4	Serum A	Peg-IFN- α 2a	30	3	10	Day 0, 3, 7, 10
C	3	Serum A	Peg-IFN- α 2a	30	3	10	Day 0, 3, 7, 10
D	3	Serum A	Peg-IFN- α 2a	30	3	10	Day 0, 3, 7, 10
A	2	Serum B	Peg-IFN- α 2a	30	3	10	Day 0, 3, 7, 10
C	2	Serum B	Peg-IFN- α 2a	30	3	10	Day 0, 3, 7, 10
A	2	Serum C	Peg-IFN- α 2a	30	3	10	Day 0, 3, 7, 10
C	2	Serum C	Peg-IFN- α 2a	30	3	10	Day 0, 3, 7, 10

*Pegasys; Chugai Pharmaceutical Co., Ltd., Tokyo, Japan.

†The *IL28B* genetic variation of the donor hepatocytes was indicated in table 2.

HCV, hepatitis C virus; peg-IFN- α , pegylated interferon α .

was performed using 2.0 μ g of total RNA (High Capacity RNA-to-cDNA kit; Applied Biosystems). Fluorescence real-time PCR analysis was performed using an ABI 7500 instrument (Applied Biosystems) and TaqMan Fast Advanced gene expression assay (Applied Biosystems). TaqMan Gene Expression Assay primer and probe sets (Applied Biosystems) are shown in the supplementary information (available online only). Relative amounts of messenger RNA, determined using a FAM-Labeled TaqMan probe, were normalised to the endogenous RNA levels of the housekeeping reference gene, glyceraldehyde-3-phosphate dehydrogenase. The delta Ct method ($2^{-(\text{delta Ct})}$) was used for quantitation of relative mRNA levels and fold induction.^{23 24}

Statistical analyses

Statistical differences were evaluated by Fisher's exact test or the χ^2 test with the Yates correction. Mice serum HCV-RNA and intrahepatic ISG expression levels were compared using the Mann-Whitney U test. Differences were considered significant if p values were less than 0.05.

RESULTS

Characteristics of the study patients

Genotypes (rs8099917) TT, TG and GG were detected in 34, 19 and one patient infected with HCV genotype 1, respectively. SVR rates were significantly higher in HCV patients with genotype TT than in those with genotype TG/GG (50% vs 11%, $p=0.012$). The initial HCV serum load was comparable between

genotypes TT and TG/GG (6.0 ± 0.7 vs 5.8 ± 0.8 log IU/ml). There were no significant differences in sex (male%, 70% vs 50%), age (55.6 ± 10.1 vs 54.7 ± 11.3 years), serum alanine aminotransferase level (100.3 ± 80.8 vs 79.3 ± 45.0 IU/L), platelet count (17.1 ± 9.0 vs $16.5\pm 5.8\times 10^4/\mu$ l) and fibrosis stages (F3/4%, 42% vs 40%) between HCV patients with the favourable (rs8099917 TT) and unfavourable (rs8099917 TG/GG) *IL28B* genotypes (table 1).

Changes in serum HCV-RNA levels in patients treated by peg-IFN- α plus ribavirin

Figure 1 shows the initial change in the serum HCV-RNA level for 14 days after peg-IFN- α plus ribavirin therapy in patients infected with HCV genotype 1 based on the genetic polymorphism near the *IL28B* gene. The immediate antiviral response (viral drop 24 h after the first IFN injection) was significantly higher in HCV patients with genotype TT than genotype TG/GG (-1.08 vs -0.39 log IU/ml, $p<0.001$). Figure 2 also shows the subsequent change in the serum HCV-RNA reduction after peg-IFN- α plus ribavirin therapy in patients infected with HCV genotype 1. Similarly, during peg-IFN- α plus ribavirin therapy, a statistically significant difference in the median reduction in serum HCV-RNA levels was noted according to the genotype (TT vs TG/GG). The median reduction in the serum HCV-RNA levels (log IU/ml) at 1, 2, 4, 8 and 12 weeks between genotypes TT and TG/GG was as follows: -1.58 vs -0.62 , $p<0.001$; -2.35 vs -0.91 , $p<0.001$;




Figure 1 Rapid reduction of median hepatitis C virus (HCV)-RNA levels (log IU/ml) at 1, 7 and 14 days between *IL28B* single nucleotide polymorphisms rs8099917 genotype TT (n=34) and TG/GG (n=20) in HCV genotype 1-infected patients treated with peg-IFN- α plus ribavirin.





Figure 2 Weekly reduction of median hepatitis C virus (HCV)-RNA levels (log IU/ml) at 1, 2, 4, 8 and 12 weeks between *IL28B* single nucleotide polymorphisms rs8099917 genotype TT (n=34) and TG/GG (n=20) in HCV genotype 1-infected patients treated with pegylated interferon α plus ribavirin.

Viral hepatitis

Figure 3 (A) The first-phase viral decline slope per day (Ph1/day) and (B) second-phase viral decline slope per week (Ph2/week) in hepatitis C virus (HCV) genotype 1-infected patients treated with pegylated interferon α plus ribavirin. The lines across the boxes indicate the median values. The hash marks above and below the boxes indicate the 90th and 10th percentiles for each group, respectively.



-3.48 vs -1.56 , $p < 0.001$; -4.53 vs -2.37 , $p < 0.01$; -4.93 vs -2.86 , $p < 0.001$. Furthermore, the initial first-phase viral decline slope per day (Ph1/day) and subsequent second-phase viral decline slope per week (Ph2/week) in TT genotype were significantly higher than in genotype TG/GG (Ph1/day 0.94 ± 0.83 vs 0.38 ± 0.40 log IU/ml, $p < 0.001$; Ph2/week 0.08 ± 0.06 vs 0.04 ± 0.03 log IU/ml, $p < 0.001$) (figure 3).

Changes in serum HCV-RNA levels in chimeric mice treated by peg-IFN- α

In order to clarify the association between *IL28B* alleles of human hepatocytes and the response to peg-IFN- α , we prepared four lines of uPA/SCID mice and four different lots of human hepatocytes containing various rs8099917, rs8103142

and rs12979860 SNPs around the *IL28B* gene (table 2). The chimeric mice were inoculated with serum samples from each HCV-1b patient, and then HCV-RNA levels had increased and reached more than 10^6 copies/ml in all chimeric mice sera at 2 weeks after inoculation. After confirming the peak of HCV-RNA in all chimeric mice, they were subcutaneously administered with four times injections of the bolus dose of peg-IFN- α for 2 weeks (table 3). Figure 4 shows the change in the serum HCV-RNA levels for 14 days during IFN injection into chimeric mice transplanted with *IL28B* favourable or unfavourable human hepatocyte genotypes. On peg-IFN- α administration, no significant difference in the median reduction in HCV-RNA levels in the serum A-infected²² chimeric mice sera was observed between favourable (n=7) and unfavourable




Figure 4 Median reduction of hepatitis C virus (HCV)-RNA levels (log copies/ml) after administering pegylated interferon α to chimeric mice having human hepatocytes containing various single nucleotide polymorphisms around the *IL28B* gene as favourable (rs8099917 TT) and unfavourable (rs8099917 TG) genotypes. Data are represented as mean \pm SD. Chimeric mice infected with a) serum A (n=7; favourable genotype, n=6; unfavourable genotype), (B) serum B (n=2, each genotype), and (C) serum C (n=2, each genotype). All serum samples were obtained from HCV-1b patients.

(n=6) *IL28B* genotypes on days 1, 3, 7 and 14 (-1.2 vs -1.3, -1.4 vs -1.4, -1.8 vs -1.7, and -2.3 vs -1.9 log copies/ml) (figure 4A). Moreover, we prepared two additional serum samples from the other HCV-1b patients (serum B and C)²¹ to confirm the influence of *IL28B* genotype in early viral kinetics during IFN treatment. After establishing persistent infection with new HCV-1b strains in all chimeric mice, they were also administered four times injections of the bolus dose of peg-IFN- α 2a for 2 weeks (figure 4B,C). In a similar fashion, no significant difference in HCV-RNA reduction in chimeric mice sera was observed between favourable and unfavourable *IL28B* genotypes.


Expression levels of ISG in chimeric mice livers

Because chimeric mice have the characteristic of severe combined immunodeficiency, the viral kinetics in chimeric mice

sera during IFN treatment could be contributed by the innate immune response of HCV-infected human hepatocytes. Therefore, ISG expression levels in mice livers transplanted with human hepatocytes were compared between favourable and unfavourable *IL28B* genotypes (figure 5).

As shown in figure 5A, ISG expression levels in mice livers were measured at 8 h and 24 h after IFN treatment. The levels of representative antiviral ISG (eg, myxovirus resistance protein A, oligoadenylate synthetase 1, RNA-dependent protein kinase) and other ISG for promoting antiviral signalling (eg, Toll-like receptor 3, retinoic acid-inducible gene 1) were significantly induced at least 8 h after treatment, and prolonged at 24 h. No significant difference in ISG expression levels in HCV-infected livers was observed between favourable and unfavourable *IL28B* genotypes. The other inoculum for persistent infection of HCV-1b also demonstrated no significant difference in ISG

Figure 5 Intrahepatic interferon (IFN)-stimulated gene (ISG) expression levels in the pegylated interferon α (peg-IFN- α)-treated chimeric mice having human hepatocytes containing homozygous favourable allele (rs8099917 TT; MA) and heterozygous unfavourable allele (rs8099917 TG; HE) were measured and expressed relative to glyceraldehyde-3-phosphate dehydrogenase (GAPDH) messenger RNA. Data are represented as mean+SD. (A) Time kinetics of ISG after administration of the peg-IFN- α in serum A-infected chimeric mice (n=3, each genotype). Comparison of ISG expression levels at (B) 8 h in serum B-infected mice and (C) 24 h in serum C-infected mice after administering peg-IFN- α (n=3, each genotype). Predesigned real-time PCR assay of *IL28B* transcript purchased from Applied Biosystems can be cross-reactive to *IL28A* transcript. *p<0.05. MxA, myxovirus resistance protein A; OAS1, oligoadenylate synthetase 1; PKR, RNA-dependent protein kinase; RIG-1, retinoic acid-inducible gene 1; TLR3, Toll-like receptor 3.



Viral hepatitis

expression levels between favourable and unfavourable *IL28B* genotypes (figure 5B,C). Interestingly, IFN- λ expression levels by treatment of peg-IFN- α were significantly induced in HCV-infected human hepatocytes harbouring the favourable *IL28B* genotype (figure 5 A–C).

DISCUSSION

Several recent studies have demonstrated a marked association between the chronic hepatitis C treatment response^{6–9} and SNP (rs8099917, rs8103142 and rs12979860) near or within the region of the *IL28B* gene, which affected the viral dynamics during peg-IFN- α plus ribavirin therapy in Caucasian, African American and Hispanic individuals.¹³

It has been reported that when patients with chronic hepatitis C are treated by IFN- α or peg-IFN- α plus ribavirin, HCV-RNA generally declines after a 7–10 h delay.²⁵ The typical decline is biphasic and consists of a rapid first phase lasting for approximately 1–2 days during which HCV-RNA may fall 1–2 logs in patients infected with genotype 1, and subsequently a slower second phase of HCV-RNA decline.²⁶ The viral kinetics had a predictive value in evaluating antiviral efficacy.¹⁴ In this study, biphasic decline of the HCV-RNA level during peg-IFN- α treatment was observed in both patients and chimeric mice infected with HCV genotype 1; however, in the first and second phases of viral kinetics, a difference between *IL28B* genotypes was observed only in HCV-infected patients; a more rapid decline in serum HCV-RNA levels after administering peg-IFN- α plus ribavirin was confirmed in patients with the TT genotype of rs8099917 compared to those with the TG/GG genotype.

On the other hand, in-vivo data using the chimeric mouse model showed no significant difference in the reduction of HCV-RNA titers in mouse serum among four different lots of human hepatocytes containing *IL28B* favourable (rs8099917 TT) or unfavourable (rs8099917 TG) genotypes, which was confirmed by the inoculation of two additional HCV strains. These results indicated that variants of the *IL28B* gene in donor hepatocytes had no influence on the response to peg-IFN- α under immunosuppressive conditions, suggesting that the immune response according to *IL28B* genetic variants could contribute to the first and second phases of HCV-RNA decline and might be critical for HCV clearance by peg-IFN- α -based therapy.

Two recent studies indeed revealed an association between the *IL28B* genotype and the expression level of hepatic ISG in human studies.^{27–28} Quiescent hepatic ISG before treatment among patients with the *IL28B* favourable genotype have been associated with sensitivity to exogenous IFN treatment and viral eradication; however, it is difficult to establish whether the hepatic ISG expression level contributes to viral clearance independently or appears as a direct consequence of the *IL28B* genotype. Another recent study addressed this question and the results suggested that there is no absolute correlation with the *IL28B* genotype and hepatic expression of ISG.²⁹ Our results on the hepatic ISG expression level in immunodeficient chimeric mice also suggested that no significant difference in ISG expression levels was observed between favourable and unfavourable *IL28B* genotypes. However, these results were not consistent with a previous report using chimeric mice that the favourable *IL28B* genotype was associated with an early reduction in HCV-RNA by ISG induction.³⁰ The reasons for the discrepancy might depend on the dose and type of IFN treatment, as well as the time point when ISG expression was examined in the liver. In addition, although IFN- λ transcript levels measured in peripheral blood mononuclear cells or liver revealed inconsistent

results in the context of an association with the *IL28B* genotype,^{7–8} our preliminary assay on the *IL28A*, *IL28B* and *IL29* transcripts in the liver first indicated that the induction of IFN- λ on peg-IFN- α administration could be associated with the *IL28B* genotype. Therefore, the induction of IFN- λ followed by immune response might contribute to different viral kinetics and treatment outcomes in HCV-infected patients, because no difference was found in chimeric mice without immune response.

It has also been reported that the mechanism of the association of genetic variations in the *IL28B* gene and spontaneous clearance of HCV may be related to the host innate immune response.¹¹ Interestingly, participants with seroconversion illness with jaundice were more frequently rs8099917 homozygous favourable allele (TT) than other genotypes (32% vs 5%, $p=0.047$). This suggests that a stronger immune response during the acute phase of HCV infection among patients with the *IL28B* favourable genotype would induce more frequent spontaneous clearance of HCV.

Taking into account both the above results in acute HCV infection and our results conducted on chimeric mice that have the characteristic of immunodeficiency, it is suggested that the response to peg-IFN- α associated with the variation in *IL28B* alleles in chronic hepatitis C patients would be composed of the intact immune system.

Acknowledgements The authors would like to thank Kyoko Ito of Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan for doing the quantification of gene-expression assays.

Contributors YT and MM conceived the study. TW and FS and YT conducted the study equally. TW and FS coordinated the analysis and manuscript preparation. All the authors had input into the study design, patient recruitment and management or mouse management and critical revision of the manuscript for intellectual content. TW, FS and YT contributed equally.

Funding This study was supported by a grant-in-aid from the Ministry of Health, Labour, and Welfare of Japan (H22-kannen-005) and the Ministry of Education, Culture, Sports, Science and Technology, Japan, and grant-in-aid for research in Nagoya City University.

Competing interests None.

Patient consent Obtained.

Ethics approval This study was conducted with the approval of each ethics committee at the Nagoya City University and Nagasaki Medical Center (see supplementary information, available online only).

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. Ray Kim W. Global epidemiology and burden of hepatitis C. *Microbes Infect* 2002;**4**:1219–25.
2. Foster GR. Past, present, and future hepatitis C treatments. *Semin Liver Dis* 2004;**24**(Suppl. 2):97–104.
3. Fried MW, Shiffman ML, Reddy KR, et al. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002;**347**:975–82.
4. Manns MP, McHutchison JG, Gordon SC, et al. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet* 2001;**358**:958–65.
5. Mihm U, Herrmann E, Sarrazin C, et al. Review article: predicting response in hepatitis C virus therapy. *Aliment Pharmacol Ther* 2006;**23**:1043–54.
6. Ge D, Fellay J, Thompson AJ, et al. Genetic variation in *IL28B* predicts hepatitis C treatment-induced viral clearance. *Nature* 2009;**461**:399–401.
7. Suppiah V, Moldovan M, Ahlenstiel G, et al. *IL28B* is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nat Genet* 2009;**41**:1100–4.
8. Tanaka Y, Nishida N, Sugiyama M, et al. Genome-wide association of *IL28B* with response to pegylated interferon-alpha and ribavirin therapy for chronic hepatitis C. *Nat Genet* 2009;**41**:1105–9.
9. Rauch A, Kutalik Z, Descombes P, et al. Genetic variation in *IL28B* is associated with chronic hepatitis C and treatment failure: a genome-wide association study. *Gastroenterology* 2010;**138**:1338–45.

10. **Tanaka Y**, Nishida N, Sugiyama M, *et al.* lambda-Interferons and the single nucleotide polymorphisms: a milestone to tailor-made therapy for chronic hepatitis C. *Hepatol Res* 2010;**40**:449–60.
11. **Thomas DL**, Thio CL, Martin MP, *et al.* Genetic variation in IL28B and spontaneous clearance of hepatitis C virus. *Nature* 2009;**461**:798–801.
12. **Grebely J**, Petoumenos K, Hellard M, *et al.* Potential role for interleukin-28B genotype in treatment decision-making in recent hepatitis C virus infection. *Hepatology* 2010;**52**:1216–24.
13. **Thompson AJ**, Muir AJ, Sulkowski MS, *et al.* Interleukin-28B polymorphism improves viral kinetics and is the strongest pretreatment predictor of sustained virologic response in genotype 1 hepatitis C virus. *Gastroenterology* 2010;**139**:120–29.
14. **Layden-Almer JE**, Layden TJ. Viral kinetics in hepatitis C virus: special patient populations. *Semin Liver Dis* 2003;**23**(Suppl. 1):29–33.
15. **Heckel JL**, Sandgren EP, Degen JL, *et al.* Neonatal bleeding in transgenic mice expressing urokinase-type plasminogen activator. *Cell* 1990;**62**:447–56.
16. **Rhim JA**, Sandgren EP, Degen JL, *et al.* Replacement of diseased mouse liver by hepatic cell transplantation. *Science* 1994;**263**:1149–52.
17. **Tateno C**, Yoshizane Y, Saito N, *et al.* Near completely humanized liver in mice shows human-type metabolic responses to drugs. *Am J Pathol* 2004;**165**:901–12.
18. **Mercer DF**, Schiller DE, Elliott JF, *et al.* Hepatitis C virus replication in mice with chimeric human livers. *Nat Med* 2001;**7**:927–33.
19. **Tsuge M**, Hiraga N, Takaishi H, *et al.* Infection of human hepatocyte chimeric mouse with genetically engineered hepatitis B virus. *Hepatology* 2005;**42**:1046–54.
20. **Kurbanov F**, Tanaka Y, Chub E, *et al.* Molecular epidemiology and interferon susceptibility of the natural recombinant hepatitis C virus strain RF1_2k/1b. *J Infect Dis* 2008;**198**:1448–56.
21. **Kurbanov F**, Tanaka Y, Matsuura K, *et al.* Positive selection of core 70Q variant genotype 1b hepatitis C virus strains induced by pegylated interferon and ribavirin. *J Infect Dis* 2010;**201**:1663–71.
22. **Inoue K**, Umehara T, Ruegg UT, *et al.* Evaluation of a cyclophilin inhibitor in hepatitis C virus-infected chimeric mice in vivo. *Hepatology* 2007;**45**:921–8.
23. **Livak KJ**, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 2001;**25**:402–8.
24. **Silver N**, Best S, Jiang J, *et al.* Selection of housekeeping genes for gene expression studies in human reticulocytes using real-time PCR. *BMC Mol Biol* 2006;**7**:33.
25. **Dahari H**, Layden-Almer JE, Perelson AS, *et al.* Hepatitis C viral kinetics in special populations. *Curr Hepat Rep* 2008;**7**:97–105.
26. **Neumann AU**, Lam NP, Dahari H, *et al.* Hepatitis C viral dynamics in vivo and the antiviral efficacy of interferon-alpha therapy. *Science* 1998;**282**:103–7.
27. **Honda M**, Sakai A, Yamashita T, *et al.* Hepatic ISG expression is associated with genetic variation in interleukin 28B and the outcome of IFN therapy for chronic hepatitis C. *Gastroenterology* 2010;**139**:499–509.
28. **Urban TJ**, Thompson AJ, Bradrick SS, *et al.* IL28B genotype is associated with differential expression of intrahepatic interferon-stimulated genes in patients with chronic hepatitis C. *Hepatology* 2010;**52**:1888–96.
29. **Dill MT**, Duong FH, Vogt JE, *et al.* Interferon-induced gene expression is a stronger predictor of treatment response than IL28B genotype in patients with hepatitis C. *Gastroenterology* 2011;**140**:1021–31.
30. **Hiraga N**, Abe H, Imamura M, *et al.* Impact of viral amino acid substitutions and host interleukin-28b polymorphism on replication and susceptibility to interferon of hepatitis C virus. *Hepatology* 2011;**54**:764–71.

Mechanism of the dependence of hepatitis B virus genotype G on co-infection with other genotypes for viral replication

T. Sakamoto,^{1,2} Y. Tanaka,¹ T. Watanabe,¹ S. Iijima,¹ S. Kani,¹ M. Sugiyama,¹ S. Murakami,¹ K. Matsuura,^{1,2} A. Kusakabe,^{1,2} N. Shinkai,^{1,2} S. Fuminaka,^{1,2} and M. Mizokami^{1,3}

¹Department of Virology & Liver unit; ²Department of Gastroenterology and Metabolism, Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan; and ³Research Center for Hepatitis and Immunology, National Center for Global Health and Medicine, Ichikawa, Japan

Received April 2012; accepted for publication September 2012

SUMMARY. Hepatitis B virus (HBV) is classified into several genotypes. Genotype G (HBV/G) is characterised by worldwide dispersion, low intragenotypic diversity and a peculiar sequence of the precore and core region (stop codon and 36-nucleotide insertion). As a rule, HBV/G is detected in co-infection with another genotype, most frequently HBV/A2. In a previous *in vivo* study, viral replication of HBV/G was significantly enhanced by co-infection with HBV/A2. However, the mechanism by which co-infection with HBV/A2 enhances HBV/G replication is not fully understood. In this study, we employed 1.24-fold HBV/A2 clones that selectively expressed each viral protein and revealed that the core protein expressing construct significantly enhanced the replication of HBV/G in Huh7 cells. The introduction of the HBV/A2 core promoter or core protein or both genomic regions into the HBV/G genome showed

that both the core promoter and core protein are required for efficient HBV/G replication. The effect of genotype on the interaction between foreign core protein and HBV/G showed that HBV/A2 was the strongest enhancer of HBV/G replication. Furthermore, Western blot analysis of Dane particles isolated from cultures of Huh7 cells co-transfected by HBV/G and a cytomegalovirus (CMV) promoter-driven HBV/A2 core protein expression construct indicated that HBV/G employed HBV/A2 core protein during particle assembly. In conclusion, HBV/G could take advantage of core proteins from other genotypes during co-infection to replicate efficiently and to effectively package HBV DNA into virions.

Keywords: co-transfection, core protein, genotype A, genotype G, hepatitis B virus, replication.

INTRODUCTION

Hepatitis B virus (HBV) infection affects more than 350 million people and is one of the major causes of acute and chronic liver disease. Acute HBV infection in adults is usually self-limiting, while chronic HBV infection can cause chronic hepatitis, liver cirrhosis or hepatocellular carcinoma [1]. As the clinical course in infected individuals depends on a complex interplay among various factors including viral, host and environmental factors, molecular characteristics of HBV including the genotype could become increasingly important in our understanding of HBV clinical implications [2].

Abbreviations: CMV, cytomegalovirus; CP, core promoter; HBeAg, hepatitis B e antigen; HBsAg, hepatitis B surface antigen; HBV, hepatitis B virus; SEAP, secreted alkaline phosphatase.

Correspondence: Masashi Mizokami, MD, PhD, The Research Center for Hepatitis and Immunology, National Center for Global Health and Medicine 1-7-1, Kohnodai, Ichikawa 272-8516, Japan. E-mail: mmizokami@hospk.ncgm.go.jp

Eight major HBV genotypes (A–H) have been identified by a sequence divergence >8% in the entire HBV genome [3,4] and have a relatively distinct geographical distribution, which may be associated with anthropological history [5]. Hepatitis B virus genotype G (HBV/G) was first described in 2000 by studies carried out in France [6]. It is usually detected during co-infection with other genotypes, most frequently with HBV/A2 [7,8]. Co-infection with HBV/C and H has also been reported [9–11]. One of the features distinguishing HBV/G from other genotypes is the 36-nucleotide (nt) insertion in its core gene [6,12]. Recent studies indicated that the 36-nt insertion increased core protein translation without enhancing mRNA abundance [13], and insertion of the 36-nt in the core region of genotypes A and D impaired genome replication, despite upregulation of core protein expression, indicating that the 36-nt insertion could alter core protein expression without altering the mRNA expression [14]. The other feature of the HBV/G genome that is unique is the possession of two stop codons in the precore region that prevents the expression of hepatitis B e antigen (HBeAg) [6,12]. Nevertheless,

some HBV/G carriers are HBeAg positive, which is explained by co-infection with an HBeAg-expressing HBV/A strain [7].

As previously reported, HBV/G monoinfection in uPA/SCID mice that had been transplanted with human hepatocytes (hereafter referred to as chimeric mice) resulted in very low level viral replication, but HBV replication increased markedly when the animals were co-infected with HBV/A2, C or H [11,15]. Furthermore, the co-infection induced more pronounced fibrosis, which concurs with findings from studies of immunosuppressed patients [16]. However, as it is still unclear how the interaction between HBV/G and other genotypes enhances the replication of HBV/G and affects the virological and clinical manifestation within an individual, we conducted *in vitro* studies using 1.24-fold HBV clones to elucidate the mechanism of HBV/G replication during co-infection.

MATERIALS AND METHODS

Plasmid constructs of HBV DNA and sequencing

Hepatitis B virus DNA was extracted from 100 μ L of serum using the QIAamp DNA blood kit (Qiagen GmbH, Hilden, Germany). Four primer sets were designed to amplify two fragments (A and B) covering the entire HBV/G genome. PCR with nested primers was performed using TaKaRa LA Taq polymerase (Takara Biochemicals, Kyoto, Japan) for 35 cycles (30 s at 95° C, 30 s at 60° C and 2 min at 72° C). The primer pairs and protocols for plasmid construction are outlined in the Supporting Information. As reported previously [17], these fragments were added to the pUC19 vector, which had been deprived of promoters (Invitrogen Corp., Carlsbad, CA, USA), by digestion with *Hind*III and *Eco*RI, resulting in the 1.24-fold HBV genome – required to transcribe the oversized pregenome and precore messenger RNA. Cloned HBV DNA sequences were confirmed with Prism BigDye (Applied Biosystems, Foster City, CA, USA) using the ABI 3100 automated sequencer.

HBV DNA mutagenesis and construct design

HBV/A2 and HBV/G clones containing the 1.24-fold HBV genome were constructed using isolates obtained from a co-infected Caucasian patient from the San Francisco cohort described in our previous study (patient #1) [7]. The study design conformed to the 1975 Declaration of Helsinki and was approved by our institutional ethics committee. Written informed consent was obtained from the patient. The HBV/A2 clones isolated from the patient's blood specimen did not possess any precore or core promoter mutations that are known to affect HBeAg expression. To study the interaction between the different genotype isolates, the following viral protein expression

constructs were prepared (outlined in Fig. 1) in HBV/A2 recombinant plasmids: HBV/A2-S, HBV/A2-core, HBV/A2-pol and HBV/A2-X and were each able to selectively translate one of the four viral proteins (the large surface, precore/core, polymerase and X proteins, respectively), whereas translation of the other three was prevented by the introduction of point mutations that produced corresponding stop codons (Fig. 1a). The following stop codons were used: (i) for surface protein: change from TTA to TAG in the 15th codon of the S gene (T198A) [18], (ii) for core protein: change from AAG to TAG in the 96th codon of the core gene (A2186T), (iii) for polymerase: change from CAACAA to TAATAA in the 283rd and 284th codons of the pol gene (C2558T/C2592T) and (iv) for X protein: change from CAA to TAA in the 7th codon of the HBx gene (C1395T) [19]. All of the above HBV/A2 recombinant plasmids possessed a TCTG motif after nucleotide position 1876, which abolished genome replication by altering the ϵ loop (CTGT to TCTG, nt 1877–1880) [20]. The 'HBV/A2-N' clone contained all six mutations and was used as an experimental negative control. All of the mutations in this study (substitutions, insertions and deletions) were created by overlapping PCR extension followed by the exchange of endonuclease enzyme-restricted fragments, as described previously [13,21].

Three cytomegalovirus (CMV) promoter-driven expression clones were constructed containing the whole core genes (not including the precore section) of HBV/G (nt 1901–2488), HBV/A2 (nt 1901–2458) and HBV/C (nt 1901–2452): CMV-HBV/G/core, CMV-HBV/A2/core and CMV-HBV/C/core, respectively (Fig. 1b).

Three replicating recombinant constructs were created by recombination of different genomic sections of HBV/G and HBV/A2 (Fig. 1c). The 'HBV-G/A2-CP' clone was a HBV/G-based construct in which the leading fragment containing the core promoter (CP) region (nt 1413–1806) was replaced with that of HBV/A2. The 'HBV-G/A2-CP+core' clone was also an HBV/G-based construct, in which the leading fragment containing the core promoter (CP), precore and core region (nt 1413–2821) of HBV/G was replaced with those of HBV/A2. The 'HBV-G/A2-core' clone was an HBV/G-based construct in which the fragment of the precore and core region (nt 1806–2821) was replaced with those of HBV/A2.

Cell culture and transfection

After 16 h of culture, Huh7 cells were transfected with 5 μ g of DNA construct per 10-cm diameter dish using the Fugene 6 transfection reagent (Promega, Madison, WI, USA) according to the manufacturer's protocol and harvested 3 days later. Transfection efficiency was measured by co-transfection with 0.5 μ g of a reporter plasmid expressing secreted alkaline phosphatase (SEAP) and normalised with subsequent SEAP measurement from culture




Fig. 1 HBV constructs (1.24-fold) and CMV-driven HBV core protein expression constructs used for the present study. CP, core promoter; BCP, basal core promoter; CMV, cytomegalovirus promoter. Stop codons for the corresponding HBV protein are indicated by crosses and arrows. All HBV/A2 recombinant plasmids consisted of the packaging-negative mutation (ϵ signal block). In three recombinant constructs between HBV/A2 and HBV/G, the corresponding recombinant genomic parts are shown by the grey bar. CMV-core constructs produce core protein without generating HBeAg in the absence of the preceding ϵ signal.

supernatant using a SEAP reporter assay kit (TOYOBO) [17]. Three experiments were conducted for each clone.

Determination of HBV markers

The expression levels of hepatitis B surface antigen (HBsAg) and HBeAg were determined by chemiluminescent enzyme immunoassay using commercial assay kits (Fujirebio Inc., Tokyo, Japan). The detection limit of the HBsAg assay is 0.05 IU/mL. HBV core-related antigen (HBcrAg) was measured in serum using a previously described chemiluminescent enzyme immunoassay [22]. The detection limit of the HBcrAg assay is 1.0 kU/mL.

Southern blot hybridisation

Southern blot hybridisation was performed with full-length probes for each genotype/subgenotype according to previously described methods [23]. In brief, cells were harvested and lysed in 1.5 mL of lysis buffer containing

50 mM Tris-HCl (pH 7.4), 1 mM EDTA and 1% NP-40. Half of the cell lysate was treated with 100 μ g/mL of RNase A and 200 μ g/mL of DNase I for 2 h at 37°C, in the presence of 6 mM Mg acetate. Then, HBV DNA was released by proteinase K digestion, extracted with phenol and precipitated with ethanol after the addition of 20 μ g of glycogen. DNA was separated on a 1.2% agarose gel, transferred to a positively charged nylon membrane (Roche Diagnostics, Germany) and hybridised with an alkaline phosphatase-labelled full-length HBV/G or HBV/A2 fragment generated with a Gene Images AlkPhos direct labelling module (GE Healthcare, UK). The detection was performed with CDP-Star, ready-to use (Roche Diagnostics). The signals were analysed by using a LAS-3000 image analyzer (Fuji Photo Film, Japan).

Western blot analysis

Serum or culture medium samples were subjected to SDS-PAGE under 15–25% polyacrylamide gel electrophoresis

conditions. The proteins in the gel were electroblotted onto a polyvinylidene difluoride membrane (Immobilon-P; Millipore, Billerica, MA, USA) at 15 V for 45 min. The membrane was then blocked and probed using alkaline phosphatase-conjugated HB50 (for HBcAg) or HB91 (for HBcrAg) monoclonal antibody [22] at room temperature for 1 h, before being washed and incubated with 5-bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium substrate solution (KPL, Gaithersburg, MD, USA) for 15 min (for HBcrAg) or 90 min (for HBcAg).

Sucrose density gradient ultracentrifugation

Aliquots (1.7 mL) of 10%, 20%, 30%, 40%, 50% or 60% (w/w) sucrose in 10 mM Tris-HCl, 150 mM NaCl and 1 mM EDTA (pH 7.5) were carefully layered in a 12-mL ultracentrifuge tube and left at room temperature for 6 h. The culture supernatant of Huh7 cells that had been co-transfected with the 1.24-fold HBV genome construct (HBV/G or HBV/A) and/or the CMV-HBV/A2-core plasmid was layered onto this sucrose gradient, and ultracentrifugation was performed at $200\,000 \times g$ for 15 h at 4°C in a Beckman Sw40Ti rotor. Fractions were collected from the top to the bottom of the gradient. The density of each fraction was calculated from its weight and volume. Each fraction was diluted 10-fold and tested for HBcrAg, HBsAg, HBeAg and HBV DNA.

Immunoprecipitation

Immunoprecipitation was carried out using magnetic beads coated with monoclonal anti-HBs from the 'Magneosphere™ MS300/Caboxyl' kit (JSR Corp., Tokyo, Japan) [24]. A 100- μL aliquot of sample was mixed with 100 μL of a magnetic bead suspension. The mixture was then incubated for 1 h at room temperature under gentle agitation and then magnetically separated. The core protein in the precipitate was analysed by Western blotting.

RESULTS

The replication of HBV/G is enhanced by HBV/A2 in co-transfection experiments

6 In this study, HBV/G and HBV/A2 genome clones (1.24-fold) were constructed from the serum of a HBV carrier that had been co-infected with HBV/G and HBV/A2. The HBV/G-d36 clone is a HBV/G genome-based construct in which the genotype-specific 36-nt insertion was deleted. We performed co-transfection with HBV/A2 and HBV/G clones and assessed virological features. Because of an over 12% sequence divergence between genotype A and G at the nucleotide level [12], the blot was hybridised successively with genotype-specific probes to DNA of each genotype. However, due to the unbiased binding of each probe at lower efficiency in Southern blot analysis [although the

replication of HBV/A2 was higher than that of HBV/G, relative value of HBV/A2 with probe G became lower (0.63), as well as the detection of HBV/G with probe A was very weak (0.24)], each probe of genotype G or A was used for hybridisation with the HBV/A2 and HBV/G clones (Fig. 2a). The density of single-strand HBV DNA detected by the genotype-specific probes in Southern blot analysis revealed that co-transfection with HBV/A2




Fig. 2 (a) Southern blot analysis for replicative activity among HBV/G montransfection, HBV/G-d36 montransfection, HBV/A2 montransfection and co-transfection with HBV/A2 and HBV/G (3 days after transfection). HBV/G-d36 clone was a deletion mutant lacking the 36-nt unique insertion in the core gene of the wild-type HBV/G clone. Hybridisation of the blot with genotype-specific probes of genotype A2 (upper) and G (lower). The density values shown at the bottom were measured to the probe-specific DNA sample. Single-stranded (SS) DNA is indicated by arrows. (b) Intracellular expression of core protein was estimated by detecting HBV-core-related antigen (HBcrAg) [22] as measured by a commercial chemiluminescent enzyme immunoassay (mean and standard deviation, $n = 3$). (c) HBsAg levels in the supernatant as detected by a commercial chemiluminescent enzyme immunoassay (mean and standard deviation, $n = 3$). All experiments were tested at least three times.

1 resulted in increased replication of the wild-type HBV/G
 2 clone, compared with monotransfection of the wild-type
 3 HBV/G (Fig. 2a). The intracellular expression of core pro-
 4 tein in the cell lysates and the expression of HBsAg in the
 5 culture supernatant were also enhanced by the co-transfec-
 6 tion with both HBV/A2 and HBV/G clones (Fig. 2b and
 7 2c). Removing the 36-nt insertion from the wild-type
 8 HBV/G genome resulted in a significant reduction in viral
 9 replication and core protein expression compared with the
 10 wild-type HBV/G clone. These results are in agreement
 11 with the observations of a previous study [13].

12 The core protein of HBV/A2 is essential for efficient 13 replication of HBV/G

14 To determine how HBV/A2 rescues HBV/G replication dur-
 15 ing co-transfection, we constructed four HBV/A2 recombi-
 16 nant plasmids that selectively expressed one of the four
 17 viral proteins, whereas translation of the other three was
 18 prevented by the introduction of stop codons (Fig. 1a). All
 19 of these plasmids were prevented from coding for the viral
 20 pregenomic RNA containing the 'packaging-negative'
 21 mutation in the ϵ signal loop to abrogate encapsidation
 22 (see Materials and methods). Huh7 cells were co-transfec-
 23 ted with the wild-type HBV/G clone and one of the four
 24 plasmids expressing a single viral protein. According to
 25 Southern blot analysis, the expression of intracellular HBV
 26 DNA was greatly increased when HBV/G was co-transfec-
 27 ted with HBV/A2-core compared with the other three
 28 expression plasmids or the experimental control (pUC19 or
 29 HBV/A2-N) (Fig. 3a). The intracellular expression of core
 30 protein in the cell lysates was also the highest when HBV/
 31 G was co-transfected with HBV/A2-core (Fig. 3b). The
 32 expression of HBsAg in the culture supernatant was only
 33 increased when HBV/G was co-transfected with the HBV/
 34 A2-S plasmid (Fig. 3c). These results indicated that the
 35 core protein translated from the HBV/A2 recombinant
 36 plasmid can enhance HBV/G replication.

37 The core protein of HBV/A2 is more effective than those 38 of HBV/C and HBV/G at promoting HBV/G replication

39 To compare the effects of genotype on the ability of the core
 40 protein to increase HBV/G replication in co-transfection
 41 experiments, we generated three genotype-specific core pro-
 42 tein expression constructs (HBV/G, HBV/A2 and HBV/C) dri-
 43 ven by the CMV promoter, which produced core protein in
 44 the absence of a preceding ϵ signal (Fig. 1c). Huh7 cells were
 45 co-transfected with HBV/G and one of the three core protein
 46 expression vectors. Southern blot analysis showed that the
 47 level of intracellular HBV DNA was highest during co-transfec-
 48 tion with CMV-HBV/A2/core, followed by CMV-HBV/G/
 49 core, and was lowest for CMV-HBV/C/core (Fig. 4a), although
 50 the expression of core protein in the cell lysates was the high-
 51 est during co-transfection with CMV-HBV/C/core, followed by




Fig. 3 (a) Southern blot analysis for replication competence of HBV/G clones co-transfected with each of the four HBV/A2 recombinant plasmids: HBV/A2-S, HBV/A2-core, HBV/A2-pol and HBV/A2-X selectively expressing only one of the four viral proteins (large surface, precore/core, polymerase or X protein, respectively). The 'HBV/A2-N' contained all the six mutations to be used as an experimental control. All of the above HBV/A2 recombinant plasmids had the 'packaging-negative' mutation in the ϵ signal to abrogate encapsidation. (b) Intracellular expression of core protein was measured as described in Fig. 2b. (c) The expression of HBsAg in the culture supernatant was detected as described in Fig. 2c.

CMV-HBV/G/core, and CMV-HBV/A2/core (Fig. 4b). As anticipated, there was no difference in the expression levels of HBsAg in any co-transfection experiment (Fig. 4c).

52 A comparison of viral replication among HBV/G and recombinant HBV/G clones

53 To examine the effects of genetic recombination and the roles of the core promoter, precore and core genomic regions in the interaction of HBV/G and HBV/A2 during co-transfection, we employed three HBV/G and HBV/A2 chimaeric replicating constructs (see Materials and methods), which are shown in Fig. 1c. After the transfection experiment, Southern blot analysis of cell lysates indicated an abundant level of DNA expression in HBV/G/A2-CP/core-transfected cells compared with those in the cells transfected with HBV/G-wild type, HBV/G/A2-CP and HBV/G/A2-core (Fig. 5a). As




Fig. 4 (a) Southern blot analysis for the expression of intracellular HBV/G DNA during co-transfection with the three core protein expression constructs for each genotype (HBV/G, HBV/A2 and HBV/C) driven by the CMV promoter, which produced core protein in the absence of a preceding ϵ signal. (b) Intracellular expression of core protein. (c) The expression of HBsAg in the culture supernatant.

shown in Fig. 5b, the highest levels of core protein (HBcrAg) expression were observed for the HBV/G/A2-CP and HBV/G/A2-CP/core-transfected cultures, which was in sharp contrast with the low levels observed in the HBV/G/A2-core and the wild-type HBV/G cultures. The discrepancy between viral replication and core production of the HBV/G/A2-CP clone might indicate insufficient virion assembly. Figure 5c shows the HBeAg levels measured in culture supernatants. The expression of HBeAg was the highest in the HBV/G/A2-CP/core culture distantly followed by that in the HBV/G/A2-core culture. The HBV/G/A2-CP and wild-type HBV/G clones expressed HBeAg protein at levels close to or below the level of detection. Nevertheless, a high HBcrAg titre was detected in the cell lysate of the HBV/G/A2-CP clone, although its DNA level was as low as that of the wild-type HBV/G clone (Fig. 5a). These results indicated that low replication of HBV/G might be explained by low synthesis of HBV/G core protein due to weak core promoter activity or dysfunction, as well as insufficient virion assembly due to the larger core protein of HBV/G (12-aa unique insertion).




Fig. 5 (a) Southern blot analysis for HBV replication among HBV/G and three chimeric replicating constructs created by recombination of different genomic sections of HBV/G and HBV/A2 (see Materials and methods). The 'HBV/G/A2-CP' clone was a HBV/G-based construct in which the fragment containing the core promoter (CP) region but not the precore or core was replaced by the corresponding sequence from HBV/A2. The 'HBV/G/A2-Core' clone was an HBV/G-based construct in which the section of the precore and core region was replaced with that of HBV/A2. For the 'HBV/G/A2-CP/core' clone, the CP, precore and core region of HBV/G were replaced with that of HBV/A2. (b) Intracellular expression of core protein. (c) Extracellular expression of HBeAg levels detected by a commercial chemiluminescent enzyme immunoassay (mean and standard deviation, $n = 3$).

Dane particles produced by HBV/G during co-transfection with HBV/A2 were packed in HBV/A2 protein

To investigate the effects of HBV/A2 core protein during HBV/G viral assembly, we tried to define whether the Dane particles in B Huh7 cells that had been co-transfected with wild-type HBV/G and the CMV-HBV/A2-core plasmid

(source of the Fig. 4 lane 2) contained HBV/A2 or HBV/G core protein. To extract the Dane particles, we employed ultracentrifugation of the culture media through a 10–60% (w/w) sucrose density gradient followed by immunoprecipitation using anti-HBs-coated magnetic beads. We thereby extracted Dane particles, which were then analysed using Western blotting. The obtained fractions were tested for HBcrAg, HBsAg and HBV DNA (Fig. 6a). HBcrAg appeared in the high-density fractions, and its levels peaked in the same fraction (fraction 22) as HBV DNA. As reported previously, the fraction in which the levels of HBV DNA and HBcrAg peaked contained Dane particles [22]. To eliminate contamination of the Dane particles with 'naked' core particles or core protein, they were specifically retrieved from sucrose high-density fraction 22 by means of immunoprecipitation using anti-HBs-coated magnetic beads. The media supernatant obtained from the culture of cells that had been subjected to CMV-HBV/A2/core clone monotransfection was also subjected to sucrose gradient ultracentrifugation using the same protocol. Sucrose high-density fraction 22, in which the HBcrAg concentration peaked, presumably contained 'naked' core particles or core protein (Fig. 6b). This fraction was collected and processed in the same manner via anti-HBs-coated magnetic bead separation and was used as negative control for this procedure (Fig. 6c, lane 4). To discriminate between HBV/G and HBV/A2 core proteins on Western blot analysis probed with anti-HBc antibody, we employed cell lysates produced from cells that had been transfected with the wild-type HBV/G clone and those produced with the HBV/A2 clone as controls. As can be seen on the Western blotting image (Fig. 6c), HBV/G-transfected cells (lane 1) produced larger proteins than the HBV/A2-transfected cells (lane 2), which can be explained by the 12-aa insertion in the core protein of HBV/G coded by its 36-nt unique insertion. Interestingly, the most saturated band associated with the Dane particles produced by HBV/G that had been co-transfected with CMV-HBV/A2/core (lane 3) was the same size as that for HBV/A2, suggesting that HBV/G competitively produces Dane particles consisting of HBV/A2 core protein during virion assembly.

DISCUSSION

HBV/G was first isolated in 2000 in France and the USA and was later found in Thailand, Japan and Mexico, indicating its global dissemination and association with specific risk groups, such as injection drug users (IDU) and men who had sex with men (MSM) [25]. Studies have also demonstrated that throughout the world, HBV/G strains possess unprecedented genetic homology and are mainly detected during co-infection with another genotype that is endemic in the area. Further studies have suggested that genotype G represents a 'replication-defective' variant of HBV that requires co-infection with another genotype to

establish a persistent infection. We and others have reported *in vitro* and *in vivo* experimental evidence of this HBV/G dependence [13–15]. The unique 36-nt insertion within core coding region increases core protein level and genome replication in genotype G but impairs replication, not core protein expression, in other genotypes [14]. These results strongly suggest the 36-nt/12-aa insertion has pleiotropic effects on core protein expression, genome replication and virion secretion [14]. To obtain clues about the mechanism by which genotype G works in combination with genotype A to effect its replication, we performed co-transfection experiments using Huh7 cells.

Using HBV/A2 viral proteins expressing plasmids, we determined that a HBV/A2 plasmid that selectively expressed core protein was capable of increasing the replication of the wild-type HBV/G (Fig. 3a). The replication of HBV/G during co-transfection was not affected by other viral elements of HBV/A2 because of the presence of the 'packaging-negative mutation' in the epsilon-coding region and stop codons preventing the translation of the other three viral proteins (the polymerase, surface and X proteins). The specific role of the core protein was further confirmed in experiments with CMV promoter-driven core expressing constructs, in which the core protein alone enhanced HBV/G replication in the absence of HBV pregenomic RNA. Interestingly, co-transfection of HBV/G with the CMV-HBV/A2/core expression construct produced the highest levels of intracellular DNA, even though this combination produced the lowest intracellular core protein level, compared with the CMV-core constructs of the other two genotypes (HBV/G and HBV/C) (Figs 4a,b). The replication of HBV/G was the highest during co-transfection with the CMV-HBV/A2/core expression construct, which agreed with the results of experiments using other genotype (HBV/D, HBV/B1) CMV-core constructs (data not shown). Thus, the core protein of HBV/A2 was confirmed to play an important role in upregulating HBV/G replication and performed this task more efficiently than the other genotypes. These experimental results might explain why HBV/A is the genotype that is most frequently found in co-infections with HBV/G [12,26].

Moreover, HBV/G core protein overexpression achieved by the co-transfection of HBV/G with CMV-HBV/G/core did not enhance replication, suggesting that HBV/G core protein is functionally defective; that is, results in insufficient viral packaging. To investigate the functional defect in the HBV/G core protein, we exchanged the core gene of the wild-type HBV/G for the corresponding gene of HBV/A2 (HBV/G/A2-Core); the introduction of the HBV/A2 core promoter together with core coding region into the HBV/G genome (HBV/G/A2-CP/core) significantly enhanced replication. However, the replication of the recombinant construct (HBV/A2 core coding region; HBV/G/A2-Core) did not differ from that of the wild-type HBV/G, suggesting that the replacement of HBV/G/A2-core alone was not