

- cancer cells. *Cancer Res* 2009;69:7160-7164.
12. Kaneda H, Arai T, Tanaka K, Tamura D, Aomatsu K, Kudo K, et al. FOXQ1 is overexpressed in colorectal cancer and enhances tumorigenicity and tumor growth. *Cancer Res* 2010;70:2053-2063.
13. Ormandy CJ, Musgrove EA, Hui R, Daly RJ, Sutherland RL. Cyclin D1, EMS1 and 11q13 amplification in breast cancer. *Breast Cancer Res Treat* 2003;78:323-335.
14. Takeda M, Arai T, Yokote H, Komatsu T, Yanagihara K, Sasaki H, et al. AZD2171 shows potent antitumor activity against gastric cancer over-expressing fibroblast growth factor receptor 2/keratinocyte growth factor receptor. *Clin Cancer Res* 2007;13:3051-3057.
15. Peters G, Brookes S, Smith R, Dickson C. Tumorigenesis by mouse mammary tumor virus: evidence for a common region for provirus integration in mammary tumors. *Cell* 1983;33:369-377.
16. Sakamoto H, Mori M, Taira M, Yoshida T, Matsukawa S, Shimizu K, et al. Transforming gene from human stomach cancers and a noncancerous portion of stomach mucosa. *Proc Natl Acad Sci U S A* 1986;83:3997-4001.
17. Takeo S, Arai H, Kusano N, Harada T, Furuya T, Kawauchi S, et al.

- Examination of oncogene amplification by genomic DNA microarray in hepatocellular carcinomas: comparison with comparative genomic hybridization analysis. *Cancer Genet Cytogenet* 2001;130:127-132.
18. Nishida N, Fukuda Y, Komeda T, Kita R, Sando T, Furukawa M, et al. Amplification and overexpression of the cyclin D1 gene in aggressive human hepatocellular carcinoma. *Cancer Res* 1994;54:3107-3110.
19. Chochi Y, Kawauchi S, Nakao M, Furuya T, Hashimoto K, Oga A, et al. A copy number gain of the 6p arm is linked with advanced hepatocellular carcinoma: an array-based comparative genomic hybridization study. *J Pathol* 2009;217:677-684.
20. Ikai I, Arii S, Ichida T, Okita K, Omata M, Kojiro M, et al. Report of the 16th follow-up survey of primary liver cancer. *Hepatol Res* 2005;32:163-172.
21. Uka K, Aikata H, Takaki S, Shirakawa H, Jeong SC, Yamashina K, et al. Clinical features and prognosis of patients with extrahepatic metastases from hepatocellular carcinoma. *World J Gastroenterol* 2007;13:414-420.

Table 1. Clinicopathological characteristics in responders of sorafenib

No.	Age (Y)	Sex	Viral status	AFP (ng/ml)	PIVKA (mAU/ml)	Stage	Primary Liver	Meta Lung	Meta Others	Histological type	Combination Treatment	Treatment Response	FGF3/FGF4 Amp
1	52	M	B	198	140	IV	2cm, x3	multi	Ad. grand	mod	(-)	PR	(-)
2	63	M	B	24	1983	III	6 cm	(-)	(-)	mod	(-)	CR	(-)
3	58	M	C	16	14	III	9cm, multi	(-)	(-)	well	(-)	PR	(-)
4	62	M	B	8	130	IV	(-)	x3	(-)	mod-poor	(-)	PR	(-)
5	47	F	C	1872	728	IV	2cm, multi	multi	(-)	poor	+ TAI	CR	(-)
6	66	M	C	290	18507*	IV	5cm	(-)	(-)	mod	(-)	CR	(-)
7	71	M	C	404100	1328	IV	5cm, multi	multi	(-)	poor	(-)	CR	(-)
8	66	M	non	49	7173	IV	(-)	x2	Pleu, LN	mod	(-)	PR	Amp
9	58	F	B	715	101	IV	11cm	multi	(-)	comb.	+ 5FU/IFN	PR	Amp
10	80	F	C	378	21	III	3cm, x3	(-)	(-)	poor, mod**	(-)	CR	Amp
11	57	M	C	46835	2730	IV	14cm, multi	multi	(-)	mod	(-)	CR	n.d.
12	77	M	B	435	71000	IV	4cm, multi	(-)	(-)	mod	(-)	PR	n.d.
13	84	M	non	5410	847000*	IV	13cm, multi	(-)	(-)	poor	(-)	PR	n.d.

non, nonB-nonC; Primary Liver, HCC in the liver; Meta Lung, lung metastasis; Meta Others, other metastatic sites; Comb.,

HCC with cholangiocarcinoma component; n.d., not done; IFN, interferon; Amp, gene amplification. *, warfarin treatment

(+), ** from two different HCC nodules.

Table 2. Clinicopathological characteristics and *FGF3/FGF4* gene amplification in responders and non-responders to sorafenib

		Responders n=13	Non-responders n=42	<i>p</i> value
Age (years)	Median	63	66	0.98
	Range	47 - 84	22 - 89	
Sex	M	10	30	0.97
	F	3	12	
Viral status	HBV	5	10	0.69
	HCV	6	16	
	B+C	0	1	
	non	2	15	
AFP (ng/ml)	Median	378	56	0.33
	Range	8 - 404100	2 - 114248	
PIVKA-II (mAU/ml)	Median	728	81	0.78
	Range	14 - 847000	11 - 147000	
Clinical Stage	II	0	1	0.73
	III	3	13	
	IV	10	28	
Primary tumor (cm)	Median	5	3	0.20
	Range	0 - 14	0 - 15	
Lung meta	(-)	6	31	0.13
	(+)	7	11	
Multiple Lung meta	5 <	8	40	0.006
	5 ≥	5	2	
Other meta	(-)	11	26	0.24
	(+)	2	16	
Histological type	Well	1	7	0.13
	Moderate	6	26	
	Poor	5	6	
	Combination	1	3	
Response	CR	6	-	N.D.
	PR	7	-	
	SD	-	16	
	PD	-	24	
	N.E	-	2	

non, nonB-nonC; Lung meta, lung metastasis; Other meta, other metastatic sites; Combination, HCC with cholangiocarcinoma component; n.d., not done. *p*

Accepted Article

values of viral status and histological type were calculated between HBV vs.

HVC and poorly differentiated vs. non-poorly differentiated.

Figure legends

Fig. 1.

A hepatocellular carcinoma that exhibited a marked response to sorafenib treatment harbors *FGF3/FGF4* gene amplification. (A) Abdominal CT images obtained upon pre-treatment (left panel) and two months after treatment (right panel). (B) Comparative genomic hybridization analysis of the HCC tumor. Paired background liver tissue was used as a reference sample. A gain of genomic copy number (> 4 copies, red) and a loss (< 0.5 copies, blue) were shown by the indicated colors. (C) Whole copy numbers of chromosome 11 are shown. A highly amplified region is described in the lower panel. (D) Western blot analysis of FGF3 (Arrow) in HCC and paired background liver samples. IB, immunoblotting.

Liver, paired background liver.

Fig. 2.

FGF3/FGF4 gene amplification is frequently observed in responders to sorafenib in HCC. (A) *FGF3/FGF4* gene amplification was determined using the TaqMan copy number assay in DNA samples obtained from 48 HCC samples that had been treated with sorafenib. *FGF3* amplification of over 5 copies was observed in

3 of the responders to sorafenib. *, CR+PR vs. SD+PD. (B) *FGF3/FGF4* gene amplification mediates the overexpression of *FGF3/FGF4* mRNA. The mRNA expression levels of *FGF3* and *FGF4* were examined in nine HCC samples that were available as frozen samples among forty eight HCC samples that were treated with sorafenib. Rel. mRNA, *target gene/GAPD* x10⁶.

Fig. 3.

Fluorescence *in situ* hybridization analysis of *FGF3*-amplified HCC. Green, *CEN11P* locus; Red, *FGF3* locus; No., sample numbers; Amp, gene amplification. High-power images are presented for a single cancer cell.


Fig. 4.

FGF3/FGF4 gene amplification in a series of HCC samples without sorafenib treatment. (A) A TaqMan copy number assay for *FGF3* and *FGF4* was used to examine DNA samples obtained from 82 surgical specimens. Human normal genomic DNA was used as a normal control. Well, well-differentiated HCC; Mod, moderately differentiated HCC; Poor, poorly differentiated HCC.

Fig. 5.

FGF3 and FGF4 overexpression and drug sensitivity to sorafenib *in vitro* and *in vivo*. (A) Growth inhibitory assay examining sorafenib in various cancer cell lines *in vitro*. The growth inhibitory effect of sorafenib was examined using an MTT assay. The IC₅₀ values of each cell line are shown in the graph. The black bars show that the IC₅₀ values were below 1 μM. Amp, gene amplification. (B) Cancer cell lines stably overexpressing *EGFP*, *FGF3* or *FGF4* were established and designated as A549/EGFP, A549/FGF3 and A549/FGF4. Western blotting confirmed that exogenously expressed FGF3 and FGF4 were secreted into the culture medium. IB, Immunoblotting; Sup., supernatant. (C) The 3T3 cells were exposed to indicated concentrations of sorafenib for 2 hours and were then stimulated with FGF4 conditioned medium for 20 minutes. (D) Mice inoculated with A549/EGFP, A549/FGF3 or A549/FGF4 (n=20 each) were treated with a low dose of sorafenib (n=10, 15 mg/kg/day, p.o.) or without (n=10, vehicle control, p.o.). **p*<0.05.


Fig. 1.



99x75mm (300 x 300 DPI)

Accept


Fig. 2.



99x75mm (300 x 300 DPI)

Accept


Fig. 3.



99x75mm (300 x 300 DPI)

Accept

Fig. 4.



99x75mm (300 x 300 DPI)

Accept

Characterization of naturally occurring protease inhibitor-resistance mutations in genotype 1b hepatitis C virus patients

Hiroko Shindo · Shinya Maekawa · Kazuki Komase · Ryota Sueki · Mika Miura · Makoto Kadokura · Kuniaki Shindo · Fumitake Amemiya · Takatoshi Kitamura · Yasuhiro Nakayama · Taisuke Inoue · Minoru Sakamoto · Shun-ichi Okada · Yasuhiro Asahina · Namiki Izumi · Masao Honda · Shuichi Kaneko · Nobuyuki Enomoto

Received: 21 April 2011 / Accepted: 22 July 2011 / Published online: 18 August 2011
© Asian Pacific Association for the Study of the Liver 2011

Abstract

Background and aims Protease inhibitor (PI)-resistant hepatitis C virus (HCV) variants may be present in substantial numbers in PI-untreated patients according to recent reports. However, influence of these viruses in the clinical course of chronic hepatitis C has not been well characterized.

Methods The dominant HCV nonstructural 3 (NS3) amino acid sequences were determined in 261 HCV genotype 1b-infected Japanese patients before pegylated interferon plus ribavirin (PEG-IFN/RBV) therapy, and investigated the patients' clinical characteristics as well as treatment responses including sustained virological response (SVR) rate. HCV-NS3 sequences were also determined in 39 non-SVR patients after completion of the therapy.

Results Four single mutations (T54S, Q80K, I153V, and D168E) known to confer PI resistance were found in 35 of 261 patients (13.4%), and double mutations (I153V plus

T54S/D168E) were found in 6 patients (2.3%). Responses to PEG-IFN/RBV therapy did not differ between patients with and without PI-resistance mutations (mutation group, SVR 48%; wild-type group, SVR 40%; $P = 0.38$). On the other hand, two mutations appeared in two non-SVR patients after PEG-IFN/RBV therapy (I153V and E168D, 5.1%).

Conclusions PI-resistance-associated NS3 mutations exist in a substantial proportion of untreated HCV-1b-infected patients. The impact of these mutations in the treatment of PIs is unclear, but clinicians should pay attention to avoid further development of PI resistance.

Keywords HCV · Protease inhibitor · Naturally occurring viral resistance mutations

Introduction

Hepatitis C virus (HCV) infects more than 170 million persons worldwide and thus represents a global health problem. At least 130 million infected individuals are chronic carriers of HCV and are at significant risk of developing liver cirrhosis and hepatocellular carcinoma [1]. The current standard treatment with pegylated interferon plus ribavirin (PEG-IFN/RBV) is complicated by frequent adverse reactions, and a sustained virologic response (SVR) can be achieved only in 50% of patients infected with the most prevalent genotype 1 [2]. In Japan, since 70% of patients are infected with intractable genotype 1b HCV, more effective treatments are urgently required.

A promising approach is the development of specifically targeted antiviral therapies for hepatitis C (STAT-C). HCV-specific protease inhibitors (PIs) target an essential step in HCV replication by blocking the nonstructural 3/4A (NS3/4A) protease-dependent cleavage of the HCV polyprotein

Electronic supplementary material The online version of this article (doi:10.1007/s12072-011-9306-7) contains supplementary material, which is available to authorized users.

H. Shindo · S. Maekawa (✉) · K. Komase · R. Sueki · M. Miura · M. Kadokura · K. Shindo · F. Amemiya · T. Kitamura · Y. Nakayama · T. Inoue · M. Sakamoto · S. Okada · N. Enomoto

First Department of Internal Medicine, University of Yamanashi, 1110, Shimokato, Chuo, Yamanashi 409-3898, Japan
e-mail: maekawa@yamanashi.ac.jp

Y. Asahina · N. Izumi
Division of Gastroenterology and Hepatology,
Musashino Red Cross Hospital, Tokyo, Japan

M. Honda · S. Kaneko
Department of Gastroenterology, Kanazawa University Graduate
School of Medicine, Kanazawa, Japan

[1]. Among these NS3/4A PIs, telaprevir, boceprevir, SCH446211, danoprevir (ITMN-191), naldaprevir (SCH900518), and TMC435 are now under clinical trials [1, 3–7]. In PROVE1 and PROVE2 studies [3, 4] undertaken in North America and Europe, the SVR rate was favorable (67 and 69%, respectively) in a triple therapy regimen including telaprevir. In addition, some studies have suggested that shortening of treatment duration may be possible for patients who achieve a rapid virologic response (RVR) [8, 9].

However the sole use of STAT-C drugs, such as PIs, promotes production and selection of drug-resistant variants in patients experiencing viral rebound during treatment [3, 10, 11] as well as in HCV replicon experiments [11, 12]. Therefore, these drugs should be used in combination with the PEG-IFN/RBV to prevent the appearance of drug-resistant variants. However, Kuntzen et al. [13] demonstrated the presence of these drug-resistant variants in high frequencies (8.6–16.2%) by population-based sequencing in patients not treated with the drugs [1, 13]. Gaudieri et al. [14] have suggested that regions of NS3 protease and NS5B polymerase are likely to be under HLA immune pressure and therapeutic selection, and that drug-resistant variants may occur naturally to escape the immune system. These observations seem quite astonishing and troubling, since a substantial number of patients may not respond to the new therapies such as STAT-C drugs.

In the present study, to assess the prevalence of NS3 mutations conferring PI resistance in HCV genotype 1b-infected Japanese patients who had not been previously treated with PIs, as well as to assess the influence of those mutations in response to PEG-IFN/RBV therapy, the dominant HCV-NS3 sequences were determined in 261 HCV-1b patients before starting the PEG-IFN/RBV therapy.

Methods

Patients

Serum samples were acquired from 261 HCV genotype 1b-infected adult Japanese patients before combination therapy with PEG-IFN (PEGINTRON[®], Schering-Plough, Tokyo, Japan) plus RBV (REBETOL[®], Schering-Plough) between 2004 and 2008 at the University of Yamanashi, Musashino Red Cross Hospital and Kanazawa University. The therapy was administered according to the standard PEG-IFN/RBV treatment protocol established for Japanese patients by a hepatitis study group of the Ministry of Health, Labor, and Welfare, Japan. Specifically, the patients were subcutaneously administered PEG-IFN α -2b, 1.5 μ g/kg body weight, once weekly and RBV 600–800 mg daily per os for 48 weeks. These patients were not infected with human immunodeficiency virus (HIV). The study was

approved by the ethics committees of all participating universities and the hospital, and the protocol conformed to the ethical guidelines of the 1975 Declaration of Helsinki as reflected in a priori approval by the Institutional Review Board at Massachusetts General Hospital. Written informed consent was obtained from each study participant.

Amplification and sequencing of full-length HCV genomes

Viral loads were determined using the Amplicor HCV RNA kit, version 2.0 (Roche Diagnostics, Tokyo, Japan) or the Cobas TaqMan test (Roche Diagnostics). HCV RNA was extracted from pretreatment serum samples by the AGPC method using Isogen (Wako, Osaka, Japan) according to the manufacturer's protocol. Complementary DNA was synthesised using Superscript II (Invitrogen, Tokyo, Japan) and random primers (Invitrogen), and then amplified by two-step nested PCR using the primers listed in Supplementary Table 1. All samples were initially denatured at 95°C for 7 min, followed by 40 cycles of amplification with denaturation at 95°C for 15 s, annealing at 55°C for 15 s, and extension at 72°C for 45 s using the BD Advantage[™] 2 PCR Enzyme system (BD Biosciences Clontech, CA, USA). PCR amplicons were directly sequenced using BigDye Terminator version 3.1 (ABI, Tokyo, Japan) and universal M13 forward/reverse primers using an ABI prism 3130 sequencer (ABI).

Sequence alignment and analysis

Sequences were determined in both directions, particularly for the ambiguous stretches, were assembled using the Vector NTI software (Invitrogen), and base-calling errors were corrected following the inspection of chromatograms. If mixed bases were detected as two different chromatogram peaks at the same residue, only the dominant base was called after evaluation of all overlapping fragments. A consensus sequence was generated from the alignment on the basis of the most common amino acid at each site.

Determination of PI resistance mutations

Multiple viral NS3 mutations were observed in amino acid positions reported to confer PI resistance among 261 patients: V36, Q41, F43, T54, V55, Q80, R109, I153, R155, A156, D168, V170, and M175. NS3 amino acid mutations with proven PI resistance in previously published studies (Table 1) were designated as resistance proven mutations (e.g., V36M/A). Mutations in the PI-resistance site not known to confer drug resistance were designated resistance unproven mutations (e.g., V36I). Patients were allocated to two groups according to the presence of PI-resistance

mutations (including resistance unproven mutations), and clinical characteristics including HCV RNA levels and responses to PEG-IFN/RBV therapy were compared. To assess the influence of PEG-IFN/RBV therapy on NS3 mutational status, posttreatment HCV-NS3 sequences in 39 of 58 non-SVR patients were also examined.

Statistical analysis

Statistical differences in the data, including all available patients' demographic, biochemic, hematologic, and virologic data such as sequence variation factors, were determined among the various groups by Student's *t* test or Mann–Whitney *U* test for numerical variables and Fisher's exact probability test for categorical variables.

Results

Prevalence of dominant PI-resistance-associated nonstructural 3 mutations in untreated patients

Figure 1 shows the frequency of substitutions in 261 patients for each of 181 NS3 protease amino acid residues

compared to the consensus sequence. A total of 41 resistance proven mutations were detected in 35 (13.4%) patients: T54S (14 patients, 5.4%), Q80K (1 patient, 0.4%), I153V (22 patients, 8.4%), D168E (4 patients, 1.5%), T54S plus I153V double mutation (4 patients, 1.5%), and I153V plus D168E double mutation (2 patients, 0.8%). The mutation number increased to 54 in 47 (18.0%) patients when resistance unproven mutations were included: V36I (2 patients, 0.8%), I153L (11 patients, 4.2%), and I153V plus V36I double mutation (2 patients, 1.5%). Double mutations were found in 7 patients (2.7%) (Table 1). Q80L was observed in 47 (18%) patients but these were excluded from consideration because a previous study demonstrated that this mutation does not confer resistance [15]. All mutations observed in this study would confer low- to moderate-level PI resistance according to previous studies [6, 15–19]. No mutations conferring high-level resistance such as R155 or A156 [11, 17, 19–22] were observed.

Clinical characteristics of patients with PI-resistance mutations

Table 2 presents the characteristics of patients classified according to the presence of PI-resistance mutations

Table 1 Prevalence of PI-resistance-associated NS3 mutations

Drug-resistance mutations described in the literature				References	Detected resistance mutations Genotype 1b (<i>N</i> = 261), (%)
NS3 residue	Resistance mutations	Drugs			
V36	A, M, L, G, C	Telaprevir, Boceprevir	[1, 3, 4, 10, 11, 19, 31, 37]	I × 2 (0.8)	
Q41	R	ITMN-191, Boceprevir	[19]		
F43	S, C	ITMN-191, Boceprevir, Telaprevir, TMC435	[15, 19]	S × 14 (5.4)	
T54	A, S	Telaprevir, Boceprevir, SCH900518	[1, 3, 10, 11, 19, 20, 31, 38]		
V55	A	Boceprevir	[1]	K × 1 (0.4)	
Q80	R, K	TMC435	[6, 15]		
R109	K	SCH446211	[17]	V × 22 (8.4), L × 11 (4.2)	
I153	V	SCH446211	[17]		
R155	K, T, I, M, G, L, S, Q	Telaprevir, Boceprevir, ITMN-191, BILN2061, TMC435	[1, 3, 4, 6, 10, 11, 15, 19, 20]	E × 4 (1.5)	
A156	S, T, V, I, G	Telaprevir, Boceprevir, ITMN-191, BILN2061, SCH446211, TMC435, SCH900518	[1, 3, 4, 10, 11, 15, 17, 19, 20, 38]		
D168	A, V, E, N, T, H	BILN2061, ITMN-191, TMC435	[6, 15, 20]	E × 4 (1.5)	
V170	A	Telaprevir, Boceprevir	[1, 19, 20]		
M175	L	Boceprevir	[39]		
Total number (%) of patients with resistance proven mutations				35 (13.4)	
Total number (%) of patients with resistance proven and unproven mutations				47 (18.0)	

Amino acid mutations conferring PI resistance in the literatures and those observed in PI-treatment-naïve patients in this study are indicated. Bold indicates resistance proven mutations, and the others indicate resistance unproven mutations

Double mutations found were as follows: V36I and I153V × 1, T54S and I153V × 4, I153V and D168E × 2

(including resistance unproven mutations). Age, sex ratio, body mass index, alanine aminotransferase (ALT) levels, serum albumin, platelet count, and fibrosis stage did not differ between the NS3 mutation and wild-type groups. No significant difference was observed between the two groups in the parameters of PEG-IFN/RBV treatment response, HCV sequence variations in interferon sensitivity determining region (ISDR), Core 70, interferon plus ribavirin resistance-determining region (IRRDR), or interleukin 28B (IL28B) single nucleotide polymorphism (SNP) (rs8099917; T/G and G/G vs. T/T) [23–30]. These clinical variables were also compared between the mutation group defined as resistance proven mutations and the wild-type group, but no notable differences were observed.

Unimpaired in vivo fitness of viral strains with resistance mutations

Because most PI-resistance mutations described till date have been associated with reduced replicative capacity of varying degrees [1, 10, 11, 13, 17, 20–22, 31, 32], we examined viral replication levels in patients with drug-resistance mutations (Fig. 2). The estimated *P* value indicated no significant difference between the mutation (median 1,500 KIU/ml) and wild-type (median 1,800 KIU/ml) groups (*P* = 0.69). The results indicate that drug-resistant HCVs were not necessarily impaired in their ability to replicate in vivo. However, patients with double mutations (*N* = 7) tended to have low viral loads (median 1,200 KIU/ml) (*P* = 0.09).

Resistance mutations and virologic response to PEG-IFN/RBV therapy

To determine the difference in virologic response to PEG-IFN/RBV therapy according to the PI mutation, frequency of HCV RNA levels below detection at 4 weeks (rapid viral response, RVR) and 12 weeks (complete early viral response, cEVR), and SVR rate (%) were investigated in

each group. The frequency of HCV RNA levels below detection at 4 and 12 weeks was 14 and 50%, respectively, in the mutation group, and was 11 and 46%, respectively, in the wild-type group. The SVR rate was 48 and 40% in the mutation and wild-type groups, respectively (*P* = 0.38). No significant difference was observed between the two groups in any of the indexes investigated (Table 2). The time-dependent viral clearance rate during PEG-IFN/RBV therapy was estimated in 133 patients including 25 patients (19%) with PI-resistance mutations available for the analysis. Kaplan–Meier analysis demonstrated that HCV clearance did not differ between the two groups with and without resistance mutations (log-rank test, *P* = 0.30) (Fig. 3).

Changes in nonstructural 3 amino acid sequence diversity during PEG-IFN/RBV therapy

Full-length NS3 protease sequences were determined in 39 non-SVR patients after PEG-IFN/RBV therapy. A single amino acid change at resistance-associated sites in two patients was observed. In one patient, isoleucine (Ile) at position 153 changed to valine (Val), and glutamic acid (Glu) changed to aspartic acid (Asp) at position 168 in the second (Fig. 4). At the nucleotide level, ATC (Ile) changed to GTC (Val) in I153V, and GAA (Glu) changed to GAC (Asp) in E168D. Both mutations were caused by one nucleotide exchange. No other changes were observed in the other 37 patients.

Discussion

Here we report that in 18% (47/261) HCV genotype 1b-infected patients who had not been previously treated with NS3 PIs, the viral genome contained dominant amino acid mutations within the NS3 PI-resistance sites. Even after confining the data to established PI-resistance mutations, the mutation rate was still significant in 13.4% (35/261). No clinical differences were observed between patients




Fig. 1 Frequency of polymorphic mutations for each of the 181 NS3 protease amino acid residues in 261 patients. Arrowheads indicate the sites reported to confer PI resistance. Dark bars denote the amino acid

variations at the resistant sites in this study. *80, we detected one resistant mutation (Q80K) and 47 (18%) non-resistant variations (Q80L) at the 80th residue

Table 2 Characteristics of patients with or without HCV genomes harboring drug-resistance mutations

Characteristics	Mutation type (<i>N</i> = 47)	Wild-type (<i>N</i> = 214)	<i>P</i> value
Patients' characteristics			
Age, median (range)	59 (46–72)	57 (19–77)	0.17
Male, no. (%)	26 (55)	112 (52)	0.70
BMI, median (range)	23.2 (15.5–31.9)	22.8 (16.1–31.9)	0.41
ALT IU/ml	81.3 ± 72.6 ^a	74.8 ± 51.9	0.93
Serum albumin g/dl	4.00 ± 0.37	4.01 ± 0.36	0.81
Platelet count × 10 ⁴ /μl	15.8 ± 4.3	14.5 ± 4.8	0.18
HCV RNA KIU/ml, median (range)	1,500 (58–6,310)	1800 (28–15,849)	0.69
Fibrosis, no. (%)			0.97
F0	0 (0)	7 (3)	
F1	23 (50)	89 (42)	
F2	9 (20)	52 (24)	
F3	9 (20)	40 (19)	
F4	5 (11)	26 (12)	
IFN pre-treatment no. (%)	15/40 (38) ^b	66/172 (38)	1.00
IL28B (rs8099917) T/G or G/G no. (%)	6/20 (30)	19/67 (28)	1.00
Response to PEG-IFN/RBV therapy			
SVR total cases no. (%)	22/46 (48)	83/210 (40)	0.38
RVR in total cases no. (%)	6/44 (14)	22/195 (11)	0.83
cEVR in total cases no. (%)	22/44 (50)	92/200 (46)	0.75
SVR 48w treatment no. (%)	16/29 (55)	55/130 (42)	0.29
End of treatment response no. (%)	26/41 (63)	123/202 (61)	0.91
HCV genome sequence variation			
ISDR mutation ≤1 no. (%)	32/46 (70)	167/210 (80)	0.21
Core70 R no. (%)	26/44 (59)	136/210 (65)	0.56
IRRDR mutation >3 no. (%)	25/38 (66)	107/190 (56)	0.34

^a Mean ± SD^b Number/total number (%)

harboring viruses with and without these mutations. Moreover, no differences were observed in the responses of either group to PEG-IFN/RBV therapy.

Recent studies reported that significant number of patients who were never treated with PI possess viral sequences with PI-resistance-associated NS3 mutations. In these studies, the prevalence of PI-resistance mutations was determined to be 8.6–16.2% [13, 14], in HCV genotype 1- and 3-infected patients in European–American populations. These patients were often coinfecting with HIV. Analysis of the public HCV databases (EuHCVdb and Los Alamos) also reported the presence of naturally occurring PI-resistance-associated NS3 mutations in worldwide isolates [33]. However, *in vivo* and *in vitro* studies demonstrated that most of the mutations observed conferred only low- to moderate-level PI resistance [7, 13, 14, 34, 35]. Regarding viral fitness, PI-resistant HCVs show lower fitness at varying degrees as revealed by *in vitro* studies [1, 10, 11, 17, 20–22, 31, 32], but HCV RNA levels in a clinical study did not differ significantly. The response to PEG-IFN/RBV therapy was almost comparable to that in HCV-infected patients without PI-resistance mutations either in HCV replicon experiments or in a clinical study of small number of treated patients [34].

The prevalence of 13.4% for PI-resistance-proven patients observed in the present study was almost comparable to the results of previous studies. Although HIV is known to increase HCV replication in coinfection with HCV [36], and HIV patients are often treated with the HIV-specific PIs, the HIV infection might not affect the natural occurrence of HCV-specific PI-resistance mutations since our studied patients were all proven to be free from coinfection with HIV infection. As shown in Table 1 and Fig. 1, I153 V (22/261, 8.4%), T54S (14/261, 5.4%), and D168E (4/261, 1.5%) were among the most prevalent PI-resistance-proven mutations in the present study. The most frequent mutation detected in our study I153V was reported to appear secondarily to the occurrence of R109K mutations in a HCV replicon system [17]. Although the role of this mutation is not understood, the I153V mutation on its own conferred SCH446211 resistance to the HCV replicon to a lesser degree [17]. Interestingly, I153V was often found in double mutations in our study, as shown in Fig. 2. This suggests analogy between *in vitro* and *in vivo* data. T54S and D168E, the other frequent mutations, have been also reported to occur as single dominant mutations in previous *in vitro* or *in vivo* studies in HCV genotype 1

Fig. 2 In vivo fitness of HCV with PI-resistance-associated NS3 mutations. HCV RNA levels were compared between patients with and without NS3 PI-resistance-associated mutations (a) and between patients with each resistance mutation (b). The estimated *P* value (Mann–Whitney *U* test) indicates no significant difference between the wild-type and other groups (wild-type vs. mutation type, wild-type vs. single mutation type, and wild-type vs. double mutation type). (Wild-type, *N* = 214; mutation type, *N* = 47; single mutation type, *N* = 40; double mutation type, *N* = 7; V36I, *N* = 2; T54S, *N* = 14; Q80K, *N* = 1; I153L, *N* = 11; I153V, *N* = 22; D168E, *N* = 4; E176A, *N* = 1; V36I + I153V, *N* = 1; T54S + I153V, *N* = 4, and I153V + D168E, *N* = 2)





Fig. 3 Comparison of virologic response to PEG-IFN/RBV therapy between HCV-infected patients with and without PI-resistance-associated NS3 mutations. Time-dependent HCV clearance rate analysis was based on serum HCV RNA positivity during PEG-IFN/RBV therapy for HCV isolates with resistance mutations or wild-


type sequences. A total of 133 patients for whom the limit of viral genome detection could be determined were analyzed. Among this group, NS3 mutations were detected in 25 patients (19%). The estimated *P* value (log-rank test) shows no significant difference between the two groups (*P* = 0.30)

infections showing moderate degrees of resistance [16, 18, 19].

Most PI-resistance mutations described to date have been associated with varying degrees of reduced replicative

capacity [10, 11, 17, 20–22, 31, 32]. In the present study, HCV RNA levels of those patients with low- to moderate-level resistance mutations were similar to those in patients in the wild-type groups, suggesting that in vitro viral fitness

Fig. 4 Appearance of PI-resistance-associated NS3 mutations during the PEG-IFN/RBV therapy. Chromatograms show part of the HCV NS3 sequence demonstrating PI-resistance mutations in two patients receiving therapy. **a** Site 153 isoleucine (Ile) (ATC) changed to valine (Val) (GTC), **b** Site 168 glutamic acid (Glu) (GAA) changed to aspartic acid (Asp) (GAC)



does not necessarily reflect in vivo viral fitness. This, however, does not rule out the possibility that some unknown compensatory viral mutations might have resulted in upregulation of reduced viral fitness. Interestingly, although the replicative capacity conferred by a single mutation seemed to be the same, the HCV RNA levels of double mutations were frequently low, suggesting that double mutations might weaken viral fitness.

In previous studies, clinical characteristics representing the state of liver disease other than HCV RNA levels were not studied in patients with PI-resistance mutations. In this study, we show that those clinical characteristics did not differ according to the presence of viral NS3 mutations. As shown in Table 2, age, sex ratio, fibrosis stage, ALT levels, serum albumin, platelet count, and past history of IFN pretreatment did not differ according to the presence of NS3 mutations. These results suggest that NS3 mutations occur independently of disease progression. Moreover, no evident differences were observed between viral and host factors known to affect IFN-based treatment responses. However, viral amino acid variations in the core and NS5A or the allelic frequency of IL28B SNPs, which were recently reported for the close relationship of responses to PEG-IFN/RBV therapy, did not differ between the two groups.

A significant outcome of the present study is the demonstration that PI-resistance mutations might not affect responses to PEG-IFN/RBV therapy. Previous in vitro studies demonstrated that HCV replicons harboring PI-resistance mutations were also sensitive to IFN treatment [31]. In addition, recent clinical studies also indicated that PI-resistance mutations were sensitive to the PEG-IFN/RBV [10, 34]. However, our analysis was more comprehensive because viral and host factors that contribute to treatment responses were simultaneously analyzed. A unique aspect of the present study is that we investigated the influence of the PEG-IFN/RBV treatment on the occurrence of new PI mutations by direct nucleotide sequencing, and were able to show that the PEG-IFN/RBV might not induce amino acid mutations.

Will the pre-existence of naturally occurring PI-resistance mutations have an influence on future treatment of HCV infections? Since new PIs are on the verge of clinical use, all clinicians should bear in mind the substantial numbers of HCV-infected patients with PI-resistance mutations. Although the degree of resistance is considered to be low or moderate in untreated patients, weak resistance might progress to more potent resistance with additional mutations, when PIs become widely used. Therefore, all clinicians need to be sufficiently prepared for the possibility of later onset of PI-resistance mutations that confer greater drug resistance and concomitant poorer responses to therapy. In SPRINT-1 study, the lead-in therapy was associated with a modestly lower rate of breakthrough than with no lead in [7]. Considering that PEG-IFN/RBV was equally effective for PI-resistant viruses, sufficient “lead-in” therapy before the administration of PIs could be an option in the forthcoming triple therapy modality.

In conclusion, we demonstrate here that PI-resistance-associated NS3 mutations exist in a substantial proportion of untreated HCV-1b-infected patients. Although the degree of resistance might not be strong, clinicians will need to consider this upon the introduction of triple therapy.

Acknowledgements This work was supported in part by a grant-in-aid scientific research fund from the Ministry of Education, Science, Sports, and Culture [grant number 20390206] and in part by a grant-in-aid from the Ministry of Health, Labor, and Welfare of Japan [grant number H19-kanen-002].

References

1. Susser S, Welsch C, Wang Y, et al. Characterization of resistance to the protease inhibitor boceprevir in hepatitis C virus-infected patients. *Hepatology* 2009;50:1709–1718
2. Hadziyannis SJ, Sette H Jr, Morgan TR, et al. Peginterferon-alpha2a and ribavirin combination therapy in chronic hepatitis C: a randomized study of treatment duration and ribavirin dose. *Ann Intern Med* 2004;140:346–355