

Five of our seven proposed hypotheses regarding construct validity could be confirmed, which indicates the satisfactory validity of the Japanese questionnaire [27]. Only the correlations between KOOS ADL and sport/recreation with the SF-36 role-emotional component were higher than 0.50, and the correlations of SF-36 role-physical versus KOOS pain and KOOS symptoms subscales were lower than expected. A remarkably weaker correlation between the latter subscore comparison was also reported for the Persian (0.38 for pain and 0.15 for symptoms) and the Singapore English and Chinese (for pain, 0.23 each and for symptoms 0.21 and 0.18, respectively) versions. These adapted questionnaires had generally lower correlations between the KOOS subscales and all three physical domains of the SF-36.

Some limitations of our study include the small sample size of 58 patients, which may be considered too low for testing reliability and validity. Nonetheless, this number lies within the quality criteria outlined by Terwee et al. [27] that proposes a minimum requirement of at least 50 patients for satisfactorily evaluating health status measures. Although it is difficult to prove, the two different methods in which the second questionnaire was distributed in this study could have some influence on the reliability of this

patient-rated survey. Lastly, further research is needed to assess the responsiveness of the Japanese KOOS. As the presented work is part of a recently initiated 5-year study focusing on patients with OA, this psychometric aspect will be evaluated in the context of this larger project.

The currently validated Japanese KOOS is a reliable and stable outcomes measure that provides a valuable basis for all clinical studies looking specifically at patient opinion concerning their knee OA. It also provides a significant contribution to the ever-increasing trend of internationally used patient self-assessed scores in Asian languages.

Acknowledgments The authors would like to acknowledge Toshi-nao Sawai and Naohiko Banno from Synthes K.K. for their administrative support, and Dr. Laurent Audigé from AO Clinical Investigation and Documentation for his statistical advice.

Conflict of interest The work of the study was supported by the AO Foundation and Synthes Japan. The authors did not receive any benefits from any commercial party related directly or indirectly to the subject of this article.

Appendix

See Table 3

Table 3 Reliability data for the individual 42 items of the Japanese Knee Injury and Osteoarthritis Outcome Score (KOOS) questionnaire

Item number ^a	Test (T1)				Retest (T2)				ICC	95% CI (ICC)
	Min	Max	Mean	SD	Min	Max	Mean	SD		
1 (S1)	0 (47%)	4 (12%)	1.05	1.34	0	4	1.22	1.28	0.73	0.58–0.83
2 (S2)	0 (38%)	4 (9%)	1.14	1.22	0	4	1.07	1.14	0.70	0.54–0.81
3 (S3)	0 (60%)	4 (2%)	0.53	0.80	0	3	0.55	0.78	0.51	0.29–0.68
4 (S4)	0 (29%)	4 (9%)	1.19	1.16	0	4	1.03	1.15	0.54	0.34–0.70
5 (S5)	0 (22%)	4 (12%)	1.66	1.33	0	4	1.83	1.27	0.78	0.65–0.86
6 (S6)	0 (26%)	3	1.21	0.99	0	4	1.19	0.98	0.71	0.55–0.81
7 (S7)	0 (12%)	4 (2%)	1.24	0.82	0	3	1.28	0.85	0.61	0.41–0.75
8 (P1)	0 (10%)	4 (31%)	2.40	1.38	0	4	2.29	1.28	0.72	0.57–0.82
9 (P2)	0 (14%)	4 (9%)	1.60	1.14	0	4	1.61	1.08	0.84	0.74–0.90
10 (P3)	0 (34%)	4 (9%)	1.12	1.20	0	4	1.02	1.18	0.90	0.84–0.94
11 (P4)	0 (24%)	4 (9%)	1.66	1.30	0	4	1.71	1.38	0.77	0.65–0.86
12 (P5)	0 (42%)	3	0.90	0.94	0	3	0.91	0.90	0.69	0.52–0.80
13 (P6)	0 (17%)	4 (7%)	1.60	1.11	0	4	1.53	1.10	0.67	0.52–0.80
14 (P7)	0 (43%)	3	0.76	0.80	0	3	0.86	0.89	0.83	0.73–0.90
15 (P8)	0 (34%)	4 (3%)	0.95	0.95	0	3	0.95	0.83	0.45	0.21–0.63
16 (P9)	0 (40%)	4 (2%)	0.86	0.89	0	3	0.76	0.84	0.77	0.64–0.86
17 (A1)	0 (22%)	4 (7%)	1.62	1.24	0	4	1.53	1.14	0.78	0.65–0.86
18 (A2)	0 (21%)	4 (7%)	1.45	1.17	0	4	1.34	1.12	0.72	0.57–0.83
19 (A3)	0 (16%)	4 (2%)	1.41	0.96	0	4	1.37	0.98	0.72	0.56–0.82
20 (A4)	0 (35%)	3	0.88	0.84	0	3	0.86	0.98	0.74	0.60–0.84
21 (A5)	0 (29%)	4 (3%)	1.19	1.08	0	4	1.21	0.99	0.72	0.57–0.82
22 (A6)	0 (48%)	4 (3%)	0.81	0.98	0	3	0.71	0.88	0.72	0.57–0.83

Table 3 continued

Item number ^a	Test (T1)				Retest (T2)				ICC	95% CI (ICC)
	Min	Max	Mean	SD	Min	Max	Mean	SD		
23 (A7)	0 (29%)	4 (3%)	1.12	1.06	0	4	0.98	0.93	0.79	0.67–0.87
24 (A8)	0 (40%)	4 (5%)	1.05	1.18	0	4	0.98	1.15	0.78	0.62–0.85
25 (A9)	0 (52%)	3	0.74	0.93	0	3	0.84	0.93	0.82	0.72–0.89
26 (A10)	0 (50%)	3	0.67	0.78	0	3	0.79	0.87	0.74	0.60–0.84
27 (A11)	0 (52%)	3	0.67	0.80	0	3	0.83	0.92	0.76	0.62–0.85
28 (A12)	0 (53%)	3	0.69	0.86	0	3	0.70	0.89	0.72	0.57–0.83
29 (A13)	0 (45%)	4 (3%)	0.81	0.96	0	4	0.79	0.99	0.83	0.72–0.89
30 (A14)	0 (81%)	2	0.24	0.54	0	3	0.31	0.63	0.70	0.54–0.81
31 (A15)	0 (76%)	2	0.26	0.48	0	2	0.36	0.61	0.55	0.34–0.71
32 (A16)	0 (17%)	4 (21%)	1.88	1.43	0	4	1.76	1.20	0.80	0.68–0.88
33 (A17)	0 (50%)	4 (9%)	0.93	1.23	0	4	0.93	1.20	0.79	0.67–0.87
34 (SP1)	0 (19%)	4 (22%)	1.91	1.44	0	4	1.76	1.35	0.84	0.74–0.90
35 (SP2)	0 (7%)	4 (29%)	2.38	1.34	0	4	2.31	1.38	0.81	0.70–0.88
36 (SP3)	0 (11%)	4 (32%)	2.33	1.44	0	4	2.17	1.49	0.82	0.71–0.89
37 (SP4)	0 (16%)	4 (22%)	2.07	1.40	0	4	1.95	1.46	0.81	0.70–0.88
38 (SP5)	0 (21%)	4 (30%)	2.10	1.54	0	4	1.95	1.38	0.81	0.70–0.88
39 (Q1)	0 (5%)	4 (38%)	2.78	1.24	0	4	2.48	1.30	0.70	0.54–0.82
40 (Q2)	0 (26%)	4 (7%)	1.46	1.22	0	4	1.25	1.13	0.71	0.56–0.82
41 (Q3)	0 (10%)	4 (18%)	1.93	1.27	0	4	1.75	1.27	0.79	0.67–0.87
42 (Q4)	0 (7%)	4 (16%)	1.95	1.18	0	4	1.90	1.17	0.82	0.71–0.89

Percentages in parentheses represent the proportion of study patients with the lowest or highest possible score for the nominated KOOS item, and provide information regarding floor and ceiling effects

SD standard deviation, ICC intraclass correlation coefficient, 95% CI 95% confidence interval

^a Each item refers to the individual 42 questions of the KOOS survey; a single item score of 0 points, best score, and 4, worst score

References

1. Beaton DE, Bombardier C, Guillemin F, Ferraz MB. Guidelines for the process of cross-cultural adaptation of self-report measures. *Spine*. 2000;25:3186–91.
2. Guillemin F, Bombardier C, Beaton D. Cross-cultural adaptation of health-related quality of life measures: literature review and proposed guidelines. *J Clin Epidemiol*. 1993;46:1417–32.
3. Felson DT, Zhang Y, Hannan MT, Naimark A, Weissman BN, Aliabadi P, Levy D. The incidence and natural history of knee osteoarthritis in the elderly. *The Framingham Osteoarthritis Study*. *Arthritis Rheum*. 1995;38:1500–5.
4. Muraki S, Oka H, Akune T, Mabuchi A, En-yo Y, Yoshida M, Saika A, Suzuki T, Yoshida H, Ishibashi H, Yamamoto S, Nakamura K, Kawaguchi H, Yoshimura N. Prevalence of radiographic knee osteoarthritis and its association with knee pain in the elderly of Japanese population-based cohorts: the ROAD study. *Osteoarthritis Cartilage*. 2009;17:1137–43.
5. Hashimoto H, Hanyu T, Sledge CB, Lingard EA. Validation of a Japanese patient-derived outcome scale for assessing total knee arthroplasty: comparison with Western Ontario and McMaster Universities osteoarthritis index (WOMAC). *J Orthop Sci*. 2003;8:288–93.
6. Akai M, Doi T, Fujino K, Iwaya T, Kurosawa H, Nasu T. An outcome measure for Japanese people with knee osteoarthritis. *J Rheumatol*. 2005;32:1524–32.
7. Kellgren JH, Jeffrey M, Ball J. Atlas of standard radiographs. Oxford: Blackwell Scientific; 1963.
8. Roos EM, Roos HP, Lohmander LS, Ekdahl C, Beynon BD. Knee Injury and Osteoarthritis Outcome Score (KOOS)—development of a self-administered outcome measure. *J Orthop Sports Phys Ther*. 1998;28:88–96.
9. Bellamy N, Buchanan WW, Goldsmith CH, Campbell J, Stitt LW. Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *J Rheumatol*. 1988;15:1833–40.
10. Roos EM, Lohmander LS. The Knee injury and Osteoarthritis Outcome Score (KOOS): from joint injury to osteoarthritis. *Health Qual Life Outcomes*. 2003;1:64.
11. Goncalves RS, Cabri J, Pinheiro JP, Ferreira PL. Cross-cultural adaptation and validation of the Portuguese version of the Knee injury and Osteoarthritis Outcome Score (KOOS). *Osteoarthritis Cartilage*. 2009;17:1156–62.
12. Ornetti P, Parratte S, Gossec L, Tavernier C, Argenson JN, Roos EM, Guillemin F, Mailliefert JF. Cross-cultural adaptation and validation of the French version of the Knee injury and Osteoarthritis Outcome Score (KOOS) in knee osteoarthritis patients. *Osteoarthritis Cartilage*. 2007;18:522–9.
13. Xie F, Li SC, Roos EM, Fong KY, Lo NN, Yeo SJ, Yang KY, Yeo W, Chong HC, Thumboo J. Cross-cultural adaptation and validation of Singapore English and Chinese versions of the Knee injury and Osteoarthritis Outcome Score (KOOS) in Asians with knee osteoarthritis in Singapore. *Osteoarthritis Cartilage*. 2006;14:1098–103.
14. Kessler S, Lang S, Puhl W, Stöve J. The Knee Injury and Osteoarthritis Outcome Score—a multifunctional questionnaire

- to measure outcome in knee arthroplasty. *Z Orthop Ihre Grenzgeb.* 2003;141:277–82.
15. de Groot IB, Favejee MM, Reijman M, Verhaar JA, Terwee CB. The Dutch version of the Knee Injury and Osteoarthritis Outcome Score: a validation study. *Health Qual Life Outcomes.* 2008;6:16.
 16. Salavati M, Mazaheri M, Negahban H, Sohani SM, Ebrahimi MR, Ebrahimi I, Kazemnejad A, Salavati M. Validation of a Persian-version of Knee injury and Osteoarthritis Outcome Score (KOOS) in Iranians with knee injuries. *Osteoarthritis Cartilage.* 2008;16:1178–82.
 17. Roos EM, Toksvig-Larsen S. Knee injury and Osteoarthritis Outcome Score (KOOS)—validation and comparison to the WOMAC in total knee replacement. *Health Qual Life Outcomes.* 2003;1:17.
 18. Verkleij SP, Luijsterburg PA, Koes BW, Bohnen AM, Bierma-Zeinstra SM. Effectiveness of diclofenac versus acetaminophen in primary care patients with knee osteoarthritis: [NTR1485], DIPA-trial: design of a randomized clinical trial. *BMC Musculoskelet Disord.* 2010;11:7.
 19. Birmingham TB, Giffin JR, Chesworth BM, Bryant DM, Litchfield RB, Willits K, Jenkyn TR, Fowler PJ. Medial opening wedge high tibial osteotomy: a prospective cohort study of gait, radiographic, and patient-reported outcomes. *Arthritis Rheum.* 2009;61:648–57.
 20. Englund M, Lohmander LS. Patellofemoral osteoarthritis coexistent with tibiofemoral osteoarthritis in a meniscectomy population. *Ann Rheum Dis.* 2005;64:1721–6.
 21. Dawson J, Fitzpatrick R, Murray D, Carr A. Questionnaire on the perceptions of patients about total knee replacement. *J Bone Joint Surg Br.* 1998;80:63–9.
 22. Murray DW, Fitzpatrick R, Rogers K, Pandit H, Beard DJ, Carr AJ, Dawson J. The use of the Oxford hip and knee scores. *J Bone Joint Surg Br.* 2007;89:1010–4.
 23. Takeuchi R, Sawaguchi T, Nakamura N, Ishikawa H, Saito T, Goldhahn S. Cross-cultural adaptation and validation of the Oxford 12-item knee score in Japanese. *Arch Orthop Trauma Surg.* 2010;131:247–54.
 24. Brazier JE, Harper R, Jones NM, O’Cathain A, Thomas KJ, Usherwood T, Westlake L. Validating the SF-36 health survey questionnaire: new outcome measure for primary care. *BMJ.* 1992;305:160–4.
 25. Gross Portney L, Watkins MP. *Foundations of clinical research—applications to practice.* New Jersey: Prentice Hall Health; 2010.
 26. Ware JEJ, Kosinski M, Dewey JE. *How to score - version 2 of the SF-36 health survey (standard & acute forms).* Rhode Island: QualityMetric Incorporated; 2001.
 27. Terwee CB, Bot SD, de Boer MR, van der Windt DA, Knol DL, Dekker J, Bouter LM, De Vet HC. Quality criteria were proposed for measurement properties of health status questionnaires. *J Clin Epidemiol.* 2007;60:34–42.
 28. Japanese Ministry of Health, Labour and Welfare. *Comprehensive survey of living conditions of the people on health and welfare.* <http://www.mhlw.go.jp/english/database/db-hss/cslc.html>. Accessed 17 Aug 2010.
 29. Edwards RR, Calahan C, Mensing G, Smith M, Haythornthwaite JA. Pain, catastrophizing, and depression in the rheumatic diseases. *Nat Rev Rheumatol.* 2011.
 30. Bellamy N, Sothorn RB, Campbell J. Rhythmic variations in pain perception in osteoarthritis of the knee. *J Rheumatol.* 1990;17:364–72.

RESEARCH ARTICLE

Open Access

Properties and usefulness of aggregates of synovial mesenchymal stem cells as a source for cartilage regeneration

Shiro Suzuki¹, Takeshi Muneta^{1,2}, Kunikazu Tsuji², Shizuko Ichinose³, Hatsune Makino⁴, Akihiro Umezawa⁴ and Ichiro Sekiya^{5*}

Abstract

Introduction: Transplantation of mesenchymal stem cells (MSCs) derived from synovium is a promising therapy for cartilage regeneration. For clinical application, improvement of handling operation, enhancement of chondrogenic potential, and increase of MSCs adhesion efficiency are needed to achieve a more successful cartilage regeneration with a limited number of MSCs without scaffold. The use of aggregated MSCs may be one of the solutions. Here, we investigated the handling, properties and effectiveness of aggregated MSCs for cartilage regeneration.

Methods: Human and rabbit synovial MSCs were aggregated using the hanging drop technique. The gene expression changes after aggregation of synovial MSCs were analyzed by microarray and real time RT-PCR analyses. *In vitro* and *in vivo* chondrogenic potential of aggregates of synovial MSCs was examined.

Results: Aggregates of MSCs cultured for three days became visible, approximately 1 mm in diameter and solid and durable by manipulation; most of the cells were viable. Microarray analysis revealed up-regulation of chondrogenesis-related, anti-inflammatory and anti-apoptotic genes in aggregates of MSCs. *In vitro* studies showed higher amounts of cartilage matrix synthesis in pellets derived from aggregates of MSCs compared to pellets derived from MSCs cultured in a monolayer. In *in vivo* studies in rabbits, aggregates of MSCs could adhere promptly on the osteochondral defects by surface tension, and stay without any loss. Transplantation of aggregates of MSCs at relatively low density achieved successful cartilage regeneration. Contrary to our expectation, transplantation of aggregates of MSCs at high density failed to regenerate cartilage due to cell death and nutrient deprivation of aggregates of MSCs.

Conclusions: Aggregated synovial MSCs were a useful source for cartilage regeneration considering such factors as easy preparation, higher chondrogenic potential and efficient attachment.

Introduction

Synovial mesenchymal stem cells (MSCs) are an attractive cell source for cartilage regeneration because of their high expansion and chondrogenic potentials [1-5]. We previously reported that more than 60% of synovial mesenchymal stem cells placed on osteochondral defects adhered to the defect within 10 minutes and promoted cartilage regeneration [6,7]. With this local adherent technique, we can transplant synovial MSCs without

scaffold. One of the disadvantages in this method is that the cell component in the suspension is invisible to the naked eye.

One of the solutions for this problem is to make aggregates of synovial MSCs [8-10]. This could enable MSCs not only to be visible but also to be heavier. Consequently, aggregates of MSCs will sink faster in the suspension medium than dispersed MSCs. The use of aggregates of MSCs may help to avoid loss of MSCs from targeted cartilage defects and improve the procedures of transplantation of synovial MSCs. However, there are still concerns; properties of synovial MSCs will be altered when synovial MSCs are aggregated. We do not know

* Correspondence: sekiya.orj@tmd.ac.jp

⁵Section of Cartilage Regeneration, Graduate School, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan
Full list of author information is available at the end of the article

whether aggregates of MSCs adhere on the cartilage defect as we expect it will, and the proper number of aggregates is unclear.

In this study, properties of aggregates of human synovial MSCs were analyzed from the standpoints of morphology, gene profile and *in vitro* chondrogenic potential. Also, the effect of transplantation of aggregates of synovial MSCs was investigated in a rabbit cartilage defect model in terms of aggregate number, cell behavior and influential factors in the *in vivo* chondrogenesis of aggregates of synovial MSCs. Finally, we demonstrated the usefulness of aggregates of synovial MSCs as a source for cartilage regeneration therapy.

Materials and methods

Isolation and culture of human synovial MSCs

This study was approved by an institutional review board of Tokyo Medical and Dental University (No.1030), and informed consent was obtained from all subjects. Human synovium was harvested from donors during anterior cruciate ligament reconstruction surgery for ligament injury and digested in a 3 mg/ml collagenase D solution (Roche Diagnostics, Mannheim, Germany) in α -minimal essential medium (α MEM) (Invitrogen, Carlsbad, CA, USA) at 37°C. After three hours, digested cells were filtered through a 70 μ m nylon filter (Becton, Dickinson and Company, Franklin Lakes, NJ, USA), and the remaining tissues were discarded. The digested cells were plated in a 150 cm² culture dish (Nalge Nunc International, Rochester, NY, USA) in complete culture medium (CCM): α MEM containing 10% fetal bovine serum (FBS; Invitrogen), 100 units/ml penicillin (Invitrogen), 100 μ g/ml streptomycin (Invitrogen), and 250 ng/ml amphotericin B (Invitrogen) and incubated at 37°C with 5% humidified CO₂. The medium was changed to remove nonadherent cells one day later and cultured for 14 days as passage 0, then replated at 100 cells/cm² in a 150 cm² culture dish, cultured for 14 days and cryopreserved as passage 1. To expand the cells, a frozen vial of the cells was thawed, plated in 60 cm² culture dishes, and incubated for four days in the recovery plate. These cells were replated at 100 cells/cm² in a 150 cm² culture dish (passage 3), and cultured for an additional 14 days. These passage 3 cells were harvested and used in this study.

Isolation and culture of rabbit synovial MSCs

This study was approved by the Animal Experimentation Committee of Tokyo Medical and Dental University (No.0120296A). Wild type skeletally mature Japanese White Rabbit and GFP transgenic rabbits [11,12] were anesthetized with an intramuscular injection of 25 mg/kg ketamine hydrochloride and with an intravenous injection of 45 mg/kg sodium pentobarbital and 150 μ g/kg medetomidine hydrochloride. Synovium was harvested aseptically

from knee joints of the rabbits, and digested in a 3 mg/ml collagenase type V in α MEM for three hours at 37°C. The digested cells were plated at 5×10^4 cells/cm² in a 150 cm² culture dish in CCM and incubated at 37°C with 5% humidified CO₂. The medium was changed to remove nonadherent cells one day later and cultured for seven days as passage 0. The cells were then trypsinized, harvested and resuspended to be used for further assays. The cells that were transplanted in animals to be sacrificed at Day 0 and Day 14 were labeled for cell tracking by the fluorescent lipophilic tracer DiI (Molecular Probes, Eugene, OR, USA). For labeling, synovial MSCs were resuspended at 1×10^6 cells/ml in α MEM without FBS and a DiI was added at a final concentration of 5 μ l/ml. After incubation for 20 minutes at 37°C with 5% humidified CO₂, the cells were centrifuged at 450 g for 5 minutes and washed twice with phosphate-buffered saline (PBS) and the cells were then resuspended in CCM and cultured in hanging drops. We already reported that these cells had characteristics of MSCs [3,6,7,11].

Preparation of aggregates of synovial MSCs

A total of 2.5×10^5 synovial MSCs were trypsinized, harvested and resuspended in 35 μ l of CCM, plated on an inverted culture dish lid. The lid was inverted and placed on a culture dish containing PBS. The cells were cultured at 37°C with 5% humidified CO₂ for three days in hanging drops.

Histology of aggregates of human synovial MSCs

Aggregates of human synovial MSCs were fixed with 2.5% glutaraldehyde in 0.1 M PBS for two hours. The aggregates were washed overnight at 4°C in the same buffer and post-fixed with 1% OsO₄ buffered with 0.1 M PBS for two hours. The aggregates were dehydrated in a graded series of ethanol and embedded in Epon 812. Semi-thin (1 μ m) sections for light microscopy were collected on glass slides and stained for 30 seconds with toluidine blue.

In vitro chondrogenic differentiation assay

A total of 2.5×10^5 human synovial MSCs cultured as a monolayer were pelleted by trypsinization and centrifugation. The pellets or aggregate of human synovial MSCs cultured for three days in hanging drops were cultured in 400 μ l chondrogenic medium consisting of high-glucose Dulbecco's modified Eagle's medium (Invitrogen) supplemented with 1,000 ng/ml BMP-7 (Stryker Biotech, Boston, MA, USA), 10 ng/ml transforming growth factor- β 3 (R&D Systems, Minneapolis, MN, USA), 100 nM dexamethasone (Sigma-Aldrich Corp., St. Louis, MO, USA), 50 μ g/ml ascorbate-2-phosphate, 40 μ g/ml proline, 100 μ g/ml pyruvate, and 1:100 diluted ITS+Premix (6.25 μ g/ml insulin, 6.25 μ g/ml transferrin, 6.25 ng/ml selenious acid, 1.25 mg/ml bovine serum

albumin, and 5.35 mg/ml linoleic acid; BD Biosciences Discovery Labware, Bedford, MA, USA). The medium was changed every 3 to 4 days for 21 days.

Histology of pellets of human synovial MSCs

The pellets were embedded in paraffin, cut into 5- μ m sections and stained with 1% Toluidine Blue. For immunohistochemistry, sections were treated with 0.4 mg/ml proteinase K (DAKO, Carpinteria, CA, USA) in Tris-HCl and normal horse serum after deparaffinization. Primary antibodies for type II collagen (Daiichi Fine Chemical, Toyama, Japan) and a secondary antibody of biotinylated horse anti-mouse IgG (Vector Laboratories, Burlingame, CA, USA) were employed. Immunostaining was detected with VECTASTAIN ABC reagent (Vector Laboratories) followed by 3,3'-diaminobenzidine staining.

Real-time RT PCR analysis

Total RNA was extracted from human synovial MSCs in a monolayer culture, aggregates of human synovial MSCs cultured for 1, 2 and 3 days, and the pellets cultured for 7, 14 and 21 days using QIAzol (Qiagen, Hilden, Germany) and the RNeasy mini kit (Qiagen). cDNA was synthesized with oligo-dT primer from total RNA using the Transcriptor High Fidelity cDNA Synthesis kit (Roche Diagnostics) according to the manufacturer's protocol. Reverse transcription (RT) was performed by 30 minutes incubation at 55°C followed by 5 minutes incubation at 85°C. Real-time PCR was performed in a LightCycler 480 instrument (Roche Diagnostics). Primer sequences and TaqMan probes are listed in Table 1. After an initial denaturation step (95°C for 10 minutes), amplification was performed for 40 cycles (95°C for 15 seconds, 60°C for 60 seconds). Relative amounts of mRNA were calculated and standardized as previously described [13,14].

DNA microarray analysis

Total RNA was extracted from human synovial MSCs in a monolayer culture, aggregates of human synovial MSCs cultured for three days. Human Genome U133 Plus 2.0 Array (GeneChip; Affymetrix, Santa Clara, CA, USA) containing the oligonucleotide probe set for more than 47,000 transcripts was used. The fluorescence intensity of each probe was quantified by using the GeneChip Analysis Suite 5.0 (Affymetrix). Gene expression data were normalized in Robust MultiChip Analysis (RMA). To analyze the data, we used hierarchical clustering using TIGR MultiExperiment Viewer (MeV) [15]. The microarray data have been deposited to the public database (GEO accession# GSE 31980).

In vivo transplantation

Under anesthesia, the left knee joint was approached through a medial parapatellar incision, and the patella

Table 1 Real time-RT PCR primer sequences

Primer name		Sequences	Probe No.
<i>β-actin</i>	forward	5'-ATTGGCAATGAGCGGTC-3'	11
	reverse	5'-TGAAGGTAGTTTCGTGGATGC-3'	
<i>BMP2</i>	forward	5'-CGGACTGCGGTCTCCTAA-3'	49
	reverse	5'-GGAAGCAGCAACGCTAGAAG-3'	
<i>SOX5</i>	forward	5'-TCTGTCCCAGCAGCGTTAG-3'	41
	reverse	5'-TGACAGCATCATGGTCATTTAAG-3'	
<i>SOX6</i>	forward	5'-GCTTCTGGACTCAGCCCTTA-3'	50
	reverse	5'-GGCCCTTAGCCTTGGTTA-3'	
<i>SOX9</i>	forward	5'-GTACCCGCACTGCACAAC-3'	61
	reverse	5'-TCGCTCTGTTCAGAAGTCTC-3'	
<i>TSG6</i>	forward	5'-CCAGATGACATCATCAGTACAGG-3'	78
	reverse	5'-CATTGCAACATATTTGATTTGGA-3'	
<i>STC1</i>	forward	5'-CCCAATCACTTCTCCAACAGA-3'	40
	reverse	5'-TGCTGACTGTGTCTTCATACA-3'	
<i>COL2A1</i>	forward	5'-GTGTCAGGGCCAGGATGT-3'	75
	reverse	5'-TCCCAGTGTACAGACACAGAT-3'	
<i>AGGRECAN</i>	forward	5'-CTGGAAGTCGTGGTAAAGG-3'	21
	reverse	5'-TCGAGGGTGTAGCGTGTAGA-3'	

was dislocated laterally. Full-thickness osteochondral defects (5 mm \times 5 mm wide, 1.5 mm deep) were created in the trochlear groove of the femur. A total of 5, 10, 20, 40 and 80 aggregates of autologous rabbit synovial MSCs (2.5×10^5 cells/aggregate) or 25 and 100 smaller aggregates of autologous rabbit synovial MSCs (1.0×10^5 cells/aggregate) suspended in PBS were transplanted to the defect. To trace the transplanted cells, DiI-labeled aggregates of autologous rabbit synovial MSCs and aggregates of allogenic synovial MSCs derived from GFP transgenic rabbit were transplanted to the defect. For the control group, the defect was left empty. All rabbits were returned to their cages after the operation and were allowed to move freely. Animals were sacrificed with an overdose of sodium pentobarbital at 1, 2, and 4 days and at 12 weeks after the operation ($n = 5$ at each time).

Macroscopic examination

The cartilage defects were examined macroscopically for color, integrity and smoothness. Osteoarthritic changes and synovitis of the knee were also investigated. Digital images were taken using an Olympus MVX10 (Olympus, Tokyo, Japan).

Histological examination and fluorescent microscopic examination

The dissected distal femurs were immediately fixed in a 4% paraformaldehyde (PFA) solution. The specimens were decalcified in 4% ethylenediamine tetraacetic acid solution, dehydrated with a gradient ethanol series and

embedded in paraffin blocks. Sagittal sections 5 μ m thick were obtained from the center of each defect and were stained with toluidine blue and Safranin O. For fluorescent microscopic examination and terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) staining, the fixed specimens were incubated at 4°C for three hours in 5%, 10%, 15% and 20% sucrose solution, respectively. After incubation, the fixed specimens were mounted on a holder. Then 30% optimal cutting temperature (OCT) (Sakura Finetek, Tokyo, Japan) in sucrose solution was added gently into the holder. The holder was frozen in hexan chilled by dry ice and stored at -80°C. Cryosections (10 μ m) were prepared with an ultracut S microtome (Reichert, Wien, Austria) and a Microm HM560 cryostat.

Histological score

Histological sections of the repaired tissue were analyzed using a grading system consisting of five categories (cell morphology, morphology, matrix staining, surface regularity, cartilage thickness and integration of donor with host), which were modified from the repaired cartilage score described by Wakitani and colleagues [16], so that overly thick, regenerated cartilage could not be overestimated [6]. The scoring was performed in a blinded manner by two observers and there was no significant interobserver difference. The ratio of the safranin-O positive area over the defect was evaluated. Zeiss AxioVison software (Carl Zeiss, Oberkochen, Germany) was used for measurement of defects and safranin-O positive areas.

In vitro viability assay

Aggregates of rabbit synovial MSCs were plated at 1 or 40 aggregates/well in 96-well plates (Nunc) in CCM, and incubated at 37°C with 5% humidified CO₂ for seven days without medium change. Aggregates were fixed in 4% PFA for TUNEL staining.

TUNEL staining

For TUNEL staining, an apoptosis *in situ* detection kit (Wako Pure Chemical Industries, Ltd, Osaka, Japan) was used. The frozen semi-thin sections were incubated with terminal deoxynucleotidyl transferase for 10 minutes at 37°C in a moist chamber. The sections were washed with 0.1 M PBS for 15 minutes. Peroxidase-conjugated antibody was then applied to the specimens at 37°C for 10 minutes in a moist chamber. The sections were developed with 3,3-diaminobenzidine and counterstained with methyl green.

Statistical analysis

Comparisons between two groups were analyzed using the Mann-Whitney U test. Comparisons between multi groups were analyzed using the Kruskal-Wallis test and

the Steel test. A *P*-value of < 0.05 was considered statistically significant.

Results

Appearance of aggregates of human synovial MSCs

Human synovial MSCs were aggregated using the hanging drop technique (Figure 1A). Three days after being cultured in the drop (Figure 1B), the aggregate, consisting of 250,000 MSCs, became approximately 1 mm in diameter (Figure 1C). The aggregate was not easily broken by manipulation. Sagittal sections of the aggregates showed heart-shape as a whole (Figure 1Da). The superficial layer was composed of spindle cells parallel to the surface, whereas the deep layer was comprised of round cells both at top and bottom of the aggregate (Figure 1Db, c). Though cells positive for TUNEL staining were observed, the number was only approximately under 5% (Figure 1Dd).

Transcriptome profile of aggregates of human synovial MSCs

To examine the sequential changes of gene expression profiles during aggregation of human synovial MSCs, microarray analyses were performed. The differences of gene profile between before and after aggregation exceeded those among donor variances (Figure 2A). The number of genes up-regulated more than five-fold was 621. The number of genes up-regulated more than 100-fold was 10, and these genes were related to hypoxia (integrin, alpha 2 (*ITGA2*), stanniocalcin 1 (*STC1*), chemokine (C-X-C motif) receptor 4 (*CXCR4*)), nutrient (*BMP2*, proprotein convertase subtilisin/kexin type 1 (*PCSK1*), secreted phosphoprotein 1 (*SPPI*), *ITGA2*, *STC1*), extracellular region (*MMP1*, *MMP3*), and cell adhesion (*SPPI*, *ITGA2*) (Table 2). The most up-regulated gene was *BMP2*, increased to 273 folds (Table 2). *STC1* was also highly up-regulated in aggregates of synovial MSCs. The number of genes down-regulated less than one-fifth was 409, and the ontology for the genes was related to cell cycle. The microarray data are available at the public database (GEO accession# GSE 31980).

To further investigate gene expressions during aggregation of human synovial MSCs, real time RT-PCR analyses were additionally used for chondrogenesis-related genes (SRY (sex determining region Y)-box (*SOX*)5, -6, -9, and *BMP2*) and anti-inflammatory genes (TNF α inducible gene 6 (*TSG-6*), and *STC-1*) in four donors. In most cases, expressions for these genes increased sequentially (Figure 2B).

In vitro chondrogenesis of aggregates of human synovial MSCs

In vitro chondrogenic ability of human synovial MSCs after hanging drop culture was compared to that of MSCs after monolayer culture (Figure 3A). Aggregates





Figure 1 Preparation and appearance of aggregates of human synovial MSCs. (A): Scheme of preparation of aggregates using hanging drop technique. **(B):** Drops hanging on the cover of 15 cm dish. **(C):** Macroscopic image of aggregate consisted of 250,000 MSCs, three days after cultured in hanging drop. **(D):** Sagittal sections of aggregates stained with toluidine blue (a, b, c) and TUNEL (d). TUNEL positive cells are indicated with arrows.



of MSCs differentiated into chondrocytes as well (Figure 3B). The wet weight of pellets derived from MSCs after hanging drop culture was heavier than that of pellets derived from MSCs after monolayer culture in all four donors at 14 or 21 days (Figure 3C). Real time RT-PCR analysis showed higher expression levels of collagen (*COL2A1*), aggrecan and *SOX9* for pellets derived from MSCs after hanging drop culture compared to MSCs after monolayer culture at 14 and 21 days (Figure 3D). Cartilage extracellular matrix synthesis and accumulation of type II collagen were confirmed by histological

analysis stained with toluidine blue and immunohistochemical analysis (Figure 3E).

***In vivo* analysis for cartilage regeneration by transplantation of aggregates of synovial MSCs in rabbits**

To examine whether transplantation of aggregates of synovial MSCs promotes cartilage regeneration, *in vivo* study was performed in rabbits. To further investigate the optimal number of aggregates consisting of 250,000 MSCs, 0 to 80 aggregates were transplanted into the defect.

Table 2 The top 10 upregulated genes in aggregates of MSCs

No.	Genebank No.	Gene name	Symbol	fold change
1	AA583044	bone morphogenetic protein 2	BMP2	273
2	NM_002421	matrix metalloproteinase 1	MMP1	205
3	NM_000439	proprotein convertase subtilisin/kexin type 1	PCSK1	179
4	M86849	gap junction protein, beta 2	GJB2	170
5	M83248	secreted phosphoprotein 1 (osteopontin)	SPP1	156
6	L27624	tissue factor pathway inhibitor 2	TFPI2	137
7	NM_002422	matrix metalloproteinase 3	MMP3	136
8	N95414	integrin, alpha 2 (CD49B)	ITGA2	129
9	AW003173	stanniocalcin 1	STC1	124
10	AJ224869	chemokine (C-X-C motif) receptor 4	CXCR4	101

The top 10 genes which increased higher in aggregates of MSCs cultured in hanging drops for three days compared with MSCs in a monolayer culture. Values are the means among three individual donors.

At 0 days, in the case of 40 and 80 transplanted aggregates, the osteochondral defects were filled with aggregates labeled with DiI macroscopically (Figure 4A).

At four weeks, in the case of 5 and 10 transplanted aggregates, the osteochondral defect was mostly covered with a thick cartilage matrix (Figure 4B, C). In the case of 20 and 40 transplanted aggregates, the defect was partially covered with cartilage matrix. In the case of 80 transplanted aggregates, the defect was filled with only fibrous tissue, which appeared to be similar to the control (Figure 4B).

At 12 weeks, in the case of 10 transplanted aggregates, the border between cartilage and bone moved up, and thickness of the regenerated cartilage became similar to the neighboring cartilage (Figure 4B, D). In the case of 5 and 20 transplanted aggregates, the bone defect was repaired, but the cartilage defect was filled partially with cartilage matrix. In the case of 40 and 80 transplanted aggregates, the osteochondral defect was poorly repaired, similar to the control (Figure 4B). Histological score was the best and the safranin-O positive area ratio was highest in the case of 10 transplanted aggregates both at 4 and 12 weeks (Figure 4E, F).

To trace MSCs, 10 aggregates of GFP positive MSCs were transplanted into the defect. At Day 1, no GFP positive aggregates could be observed in the knee joint except the defects with a fluorescent stereomicroscope. Histologically, aggregates changed their forms but have not fused yet (Figure 5A). At four weeks, the defect was filled with cartilage matrix and the GFP positive cells were still observed both at the bottom and the center of the regenerated cartilage (Figure 5B). Regenerated cartilage consisted of both GFP positive cells and GFP negative cells.

Influences of cell number per aggregate and of aggregate number for transplantation

Cell number per aggregate as well as aggregate number may be a factor affecting properties of the aggregates.

To answer this question, 25 or 100 aggregates, in which an aggregate consisted of 100,000 MSCs, were transplanted into the osteochondral defect.

At four weeks, in the case of 25 transplanted aggregates, the defect was fully filled with cartilage matrix (Figure 6A), in which the result was different from the case of 20 or more aggregates, in which an aggregate consisted of 250,000 MSCs. In the case of 100 transplanted aggregates, the defect was filled with fibrous tissue, and the histological score was inferior and the safranin-O positive area ratio was smaller. (Figure 6B, C).

Influences of aggregate number on viability of MSCs

To clarify why transplantation of aggregates over a certain number resulted in poor outcome, viability of cells was first examined by TUNEL staining. Compared to the case of 10 transplanted aggregates, much more TUNEL positive cells could be observed in the case of 80 transplanted aggregates (Figure 6D).

Another factor might be a nutrient deprivation and *in vitro* analyses using aggregates of rabbit synovial MSCs were performed. Seven days after 1 or 40 aggregates were cultured in a well of 96-well plates, the medium color changed to yellow in the case of 40 aggregates, while the color remained red in the case of only 1 aggregate (Figure 6E). TUNEL positive cells were much higher in the case of 40 aggregates than in the case of only 1 aggregate.

Discussion

In this study, to form aggregates of synovial MSCs, the hanging drop technique was used [8-10]. This is a simple method; expensive or specific tools are not required. Three days after cultured in the drop, the aggregate, consisting of 250,000 MSCs, became approximately 1 mm in diameter, large enough to be visible and solid enough to aspirate with a pipette. Aggregates of MSCs sank faster in the suspension medium than dispersed




Figure 3 *In vitro* chondrogenic ability of human synovial MSCs after hanging drop culture **(A):** Scheme for the analyses. **(B):** Macroscopic images of pellets derived from aggregates of MSCs and those of pellets derived from MSCs in a monolayer culture. **(C):** Wet weight in four individual donors. Values are the means with standard deviation (SD) ($P < 0.05$ by the Mann-Whitney U test). **(D):** Expressions of chondrogenesis-related genes by RT-PCR analyses. Values are the means with SD among four donors. The fold changes of *SOX9* and *AGGRECAN* expression levels were shown when the gene expression levels at Day 0 were normalized as 1. The fold changes of *COL2A1* expression levels were shown when the gene expression levels in MSCs in monolayer at Day 7 were normalized as 1 because *COL2A1* expression level at Day 0 was undetectable. **(E):** Histological sections of pellets stained with toluidine blue and immunohistochemical analysis for type II collagen.




Figure 4 Cartilage regeneration by transplantation of aggregates of synovial MSCs in rabbits. (A): Macroscopic observation of osteochondral defects one minute after transplantation of indicated number of aggregates of MSCs. The aggregate consisted of 250,000 MSCs, labeled with Dil for visualization. (B): Macroscopic and histological observation. For histologies, sagittal sections were stained with safranin-O (SO) and toluidine blue (TB). (C, D): Magnified histology of the indicated area. (E): Histological score. Values are the means with SD. ($n = 5$; $P < 0.05$ by the Kruskal-Wallis test and the Steel test). (F): Ratio of the safranin-O positive area to the defect area. Values are the means with SD. ($n = 5$; $P < 0.05$ by the Kruskal-Wallis test and the Steel test).




Figure 5 Transplantation of 10 aggregates of synovial MSCs derived from a GFP rabbit. (A): Sagittal sections of osteochondral defect under fluorescence for GFP at one day. **(B):** Macroscopic and histological observation four weeks after transplantation. Nuclei were shown as blue in higher magnified pictures.


MSCs and helped to avoid loss of MSCs from targeted cartilage defect. The use of aggregates was practically convenient for transplantation of MSCs.

In the previous report, the number of apoptotic or necrotic cells was greater in aggregates prepared with 100,000 or 250,000 human bone marrow MSCs, which was examined by flow cytometry, measuring propidium iodide uptake and annexin V labeling [10]. We examined the viability of aggregates of MSCs by TUNEL staining and confirmed that cells positive for TUNEL staining were observed; the number was small compared to the previous report. This difference may have been due to the difference of methods. Microarray analysis showed up-regulation of genes with ontology for regulation of cell death. The microarray data are available at the public database (GEO accession# GSE 31980). These results suggest that aggregation of 250,000 MSCs affect the viability of cells. However, we thought that aggregates of MSCs could be used as a source for cartilage regeneration because most cells which are cultured in drops for three days are viable.

Aggregation of synovial MSCs changed the gene expression profile dramatically without any special tools

or chemical factors. This is possibly due to environmental changes, including cell-to-cell contact, hypoxic condition and low nutrient condition. Aggregation of human synovial MSCs increased expressions of several chondrogenesis-related genes and the most up-regulated gene was BMP2, which was also up-regulated in bone marrow MSCs [8,10].

In this study, we compared *in vitro* chondrogenesis potential of synovial MSCs after hanging drop culture with that of MSCs after monolayer culture. We used 1,000 ng/ml BMP7 for *in vitro* chondrogenic differentiation assay. We previously examined the dose effect of BMP6 between 0 to 500 ng/ml for *in vitro* chondrogenesis of bone marrow MSCs. Cartilage pellets increased in size along with the concentration of BMP6, and a maximal effect was at 500 ng/ml [17]. Our preliminary experiments showed that 1,000 ng/ml BMP6 induced larger cartilage pellets than 500 ng/ml BMP6 in bone marrow and synovial MSCs. We obtained similar results with BMP7. Real time RT-PCR analysis showed higher expression levels of COL2A1, aggrecan and SOX9 for pellets derived from MSC-aggregates after hanging drop culture compared to those of MSCs in a monolayer culture.



Furthermore, the wet weight of pellets derived from MSC-aggregates after hanging drop culture was heavier than that of pellets derived MSCs in a monolayer culture. These indicate that chondrogenic potential increased in aggregates of MSCs after hanging drop culture.

In this study, we used an osteochondral defect model of rabbits, which have a higher, self-renewal capacity than bigger animals and humans. Therefore, the results obtained here should be critically evaluated. However,

we prepared negative controls, which healed poorly at 4 and 12 weeks. We previously confirmed that the osteochondral defect created in the trochlear groove of the femur, similar to this study, was not repaired without any treatments 24 weeks after surgery [6]. These findings indicate that this rabbit model is useful to evaluate the effects of the treatments for cartilage regeneration.

For *in vivo* analysis of cartilage regeneration by transplantation of aggregates of synovial MSCs in rabbits,

successful cartilage regeneration was observed in the cases of a relatively small number of transplanted aggregates of MSCs, and the worst results were observed when the highest number of aggregates of MSCs was transplanted. These results were not what we expected, because we previously reported that better cartilage regeneration was obtained when higher cell densities of MSCs were embedded in collagen gel [3].

Why were poor results obtained when more than a certain number of aggregates were transplanted? We listed three possible reasons. First, nutrition to maintain transplanted MSCs was depleted and the environment around transplanted MSCs worsened when too many aggregates were transplanted. As shown in Figure 6E, in the case of 40 aggregates that were cultured for seven days in a well of 96-well plates, medium color changed to yellow. This means that adjustment of pH could not be controlled. Second, TUNEL positive cells increased when too many aggregates were transplanted. The number of TUNEL positive cells was higher when too many aggregates were transplanted (Figure 6D) than before transplantation (Figure 1D) and after a suitable number of aggregates were transplanted (Figure 6D). Third, transplantation of too many aggregates prevented chondro-progenitor cells from moving to the osteochondral defect from bone marrow and from synovial fluid.

We confirmed that transplanted aggregates of synovial MSCs were directly differentiated into chondrocytes by transplanting MSCs derived from GFP transgenic rabbit. This result suggests that aggregates of synovial MSCs were involved in the reparative process. However, as shown in Figure 5B, in the case of aggregates of GFP positive MSCs being transplanted, regenerated cartilage consisted of both GFP positive cells and GFP negative cells. MSCs existed in synovial fluid [18] and these MSCs contributed to the repair of cartilage injury [6,19]. These results suggest that some host MSCs were also involved in the reparative process. In addition, host MSCs may have been involved in the anti-inflammatory process. In our rabbit osteochondral defect model, inflammation like a synovitis was not severe even in the control group. Therefore, we could not confirm the anti-inflammatory effect of MSCs. It would be interesting to investigate the anti-inflammatory effect of transplantation of aggregates of synovial MSCs and host MSCs in other arthritis models.

As previously reported, in bone marrow MSCs [10], aggregates of human synovial MSCs expressed anti-inflammatory genes *TSG6* and *STC1*. *TSG6* is secreted by synoviocytes, mononuclear cells and chondrocytes under inflammatory conditions and has an anti-inflammatory effect. Overexpression of *TSG6* or administration of recombinant *TSG6* inhibited inflammation and joint destruction in a murine collagen induced arthritis model

[20-23]. *STC1* is reported to have an anti-apoptotic effect as well as an anti-inflammatory effect [24,25]. However, their roles in joint homeostasis are unknown.

In this study, transplantation of low numbers of aggregates, in other words, low density of aggregates to the volume of the cartilage defect, showed better regeneration (Figures 4 and 6). This is favorable for clinical application. We have performed clinical trials of autologous human synovial MSCs transplantation for cartilage defects. In the experiences of 12 patients, approximately 50 million synovial MSCs at passage 0 were transplanted for approximately 280 mm² cartilage defects (unpublished data). In a rabbit model, we transplanted synovial MSC-aggregates into the osteochondral defects without any loss of cells, and 10 MSC-aggregates (2.5 × 10⁶ cells) per 25 mm² defects were needed for better cartilage regeneration. According to these data, we can prepare a sufficient amount of human synovial MSCs at passage 0.

In this study, we did not use scaffolds for transplantation of aggregates of synovial MSCs. We were able to adhere aggregates of synovial MSCs on the osteochondral defect without scaffolds; however, the use of scaffolds or materials to improve survival of transplanted cells is attractive. One of the methods is the use of a fibrin glue, which has an effect of improving survival of transplanted cells [26]. In addition, cell transplantation of MSCs with a fibrin glue can probably be performed under arthroscopic surgery. Further studies are needed to improve cell transplantation procedures.

Conclusion

Aggregated synovial MSCs were a useful source for cartilage regeneration considering such factors as easy preparation, higher chondrogenic potential and efficient attachment.

Abbreviations

αMEM: α-minimal essential medium; *BMP*: bone morphogenetic protein; CCM: complete culture medium; *COL*: collagen; *CXCR4*: chemokine (C-X-C motif) receptor 4; EDTA: ethylenediaminetetraacetate; FBS: fetal bovine serum; GFP: green fluorescent protein; *GJB2*: gap junction protein, beta 2; *ITGA2*: integrin, alpha 2; MeV: MultiExperiment Viewer; MMP: matrix metalloproteinase; MSC: mesenchymal stem cell; OCT: optimal cutting temperature; PBS: phosphate-buffered saline; *PCSK1*: proprotein convertase subtilisin/kexin type 1; PFA: paraformaldehyde; RMA: Robust MultiChip Analysis; RT: reverse transcription; SD: standard deviation; SO: safranin-O; *SOX*: SRY (sex determining region Y)-box; *SPP1*: secreted phosphoprotein 1; *STC1*: stanniocalcin 1; TB: toluidine blue; *TFPI2*: tissue factor pathway inhibitor 2; *TNF*: tumor necrosis factor; *TSG6*: TNFα inducible gene 6; TUNEL: terminal deoxynucleotidyl transferase dUTP nick end labeling.

Acknowledgements

This study was supported by "the Project for Realization of Regenerative Medicine" and "the Global Center of Excellence (GCOE) Program" by the Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan to IS, grants from the Japanese Ministry of Education Global Center of Excellence Program, International Research Center for Molecular Science in

Tooth and Bone Diseases to TM, and from a Health and Labor Sciences Research Grant, Research on Regenerative Medicine for Clinical Application to IS. Recombinant human BMP-7 was distributed by Stryker Biotech. We thank Miyoko Ojima for her expert help with histology and Izumi Nakagawa for management of our laboratory.

Author details

¹Section of Orthopedic Surgery, Graduate School, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan. ²Global Center of Excellence Program, International Research Center for Molecular Science in Tooth and Bone Diseases, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan. ³Instrumental Analysis Research Center, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan. ⁴Department of Reproductive Biology, National Research Institute for Child Health and Development, 2-10-1 Okura, Setagaya-ku, Tokyo 157-8535, Japan. ⁵Section of Cartilage Regeneration, Graduate School, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan.

Authors' contributions

SS participated in the design of the study, carried out the animal experiments, analyzed the results and drafted the manuscript. TM participated in the design of the study and provided the administrative and financial support. KT participated in the design of the study. SI helped with histological analysis. HM and AU carried out the microarray analysis and participated in the evaluation of the results. IS participated in the design of the study, provided the financial support and completed the final manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 24 November 2011 Revised: 30 April 2012

Accepted: 7 June 2012 Published: 7 June 2012

References

1. Sakaguchi Y, Sekiya I, Yagishita K, Muneta T: Comparison of human stem cells derived from various mesenchymal tissues: superiority of synovium as a cell source. *Arthritis Rheum* 2005, **52**:2521-2529.
2. Shirasawa S, Sekiya I, Sakaguchi Y, Yagishita K, Ichinose S, Muneta T: *In vitro* chondrogenesis of human synovium-derived mesenchymal stem cells: optimal condition and comparison with bone marrow-derived cells. *J Cell Biochem* 2006, **97**:84-97.
3. Koga H, Muneta T, Nagase T, Nimura A, Ju YJ, Mochizuki T, Sekiya I: Comparison of mesenchymal tissues-derived stem cells for *in vivo* chondrogenesis: suitable conditions for cell therapy of cartilage defects in rabbit. *Cell Tissue Res* 2008, **333**:207-215.
4. Nimura A, Muneta T, Koga H, Mochizuki T, Suzuki K, Makino H, Umezawa A, Sekiya I: Increased proliferation of human synovial mesenchymal stem cells with autologous human serum: comparisons with bone marrow mesenchymal stem cells and with fetal bovine serum. *Arthritis Rheum* 2008, **58**:501-510.
5. Koga H, Engebretsen L, Brinckmann JE, Muneta T, Sekiya I: Mesenchymal stem cell-based therapy for cartilage repair: a review. *Knee Surg Sports Traumatol Arthrosc* 2009, **17**:1289-1297.
6. Koga H, Shimaya M, Muneta T, Nimura A, Morito T, Hayashi M, Suzuki S, Ju YJ, Mochizuki T, Sekiya I: Local adherent technique for transplanting mesenchymal stem cells as a potential treatment of cartilage defect. *Arthritis Res Ther* 2008, **10**:R84.
7. Shimaya M, Muneta T, Ichinose S, Tsuji K, Sekiya I: Magnesium enhances adherence and cartilage formation of synovial mesenchymal stem cells through integrins. *Osteoarthritis Cartilage* 2010, **18**:1300-1309.
8. Potapova IA, Gaudette GR, Brink PR, Robinson RB, Rosen MR, Cohen IS, Doronin SV: Mesenchymal stem cells support migration, extracellular matrix invasion, proliferation, and survival of endothelial cells *in vitro*. *Stem Cells* 2007, **25**:1761-1768.
9. Potapova IA, Brink PR, Cohen IS, Doronin SV: Culturing of human mesenchymal stem cells as three-dimensional aggregates induces functional expression of CXCR4 that regulates adhesion to endothelial cells. *J Biol Chem* 2008, **283**:13100-13107.
10. Bartosh TJ, Ylostalo JH, Mohammadipoor A, Bazhanov N, Coble K, Claypool K, Lee RH, Choi H, Prockop DJ: Aggregation of human mesenchymal stromal cells (MSCs) into 3D spheroids enhances their antiinflammatory properties. *Proc Natl Acad Sci USA* 2010, **107**:13724-13729.
11. Miyamoto T, Muneta T, Tabuchi T, Matsumoto K, Saito H, Tsuji K, Sekiya I: Intradiscal transplantation of synovial mesenchymal stem cells prevents intervertebral disc degeneration through suppression of matrix metalloproteinase-related genes in nucleus pulposus cells in rabbits. *Arthritis Res Ther* 2010, **12**:R206.
12. Takahashi R, Kuramochi T, Aoyagi K, Hashimoto S, Miyoshi I, Kasai N, Hakamata Y, Kobayashi E, Ueda M: Establishment and characterization of CAG/EGFP transgenic rabbit line. *Transgenic Res* 2007, **16**:115-120.
13. Niikura T, Hak DJ, Reddi AH: Global gene profiling reveals a downregulation of BMP gene expression in experimental atrophic nonunions compared to standard healing fractures. *J Orthop Res* 2006, **24**:1463-1471.
14. Willems E, Leyns L, Vandesompele J: Standardization of real-time PCR gene expression data from independent biological replicates. *Anal Biochem* 2008, **379**:127-129.
15. Saeed AI, Bhagabati NK, Braisted JC, Liang W, Sharov V, Howe EA, Li J, Thiagarajan M, White JA, Quackenbush J: TM4 microarray software suite. *Methods Enzymol* 2006, **411**:134-193.
16. Wakitani S, Goto T, Pineda SJ, Young RG, Mansour JM, Caplan AI, Goldberg VM: Mesenchymal cell-based repair of large, full-thickness defects of articular cartilage. *J Bone Joint Surg Am* 1994, **76**:579-592.
17. Sekiya I, Colter DC, Prockop DJ: BMP-6 enhances chondrogenesis in a subpopulation of human marrow stromal cells. *Biochem Biophys Res Commun* 2001, **284**:411-418.
18. Morito T, Muneta T, Hara K, Ju YJ, Mochizuki T, Makino H, Umezawa A, Sekiya I: Synovial fluid-derived mesenchymal stem cells increase after intra-articular ligament injury in humans. *Rheumatology (Oxford)* 2008, **47**:1137-1143.
19. Koga H, Muneta T, Ju YJ, Nagase T, Nimura A, Mochizuki T, Ichinose S, von der Mark K, Sekiya I: Synovial stem cells are regionally specified according to local microenvironments after implantation for cartilage regeneration. *Stem Cells* 2007, **25**:689-696.
20. Wisniewski HG, Hua JC, Poppers DM, Naime D, Vilcek J, Cronstein BN: TNF/IL-1-inducible protein TSG-6 potentiates plasmin inhibition by inter-alpha-inhibitor and exerts a strong anti-inflammatory effect *in vivo*. *J Immunol* 1996, **156**:1609-1615.
21. Maier R, Wisniewski HG, Vilcek J, Lotz M: TSG-6 expression in human articular chondrocytes. Possible implications in joint inflammation and cartilage degradation. *Arthritis Rheum* 1996, **39**:552-559.
22. Mindrescu C, Dias AA, Olszewski RJ, Klein MJ, Reis LF, Wisniewski HG: Reduced susceptibility to collagen-induced arthritis in DBA/1J mice expressing the TSG-6 transgene. *Arthritis Rheum* 2002, **46**:2453-2464.
23. Mindrescu C, Thorbecke GJ, Klein MJ, Vilcek J, Wisniewski HG: Amelioration of collagen-induced arthritis in DBA/1J mice by recombinant TSG-6, a tumor necrosis factor/interleukin-1-inducible protein. *Arthritis Rheum* 2000, **43**:2668-2677.
24. Block GJ, Ohkouchi S, Fung F, Frenkel J, Gregory C, Pochampally R, DiMattia G, Sullivan DE, Prockop DJ: Multipotent stromal cells are activated to reduce apoptosis in part by upregulation and secretion of stanniocalcin-1. *Stem Cells* 2009, **27**:670-681.
25. Huang L, Garcia G, Lou Y, Zhou Q, Truong LD, DiMattia G, Lan XR, Lan HY, Wang Y, Sheikh-Hamad D: Anti-inflammatory and renal protective actions of stanniocalcin-1 in a model of anti-glomerular basement membrane glomerulonephritis. *Am J Pathol* 2009, **174**:1368-1378.
26. Guo HD, Wang HJ, Tan YZ, Wu JH: Transplantation of marrow-derived cardiac stem cells carried in fibrin improves cardiac function after myocardial infarction. *Tissue Eng Part A* 2011, **17**:45-58.

doi:10.1186/ar3869

Cite this article as: Suzuki et al.: Properties and usefulness of aggregates of synovial mesenchymal stem cells as a source for cartilage regeneration. *Arthritis Research & Therapy* 2012 **14**:R136.

Arthroscopic, histological and MRI analyses of cartilage repair after a minimally invasive method of transplantation of allogeneic synovial mesenchymal stromal cells into cartilage defects in pigs

TOMOMASA NAKAMURA¹, ICHIRO SEKIYA², TAKESHI MUNETA^{1,3}, DAISUKE HATSUSHIKA¹, MASAFUMI HORIE¹, KUNIKAZU TSUJI³, TATSUO KAWARASAKI⁴, ATSUYA WATANABE⁵, SHUJI HISHIKAWA⁶, YASUHIRO FUJIMOTO⁶, HOZUMI TANAKA⁶ & EIJI KOBAYASHI⁷

¹Section of Orthopedic Surgery, Graduate School, Tokyo Medical and Dental University, Tokyo, Japan, ²Section of Cartilage Regeneration, Graduate School, Tokyo Medical and Dental University, Tokyo, Japan, ³Global Center of Excellence Program, International Research Center for Molecular Science in Tooth and Bone Diseases, Tokyo Medical and Dental University, Tokyo, Japan, ⁴Swine and Poultry Research Center, Shizuoka Prefectural Research Institute of Animal Industry, Shizuoka, Japan, ⁵Department of Orthopedic Surgery, Teikyo University Chiba Medical Center, Chiba, Japan, ⁶Center for Development of Advanced Medical Technology, Jichi Medical University, Tochigi, Japan, and ⁷Division of Development of Advanced Treatment, Jichi Medical University, Tochigi, Japan

Abstract

Background aims. Transplantation of synovial mesenchymal stromal cells (MSCs) may induce repair of cartilage defects. We transplanted synovial MSCs into cartilage defects using a simple method and investigated its usefulness and repair process in a pig model. **Methods.** The chondrogenic potential of the porcine MSCs was compared *in vitro*. Cartilage defects were created in both knees of seven pigs, and divided into MSCs treated and non-treated control knees. Synovial MSCs were injected into the defect, and the knee was kept immobilized for 10 min before wound closure. To visualize the actual delivery and adhesion of the cells, fluorescence-labeled synovial MSCs from transgenic green fluorescent protein (GFP) pig were injected into the defect in a subgroup of two pigs. In these two animals, the wounds were closed before MSCs were injected and observed for 10 min under arthroscopic control. The defects were analyzed sequentially arthroscopically, histologically and by magnetic resonance imaging (MRI) for 3 months. **Results.** Synovial MSCs had a higher chondrogenic potential *in vitro* than the other MSCs examined. Arthroscopic observations showed adhesion of synovial MSCs and membrane formation on the cartilage defects before cartilage repair. Quantification analyses for arthroscopy, histology and MRI revealed a better outcome in the MSC-treated knees than in the non-treated control knees. **Conclusions.** Leaving a synovial MSC suspension in cartilage defects for 10 min made it possible for cells to adhere in the defect in a porcine cartilage defect model. The cartilage defect was first covered with membrane, then the cartilage matrix emerged after transplantation of synovial MSCs.

Key Words: cartilage repair, mesenchymal stromal cells, pig, synovium

Introduction

Cartilage injuries are a common clinical problem and if left untreated may cause osteoarthritis, one of the leading causes of disability (1). Stem cell therapy for cartilage repair may be one possible strategy for improvement of cartilage injury. The candidate therapeutic cells are mesenchymal stromal cells (MSCs), which can be isolated from various mesenchymal tissues (2,3). We have reported previously the superiority of human synovial-derived MSCs for

cartilage repair (4–6) and *in vitro* expansion with autologous human serum (7).

Various methods have been used to transplant MSCs into cartilage defects, such as intra-articular injection (8,9) and the use of scaffolds (10). We have demonstrated recently that leaving the knee immobilized for 10 min immediately after delivering a suspension of synovial MSCs into the defect results in approximately 60% of the cells adhering to the defect to promote cartilage repair in rabbits (11).

Correspondence: Ichiro Sekiya, MD, PhD, Section of Cartilage Regeneration, Graduate School, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo, 113-8519 Japan. E-mail: sekiya.orj@tmd.ac.jp. Eiji Kobayashi, MD, PhD, Center for Development of Advanced Medical Technology, Jichi Medical University, 3311-1 Yakushiji, Shimotsuke, Tochigi 329-0498, Japan. E-mail: eijikoba@jichi.ac.jp

(Received 21 November 2011; accepted 5 December 2011)

ISSN 1465-3249 print/ISSN 1477-2566 online © 2012 Informa Healthcare
DOI: 10.3109/14653249.2011.638912

This 'local adherent technique' can be performed less invasively and without scaffolds compared with other methods.

We hypothesized that this method will also be useful in animals that are more closely related to humans. The purpose of the present study was to examine the usefulness of the local adherent technique with synovial MSCs in pigs. The knee joints of pigs are similar to those of humans in terms of size (12) and cartilage-specific properties (13). In this study, synovial MSCs were transplanted into the cartilage defect of pigs using the local adherent technique, and repaired cartilage was examined sequentially arthroscopically, histologically and by delayed gadolinium-enhanced magnetic resonance imaging of cartilage (dGEMRIC) (14,15).

Methods

Animals

All experiments were conducted in accordance with the institutional guidelines for the care and use of experimental animals of the Tokyo Medical and Dental University (Tokyo, Japan) and Jichi Medical University (Tochigi, Japan). Nine male and six female Mexican hairless pigs (National Livestock Breeding Center, Ibaraki, Japan) were used. They were 13 months old, on average 33.5 kg in weight, and skeletally mature, with the growth plates closed. All pigs were bred under specific pathogen-free conditions and had free access during the study period to food and water in a post-operative care cage (400 mm in width, 1210 mm in length and 1090 mm in height). One wild-type pig and one transgenic green fluorescent protein (GFP) pig (16) were used as donors for synovial MSC for transplantation. Two other pigs were also used as sources for MSCs for *in vitro* proliferation and differentiation assays. These four pigs were euthanized on the day when the tissues were harvested. Twelve other wild-type pigs were used as recipients. For GFP observation, two pigs were euthanized on the day MSCs were transplanted, and for observation of 1,1'-dioctadecyl-3,3,3',3'-tetramethylindocarbocyanine perchlorate (DiI; Molecular Probes, Eugene, OR, USA) two pigs were euthanized at 7 days after transplantation. For arthroscopic, histological and MRI analyses, three pigs were euthanized at 1 month, and five pigs were euthanized at 3 months, after transplantation.

Cell isolation and culture

Synovial tissue was harvested from the suprapatellar pouch, which overlays the non-cartilaginous areas of the femur, through an arthrotomy of the knee. The

tissue was digested in 3 mg/mL collagenase D solution (Roche Diagnostics, Mannheim, Germany) in α -minimal essential medium (α MEM; Invitrogen, Carlsbad, CA, USA) at 37°C for 3 h, filtered through a 70- μ m nylon filter (Becton-Dickinson and Co., Franklin Lakes, NJ, USA) and the nucleated cells plated in a 150-cm² culture dish (Nalge Nunc International, Rochester, NY, USA) in complete culture medium [α MEM containing 10% fetal bovine serum (FBS), 100 U/mL penicillin, 100 μ g/mL streptomycin and 250 ng/mL amphotericin B (all from Invitrogen)] and incubated at 37°C with 5% humidified CO₂. The medium was changed to remove non-adherent cells every 4–5 days and then cultured for 14 days as passage 0 without refeeding. To cryopreserve the cells, they were resuspended at a concentration of 2×10^6 cells/mL in α MEM with 5% dimethylsulfoxide (Wako, Osaka, Japan) and 10% FBS. Aliquots of 2 mL were frozen slowly in a Cryo 1°C freezing container (Nalge Nunc International) and cryopreserved at –80°C. To expand the cells, a frozen vial of the cells was thawed, plated in 60-cm² culture dishes, and incubated for 4 days. Then the cells were replated at 5×10^5 cells/150-cm² culture dish (passage 2) and cultured for an additional 14 days. The nucleated cells derived from periosteum, muscle and adipose tissue were isolated and expanded in the same manner as those from synovium.

Bone marrow was aspirated from the tibial tuberosity. Periosteum was peeled off from the tibia. Muscle was obtained from the quadriceps. Adipose tissue was prepared from the subcutaneous fat around the knee. Nucleated cells from the bone marrow were isolated with a density gradient (Ficoll-Paque; Amersham Biosciences, Uppsala, Sweden).

Colony-formation assay

Nucleated cells derived from synovium were plated at 0.5, 5, 50 and 500×10^3 cells/60-cm² dish, cultured for 14 days, and stained with crystal violet. The optimal initial cell density was determined based on the following criteria: (a) the colony size was not affected by contact inhibition, and (b) the greatest number of colonies was obtained. We then harvested the cells plated at optimal densities from the remaining dishes and expanded them as mentioned above.

In vitro proliferation assay

Synovial MSCs were plated at 5×10^3 cells/60-cm² dish in complete culture medium and passaged every 14 days. Cells from each passage were harvested and counted with a hemocytometer, and the total accumulated cell number was calculated.

In vitro differentiation assay

For chondrogenesis, 250 000 cells were placed in a 15-mL polypropylene tube (Becton-Dickinson and Co.) and centrifuged at 450 *g* for 10 min. The pellets were cultured in chondrogenesis medium consisting of high-glucose Dulbecco's modified Eagle's medium (Invitrogen) supplemented with 1 µg/mL bone morphogenetic protein (BMP)-7 (Stryker Biotech, Hopkinton, MA, USA), 10 ng/mL transforming growth factor (TGF)-β3 (R&D Systems, Minneapolis, MN, USA), 100 nM dexamethasone (Sigma-Aldrich Corp., St Louis, MO, USA), 50 µg/mL ascorbate-2-phosphate, 40 µg/mL proline, 100 µg/mL pyruvate and 1:100 diluted ITS + Premix (6.25 µg/mL insulin, 6.25 µg/mL transferrin, 6.25 ng/mL selenious acid, 1.25 mg/mL bovine serum albumin and 5.35 mg/mL linoleic acid; BD Biosciences Discovery Labware, Bedford, MA, USA). For microscopy, the pellets were embedded in paraffin, cut into 5-µm sections, and stained with toluidine blue (17–19).

For adipogenesis, cells were cultured in adipogenic medium, which consisted of complete medium supplemented with 100 nM dexamethasone (Sigma-Aldrich Corp.), 0.5 mM isobutyl-methylxanthine (Sigma-Aldrich Corp.) and 50 µM indomethacin (Wako), for 21 days. The adipogenic cultures were fixed in 4% paraformaldehyde and then stained with fresh Oil Red O solution (20).

For calcification, cells were cultured in calcification medium, which consisted of a complete medium of 1 nM dexamethasone, 20 mM β-glycerol phosphate (Wako) and 50 µg/mL ascorbate-2-phosphate (Sigma-Aldrich Corp.), for 21 days. The cells were fixed in 4% paraformaldehyde and stained with 0.5% Alizarin Red solution (21).

DiI labeling

Synovial MSCs were resuspended at 1×10^6 cells/mL in αMEM without FBS, and a fluorescent lipophilic tracer, DiI, was added at a final concentration of 5 µL/mL. After incubation for 20 min at 37°C and two washings with phosphate-buffered saline (PBS), DiI-labeled cells were resuspended in 100 µL culture medium (22).

Experimental set-up

The first pig was used for anatomical study and harvesting mesenchymal tissues to stock the MSCs for further analyses. When pigs for the *in vivo* study were prepared, cryopreserved synovial MSCs were thawed and expanded 2 weeks before transplantation. On the day of transplantation surgery, all colony-forming

cells were harvested and suspended in 100 µL culture medium and transplanted as described. Four pigs were used for an early adhesion assay with transplantation of GFP porcine synovial MSCs ($n = 2$) (Figure 2A) and DiI-labeled MSCs ($n = 2$). Other pigs were analyzed by arthroscopy every month, and two pigs were sacrificed at 1 month after treatment for histological, macroscopical and MRI analyses. Five pigs were sacrificed at 3 months after treatment and analyzed by histology and MRI (Figure 2D).

Transplantation of synovial MSCs into the cartilage defects

All pigs underwent general anesthesia, and the medial femoral condyle was approached through a medial parapatellar incision. Full-thickness osteochondral defects (8 × 8 mm square and 2 mm deep; approximately 1.5 mm cartilaginous and 0.5 mm bony part) were created with various sizes of drills in the weight-bearing area of the medial femoral condyles in both knees, 10 mm below the terminal ridge. When the defects were created, bleeding was not observed, and a procedure to stop bleeding from the bottom of the defect was not required.

The right knee of each pig was treated with MSCs and the left knee served as a vehicle internal control. The MSCs were harvested and collected from the culture dishes several hours before transplantation, and harvested MSCs were suspended in a 50-mL conical tube containing 40 mL culture medium. Just before the transplantation, the tube was centrifuged for 5 min at 1500 r.p.m., and the supernatant was removed. Centrifuged MSCs were suspended in 100 µL culture medium. The transplanted cell number was a maximum of 5.3×10^7 , a minimum of 2.2×10^7 , and on average 3.8×10^7 .

The cartilage defect was faced upward, and its position was held manually. A suspension of prepared MSCs in 100 µL culture medium was placed into the defect through an 18-gauge needle. Culture medium alone (100 µL) was placed into the defects in the left knee in the same manner. After 10 min, the incisions were closed without washing the inside of the knee joint. After the anesthetic wore off, the pigs were allowed to walk freely without fixation. To reduce the risk of infection, we avoided the use of an immune suppressor.

For euthanasia, an overdose intravenous injection of KCl was used under adequately deep general anesthesia. For macroscopic analyses, all samples at 1 month ($n = 3$) and 3 months ($n = 5$) were evaluated with the International Cartilage Repair Society (ICRS) macroscopic score (23) (see the supplementary tables).

Arthroscopy

All knees were observed with arthroscopy (Linvatec 8180A camera console surgical video equipment, with LIS8430 for the light source; Zimmer Inc., Warsaw, IN, USA) at 1, 2 and 3 months after transplantation. An arthroscope, a probe and a shaver system were inserted through longitudinal incisions at the medial and lateral sides of the patella tendon. All arthroscopic observations were evaluated by Oswestry arthroscopy score (23) (see the supplementary tables). For arthroscopic observation of GFP MSCs, a newly developed fluorescence arthroscope (Olympus Medical Systems Corp., Tokyo, Japan) was used.

Histological analyses

The samples were cut into a thickness of a 15 mm square with 5 mm containing a defect, fixed in 4% paraformaldehyde, and decalcified with 0.5 M ethylene diamine tetra acetic acid (EDTA; pH 7.5) for 3 days at 4°C. Paraffin sections were stained with Safranin O. All samples were evaluated with a modified Wakitani score (11) (see the supplementary tables).

dGEMRIC

Before histological analyses, medial femoral condyles were collected and pre-contrast MRI was performed. An MRI system at 1.5 Tesla (Signa HDx; GE Healthcare, Chalfont St Giles, UK) was used with a custom-made micro-imaging coil. Each specimen was pre-treated with 0.5 mm gadopentate dimeglumine (Gd-DTPA²⁻; Magnevist®; Schering, Berlin, Germany) in 0.9% normal saline overnight at 4°C with continuous stirring. The next day the samples were removed from refrigeration, and post-contrast MRI was performed at room temperature. R1 was defined as the reciprocal of the T1 value. The R1 measurement was performed using a fast-spin echo inversion-recovery (FSE-IR) sequence (2400 ms repetition time, 18 ms echo time, six inversion times of 50–2000 ms, 30×30 mm field of view, 1.0-mm section thickness, 512×512 matrix). The difference between the pre-Gd-enhanced R1 value and the post-Gd enhanced R1 value ($\Delta R1$) indicated the glycosaminoglycan (GAG) concentration (14). Color-coded $\Delta R1$ -calculated heat maps of the cartilage were generated using MATLAB (Mathworks, Natick, MA, USA) with a mono-exponential curve fit. Blue represents a high content of GAG, and red a low content. For R1 measurements, the region of interest (ROI) for repaired tissue was defined as the area where both sides were connected between native

and repaired cartilage; the bottom was the interface between bone and repaired cartilage, and the top was the superficial surface of the repaired cartilage. The ROI for native cartilage was drawn over the full-thickness weight-bearing areas of the femoral condyle at both sides of the repair site, about 3 mm from the lateral edge of the repair site (14,15).

Statistical analyses

To assess differences, Wilcoxon rank-sum tests were used except for MRI analysis. For MRI analysis, the paired *t*-test was used. A value of $P < 0.05$ was considered significant.

Results

Characteristics of porcine synovial cells as MSCs

The initial cell-plating density to produce the optimal colony number was determined to be 5×10^3 cells/60-cm² dish (Figure 1A). Three cell lineages derived from three different pigs maintained their proliferation potential over 20 passages (Figure 1B). Colony-forming cells derived from porcine synovium displayed a trilineage potential, differentiating into chondrocytes and adipocytes, and osteocytes, when cultured in their respective differentiation media (Figure 1C). *In vitro* chondrogenesis assays demonstrated that cartilage pellets of colony-forming cells derived from synovium were the heaviest among those derived from the other mesenchymal tissues (Figure 1D). These results indicated that colony-forming cells derived from porcine synovium had similar characteristics to those of MSCs, and the highest chondrogenic potential compared with cells derived from the other tissues examined.

Local adherent technique for transplantation of MSCs

After expanding for 14 days (Figure 2A), colony-forming cells derived from synovium of the transgenic GFP pig expressed GFP (Figure 2B). A drop of MSC suspension through a needle (Figure 2Ci) could be detected with the GFP arthroscopy system (Figure 2Cii). After placement of the MSC suspension for 10 min, the bottom of the cartilage defect looked foggy (Figure 2Ciii) and GFP MSCs were still detected in the cartilage defect (Figure 2Civ), even though the irrigation fluid was flushed from the tip of the arthroscope (see the supplementary movies). DiI-labeled MSCs were also traced (Figure 2D, E) and remained in the cartilage defect at 7 days (Figure 2F), but they could not be found at 1 and 3 months.