

human OCT4 promoter was digested with HindIII and AgeI, and then inserted into pEGFP-1 (Clontech).

2.2. Maintenance and transfection of hESCs

The hESC lines (KhES-1, KhES-3, and WA09 (H9)) were routinely cultured as described previously [6–8] on mitomycin C-treated MEF feeder cells in hESC medium consisting of DMEM/F12 (D-6421; Sigma) supplemented with 20% KnockOut Serum Replacement (KSR; Invitrogen), 0.1 mM non-essential amino acids (Sigma), 2 mM L-glutamine (Sigma), 0.1 mM 2-mercaptoethanol (Sigma), and 5 ng/ml bFGF (Wako, Japan). hESC medium without bFGF was used as hESC-basal medium (BM) in this study.

For feeder-free culture, hESCs were incubated with 2 mg/ml dispase (Invitrogen) in DMEM/F12 at 37 °C for 10 min, and then detached using a cell scraper. Small clumps of hESCs were seeded onto Matrigel-coated plates in CM or hESC-BM with each compound. The plates were pre-coated with 0.2 mg/ml Matrigel (growth factor reduced; BD Biosciences) at 4 °C overnight. The medium was removed, and the plates were washed with DMEM/F-12 to remove unbound Matrigel and then warmed to room temperature before use. CM was prepared as described previously [9] with the addition of 5 ng/ml bFGF.

Each compound, including small molecules from the Prestwick Chemical library, was added to hESC-BM at the indicated concentrations in the presence of 0.1% DMSO. Ethopropazine, promazine, retinoic acid (RA), PD98059, and trimipramine were purchased from Sigma, methotrimeprazine from Aurora Fine Chemicals LLC, and trimiprazine from the United States Pharmacopeial Convention.

Cell lines carrying hOCT4pro-EGFP were established by transfection with the hOCT4pro-EGFP reporter plasmid. Before transfection, KhES-1 cells were seeded onto Matrigel-coated 100 mm tissue culture dishes in CM. ApaI-linearized hOCT4pro-EGFP plasmid was transfected into KhES-1 cells using Fugene HD (Roche Diagnostics) according to the manufacturer's instructions. G418 selection (100 µg/ml) was applied at 24 h after transfection. After about 14 days of selection, the surviving colonies were picked up individually and expanded as clones.

2.3. High-content screening

For HCS, cells cultured on feeder cells were treated with a CTK solution consisting of 1 mg/ml collagenase IV (Invitrogen), 0.25% trypsin (Invitrogen), 1 mM CaCl₂ and 20% KSR, and then detached as small clumps [6]. The cells were seeded in hESC-BM into 96-well plates (Greiner Bio-One), and from the following day, hESC-BM containing 2 µg/ml (~5 µM) of each small molecule from the Prestwick Chemical library (in the presence of 0.1% DMSO) was changed daily. Each compound was assessed in quadruplicate. Control wells containing 0.1% DMSO in hESC-BM were included on each plate. After 6 days of culture, cells were fixed with 4% paraformaldehyde/PBS, washed with PBS, and then stained with 1 µg/ml 4',6-diamidino-2-phenylindole (DAPI; Invitrogen). After three washes with PBS, fluorescence images were acquired by an ArrayScan-VTI System (Cellomics) and processed with the Target Activation BioApplication (Cellomics). To normalize data, a z-score based on the EGFP fluorescence intensity was calculated for each medium using the negative control medium (hESC-BM containing 0.1% DMSO).

2.4. Semi-quantitative PCR

Total RNA was extracted using an RNeasy Micro Kit (Qiagen), and then 0.5–1 µg total RNA was reverse transcribed with an Omniscript RT Kit (Qiagen) according to the manufacturer's instructions. For semi-quantitative PCR analysis, PCR was

performed with TaKaRa ExTaq (TaKaRa, Japan). PCRs were optimized to allow semi-quantitative comparisons within the log phase of amplification. PCR products were separated on 2% agarose gels and visualized by ethidium bromide staining. Gene-specific primers are listed in Supplementary Table S1.

2.5. Teratoma formation assay

Approximately 2×10^6 cells were injected into the testes of severe combined immunodeficiency (SCID) mice (CLEA Japan). After 8 weeks, teratomas were surgically dissected from the mice, and then fixed in 4% paraformaldehyde/PBS. Samples were embedded in paraffin, sectioned at 5 µm and processed for hematoxylin and eosin staining. Animal protocols were approved by the Institutional Board on Animal Care at Kyoto University.

2.6. Statistical analysis

Data are shown as the average ± standard deviation (SD). Statistical significance was assessed using the Student's *t*-test. The probability level accepted for significance was $P < 0.05$.

2.7. Other methods

See Supplementary Methods for remaining methods including flow cytometric analysis, immunocytochemistry, Karyotype analysis, the EdU incorporation assay, the TUNEL assay and *in vitro* differentiation assay.

3. Results

3.1. High-throughput chemical screening to identify promotion of hESC self-renewal

To carry out a screen, we first established a human OCT4-GFP reporter system in hESCs, which contained –3917 to –1 base pairs relative to the transcriptional start site [5] (Fig. 1A). OCT4 is highly expressed in hESCs and downregulated upon differentiation. We initially isolated a 3D6 hESC clone, which showed GFP expression in an undifferentiated state, and was morphologically indistinguishable from the parental KhES1 cells (Fig. 1A). Furthermore, GFP expression in the cells was lost upon differentiation by 5 days of 10 µM RA treatment, as indicated by both fluorescence microscopy (Supplementary Fig. S1A) and flow cytometric analysis (Supplementary Fig. S1B). Moreover, flow cytometric analysis showed that OCT4-GFP expression was well correlated with OCT4 expression.

Undifferentiated 3D6 hESCs were seeded onto Matrigel-coated 96-well plates at a density of 3000 cells/well in hESC-BM. After overnight incubation, a compound from the chemical library, as described in Section 2, was added to each well (2 µg/ml, ~5 µM). Medium containing compounds was changed daily for a further 5 days of incubation. Cells were analyzed for GFP expression using an Arrayscan VTI system (Cellomics).

Before performing our screening, to confirm whether the intensity of GFP fluorescence indicated the status of hESCs, we used CM, bFGF and mTeSR1 medium (StemCell Technologies) as controls for promotion of self-renewal, and RA or PD98059 as controls for induction of differentiation (Fig. 1B, and data not shown). The *z'*-factor is a parameter in statistics to assess the performance in high-throughput screening [10]. A *z'*-factor of >0.5 was routinely obtained using this assay system, thereby supporting our conclusion that the intensity of GFP fluorescence using the hOCT4pro-EGFP reporter gene system could reliably identify small molecules that maintained hESCs in an undifferentiated state.




Fig. 1. High-content screening system and characterization of hit compounds. (A) (Left) Vector map of hOCT4pro-EGFP. (Right) Phase contrast and fluorescence images of parental KhES-1 and hOCT4pro-EGFP transfectant (3D6 clone) hESCs cultured on MEF feeders. Bar indicates 200 μ m. (B) (upper) Representative fluorescence image of EGFP in 3D6 clones cultured in various media (CM, hESC-BM containing 0.1% DMSO, and hESC-BM containing 10 μ M RA) for 5 days. Bar indicates 500 μ m. (lower) Calculation of GFP intensity in upper fluorescence images. A.U. corresponds to arbitrary unit. Data are presented as the means \pm SD. (C) Five of the 18 hit small molecules from the 1120-chemical library. The small molecules shared a z-score of higher than 2 SD, as well as structural and physiological similarities. (D) Representative fluorescence images of 3D6 hESCs cultured in hESC-BM with the five hit compounds. Expression of markers of the undifferentiated state was observed by EGFP reporter gene expression driven by the hOCT4 promoter (hOCT4pro-EGFP; green) in immunocytochemistry. Cells were identified by DAPI (blue). Bar indicates 500 μ m. CM, conditioned medium. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Using this assay, we screened 1120 compounds from Prestwick libraries, and 18 hit compounds (1.6% of the total chemicals screened; listed in Supplementary Table S2) that had effects on the self-renewal of hESCs were identified by exhibiting higher OCT4-GFP expression by more than two SDs. Interestingly, we found that five out of the 18 chemicals were neurotransmitter antagonists that may share structural and physiological similarities, namely methotrimeprazine, trimipramine, trimeprazine, ethopropazine, and promethazine (Fig. 1C and D). In this study, we used trimipramine and ethopropazine for subsequent experiments. HCS using OCT4 immunostaining further showed that almost of the cells exhibited OCT4^{high} in CM as a control for the undifferentiated state (98.9%). In contrast, DMSO strongly decreased the OCT4^{high} population to 7.6%, whereas trimipramine and ethopropazine were effective for recovery of the OCT4^{high} population to 41.0% and 16.1%, respectively (Fig. 2A). The effect became more noticeable over several passages. Both trimipramine and ethopropazine maintained KhES-3 and WA09 hESCs in an undifferentiated state, whereas DMSO treatment resulted in continuous differentiation of the cells (Fig. 2B). Although we initially screened these chemicals using KhES-1 hESCs, the cells were differentiated even in the presence of the chemicals after several passages (data not shown). In addition, OCT4-GFP or OCT4 expression in the cells, which was

increased by the addition of chemicals to the medium, was still lower than that in cells cultured in CM (Figs. 1D and 2A).

3.2. Effect of trimipramine and ethopropazine on long-term culture of hESCs

We found that two hESC lines (KhES-3 and WA09) cultured on Matrigel in hESC-BM containing trimipramine or ethopropazine proliferated for at least 20 passages without bFGF in the medium. Under these culture conditions, the cells showed compact colonies (Fig. 3A), which were similar to the undifferentiated colonies cultured in CM generally used as gold standard. We karyotyped 50 randomly selected cells by G-banding of both cell lines after 20 passages. Cells cultured with trimipramine or ethopropazine had normal karyotypes (Fig. S2A). Furthermore, RT-PCR analysis revealed that hESCs maintained high expression levels of pluripotency markers including OCT4 and NANOG, while the expression of differentiation marker genes was suppressed (Fig. 3B). Immunofluorescence analysis showed that the cells maintained the expression of pluripotency markers OCT4, NANOG and SOX2 (Fig. 3C). Flow cytometric analysis further indicated that the majority of hESCs cultured with these chemicals expressed pluripotency markers SSEA-3, SSEA-4, TRA-1-60, TRA-1-81, and OCT4 with only




Fig. 2. Small molecules support short-term culture of hESCs without bFGF in the medium. (A) Histograms of the intensity of OCT4 immunostaining in hESCs treated with CM, DMSO, or small chemicals. (B) Morphology of KhES-3 and WA09 hESCs cultured in hESC-BM with trimipramine or ethopropazine after several passages, respectively. DMSO caused differentiation of hESCs, whereas CM and small molecule-containing media maintained the undifferentiated state of hESCs.

low-level of expression of the differentiation marker SSEA-1 (Fig. 3D). Thus, trimipramine and ethopropazine may become substitutes for bFGF to maintain hESCs in an undifferentiated state. Furthermore, we found that trimipramine and ethopropazine treatments resulted in a similar proliferation rate of both hESC lines compared with those cultured in CM (Fig. 3E). We could not detect significant differences between trimipramine or ethopropazine in hESC-BM and CM in terms of cell cycling using a 5-ethyl-2'-deoxyuridine (EdU) uptake assay (S-phase) (Supplementary Fig. S2B). In addition, the chemicals did not induce apoptosis of hESCs using the TUNEL assay, although they showed insignificantly increased apoptosis of KhES-3 hESCs compared with those cultured in CM (Supplementary Fig. S2C). Together, we concluded that hESCs cultured with trimipramine or ethopropazine showed similar behavior compared with those cultured in CM.

Finally, we confirmed the pluripotency of hESCs that had been expanded in the presence of trimipramine or ethopropazine by

examining their ability to form teratomas comprised of all three germ layers including neuroepithelium (ectoderm), intestinal epithelium (endoderm) and cartilage (mesoderm) (Fig. 4A). To assess the pluripotency of hESCs *in vitro*, the cells were cultured in a differentiation medium described in the Supplementary Methods. We found that hESCs could differentiate into ectoderm (β -III tubulin), mesoderm (α -smooth muscle actin (SMA)) and endoderm (α -fetoprotein (AFP)) as determined by immunohistochemical analyses (Fig. 4B).

4. Discussion

Small molecules are different from proteins in terms of molecular size, and can penetrate multi-layer tissues easily. Therefore, small molecules are expected to be more efficient for maintenance of the undifferentiated state of hESCs and hiPSCs to subsequently




Fig. 3. Small molecules support long-term culture of undifferentiated hESCs without bFGF in the medium. (A) Phase contrast micrographs of KhES-3 and WA09 hESCs cultured in hESC-BM with trimipramine or ethopropazine after 25 passages. Bar indicates 200 μm . (B) Expression of pluripotency marker genes (*OCT4* and *NANOG*) and differentiation marker genes (ectoderm: *LHX2* and *DLK1*; endoderm: *GATA6* and *GATA4*; mesoderm: *CDH5* and *FOXF1*; trophoctoderm: *CDX2* and *CGA*) was analyzed by RT-PCR. (C) Immunocytochemical detection of pluripotency marker genes (*OCT4*, *SOX2*, and *NANOG*) in KhES-3 and WA09 hESCs cultured in hESC-BM with trimipramine or ethopropazine after 40 passages. Bar indicates 100 μm . (D) Flow cytometric analysis of markers of the undifferentiated state (*OCT4*, *SSEA-3*, *SSEA-4*, *TRA-1-60*, and *TRA-1-81*) and a differentiation marker (*SSEA-1*) after 40 passages of culture in CM or hESC-BM with trimipramine or ethopropazine. (E) Growth rates of KhES-3 and WA09 hESCs cultured in CM or hESC-BM with trimipramine or ethopropazine during 4 days of culture after seeding. Cell numbers were counted every 24 h. The data represent means \pm SD.

induce specific differentiation. For example, Y-27632 has been identified as effective for maintaining the survival of dissociated hiPSCs [11]. Several small chemicals have also been identified to promote the generation of iPSCs from somatic cells [12–16]. Moreover, (–)-indolactam V has been found to promote pancreatic differentiation of human pluripotent cells [17]. Here, we show that some neurotransmitter antagonists support the expansion of undifferentiated hESCs without bFGF in the medium.

Although a previous study using a HCS system identified small molecules that regulate undifferentiated proliferation of hESCs, the small molecules used in the study cannot support long-term

culture [18]. In this study, we first addressed following points to develop a more reliable HCS system for hESCs. Previous HCS systems evaluate the undifferentiated state of hESCs by the expression of *OCT4* or *SSEA-3* using immunohistochemistry [18–20]. Instead, we used a reporter gene system consisting of GFP regulated by the h*OCT4* promoter, which enables evaluation of *OCT4* expression in cells directly without immunostaining procedures. Furthermore, we assessed not only the score of the effect of each small molecule, but also their similarities in terms of structure and pharmacological effects for hit compounds previously proposed by Lukaszewicz et al. [21]. This strategy may enable more reliable HCS for the




Fig. 4. Small molecules sustain the pluripotency of KhES-3 and WA09 hESCs without bFGF in the medium. (A) Characterization of teratomas derived from KhES-3 and WA09 hESCs cultured with trimipramine or ethopropazine. Hematoxylin and eosin staining of paraffin-embedded teratoma sections identified hESC differentiation into various tissues including neural pigment (ectoderm), a gut-like tube (endoderm), and cartilage (mesoderm). Bar indicates 100 μ m. (B) Immunostaining for markers of the three germ layers in differentiating KhES-3 and WA09 hESCs cultured with hit compounds after 40 passages: ectoderm (β III-tubulin), endoderm (AFP), and mesoderm (α -SMA).

identification of small molecules that drive hESC self-renewal as a replacement for bFGF.

In this study, we show that trimipramine and ethopropazine permit long-term hESC self-renewal without bFGF in the medium. We examined three hESCs lines (KhES-1, KhES-3, and WA09) and found that KhES-3 and WA09, but not KhES-1, hESCs successfully maintain undifferentiated expansion in the presence of these small molecules. Indeed, as indicated in Figs. 1D and 2A, OCT4 expression increased in the presence of the small molecules compared with that of DMSO, but OCT4 expression was still lower than that in CM. Therefore, the small molecules were not fully effective for the maintenance of undifferentiated proliferation of some hESC lines. We examined higher concentrations of the small molecules, but they showed cytotoxicity at such concentrations (data not shown). Taken together, further improvement of the small molecules to reduce their cytotoxicity while increasing the self-renewal of many hESCs lines needs to be achieved for complete replacement of bFGF in media.

The FGF pathway is activated by FGF ligands binding to FGF receptors, which in turn may trigger activation of various downstream signaling pathways including MAP kinase, PI3K kinase/AKT and PLC/PKC pathways [22]. However, the molecular mechanism by which bFGF promotes undifferentiated expansion of hESCs is still unclear. A MAP kinase kinase (MEK) inhibitor, PD0325901, is reported to be effective for hESC maintenance [23]. Activation of the PLC/PKC pathway may induce hESC differentiation [24]. PIK3/AKT signaling activity has been reported to participate in undifferentiated proliferation by suppression of MAP and canonical Wnt signaling pathways [25]. Therefore, the PIK3/AKT pathway may be a potential target of bFGF signaling. However, it remains unclear whether such small molecules related to PIK3/AKT pathways are ideal replacements for bFGF. Thus, small molecules related to growth factors and their signaling pathways have not been fully identified for the replacement of bFGF to expand undifferentiated hESCs in culture.

In this study, we showed that trimipramine and ethopropazine permit long-term hESC self-renewal without bFGF in the medium. Further studies will clarify the molecular mechanisms of trimipramine and ethopropazine, which enable long-term expansion of hESCs, including the interactions between these chemicals and the FGF or PIK3/AKT pathways.

For feeder-free culture systems, CM is usually used as the current gold standard. In this study, we showed that trimipramine and ethopropazine treatments resulted in similar growth curves and cell cycles as those of hESCs cultured in CM. Thus, the small molecules enabled similar performance compared with that of CM for the expansion of undifferentiated hESCs.

In conclusion, we established a HCS platform to identify small chemicals that promote hESC self-renewal. Using the screening system, we identified trimipramine and ethopropazine as novel

small molecules that promote long-term hESC self-renewal without bFGF in the medium. This screening system may help to identify more small molecules to achieve robust proliferation of hESCs and hiPSCs in low-cost and growth factor-free medium.

Acknowledgments

We thank Ms. Mari Hamao and Ms. Ritsuko Goto for their valuable technical assistance and the members of the laboratories of N. Nakatsuji for insightful discussions. We also thank Mr. Yoshishiro Koyama for his valuable technical assistance with the HCS analysis. This work was supported by the New Energy and Industrial Technology Development Organization (NEDO) of Japan, and partially supported by a Grant-in-Aid for Scientific Research (to E.K.) and the “Funding Program for Next Generation World-Leading Researchers” (to M.U.). H.K. was supported, in part, as a predoctoral fellow for the Global COE Program “Center for Frontier Medicine” by the Ministry of Education, Culture, Sports, Science, and Technology (MEXT), Japan. iCeMS is supported by the World Premier International Research Center Initiative (WPI), MEXT, Japan.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.bbrc.2013.03.061>.

References

- [1] J.A. Thomson, J. Itskovitz-Eldor, S.S. Shapiro, M.A. Waknitz, J.J. Swiergiel, V.S. Marshall, J.M. Jones, Embryonic stem cell lines derived from human blastocysts, *Science* 282 (1998) 1145–1147.
- [2] K. Takahashi, K. Tanabe, M. Ohnuki, M. Narita, T. Ichisaka, K. Tomoda, S. Yamanaka, Induction of pluripotent stem cells from adult human fibroblasts by defined factors, *Cell* 131 (2007) 861–872.
- [3] J. Yu, M.A. Vodyanik, K. Smuga-Otto, J. Antosiewicz-Bourget, J.L. Frane, S. Tian, J. Nie, G.A. Jonsdottir, V. Ruotti, R. Stewart, Slukvin II, J.A. Thomson, Induced pluripotent stem cell lines derived from human somatic cells, *Science* 318 (2007) 1917–1920.
- [4] V. Akopian, P.W. Andrews, S. Beil, N. Benvenisty, J. Brehm, M. Christie, A. Ford, V. Fox, P.J. Gokhale, L. Healy, F. Holm, O. Hovatta, B.B. Knowles, T.E. Ludwig, R.D. McKay, T. Miyazaki, N. Nakatsuji, S.K. Oh, M.F. Pera, J. Rossant, G.N. Stacey, H. Suemori, Comparison of defined culture systems for feeder cell free propagation of human embryonic stem cells, *In Vitro Cell Dev. Biol. Anim.* 46 (2010) 247–258.
- [5] L. Gerrard, D. Zhao, A.J. Clark, W. Cui, Stably transfected human embryonic stem cell clones express OCT4-specific green fluorescent protein and maintain self-renewal and pluripotency, *Stem Cells* 23 (2005) 124–133.
- [6] H. Suemori, K. Yasuchika, K. Hasegawa, T. Fujioka, N. Tsuneyoshi, N. Nakatsuji, Efficient establishment of human embryonic stem cell lines and long-term maintenance with stable karyotype by enzymatic bulk passage, *Biochem. Biophys. Res. Commun.* 345 (2006) 926–932.
- [7] T. Miyazaki, S. Futaki, K. Hasegawa, M. Kawasaki, N. Sanzen, M. Hayashi, E. Kawase, K. Sekiguchi, N. Nakatsuji, H. Suemori, Recombinant human laminin isoforms can support the undifferentiated growth of human embryonic stem cells, *Biochem. Biophys. Res. Commun.* 375 (2008) 27–32.

- [8] T. Miyazaki, S. Futaki, H. Suemori, Y. Taniguchi, M. Yamada, M. Kawasaki, M. Hayashi, H. Kumagai, N. Nakatsuji, K. Sekiguchi, E. Kawase, Laminin E8 fragments support efficient adhesion and expansion of dissociated human pluripotent stem cells, *Nat. Commun.* 3 (2012) 1236.
- [9] C. Xu, M.S. Inokuma, J. Denham, K. Golds, P. Kundu, J.D. Gold, M.K. Carpenter, Feeder-free growth of undifferentiated human embryonic stem cells, *Nat. Biotechnol.* 19 (2001) 971–974.
- [10] J.H. Zhang, T.D. Chung, K.R. Oldenburg, A simple statistical parameter for use in evaluation and validation of high throughput screening assays, *J. Biomol. Screen* 4 (1999) 67–73.
- [11] K. Watanabe, M. Ueno, D. Kamiya, A. Nishiyama, M. Matsumura, T. Wataya, J.B. Takahashi, S. Nishikawa, K. Muguruma, Y. Sasai, A ROCK inhibitor permits survival of dissociated human embryonic stem cells, *Nat. Biotechnol.* 25 (2007) 681–686.
- [12] Y. Shi, J.T. Do, C. Desponts, H.S. Hahm, H.R. Scholer, S. Ding, A combined chemical and genetic approach for the generation of induced pluripotent stem cells, *Cell Stem Cell* 2 (2008) 525–528.
- [13] T. Lin, R. Ambasudhan, X. Yuan, W. Li, S. Hilcove, R. Abujarour, X. Lin, H.S. Hahm, E. Hao, A. Hayek, S. Ding, A chemical platform for improved induction of human iPSCs, *Nat. Methods* 6 (2009) 805–808.
- [14] D. Huangfu, R. Maehr, W. Guo, A. Eijkelenboom, M. Snitow, A.E. Chen, D.A. Melton, Induction of pluripotent stem cells by defined factors is greatly improved by small-molecule compounds, *Nat. Biotechnol.* 26 (2008) 795–797.
- [15] C.A. Lyssiotis, R.K. Foreman, J. Staerk, M. Garcia, D. Mathur, S. Markoulaki, J. Hanna, L.L. Lairson, B.D. Charette, L.C. Bouchez, M. Bollong, C. Kunick, A. Brinker, C.Y. Cho, P.G. Schultz, R. Jaenisch, Reprogramming of murine fibroblasts to induced pluripotent stem cells with chemical complementation of Klf4, *Proc. Natl. Acad. Sci. USA* 106 (2009) 8912–8917.
- [16] Z. Li, T.M. Rana, A kinase inhibitor screen identifies small-molecule enhancers of reprogramming and iPSC cell generation, *Nat. Commun.* 3 (2012) 1085.
- [17] S. Chen, M. Borowiak, J.L. Fox, R. Maehr, K. Osafune, L. Davidow, K. Lam, L.F. Peng, S.L. Schreiber, L.L. Rubin, D. Melton, A small molecule that directs differentiation of human ESCs into the pancreatic lineage, *Nat. Chem. Biol.* 5 (2009) 258–265.
- [18] S.C. Desbordes, D.G. Placantonakis, A. Ciro, N.D. Socci, G. Lee, H. Djaballah, L. Studer, High-throughput screening assay for the identification of compounds regulating self-renewal and differentiation in human embryonic stem cells, *Cell Stem Cell* 2 (2008) 602–612.
- [19] R. Damoiseaux, S.P. Sherman, J.A. Alva, C. Peterson, A.D. Pyle, Integrated chemical genomics reveals modifiers of survival in human embryonic stem cells, *Stem cells* 27 (2009) 533–542.
- [20] I. Barbaric, M. Jones, D.J. Harley, P.J. Gokhale, P.W. Andrews, High-content screening for chemical modulators of embryonal carcinoma cell differentiation and survival, *J. Biomol. Screen* 16 (2011) 603–617.
- [21] A.I. Lukaszewicz, M.K. McMillan, M. Kahn, Small molecules and stem cells. Potency and lineage commitment: the new quest for the fountain of youth, *J. Med. Chem.* 53 (2010) 3439–3453.
- [22] K. Dorey, E. Amaya, FGF signalling: diverse roles during early vertebrate embryogenesis, *Development* 137 (2010) 3731–3742.
- [23] H. Tsutsui, B. Valamehr, A. Hindoyan, R. Qiao, X. Ding, S. Guo, O.N. Witte, X. Liu, C.M. Ho, H. Wu, An optimized small molecule inhibitor cocktail supports long-term maintenance of human embryonic stem cells, *Nat. Commun.* 2 (2011) 167.
- [24] X. Feng, J. Zhang, K. Smuga-Otto, S. Tian, J. Yu, R. Stewart, J.A. Thomson, Protein kinase C mediated extraembryonic endoderm differentiation of human embryonic stem cells, *Stem cells* 30 (2012) 461–470.
- [25] A.M. Singh, D. Reynolds, T. Cliff, S. Ohtsuka, A.L. Mattheyses, Y. Sun, L. Menendez, M. Kulik, S. Dalton, Signaling network crosstalk in human pluripotent cells: a Smad2/3-regulated switch that controls the balance between self-renewal and differentiation, *Cell Stem Cell* 10 (2012) 312–326.

ARTICLE

Received 30 Jul 2012 | Accepted 29 Oct 2012 | Published 4 Dec 2012

DOI: 10.1038/ncomms2231

OPEN

Laminin E8 fragments support efficient adhesion and expansion of dissociated human pluripotent stem cells

Takamichi Miyazaki¹, Sugiko Futaki², Hirofumi Suemori¹, Yukimasa Taniguchi², Masashi Yamada², Miwa Kawasaki², Maria Hayashi², Hideaki Kumagai¹, Norio Nakatsuji^{3,4}, Kiyotoshi Sekiguchi² & Eihachiro Kawase¹

Human embryonic stem cells (hESCs) and induced pluripotent stem cells (hiPSCs) have the potential to provide an infinite source of tissues for regenerative medicine. Although defined xeno-free media have been developed, culture conditions for reliable propagation of hESCs still require considerable improvement. Here we show that recombinant E8 fragments of laminin isoforms (LM-E8s), which are the minimum fragments conferring integrin-binding activity, promote greater adhesion of hESCs and hiPSCs than do Matrigel and intact laminin isoforms. Furthermore, LM-E8s sustain long-term self-renewal of hESCs and hiPSCs in defined xeno-free media with dissociated cell passaging. We successfully maintained three hESC and two hiPSC lines on LM-E8s in three defined media for 10 passages. hESCs maintained high level expression of pluripotency markers, had a normal karyotype after 30 passages and could differentiate into all three germ layers. This culture system allows robust proliferation of hESCs and hiPSCs for therapeutic applications.

¹Department of Embryonic Stem Cell Research, Institute for Frontier Medical Sciences, Kyoto University, 53 Kawahara-cho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan. ²Laboratory of Extracellular Matrix Biochemistry, Institute for Protein Research, Osaka University, 3-2 Yamadaoka, Suita, Osaka 565-0871, Japan. ³Department of Development and Differentiation, Institute for Frontier Medical Sciences, Kyoto University, 53 Kawahara-cho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan. ⁴Institute for Integrated Cell-Material Sciences (WPI-iCeMS), Kyoto University, Ushinomiya-cho, Yoshida, Sakyo-ku, Kyoto 606-8501, Japan. Correspondence should be addressed to E.K. (email: kawase8@frontier.kyoto-u.ac.jp) for general correspondence or K.S. (email: sekiguch@protein.osaka-u.ac.jp) for recombinant laminin products.

Human embryonic stem cells (hESCs) and induced pluripotent stem cells (hiPSCs) have great potential for a wide range of applications in regenerative medicine, and as tools for human disease modelling and drug discovery. However, defined xeno-free and feeder-free systems have not been fully established for the realization of these applications. Several types of defined medium have been developed, which consist of known components and avoid the use of animal products^{1–5}. However, the widespread use of Matrigel as a culture substrate is potentially problematic⁶. Matrigel is not an optimal substrate because it is derived from mouse Engelbreth–Holm–Swarm tumours and contains many unknown components. Additionally, hESCs usually propagate as colonies in culture, and their passaging involves either enzymatic dissociation by gently pipetting or manual microdissection of the cells⁷. hESC colonies grow as a monolayer and sustain an undifferentiated state, whereas the formation of large aggregates during passaging accelerates their spontaneous differentiation. Thus, it is somewhat difficult to precisely control appropriate dissociation of hESCs while passaging. As a result, conventional colony-based protocols often have considerable variations in culture outcomes. One method to overcome such difficulties might be complete dissociation of hESCs by an optimized and standardized protocol. However, complete dissociation of hESC clumps into single cells causes extensive cell death, and the survival is generally <1% (refs 8,9). Furthermore, complete cell dissociation is thought to be associated with an increased risk of karyotypic abnormalities^{10,11}. In this study, we aimed to identify an optimal substrate with the greatest adhesive property for the culture of completely dissociated hESCs and hiPSCs, and we further examined whether it could promote robust propagation of these cells while retaining a normal karyotype.

Although there are reports of the replacement of Matrigel with defined substrates, such as small peptides, to allow sustained hESC self-renewal in defined or xeno-free media^{3–5,12–18}, considerable improvements and refinements of these substrates are still needed. Laminin, the main component of Matrigel, is a heterotrimeric glycoprotein composed of three covalently linked chains that are termed α , β and γ ¹⁹. Thus far, 15 laminin isoforms formed by various combinations of each chain ($\alpha 1$ – $\alpha 5$, $\beta 1$ – $\beta 3$ and $\gamma 1$ – $\gamma 3$) have been identified. We have previously shown that human recombinant laminin isoforms (laminin-511, -332 and -111) support hESC proliferation in an undifferentiated state, although laminin-111 shows weak cell adhesive properties²⁰. Laminin-511, the first and only alternative substrate with a greater adhesive property for hESCs than that provided by Matrigel, also permits sustained self-renewal of hESCs in xeno-free culture conditions¹². Typically, procedures for the isolation of native pure laminins from tissues are arduous or highly impractical. Recently, some recombinant laminins have been produced using mammalian expression systems^{21–25}. However, full-length laminins are large heterotrimeric proteins (Fig. 1a), and mass production of these recombinant proteins is still very difficult. Therefore, we examined whether recombinant laminin fragments could support adhesion and undifferentiated proliferation of hESCs and hiPSCs. These fragments are obviously smaller molecules and have a much higher yield compared with that of the full-length protein.

Laminin isoforms bind to multiple receptors on the cell surface, including integrins, syndecans and dystroglycans, of which integrins serve as the major adhesion receptors of hESCs²⁶. Laminin-511, -332 and -111 share the characteristic of specific high affinity for $\alpha 6\beta 1$ integrin²⁷. Using function-blocking antibodies, $\alpha 6\beta 1$ integrin has been identified as the major cell adhesion receptor for both Matrigel and laminin-511 (refs 12,28). Furthermore, $\alpha 6\beta 1$ integrin is abundantly expressed by both

hESCs²⁰ and hiPSCs (Supplementary Fig. S1). Thus, we examined whether the integrin-binding site in laminin is solely necessary for the maintenance of hESCs and hiPSCs.

Laminin E8 fragments (LM-E8s) are truncated proteins composed of the C-terminal regions of the α , β and γ chains. These truncated proteins contain the active integrin-binding site comprising the laminin globular 1–3 domains of the α chain²¹ and the glutamate residue in the C-terminal tail of the γ chain²⁹, but lack other activities such as the heparin/heparan sulphate-binding activity, which are associated with intact laminins²¹. Therefore, LM-E8s serve as a functionally minimal form that retains the full capability for binding to $\alpha 6\beta 1$ integrin³⁰. When LM-E8s are separated into an α chain and a disulphide-bonded dimer of β and γ chains, they lose their cell adhesive activity³¹, underscoring the requirement for all three chains for their integrin-binding activity. Thus, the integrin-binding site of laminin appears to be a composite epitope composed of multiple polypeptides³², thereby making it difficult to recapitulate the integrin-binding site with synthetic peptides modelled on individual laminin subunits.

In this study, we produced two recombinant LM-E8s, LM511-E8 from laminin-511 and LM332-E8 from laminin-332, both of which have high cell adhesive properties (Fig. 1a), and examined their capabilities as adhesion substrates for hESCs and hiPSCs. Both fragments display stronger adhesion affinities for hESCs and hiPSCs than those of other substrates. Furthermore, LM-E8s enable robust propagation of hESCs and hiPSCs in an undifferentiated state in cultures with defined and xeno-free media using dissociated cell passaging.

Results

LM-E8s maximize hESC and hiPSC adhesion. Extracellular matrix (ECM) proteins vary considerably in their affinities for receptors, and in protein size and charge. Consequently, different proteins vary in their optimal use as a cell culture support matrix. To determine whether LM511-E8 and LM332-E8 were suitable for hESC and hiPSC adhesion, we first compared the adhesive properties of completely dissociated H9 hESCs and iPSC(IMR90)-1 cells in mTeSR1 medium on matrices containing LM511-E8 or LM332-E8 with those on other matrices. Cell adhesion on all matrices showed dose-dependent increases, but with different maxima: $15 \mu\text{g cm}^{-2}$ for Matrigel; $3\text{--}6 \mu\text{g cm}^{-2}$ for intact laminin-511 and -332; and $1.5 \mu\text{g cm}^{-2}$ for LM511-E8 and LM332-E8 (Fig. 1b and Supplementary Figs S2a and S3a). In agreement with a previous report¹², adhesion to intact laminin-511 or -332 was higher than that to Matrigel. However, we found that LM511-E8 and LM332-E8 had significantly higher adhesive properties than those of intact laminin-511 and -332. We therefore used LM511-E8 and LM332-E8 at a coating concentration of $1.5 \mu\text{g cm}^{-2}$ in subsequent experiments.

To confirm that LM511-E8 and LM332-E8 were recognized by $\alpha 6\beta 1$ integrin on the cell surface, we used integrin function-blocking antibodies to investigate whether $\alpha 6$ and/or $\beta 1$ integrins were required for the adhesion of H9 hESCs to LM-E8-coated substrata in mTeSR1 medium (Fig. 1c; Statistical results of multiple comparisons are shown in Supplementary Tables S1a and S1b). Cell attachment to both LM-E8s was significantly reduced by the addition of antibodies against $\alpha 6$ and/or $\beta 1$ integrins. We further confirmed that an inactive form of LM511-E8 (LM511(EQ)), in which the glutamic acid residue (E) at the third position from the C-terminus in the γ chain was replaced by a glutamine residue (Q)²⁹, showed dramatic abrogation of cell adhesion (Fig. 1c). Taken together, these data showed that $\alpha 6\beta 1$ integrin mediated the adhesion of completely dissociated hESCs to LM511-E8 and LM332-E8.




Figure 1 | Validation of LM-E8s for H9 hESC attachment. (a) Schematic representations of the two LM-E8s and the original intact laminin isoforms. A 6xHis-tag and FLAG-tag were attached to the N-termini of the α and γ chains, respectively, of LM-E8s to facilitate the purification of recombinant LM-E8s. The molecular weights of individual proteins are shown in parentheses. (b) Dose-response adhesion curves of H9 hESCs on various ECMs. LM-E8s at $1.5 \mu\text{g cm}^{-2}$ showed higher efficiency for supporting the adhesion of dissociated cells than did other matrices. The means of absorbance (OD570) represent the relative number of adherent cells normalized against the values at the maximum effect on Matrigel, which was arbitrarily set as 1. Error bars show the s.e.m. of five independent assays, except for vitronectin, which shows the average of duplicate assays. (c) Inhibition of integrins on LM-E8s. Completely dissociated H9 hESCs were incubated with an integrin-specific blocking antibody for 30 min in mTeSR1 medium. LM511(EQ); an inactive form of LM511-E8. Error bars indicate the s.d. of five individual assays. $**P < 0.05$; ns, non-significant; two-way ANOVA or Tukey's test. (d) Seeding density-dependent adhesion of H9 hESCs on LM-E8s. LM-E8s showed a higher cell adhesive activity for completely dissociated H9 hESCs than did Matrigel. Completely dissociated cells were incubated on LM-E8s in mTeSR1 medium for 6 h with or without Y-27632. Data represent the means \pm s.d. of three experiments. $**P < 0.05$; ns, non-significant, two-tailed Student's *t*-test.

Both LM332-E8 and LM511-E8 showed higher adhesion for completely dissociated hESCs or hiPSCs than did Matrigel at various cell densities (Fig. 1d and Supplementary Fig. S3b). Furthermore, using the optimal cell density (more than 50,000 cells per cm^2), we observed attachment of 40–70% of cells to LM-E8s, which raised the possibility of passaging cultures by completely dissociating hESCs and hiPSCs.

It has been previously shown that the Rho-associated protein kinase (ROCK) inhibitor Y-27632 increases the adhesion of dissociated hESCs³³. However, the effect of Y-27632 on hESC and hiPSC cultures in LM-E8-coated plates was unclear (Fig. 1d and Supplementary Fig. S3b). Y-27632 pretreatment before complete dissociation did not significantly improve the attachment of dissociated H9 hESCs on LM-E8s (Supplementary Fig. S2b). We therefore concluded that the addition of Y-27632 to the medium was unnecessary for cultures using LM-E8s, unlike attachment onto Matrigel, fibronectin, vitronectin and Synthemax that has a synthetic culture surface consisting of short peptides derived from vitronectin (Supplementary Fig. S2c). To focus on the effectiveness of LM-E8s, we did not use Y-27632 in subsequent experiments. In addition, we selected 50,000 cells per cm^2 as the seeding density for passaging hESCs and hiPSCs.

LM-E8s support vigorous proliferation in defined media. We next examined whether LM-E8s supported undifferentiated

growth of completely dissociated hESCs and hiPSCs in mTeSR1 medium. H9 hESC colonies dissociated by a conventional method showed superior attachment to LM-E8s (Fig. 2a). Likewise, completely dissociated H9 hESCs showed strong adherence on LM-E8s. Cells quickly adhered and spread on LM-E8-coated substrates, with extending lamellipodium-like protrusions (Fig. 2b and Supplementary Movies 1 and 2). H9 hESCs adhered to LM-E8s promptly recovered cell–cell contacts, formed highly dense colonies (Fig. 2c), and showed steady proliferation similar to that of conventional colony cultures (Fig. 2d). Similarly, completely dissociated iPS(IMR90)-1 cells also exhibited steady proliferation on LM-E8 fragments (Supplementary Fig. S3c). In contrast, completely dissociated H9 hESCs displayed weak adhesion on Matrigel or considerably poor adhesion on fibronectin and vitronectin, although H9 hESC colonies normally adhere to these substrates (Fig. 2a). Most dissociated cells on Matrigel, fibronectin and vitronectin remained as single cells and rapidly underwent cell death (Fig. 2b and Supplementary Movies 3–5). In agreement with the cell adhesion and survivability on LM-E8s, molecular analysis revealed that LM-E8s stimulated the signalling pathways downstream of integrins in hESCs and iPSCs. AKT, FAK and ERK1/2, signalling molecules closely associated with cell survival and migration³⁴, were activated to a greater extent in cells cultured on LM-E8s than in cells cultured on fibronectin or vitronectin (Fig. 2e). Y-27632 treatment abrogated MLC2 phosphorylation regardless of culture substrates, indicating that




Figure 2 | LM-E8s support undifferentiated proliferation in defined medium. (a) Distinct cell survival of H9 hESCs depending on ECMs. Data represent the numbers of live cells at 24 h after seeding 3.75×10^4 cells as single cells (single) or colonies (colony). (b) Serial phase-contrast images of H9 hESCs on LM-E8s or Matrigel in mTeSR1 medium at initial attachment. Images at the bottom right show an enlarged view of enclosed boxes. Note that dissociated H9 hESCs promptly extended and reached neighbouring cells on LM-E8s, but failed cellular extension on Matrigel. See also Supplementary Movies 1–5. (c) Phase-contrast images of H9 hESCs in mTeSR1 medium on LM-E8s or Matrigel during the growth phase. (d) Short-term growth curves of H9 hESCs on LM-E8s in mTeSR1 medium. Dissociated H9 hESCs on LM-E8s proliferated and had a similar growth rate as that in conventional colony culture during logarithmic growth. Data represent the means \pm s.d. of three individual experiments. (e) Detection of phosphorylated proteins associated with cell adhesion by western blot analysis. Cell lysates were prepared from H9 hESCs incubated for 1 h on each substrate with or without Y-27632. Sus, dissociated H9 hESCs incubated for 1 h in suspension; FN, fibronectin; VN, vitronectin. (f) Long-term growth curves of H9 hESCs on LM-E8s in mTeSR1 medium. The numbers of seeded cells at the initial passage were converted to 1, and the counted cell numbers at each passage are shown as cumulative cell numbers. Note that dissociation culture on LM-E8s resulted in a more than 200-fold increase in cell numbers relative to conventional colony culture over 1 month. Scale bars, 200 μ m.

the signalling pathways promoting cell survival on LM-E8s were not the same as those targeted by Y-27632. Once H9 hESCs adhered on each culture substrate, the cells similarly proliferated at an invariable rate that was regardless of the culture substrate, cluster size and Y-27632 treatment (Fig. 2d and Supplementary Fig. S2d). Therefore, the expansion efficiency was simply dependent on the initial adhesion efficiency and passaging ratio. LM-E8s allowed a higher passaging ratio during subculture, which was approximately 1:10 compared with 1:4 for conventional colony culture. Indeed, LM-E8s yielded more than

200-fold of the number of cells compared with that obtained by conventional culture over a 1-month period (Fig. 2f). Thus, LM-E8s enabled rapid expansion of completely dissociated hESCs and hiPSCs by supporting robust adhesion and rapid cell expansion.

Using our culture protocol with LM-E8s and two chemically defined media (mTeSR1 and StemPro), three hESC lines (H9, HES3 and KhES-1) and two hiPSC lines (iPS(IMR90)-1 and 253G1) showed robust proliferation for at least 10 serial passages. Flow cytometric analysis further indicated that the majority of dissociated hESCs and hiPSCs cultured on LM-E8s




Figure 3 | Flow cytometric analysis of hESC or hiPSC lines. Three hESC lines (H9, HES3 and KhES-1) and two hiPSC line (iPS(IMR90)-1 and 253G1) were cultured on LM-E8s after complete dissociation. These cells maintained higher expression levels of the undifferentiated markers, compared with that on Matrigel with either colony or complete dissociation culture. Undifferentiated markers: SSEA-3, SSEA-4, Tra-1-60, Tra-1-81, GCTM2 and Tra-2-54. Differentiation marker: SSEA-1. Data were collected from cells cultured for 10 passages. The percentages of positive cells are indicated on the graph.

expressed pluripotency markers SSEA-3, SSEA-4, Tra-1-60, Tra-1-81, GCTM2 and Tra-2-54, with only low level of expression of the differentiation marker SSEA-1 (Fig. 3). It has been reported that N2B27 medium is less robust for hESC maintenance, compared with two other media², but almost all hESCs and hiPSCs, except for KhES-1, cultured on LM-E8s showed undifferentiated growth in N2B27 medium. We karyotyped 50 randomly selected cells after G-banding of each cell line for all experiments at passage 10. Cells propagated on LM-E8s after complete dissociation had normal karyotypes (Table 1 and Supplementary Fig. S4a, b). Using multiplex fluorescence *in situ* hybridization (M-FISH) analysis, we showed that completely dissociated H9 hESCs and iPS(IMR90)-1 cells had a normal karyotype after 30 and 20 passages on LM-E8s, respectively (Supplementary Fig. S4c–f).

To assess the pluripotency of hESCs that had been cultured on LM-E8s in mTeSR1 medium for 10 passages, the cells were treated with the growth factors described in the Methods. We found that hESCs could differentiate into ectoderm (β III tubulin), mesoderm (α -smooth muscle actin) or endoderm (α -fetoprotein), as determined by immunohistochemical analyses (Fig. 4a). Furthermore, reverse transcriptase PCR (RT-PCR) analysis showed that, when hESCs were placed in suspension culture, they formed embryoid bodies with increased expression of gene markers for all three germ layer lineages after 10 days of

incubation (Fig. 4b). Similar results were obtained for iPS (IMR90)-1 cells cultured on LM-E8s (Supplementary Fig. S5). We further confirmed that hESCs cultured on LM-E8s in mTeSR1 medium for 25 passages formed teratomas consisting of all three germ layers, including neuroepithelium (ectoderm), intestinal epithelium (endoderm) and cartilage (mesoderm) (Fig. 4c).

Thus, our new culture method using completely dissociated cells on LM-E8s supported long-term maintenance and efficient cell expansion of hESCs and hiPSCs in an undifferentiated state.

LM-E8s support vigorous proliferation in xeno-free medium.

The final goal of this study was to characterize LM-E8s as substrates that support hESCs in a sustainable undifferentiated state under a xeno-free culture condition. We cultured completely dissociated H9 hESCs and 253G1 iPSCs on LM511-E8 or LM332-E8 in TeSR2 medium that is chemically defined and animal protein free. Dissociated H9 hESCs tightly adhered on LM-E8s in TeSR2 medium (Fig. 5a). Similarly to the results described for defined media, we found that H9 hESCs and 253G1 iPSCs proliferated robustly for more than 30 and 10 passages, respectively. Flow cytometric analysis also confirmed that the cells retained their undifferentiated state (Fig. 5b and Supplementary Fig. S6a). RT-PCR analysis of H9 hESCs and 253G1 iPSCs

Table 1 | Summary of karyotypic analysis.

Cell line	Medium	Colony culture	Matrigel	LM332-E8	LM511-E8
H9	mTeSR1	Normal*	Normal*	Normal	Normal*
iPS (IMR90)-1	mTeSR1	Normal*	Normal*	Normal*	Normal*
HES3	mTeSR1	Normal*	Normal*	Normal*	Normal
KhES-1	mTeSR1	46, xx, add(13)[1]	46, xx, add(13)[13]	Normal*	Normal*
253G1	mTeSR1	Normal	Normal	Normal	Normal
H9	StemPro	Normal*	Normal*	Normal*	Normal*
iPS (IMR90)-1	StemPro	Normal	Normal	Normal	Normal*
HES3	StemPro	Normal*	Normal	Normal*	Normal*
KhES-1	StemPro	Normal	Normal*	Normal*	Normal*
253G1	StemPro	Normal	Normal	Normal	Normal
H9	N2B27	Normal	Normal	Normal	Normal
iPS (IMR90)-1	N2B27	Normal*	Normal*	Normal*	Normal*
HES3	N2B27	Normal*	Normal	Normal*	Normal
KhES-1	N2B27	Normal	Normal	Normal	Normal
253G1	N2B27	Normal	Normal	Normal	Normal

The table lists the abnormalities found in 50 randomly selected G-banded cells from tenth passage cultures.
 *Indicates that a single defect of one chromosome was observed, suggesting that it may have been due to technical/procedural issues.




Figure 4 | LM-E8s sustain pluripotency in defined medium. (a) Immunostaining for markers of the three germ layers in differentiating H9 hESCs cultured on LM-E8s in mTeSR1 medium after 10 passages: ectoderm (β III tubulin), endoderm (alpha-fetoprotein (AFP)), and mesoderm (α -smooth muscle actin (SMA)). (b) RT-PCR analysis of marker genes for the differentiation of embryoid bodies. H9 hESCs were cultured in mTeSR1 medium for 10 passages on LM-E8s after complete dissociation, or on Matrigel after colony dissociation. Cells were then allowed to form embryoid bodies for 10 days. (c) Teratomas consisting of the three germ layers developed following the transfer of H9 hESCs cultured on LM-E8s after 25 passages in mTeSR1 medium. Cells were injected into the testes of SCID mice. After 8 weeks, the fate of the cells was analysed. Haematoxylin and eosin staining showed differentiation into various tissues including neuroepithelium (ectoderm), intestinal epithelium (endoderm) and cartilage (mesoderm). Scale bars, 200 μ m.

cultured on LM-E8s after complete dissociation showed that the expression of pluripotency markers, including *NANOG* and *OCT4*, remained high, whereas the expression of differentiation lineage markers was suppressed (Fig. 5c and Supplementary

Fig. S6b). H9 hESCs and 253G1 iPSCs had a normal karyotype after 35 and 10 serial passages, as indicated by G-band analysis (Supplementary Fig. S7a–d), and M-FISH analysis corroborated these findings (Supplementary Fig. S7e,f). In addition, we




Figure 5 | LM-E8s sustain the pluripotency in xeno-free TeSR2 medium. (a) Phase-contrast images of H9 hESCs. Dissociated H9 hESCs in TeSR2 medium reformed clusters (24 h) and actively proliferated (72 h). (b) Flow cytometric profiles of H9 hESCs cultured on LM-E8s for 34 passages after complete dissociation, or on Matrigel for 30 passages after colony dissociation. H9 hESCs maintained expression of markers of the undifferentiated state. Numbers indicate percentages of cells that were positive for surface markers. (c) RT-PCR analysis of the expression of pluripotency marker genes in H9 hESCs cultured in TeSR2 medium. H9 hESCs cultured for 30 passages on LM-E8s after complete dissociation maintained high-level expression of pluripotency marker genes, while differentiation lineage marker genes were suppressed. Embryoid bodies derived from H9 hESCs cultured as colonies in mTeSR1 medium were used as the control to assess differentiation. (d) Teratomas consisting of the three germ layers developed from H9 hESCs cultured on LM-E8s for 34 passages in TeSR2 medium after complete dissociation. Cells were injected into the testes of SCID mice. After 8 weeks, the fate of the cells was analysed. Haematoxylin and eosin staining showed differentiation into various tissues including neuroepithelium (ectoderm), intestinal epithelium (endoderm) and cartilage (mesoderm). Scale bars, 200 μ m.

confirmed the pluripotency of hESCs cultured on LM-E8s by their ability to form teratomas comprising all three germ layers (Fig. 5d).

Discussion

Stem cell–ECM interactions are important for maintaining stem cell adhesion, survival and self-renewal both *in vivo* and *in vitro*. The development of an ECM that sustains hESC and hiPSC cultures still requires substantial effort, particularly to avoid the use of feeder cells. In this study, we have demonstrated that LM-E8s support stronger adhesion of hESCs and hiPSCs than do other substrates, including Matrigel and intact laminin-511. We showed that LM-E8s enabled rapid and steady expansion of these cells. Furthermore, the cells cultured on LM-E8s for more than 30 passages in defined xeno-free media maintained an undifferentiated state and retained a normal karyotype and pluripotency. The adhesion of these cells to LM-E8s is primarily $\alpha 6\beta 1$ integrin-dependent, because a function-blocking antibody against $\alpha 6\beta 1$ integrin strongly inhibits the cell adhesion. Furthermore, LM511(EQ), a mutant LM-E8 that cannot bind to $\alpha 6\beta 1$ integrin owing to a single amino-acid substitution²⁹, does not support the

adhesion of hESCs. Thus, we believe that the interaction of LM-E8s with $\alpha 6\beta 1$ integrin is the principal route for binding to hESCs and promotion of proliferation in an undifferentiated state.

Although laminin-511 was shown to be a potential substrate for feeder-free culture of hESCs, intact laminin-511 is a large heterotrimeric protein of 780 kDa (Fig. 1a) and requires three independent transfections of expression vectors encoding the α , β and γ chains, each of which is larger than 10 kb, for recombinant protein production. It is therefore an arduous protocol to produce recombinant laminin-511 in sufficient quantities for the clinical application of hESCs. Here, we have shown that LM-E8s, the minimum structure harbouring the full integrin-binding activity of laminins, are superb substrates for the long-term culture of hESCs, with a substantial advantage over intact laminin-511 and laminin-332. LM-E8s are much smaller and easier to produce recombinantly than intact laminins. Indeed, the smaller size of LM-E8s significantly improves recombinant protein expression, allowing LM-E8s to yield more than five times more recombinant protein than intact laminin-511 (see Methods). In addition, LM-E8s can be easily purified using 6xHis- and FLAG-tags attached to the α and γ chains, respectively (Fig. 1a). This method reduces the time required

for purification and increases the purity of recombinant proteins, compared with intact laminin-511. These advantages endow LM-E8s with a greater potential for scaled-up production, compared with intact laminins. Importantly, LM-E8s have a higher adhesion affinity for hESCs than do intact laminin-511 and laminin-332, although the reasons for this property are yet to be fully determined. The smaller size of LM-E8s may allow them to pack more closely to each other on the substrate and thereby increases the adhesiveness of the substrate for hESC attachment by either increasing the ligand-binding affinity of integrins and/or enhancing the signalling events triggered by integrin clustering.

Several groups have reported that vitronectin and one of its variants (VTN-NC) supports the maintenance of hESCs via $\alpha V\beta 5$ integrin^{4,13}. The binding affinity of vitronectin to $\alpha V\beta 5$ integrin was 10-fold lower than that of LM-E8 to $\alpha 6\beta 1$ integrin (Yuya Sato, K.S., unpublished data), suggesting that vitronectin is potentially inferior for hESC adhesion compared with LM-E8s. Indeed, although hESCs and hiPSCs have been cultured as colonies on fibronectin and vitronectin^{1,13,35,36}, hESCs and hiPSCs that had been completely dissociated showed less adhesion (Fig. 2a). It has been reported that AKT, ERK1/2 and kinases interacting with FAK are highly phosphorylated in human pluripotent stem cells³⁷. Furthermore, it is well known that AKT and FAK, stimulated via integrin activation, perform numerous cell survival functions³⁴. The phosphorylation of AKT, FAK and ERK1/2 confirms that LM-E8s have the potential to promote the survival of completely dissociated hESCs by robust adhesion via integrins, whereas vitronectin and fibronectin are unfavourable for culturing of completely dissociated hESCs and hiPSCs. The ROCK inhibitor Y-27632 has been shown to increase the survival of dissociated hESCs³³. Tsutsui *et al.*⁵ reported maintenance of hESCs on a fibronectin-coated surface by dissociated cell passaging, but their method requires a high dose of Y-27632 for cell attachment to fibronectin. It should be noted that a high concentration of Y-27632 increases toxicity and leads to a reduction of hESC numbers in some cases³⁸. Additionally, the maintenance of hESCs on fibronectin requires constant administration of Y-27632, even after cell attachment. In contrast, our method using LM-E8s does not require Y-27632 administration owing to sufficient stimulation of cell survival from the culture substrate. This method might be advantageous not only for supporting cell expansion but also for use in drug screening studies. This is because LM-E8s not only improve the culture condition to avoid transient adverse effects of untargeted chemicals, such as Y-27632, but also allow uniform seeding of routine numbers of hESCs and hiPSCs. Therefore, LM-E8s are an appropriate culture substrate for the culture of completely dissociated hESCs and hiPSCs. Although LM-E8s ensured hESC survival just after cell seeding, single hESCs exhibited poor survival over prolonged time periods, even on LM-E8s. It is known that E-cadherin signalling, which arises from cell–cell contact, has a key role in the self-renewal of hESCs as well as the generation of hiPSCs^{39–41}. LM-E8 signalling may be insufficient for prolonged survival of single hESCs, but LM-E8s enabled hESCs to actively migrate, and the cells aggregated promptly and restored cell–cell contacts to form small colonies within 24 h of culture initiation (Figs 2b, c and 5a). FAK is known to perform multiple functions and regulates cell migration⁴². The increased phosphorylation of FAK is indicative of active migration of dissociated hESCs on LM-E8s. It has been recently reported that cell–cell contact promotes hESC survival, and Y-27632 treatment accelerates the migration of hESCs to reconstruct cell–cell contacts⁴³. Here, we show that LM-E8s also activated dissociated hESC migration, but the mechanism is different from that of Y-27632 treatment, as shown by the MLC2 phosphorylation analysis. Collectively, LM-E8s support not only

strong adhesion to hESCs and hiPSCs but also promote cell–cell contact by inducing their migration ability, thereby ensuring cell survive and proliferation. Thus, our new protocol improves the adhesion of hESCs and hiPSCs and allows efficient propagation in a simple and rapid manner.

In conclusion, we demonstrate a defined xeno- and feeder-free culture system for hESCs and hiPSCs. LM-E8s derived from laminin isoforms support increased adhesion, survival, self-renewal and differentiation of hESCs. This simple culture system should facilitate bulk proliferation of hESCs and hiPSCs under defined xeno-free conditions. We believe that our protocol will be useful for clinical applications of hESCs and hiPSCs in compliance with good manufacturing practice and/or current good tissue practice.

Methods

Preparation of ECMs. Recombinant human laminins, laminin-111, -332 and -511 and LM-E8s (LM322-E8 and LM511-E8) were produced using a FreeStyle 293 Expression System (Invitrogen) as described previously^{21,29,30}. Laminin-111, -332 and -511 and LM-E8s were filtered through 0.22- μm -pore-sized Ultrafree-CL Centrifugal Filter Units (Millipore) for sterilization. Using these methods, we purified an average of 0.74 mg intact LM511 ($n = 5$) and 4.1 mg LM511-E8 ($n = 4$) from 1 l of culture supernatant. Thus, the yield was much higher for LM511-E8 (5.5-fold in terms of weight, or 27.7-fold per mol) than for intact laminin. Furthermore, purification of intact laminin-511 required 1–2 weeks, whereas purification of LM-E8s required only 2–3 days.

Human fibronectin and vitronectin were purified from plasma, as described elsewhere^{44,45}. Matrigel (Growth Factor Reduced) was purchased from BD Biosciences.

Coating of matrices. For the dose–response binding assay, 96-well flat-bottomed plates (non-treated, BD Falcon) were pre-coated with each of the matrices at a concentration of 0–25 $\mu\text{g cm}^{-2}$ for 3 h at room temperature just before cell seeding. For other experiments, we used 48-well plates (non-treated, BD Falcon) pre-coated with LM-E8s, fibronectin, or vitronectin at 1.5 $\mu\text{g cm}^{-2}$ or Matrigel at 25 $\mu\text{g cm}^{-2}$ for 3 h at room temperature for the culture of hESCs and hiPSCs. For conventional colony passaging as an experimental control, 12-well plates were coated with Matrigel at 25 $\mu\text{g cm}^{-2}$. Coated dishes were not rinsed with buffer before use. To dilute the ECMs, Dulbecco's PBS was used for laminin-111, -332 and -511, LM-E8s, fibronectin and vitronectin. DMEM/F12 was used to dilute Matrigel.

Preparation of defined media. Commercially available defined media, mTeSR1 (mTeSR1; Stem Cell Technologies) and StemPro hESC SFM (StemPro, Invitrogen) and xeno-free medium, TeSR2 (TeSR2; Stem Cell Technologies), were used according to the manufacturer's instructions and freshly prepared every day. N2B27 medium was prepared daily, according to previous reports^{1,2}, and was composed of DMEM/F12 (Sigma, D6421), 1 \times N2 supplement (Invitrogen), 1 \times B27 supplement (Invitrogen), 1 mM non-essential amino acids (Sigma, M7145), 1 mM L-glutamine (Sigma, G7513), 0.1 mM 2-mercaptoethanol (Invitrogen) and 100 ng ml⁻¹ FGF-2 (Wako, Japan, 060-04543).

Cell culture. hESC lines, KhES-1 at passage 22 and HES3 at passage 100, and hiPSC line 253G1 at passage 50 were maintained as previously described on mitomycin-C-treated mouse embryonic fibroblast (MEF) feeder cells⁴⁶. The hESC line H9 (WA09) at passage 36, hiPSC line iPS (IMR90) clone 1 (iPS(IMR90)-1) at passage 80 and iPS (foreskin) line at passage 64 were purchased from the National Stem Cell Bank. Cells were maintained in accordance with National Stem Cell Bank protocols, except for the use of mitomycin-C-treated MEFs from the ICR strain at a density of 3.2×10^4 cells per cm².

Experimental control cells were obtained from conventional colony cultures that were maintained as follows. Near-confluent cells were incubated with 2 mg ml⁻¹ dispase in DMEM/F12 at 37 °C for 2 min, and then rinsed twice with DMEM/F12. After addition of defined medium, weakly adherent colonies were detached using a cell scraper. Cells were collected and centrifuged at 200 g for 3 min at 4 °C. Small colonies were then passaged into Matrigel-coated vessels at a ratio of 1:4. Colonies showing morphological differentiation were manually removed under a microscope during each passage, except at the time of analysis. Completely dissociated cells were cultured after preculture in the appropriate defined medium for one passage on Matrigel-coated dishes to remove MEF feeders. This stage was defined as passage 0. Near-confluent cell clusters were treated with 4.8 mM EDTA/PBS for 2–5 min at room temperature and then TrypLE Select (Invitrogen) for 1 min at 37 °C. Cells in defined medium were pipetted to disperse the cells completely. After centrifugation at 200 g for 3 min at 4 °C, cells were seeded at the appropriate density (5×10^4 cells per cm² for passaging or as indicated) into new culture vessels

precoated with each matrix. The medium was changed daily for all experiments. hESCs and hiPSCs on LM-E8s were subcultured every 4–6 days, and those on Matrigel every 6–8 days.

Time-lapse recording. For time-lapse recording during initial attachment, completely dissociated H9 hESCs were seeded onto LM-E8-, fibronectin-, vitronectin- or Matrigel-coated 35-mm dishes at 2.5×10^4 cells per cm^2 in mTeSR1 medium. Recording began at 5 min after cell seeding and continued for 24 h. Movies were edited using AxioVisio software.

Cell adhesion assay. Dose–response binding assays were performed using a protocol described elsewhere¹³, with some modification. Briefly, hESCs or hiPSCs were cultured for one passage on Matrigel in mTeSR1 medium to remove feeder cells, and then the cells were detached using TrypLE Select, collected and counted. A total of 5×10^4 cells were seeded into each well of 96-well plates coated with the matrices described above. After 6 h of incubation, non-adherent cells were removed by rinsing the plates twice with DMEM/F12, and the remaining adherent cells were fixed with 10% formalin for 15 min, and postfixed in 100% ethanol for 5 min at room temperature. Cells were then stained with 0.4% crystal violet in methanol for 5 min at room temperature. After washing extensively with demineralized water, the remaining cells were solubilized by adding 100 μl of 1% SDS, and the optical density at 570 nm was measured by a multiwell plate reader. For the functional inhibition assay of integrins, H9 hESCs were seeded into each well of 96-well plates coated with the various matrices including LM511(EQ) in the presence or absence of $2 \mu\text{g ml}^{-1}$ integrin-blocking antibodies, anti- $\alpha 6$ integrin (R&D systems) and/or anti- $\beta 1$ integrin (R&D systems). After 30 min of incubation, adherent cells were screened as described above for the dose–response assay.

For the seeding density assay, hESCs or hiPSCs were precultured in mTeSR1 medium on Matrigel-coated vessels for one passage to remove feeder cells. Cells were detached using an EDTA solution and TrypLE Select as described above. Cells were seeded into 48-well plates coated with LM-E8s or Matrigel. After 6 h of incubation, non-adherent cells were rinsed twice with prewarmed DMEM/F12, and the remaining adherent cells were detached using 0.25% Trypsin/EDTA. Viable cells were counted by trypan blue exclusion.

For the proliferation assay, dissociated single cells were seeded at a density of 4×10^4 cells per cm^2 , and viable cells were counted every 24 h as described above. In parallel, colonies were seeded on 24-well plates coated with Matrigel at the same cell density.

To evaluate Y-27632 pretreatment, H9 hESCs cultured on Matrigel in mTeSR1 medium were incubated with 10 μM Y-27632 (Wako, 253-00513) for 1 h just before cell detachment. Together with the cultures without Y-27632 pretreatment, cells were detached by EDTA and TrypLE Select treatment and then seeded in multiwell plates coated with substrates in mTeSR1 medium containing 10 μM Y-27632.

To evaluate the effect of Y-27632 treatment on cell proliferation, H9 hESCs after complete dissociation or as colonies were seeded at a density of 5×10^4 cells per cm^2 with or without 10 μM Y-27632 in 48-well plates coated with each substrate. Viable cells were counted every 24 h. To evaluate a derivation of vitronectin, a Synthemax-R Surface 6-well plate (Corning) was used as a substrate.

Flow cytometric analysis. Cells were dissociated by treatment with 4.8 mM EDTA/PBS for 2–5 min, followed by further treatment with a 0.05% trypsin/EDTA solution for 1 min. After two washes with DMEM/10% FBS and one wash with staining buffer (0.1% BSA/PBS), 1×10^5 cells were incubated for 30 min at 4 °C with primary antibodies diluted in staining buffer. Cells were then rinsed twice with staining buffer and incubated for 30 min at 4 °C with secondary antibodies diluted in staining buffer. Then, cells were rinsed twice with staining buffer, and counterstained with propidium iodide just before analysis.

Fluorescence intensities were analysed on a FACScalibur flow cytometer (Becton Dickinson), and the FL-1 positive ratios in the propidium iodide-negative region were monitored using Cell Quest software. The following primary antibodies were used: SSEA-3 ($2 \mu\text{g ml}^{-1}$; Developmental Studies Hybridoma Bank (DSHB)), SSEA-4 ($1 \mu\text{g ml}^{-1}$; DSHB), Tra-1-60 ($1 \mu\text{g ml}^{-1}$; Chemicon), Tra-1-81 ($1 \mu\text{g ml}^{-1}$; Chemicon), GCTM2 ($1 \mu\text{g ml}^{-1}$; Chemicon), Tra-2-54 ($1 \mu\text{g ml}^{-1}$, DSHB), and SSEA-1 ($2 \mu\text{g ml}^{-1}$; DSHB). Anti-mouse Ig/FITC conjugated ($10 \mu\text{g ml}^{-1}$; Becton Dickinson) and anti-rat IgM/Alexa 488 conjugated ($1 \mu\text{g ml}^{-1}$; Molecular Probes) were used as secondary antibodies.

Karyotyping. hESCs and hiPSCs were treated with 100 ng ml^{-1} colcemid (Invitrogen) for 2–3 h. After dissociation in 0.25% trypsin/EDTA solution, cells were treated with a hypotonic solution and then fixed in Carnoy's solution. Cells were spread onto glass slides and stained with Giemsa. Chromosome spreads were analysed by randomly counting 50 cells using the Ikaros Karyotyping System (META system). M-FISH analysis was performed using the 24XCyte Human Multicolor FISH (mFISH) Probe Kit (META system) following the manufacturer's instructions, and analysed by ISIS software (META system).

Statistical analysis. The statistical significance of differences was determined by the two-tailed Student's *t*-test for two group comparisons, or two-way analysis of

variance (two-way ANOVA) for multiple comparisons followed by Tukey's test. Differences with a value of $P < 0.05$ were considered significant.

Other methods. See Supplementary Methods and Supplementary Table S2 for remaining methods, including semi-quantitative RT–PCR, the differentiation assay, immunofluorescence staining and immunoblotting.

References

- Liu, Y. *et al.* A novel chemical-defined medium with bFGF and N2B27 supplements supports undifferentiated growth in human embryonic stem cells. *Biochem. Biophys. Res. Commun.* **346**, 131–139 (2006).
- Akopian, V. *et al.* Comparison of defined culture systems for feeder cell free propagation of human embryonic stem cells. *In Vitro Cell Dev. Biol. Anim.* **46**, 247–258 (2010).
- Ludwig, T. E. *et al.* Derivation of human embryonic stem cells in defined conditions. *Nature Biotech.* **24**, 185–187 (2006).
- Chen, G. *et al.* Chemically defined conditions for human iPSC derivation and culture. *Nat. Methods* **8**, 424–429 (2011).
- Tsutsui, H. *et al.* An optimized small molecule inhibitor cocktail supports long-term maintenance of human embryonic stem cells. *Nat. Commun.* **2**, 167 (2011).
- Baker, M. Stem cells in culture: defining the substrate. *Nat. Methods* **8**, 293–297 (2011).
- Thomson, J. A. *et al.* Embryonic stem cell lines derived from human blastocysts. *Science* **282**, 1145–1147 (1998).
- Amit, M. *et al.* Clonally derived human embryonic stem cell lines maintain pluripotency and proliferative potential for prolonged periods of culture. *Dev. Biol.* **227**, 271–278 (2000).
- Pyle, A. D., Lock, L. F. & Donovan, P. J. Neurotrophins mediate human embryonic stem cell survival. *Nature Biotech.* **24**, 344–350 (2006).
- Draper, J. S. *et al.* Recurrent gain of chromosomes 17q and 12 in cultured human embryonic stem cells. *Nature Biotech.* **22**, 53–54 (2004).
- Mitalipova, M. M. *et al.* Preserving the genetic integrity of human embryonic stem cells. *Nature Biotechnol.* **23**, 19–20 (2005).
- Rodin, S. *et al.* Long-term self-renewal of human pluripotent stem cells on human recombinant laminin-511. *Nature Biotechnol.* **28**, 611–615 (2010).
- Braam, S. R. *et al.* Recombinant vitronectin is a functionally defined substrate that supports human embryonic stem cell self-renewal via $\alpha 5 \beta 1$ integrin. *Stem Cells* **26**, 2257–2265 (2008).
- Melkounian, Z. *et al.* Synthetic peptide-acrylate surfaces for long-term self-renewal and cardiomyocyte differentiation of human embryonic stem cells. *Nature Biotechnol.* **28**, 606–610 (2010).
- Villa-Diaz, L. G. *et al.* Synthetic polymer coatings for long-term growth of human embryonic stem cells. *Nature Biotech.* **28**, 581–583 (2010).
- Mei, Y. *et al.* Combinatorial development of biomaterials for clonal growth of human pluripotent stem cells. *Nat. Mater.* **9**, 768–778 (2010).
- Klim, J. R., Li, L., Wrighton, P. J., Piekarczyk, M. S. & Kiessling, L. L. A defined glycosaminoglycan-binding substratum for human pluripotent stem cells. *Nat. Methods* **7**, 989–994 (2010).
- Saha, K. *et al.* Surface-engineered substrates for improved human pluripotent stem cell culture under fully defined conditions. *Proc. Natl. Acad. Sci. USA* **108**, 18714–18719 (2011).
- Miner, J. H. & Yurchenco, P. D. Laminin functions in tissue morphogenesis. *Annu. Rev. Cell Dev. Biol.* **20**, 255–284 (2004).
- Miyazaki, T. *et al.* Recombinant human laminin isoforms can support the undifferentiated growth of human embryonic stem cells. *Biochem. Biophys. Res. Commun.* **375**, 27–32 (2008).
- Ido, H. *et al.* Molecular dissection of the α -dystroglycan- and integrin-binding sites within the globular domain of human laminin-10. *J. Biol. Chem.* **279**, 10946–10954 (2004).
- Smirnov, S. P. *et al.* Contributions of the LG modules and furin processing to laminin-2 functions. *J. Biol. Chem.* **277**, 18928–18937 (2002).
- Doi, M. *et al.* Recombinant human laminin-10 ($\alpha 5 \beta 1 \gamma 1$). Production, purification, and migration-promoting activity on vascular endothelial cells. *J. Biol. Chem.* **277**, 12741–12748 (2002).
- Kortessmaa, J., Yurchenco, P. & Tryggvason, K. Recombinant laminin-8 ($\alpha 4 \beta 2 \gamma 1$). Production, purification, and interactions with integrins. *J. Biol. Chem.* **275**, 14853–14859 (2000).
- Kariya, Y. *et al.* Efficient expression system of human recombinant laminin-5. *J. Biochem.* **132**, 607–612 (2002).
- Vuoristo, S. *et al.* Laminin isoforms in human embryonic stem cells: synthesis, receptor usage and growth support. *J. Cell Mol. Med.* **13**, 2622–2633 (2009).
- Nishiuchi, R. *et al.* Ligand-binding specificities of laminin-binding integrins: a comprehensive survey of laminin-integrin interactions using recombinant $\alpha 3 \beta 1$, $\alpha 6 \beta 1$, $\alpha 7 \beta 1$ and $\alpha 6 \beta 4$ integrins. *Matrix Biol.* **25**, 189–197 (2006).
- Meng, Y. *et al.* Characterization of integrin engagement during defined human embryonic stem cell culture. *FASEB J.* **24**, 1056–1065 (2010).

29. Ido, H. *et al.* The requirement of the glutamic acid residue at the third position from the carboxyl termini of the laminin gamma chains in integrin binding by laminins. *J. Biol. Chem.* **282**, 11144–11154 (2007).
30. Taniguchi, Y. *et al.* The C-terminal region of laminin beta chains modulates the integrin binding affinities of laminins. *J. Biol. Chem.* **284**, 7820–7831 (2009).
31. Deutzmann, R. *et al.* Cell adhesion, spreading and neurite stimulation by laminin fragment E8 depends on maintenance of secondary and tertiary structure in its rod and globular domain. *Eur. J. Biochem.* **191**, 513–522 (1990).
32. Takagi, J. Structural basis for ligand recognition by integrins. *Curr. Opin. Cell Biol.* **19**, 557–564 (2007).
33. Watanabe, K. *et al.* A ROCK inhibitor permits survival of dissociated human embryonic stem cells. *Nature Biotech.* **25**, 681–686 (2007).
34. Vachon, P. H. Integrin signaling, cell survival, and anoikis: distinctions, differences, and differentiation. *J. Signal Transduct.* **2011**, 738137 (2011).
35. Prowse, A. B. *et al.* Long term culture of human embryonic stem cells on recombinant vitronectin in ascorbate free media. *Biomaterials* **31**, 8281–8288 (2010).
36. Lu, J., Hou, R., Booth, C. J., Yang, S. H. & Snyder, M. Defined culture conditions of human embryonic stem cells. *Proc. Natl Acad. Sci. USA* **103**, 5688–5693 (2006).
37. Phanstiel, D. H. *et al.* Proteomic and phosphoproteomic comparison of human ES and iPS cells. *Nat. Methods* **8**, 821–827 (2011).
38. Andrews, P. D. *et al.* High-content screening of feeder-free human embryonic stem cells to identify pro-survival small molecules. *Biochem. J.* **432**, 21–33 (2010).
39. Xu, Y. *et al.* Revealing a core signaling regulatory mechanism for pluripotent stem cell survival and self-renewal by small molecules. *Proc. Natl Acad. Sci. USA* **107**, 8129–8134 (2010).
40. Redmer, T. *et al.* E-cadherin is crucial for embryonic stem cell pluripotency and can replace OCT4 during somatic cell reprogramming. *EMBO Rep.* **12**, 720–726 (2011).
41. Chen, T. *et al.* E-cadherin-mediated cell-cell contact is critical for induced pluripotent stem cell generation. *Stem Cells* **28**, 1315–1325 (2010).
42. Schaller, M. D. Cellular functions of FAK kinases: insight into molecular mechanisms and novel functions. *J. Cell Sci.* **123**, 1007–1013 (2010).
43. Li, L. *et al.* Individual cell movement, asymmetric colony expansion, rho-associated kinase, and E-cadherin impact the clonogenicity of human embryonic stem cells. *Biophys. J.* **98**, 2442–2451 (2010).
44. Sekiguchi, K. & Hakomori, S. Domain structure of human plasma fibronectin. Differences and similarities between human and hamster fibronectins. *J. Biol. Chem.* **258**, 3967–3973 (1983).
45. Yatohgo, T., Izumi, M., Kashiwagi, H. & Hayashi, M. Novel purification of vitronectin from human plasma by heparin affinity chromatography. *Cell Struct. Funct.* **13**, 281–292 (1988).
46. Suemori, H. *et al.* Efficient establishment of human embryonic stem cell lines and long-term maintenance with stable karyotype by enzymatic bulk passage. *Biochem. Biophys. Res. Commun.* **345**, 926–932 (2006).

Acknowledgements

We thank Ms Mari Hamao and Ms Ritsuko Goto for their valuable technical assistance and the members of the laboratories of N.N. and K.S. for insightful discussions. This work was supported by the New Energy and Industrial Technology Development Organization (NEDO) of Japan, and partially supported by the Coordination, Support and Training Programme for Translational Research of the Ministry of Education, Culture, Sports, Science and Technology of Japan (to K.S.).

Author contributions

T.M., K.S. and E.K. designed research, T.M., S.F., Y.T., M.Y., M.K., M.H. and H.K. performed research, T.M., H.S., N.N., K.S. and E.K. analysed data and T.M., K.S. and E.K. wrote the paper.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://ngp.nature.com/reprintsandpermissions/>

How to cite this article: Miyazaki, T. *et al.* Laminin E8 fragments support efficient adhesion and expansion of dissociated human pluripotent stem cells. *Nat. Commun.* **3**:1236 doi: 10.1038/ncomms2231 (2012).




This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Erratum: Laminin E8 fragments support efficient adhesion and expansion of dissociated human pluripotent stem cells

Takamichi Miyazaki, Sugiko Futaki, Hirofumi Suemori, Yukimasa Taniguchi, Masashi Yamada, Miwa Kawasaki, Maria Hayashi, Hideaki Kumagai, Norio Nakatsuji, Kiyotoshi Sekiguchi & Eihiro Kawase

Nature Communications 3:1236 doi: 10.1038/ncomms2231 (2012); Published 4 Dec 2012; Updated 29 Jul 2013

In the key to Fig. 3 of this Article, labels for the colours representing SSEA-1 and Tra-2-54 were inadvertently switched during the production process. The correct version of the figure appears below.



RESEARCH COMMUNICATION

The SMAD2/3 corepressor SNON maintains pluripotency through selective repression of mesendodermal genes in human ES cells

Norihiro Tsuneyoshi,¹ Ee Kim Tan,¹
Akila Sadasivam,¹ Yogavalli Poobalan,¹
Tomoyuki Sumi,^{2,3} Norio Nakatsuji,^{4,5}
Hirofumi Suemori,² and N. Ray Dunn^{1,6}

¹Institute of Medical Biology, A*STAR (Agency for Science, Technology, and Research), Singapore 138648, Singapore;

²Department of Embryonic Stem Cell Research, Institute for Frontier Medical Sciences, Kyoto University, Kyoto 606-8507, Japan; ³Department of Developmental Biology, Graduate School of Medical Sciences, Kyushu University, Fukuoka 812-8582, Japan; ⁴Department of Development and Differentiation, Institute for Frontier Medical Sciences, Kyoto University, Kyoto 606-8507, Japan; ⁵Institute for Integrated Cell-Material Sciences (WPI-iCeMS), Kyoto University, Kyoto 606-8501, Japan

Activin/Nodal signaling via SMAD2/3 maintains human embryonic stem cell (hESC) pluripotency by direct transcriptional regulation of *NANOG* or, alternatively, induces mesoderm and definitive endoderm (DE) formation. In search of an explanation for these contrasting effects, we focused on SNON (SKIL), a potent SMAD2/3 corepressor that is expressed in hESCs but rapidly down-regulated upon differentiation. We show that SNON predominantly associates with SMAD2 at the promoters of primitive streak (PS) and early DE marker genes. Knockdown of SNON results in premature activation of PS and DE genes and loss of hESC morphology. In contrast, enforced SNON expression inhibits DE formation and diverts hESCs toward an extraembryonic fate. Thus, our findings provide novel mechanistic insight into how a single signaling pathway both regulates pluripotency and directs lineage commitment.

Supplemental material is available for this article.

Received July 23, 2012; revised version accepted September 28, 2012.

Since the isolation of human embryonic stem cells (hESCs) more than a decade ago, numerous extrinsic signals have been implicated in the maintenance of their cardinal features of pluripotency and indefinite self-renewal (Pera and Tam 2010). These include growth factors belonging to the TGF β , FGF, WNT, and IGF families, which are often supplemented with culture

medium containing serum, feeder cell-conditioned medium, and/or ill-defined knockout serum replacement (Price et al. 1998). These various and complex platforms have obscured the identification of the minimal complement of extracellular cues that govern the intrinsic transcriptional regulators of pluripotency and self-renewal. Recently, however, it was shown that either recombinant TGF β 1 or the related ligand Activin A in combination with FGF2 can maintain hESCs in chemically defined and feeder-free conditions partly through direct regulation of the core pluripotency factor gene *NANOG* (Xu et al. 2008; Vallier et al. 2009).

Both TGF β and Activin signals are transduced intracellularly by phosphorylation of the cytoplasmic effector proteins SMAD2 and SMAD3 (Schmierer and Hill 2007). Once activated, they partner with SMAD4, translocate to the nucleus, and associate with tissue-specific transcription factors to regulate batteries of target genes. Chromatin immunoprecipitation (ChIP) experiments show that SMAD2/3 bind the *NANOG* promoter, and luciferase reporter assays confirm that this SMAD-binding element (SBE) is crucial for *NANOG* transcriptional activity (Xu et al. 2008; Vallier et al. 2009). Recently, SMAD2/3 ChIP combined with next-generation sequencing (ChIP-seq) revealed that, in addition to *NANOG*, SMAD2/3 occupy a broad number of genes both expressed in hESCs and associated with the maintenance of pluripotency (Brown et al. 2011; Kim et al. 2011). These include the additional core pluripotency factors *OCT4* and *SOX2* as well as other key regulators, such as *SUZ12*, *UTF1*, *TERT*, *DPPA4*, and *LIN28A*. Taken together, these findings emphasize the importance of continuous TGF β /Activin signaling for the maintenance of hESC pluripotency. However, it is a third divergent TGF β -related ligand, Nodal, that carries out this fundamental activity in vivo.

Nodal transcripts identify the pluripotent cells within the mouse blastocyst inner cell mass (ICM) and persist in these cells as they extensively reorganize into the cup-shaped epiblast epithelium after implantation (Mesnard et al. 2006; Arnold and Robertson 2009). In the absence of *Nodal*, null mutant embryos down-regulate *Oct4*, *Nanog*, and *Foxd3*, which are normally uniformly expressed throughout the epiblast; prematurely and ectopically up-regulate neural markers; and fail to gastrulate (Camus et al. 2006; Mesnard et al. 2006). With the onset of gastrulation, *Nodal* expression resolves to the posterior primitive streak (PS), where epiblast cells ingress and emerge as mesoderm and definitive endoderm (DE) (Arnold and Robertson 2009). Extensive genetic studies have shown that the highest levels of *Nodal* specify DE, while the assorted mesodermal lineages are patterned by intermediate to lower *Nodal* levels (Arnold and Robertson 2009). Thus, there is a remarkable conversion in a narrow developmental time span in how *Nodal* signals are interpreted intracellularly—from maintenance of pluripotency within the ICM/epiblast to inducer of differentiation within the PS. This raises the intriguing question of how *Nodal*/Smad2/3 signals in the ICM/epiblast engage only the pluripotency machinery and avert precocious activation of differentiation genes. This question also stringently applies to hESCs, which share many defining characteristics with the mouse epiblast and its derivative cell lines (termed epiblast stem cells [EpiSCs]) (Pera and Tam 2010),

[*Keywords*: Activin/Nodal; SNON; human embryonic stem cells; mesendoderm; pluripotency; repressor]

⁶Corresponding author

E-mail ray.dunn@imb.a-star.edu.sg

Article is online at <http://www.genesdev.org/cgi/doi/10.1101/gad.201772.112>.

Tsuneyoshi et al.

as the Nodal analog Activin A both regulates pluripotency and is nearly universally used to initiate mesendoderm differentiation *in vitro* to produce a variety of lineage-specific cell types (Sulzbacher et al. 2009).

We reasoned that one explanation for these contrasting effects is the existence of discriminating nuclear repressor proteins. Moreover, the expression of such a repressor would likely be under the direct control of the core pluripotency factors. Previous ChIP combined with DNA microarrays (ChIP-chip) identified numerous active and inactive genes that are triply bound by OCT4, SOX2, and NANOG in hESCs (Boyer et al. 2005). One of these is the SKI-related proto-oncogene *SNON* (*SKIL*) that encodes a potent negative regulator of TGF β -related signaling (Deheuninck and Luo 2009; Zhu and Luo 2012). *SNON* physically interacts with SMAD2, SMAD3, and SMAD4, disrupting their assembly into an activated heteromeric complex (Stroschein et al. 1999; Wu et al. 2002). In addition, *SNON* prevents SMAD2 and SMAD3 from associating with the transcriptional coactivator p300/CBP and recruits a repressor complex comprised of nuclear receptor corepressor (N-CoR) and histone deacetylase (HDAC) (Schmierer and Hill 2007). Thus, the presence of *SNON* at a particular SBE ensures transcriptional repression. Here, we found that *SNON* protein is highly abundant in hESCs and specifically enriched at differentiation gene promoters. Loss of *SNON* in hESCs prematurely activates PS and early DE genes and destabilizes pluripotency, while constitutive *SNON* expression restrains differentiation into DE. Taken together, our results identify *SNON* as a key component of the nuclear machinery that safeguards pluripotency through direct repression of differentiation genes downstream from Activin/Nodal.

Results and Discussion

SNON is abundant in hESCs and is targeted to the proteasome during early differentiation

We first sought to confirm the expression of *SNON* in pluripotent hESCs and examine its regulation during DE formation *in vitro*. Differentiation was carried out according to the schematic in Figure 1A (Teo et al. 2012) and, expectedly, resulted in the loss of pluripotency markers (*NANOG*, *OCT4*, and *SOX2*) and up-regulation of PS (*WNT3*, *FGF8*, *EOMES*, and *MIXL1*), mesendoderm (*GSC* and *LHX1*), and DE (*FOXA2* and *SOX17*) markers by day 5 (Supplemental Fig. 1A). These data were confirmed by Western analysis, immunohistochemistry, and FACS (Supplemental Fig. 1B–D). We observed that *SNON* transcript levels are robust in undifferentiated HES3 and on day 1 of differentiation but drop considerably by day 3 and decline further on day 5 (Fig. 1B).

This decrease in *SNON* levels *in vitro* closely mirrors the expression dynamics of *SnoN* in the early mouse embryo. Prior to gastrulation, *SnoN* transcripts label the pluripotent epiblast epithelium (Supplemental Fig. 1E). Mouse ESCs and EpiSCs also express *SnoN* (Supplemental Fig. 1F). With the onset of gastrulation, *SnoN* transcript levels are conspicuously down-regulated in the region of the incipient PS and continue to decline posteriorly as the PS lengthens (Supplemental Fig. 1E). Nascent mesoderm and DE also lack *SnoN* expression. By the late PS stage (embryonic day 7.5 [E7.5]), *SnoN* transcripts resolve to the anterior epiblast, but shortly




Figure 1. *SNON* expression in hESCs is controlled by SMAD2/3 and OCT4, SOX2, and NANOG. (A) hESCs were differentiated into DE for 5 d according to the schematic. Activin A and BMP4 (both at 50 ng/mL) were added on day 0. Only Activin A was replenished on day 3. (B) *SNON* expression levels during DE differentiation by qPCR. Data were normalized against *GUSB* and are shown relative to undifferentiated hESCs (= 1.0). (C) Western blot analysis for *SNON* protein levels during DE differentiation. (D) OCT4 and *SNON* immunofluorescence on undifferentiated HES3 (hESCs). Bar, 100 μ m. (E) Luciferase reporter assays of *SNON* promoter activity in HES3. Black shading indicates specific mutation of the OCT4/SOX2/NANOG-binding sites in the 5' distal enhancer or of the four SBEs proximal to the *SNON* transcriptional start site.

thereafter, this domain has largely disappeared. These expression data suggest that loss of the Smad2/3 corepressor *SnoN* is an obligate step in the production of PS-derived lineages.

At the protein level, *SNON* is abundant in undifferentiated HES3 but, in contrast to our quantitative PCR (qPCR) data, drastically reduced by day 1 (Fig. 1C,D). SKI, which is closely related to *SNON*, is also present in hESCs, but its levels fluctuate negligibly between days 1 and 5 (Supplemental Fig. 1B,G). Several E3 ubiquitin ligases previously shown to target *SNON* to the proteasome are expressed in hESCs and during differentiation, including SMURF2, the RING domain-containing protein ARKADIA (RNF111), and the anaphase-promoting complex/cyclosome (APC/C) (Deheuninck and Luo 2009). *SMURF2* levels are up-regulated on day 1 and then plateau, while expression of *ARKADIA* and two APC/C subunits—*CDH1* (*FZR1*) and *ANAPC2*—increase gradually (Supplemental Fig. 1G). Similar results were observed by Western for SMURF2 and ARKADIA (Supplemental Fig. 1B). Other APC/C components, including *ANAPC11*, *CDC16*, and *CDC27*, were equally expressed at all time points (Supplemental Fig. 1G).

To confirm that the rapid loss of *SNON* protein is indeed due to proteasomal degradation, feeder-free HES3 cells were treated with two proteasome inhibitors: MG132 and lactacystin. When HES3 cells are differentiated in the presence of MG132 or lactacystin, *SNON* protein levels expectedly persist on day 1 (Supplemental Fig. 1H). Taken together, these data show that *SNON* expression is tightly linked to pluripotency and that *SNON* is targeted to the proteasome at the onset of DE formation.

SNON represses mesendoderm genes in hESCs

SNON expression in hESCs is controlled by *SMAD2/3* and *OCT4*, *SOX2*, and *NANOG*

ChIP–chip, recent ChIP–seq, and phylogenetic alignments reveal a series of conserved *SMAD2/3*- and *OCT4/SOX2/NANOG*-binding regions within the ~5-kb *SNON* promoter/enhancer (Fig. 1E; Supplemental Fig. 1I; Boyer et al. 2005; Zhu et al. 2005; Brown et al. 2011; Kim et al. 2011). We investigated the contribution of each of these elements to *SNON* regulation in undifferentiated hESCs using luciferase reporter constructs. Serial deletion analysis showed that the distal 5' *OCT4/SOX2/NANOG*-binding region is essential for robust reporter activity (Supplemental Fig. 1I). The more proximal *NANOG* and *SMAD2/3* elements contribute modestly to the regulation of *SNON* when compared with the –3.6-kb construct (Supplemental Fig. 1I). We also generated a series of –3.6-kb reporters in which *OCT4/SOX2/NANOG*- and *SMAD2/3*-binding sites were individually mutated (Fig. 1E; Supplemental Table 1). Consistent with our deletion studies, the mutated *OCT4/SOX2/NANOG* (mtO/S/N) construct in which distal *OCT4/SOX2/NANOG* binding is specifically disrupted shows significantly reduced (~13-fold) reporter activity (Fig. 1E). In contrast, mutating all four SBEs (mtSBEs) diminishes *SNON* reporter activity roughly twofold (Fig. 1E). Taken together, these data establish that *SNON* is TGF β -responsive and that *OCT4/SOX2/NANOG* serve as critical upstream regulators of *SNON* expression in hESCs.

SNON knockdown destabilizes pluripotency

To analyze *SNON* function, we first performed tetracycline (Tet)-inducible shRNA-mediated knockdown of *SNON* in S4TR5 cells, a derivative of the Shef4 hESC line that constitutively expresses the nuclear Tet repressor (TetRnls) (Zafarana et al. 2009). Two independent and previously validated human *SNON*-shRNA target sequences—shSNON(954) and shSNON(1307)—were transfected into S4TR5 cells (Sarker et al. 2005; Zhu et al. 2007). After selection, 11 of 12 clones for shSNON(954) and 10 of 12 clones for shSNON(1307) showed ~80% *SNON* knockdown by qPCR after 3 d of doxycycline (Dox) treatment (data not shown). Two clones, designated shSNON(954) and shSNON(1307), were chosen for further study. Both were indistinguishable by morphology (Supplemental Fig. 2A) and showed *OCT4*, *SOX2*, and *NANOG* levels comparable with the parental S4TR5 line by qPCR and Western (Supplemental Fig. 2B,C). Dox treatment resulted in a significant knockdown of *SNON* transcripts on day 1 (Fig. 2A,B). Interestingly, both shSNON clones lost their compact colony morphology after 5 d of continuous culture in Dox and appeared differentiated (Supplemental Fig. 2A). This was accompanied by a specific increase in PS (*BRACHYURY*, *MIXL1*, and *EOMES*), mesendoderm (*GSC*), and early DE (*FOXA2* and *SOX17*) markers beginning around day 3 (Fig. 2B; Supplemental Fig. 2B). Neuroectodermal gene expression was not above background (Supplemental Fig. 2B) and, importantly, levels of *OCT4*, *SOX2*, and *NANOG* remained unchanged (Supplemental Fig. 2B,C). In addition, loss of *SNON* did not impact *SKI*, *SMURF2*, *SMAD2/pSMAD2*, or *SMAD3/pSMAD3* levels, suggesting that the genes encoding these proteins are not regulated by the *SNON* corepressor in hESCs (Supplemental Fig. 2C). Consistent with our qPCR data, Western analysis




Figure 2. *SNON* knockdown induces differentiation in hESCs. (A) Schematic of the experimental design using two stable DOX-inducible S4TR5 hESC lines—shSNON(954) and shSNON(1307)—where DOX was replenished daily for 5 d. (B) *SNON*, *FOXA2*, and *SOX17* expression analysis by qPCR. RNA was collected on days 0–5 according to A. Data were normalized against *GUSB* and are depicted relative to S4TR5 (DOX–) at day 0 (= 1.0). *P*-values were calculated according to the Student's *t*-test. (*) *P* < 0.05. (C) Western blot analysis for *SNON*, *FOXA2*, and *SOX17*.

revealed that *SNON* is hardly detectable on day 1, and low levels of *FOXA2* were first detected on day 2, followed by *SOX17* on day 5 (Fig. 2C; Supplemental Fig. 2D). Immunofluorescence also confirmed loss of *SNON*, persistence of *OCT4*, and up-regulation of *SOX17* and *FOXA2* on day 5 (Supplemental Fig. 2E,F).

To provide additional evidence that loss of *SNON* results in differentiation, we performed a replating experiment whereby S4TR5 cells and both knockdown clones were passaged after 5 d of Dox treatment and cultured for four additional days in Dox. *OCT4*-DAB staining was used to identify hESC colonies and clearly revealed the dramatically reduced plating efficiency of the shSNON knockdown clones after passaging (Supplemental Fig. 2G). Taken together, our results show that diminished *SNON* levels in hESCs destabilizes pluripotency through precocious activation of mesendodermal genes.

SNON overexpression blocks differentiation into DE and mesoderm

We next performed the reciprocal experiment by overexpressing *SNON* in hESCs. After repeated attempts, we were unable to isolate HES3 subclones with levels greater than endogenous *SNON* (data not shown). We therefore chose to overexpress a stable truncated form of *SNON*—*SNON*(1–366)—that retains its ability to bind *SMAD2/3* and is sufficient for transcriptional repression but lacks the C-terminal lysine residues critical for ubiquitination and proteasomal degradation (Stroschein et al. 1999, 2001). Two *SNON*(1–366) clones (#1 and #9) were isolated that produce roughly equivalent levels of *SNON*(1–366) (Fig. 3A) and remain undifferentiated, with *OCT4* immunoreactivity comparable with wild-type HES3 (Supplemental Fig. 3A).

Clones #1 and #9 were then subjected to DE differentiation according to the schematic in Figure 1A, along with two controls: wild-type HES3 and HES3 harboring pCAG-IRES-puro (vector control). *NANOG*, *OCT4*, *SOX2*, and *DPPA4* levels declined with similar kinetics in controls