

2.89 (t, 16H,  $J=6$  Hz), 3.26–3.32 (m, 24H), 4.06 (t, 16H,  $J=5$  Hz), 5.14 (br, 8H).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta=28.3, 33.4, 39.4, 43.3, 49.3, 54.8, 63.5, 78.0, 79.2, 155.6, 169.6, 171.9$ . MS (FAB):  $m/z$  2150  $[\text{M}+\text{H}]^+$ . MS (MALDI):  $m/z$  calcd for  $\text{C}_{98}\text{H}_{164}\text{N}_{12}\text{O}_{40}$   $[\text{M}+\text{H}]^+$ : 2150.117; found: 2150.650.

**5.4.2. Synthesis of Boc-G2 7.** Compound **5** (112 mg, 0.05 mmol) was added to TFA (2.0 ml) and stirred for 2 h. The concentrated compound was reprecipitated with ethyl acetate and lyophilized. The deprotected compound (100 mg, 0.04 mmol) was dissolved in  $i$ PrOH (0.4 ml), and **1** (1.5 g, 7.07 mmol) and triethylamine (123  $\mu\text{l}$ , 0.88 mmol) were added. After stirring for 7 days at 45 °C under a nitrogen atmosphere, the reaction mixture was extracted with ethyl acetate and dried over  $\text{Na}_2\text{SO}_4$ . Purification of the concentrated compound was carried out by column chromatography on silica gel using ethyl acetate and acetone as eluent. A yellowish gum; trace.

## 5.5. Synthesis of benzyl ester glycine dendrons

**5.5.1. Synthesis of BnGD Boc-G1 8.** Glycine benzyl ester *p*-toluenesulfonate (2.0 g, 5.92 mmol) was dissolved in  $\text{CH}_3\text{CN}$  (10 ml), and **1** (3.8 g, 17.78 mmol) and triethylamine (1.6 ml, 11.86 mmol) were added. After stirring for 4 days at 60 °C under a nitrogen atmosphere, the reaction mixture was extracted with ethyl acetate and dried over  $\text{Na}_2\text{SO}_4$ . Purification of the concentrated compound was carried out by column chromatography on silica gel using ethyl acetate and hexane as eluent. A pale yellowish oil; 3.3 g, 93%. FTIR:  $\nu$  ( $\text{cm}^{-1}$ ) 3730, 3592, 3368, 3064, 2977, 1958, 1712.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta=1.43$  (s, 18H), 2.46 (t, 4H,  $J=7$  Hz), 2.98 (t, 4H,  $J=7$  Hz), 3.33–3.39 (m, 4H), 3.46 (s, 2H), 4.12 (t, 4H,  $J=5$  Hz), 5.13 (m, 4H), 7.34 (s, 5H).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta=28.4, 33.4, 39.6, 49.6, 54.5, 63.7, 66.3, 79.4, 128.2, 128.3, 128.5, 135.4, 155.7, 170.8, 172.1$ . MS (FAB):  $m/z$  596  $[\text{M}+\text{H}]^+$ . HRMS (FAB):  $m/z$  calcd for  $\text{C}_{29}\text{H}_{45}\text{N}_3\text{O}_{10}$   $[\text{M}+\text{H}]^+$ : 596.3105; found: 596.3177.

**5.5.2. Synthesis of BnGD Boc-G2 9.** Compound **8** (721 mg, 1.21 mmol) was added to TFA (3 ml) and stirred for 2 h. The concentrated compound was lyophilized. The deprotected compound (916 mg, 1.24 mmol) was dissolved in  $\text{CH}_3\text{CN}$  (14.1 ml), and **1** (6.1 g, 28.31 mmol) and triethylamine (1.2 ml, 8.49 mmol) were added. After stirring for 5 days at 45 °C under a nitrogen atmosphere, the reaction mixture was extracted with ethyl acetate and dried over  $\text{Na}_2\text{SO}_4$ . Purification of the concentrated compound was carried out by column chromatography on silica gel using ethyl acetate and hexane as eluent. A yellowish gum; 577 mg, 37%. FTIR:  $\nu$  ( $\text{cm}^{-1}$ ) 3730, 3372, 2976, 2011, 1957, 1715.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta=1.37$  (s, 36H), 2.39 (t, 12H,  $J=7$  Hz), 2.64 (t, 4H,  $J=6$  Hz), 2.75 (t, 8H,  $J=7$  Hz), 2.92 (t, 4H,  $J=7$  Hz), 3.27–3.33 (m, 8H), 3.40 (s, 2H), 4.02–4.08 (m, 12H), 5.06 (s, 2H), 5.17 (br, 4H), 7.28 (s, 5H).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta=28.3, 32.7, 33.2, 39.4, 49.3, 49.6, 51.8, 54.4, 62.0, 63.5, 66.0, 79.2, 128.0, 128.3, 135.3, 155.5, 170.6, 171.8, 171.9$ . MS (FAB):  $m/z$  1256.6  $[\text{M}+\text{H}]^+$ . HRMS (FAB):  $m/z$  calcd for  $\text{C}_{59}\text{H}_{97}\text{N}_7\text{O}_{22}$   $[\text{M}+\text{H}]^+$ : 1256.6687; found: 1256.6769. MS (MALDI):  $m/z$  calcd for  $\text{C}_{59}\text{H}_{97}\text{N}_7\text{O}_{22}$   $[\text{M}+\text{H}]^+$ : 1256.669; found: 1256.432.

**5.5.3. Synthesis of BnGD Boc-G3 10.** Compound **9** (535 mg, 0.43 mmol) was added to TFA (2 ml) and stirred for 2 h. The concentrated compound was lyophilized. The deprotected compound (631 mg, 0.38 mmol) was dissolved in  $\text{CH}_3\text{CN}$  (3.8 ml), and **1** (3.3 g, 15.24 mmol) and triethylamine (740  $\mu\text{l}$ , 5.33 mmol) were added. After stirring for 7 days at 45 °C under a nitrogen atmosphere, the reaction mixture was extracted with ethyl acetate and dried over  $\text{Na}_2\text{SO}_4$ . Purification of the concentrated compound was carried out by column chromatography on silica gel using ethyl acetate and

acetone as eluent. A brownish gum; 212 mg, 22%. FTIR:  $\nu$  ( $\text{cm}^{-1}$ ) 3726, 3593, 3371, 2977, 2837, 2013, 1729.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta=1.35$  (s, 72H), 2.38 (t, 28H,  $J=6$  Hz), 2.63 (t, 12H,  $J=5$  Hz), 2.74 (t, 24H,  $J=6$  Hz), 2.89 (t, 4H,  $J=7$  Hz), 3.27–3.32 (m, 16H), 3.37 (s, 2H), 4.01–4.07 (m, 28H), 5.04 (s, 2H), 5.19 (br, 8H), 7.27 (s, 5H).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta=28.3, 32.5, 32.7, 33.2, 39.4, 49.3, 49.4, 49.6, 51.7, 51.8, 54.5, 62.0, 62.1, 63.4, 66.0, 79.1, 127.9, 128.0, 128.3, 135.4, 155.5, 170.6, 171.7, 171.8, 171.8$ . MS (MALDI):  $m/z$  calcd for  $\text{C}_{119}\text{H}_{201}\text{N}_{15}\text{O}_{46}$   $[\text{M}+\text{H}]^+$ : 2578.943; found: 2576.640.

## 5.6. Synthesis of amide glycine dendrimers

**5.6.1. Synthesis of AG Boc-G1 11.** Compound **8** (150 mg, 0.25 mmol) was dissolved in  $\text{CH}_3\text{CN}$  (2.5 ml) and PdOH/C (150 mg) was added. After stirring for 2 h under a hydrogen atmosphere, the reaction mixture was filtered. The concentrated compound was dissolved in  $\text{CH}_3\text{CN}$  (2.5 ml), and **4** (37 mg, 0.042 mmol), PyBOP (109 mg, 0.21 mmol), and triethylamine (47  $\mu\text{l}$ , 0.34 mmol) were added. After stirring for 1 day at room temperature under a nitrogen atmosphere, the reaction mixture was extracted with ethyl acetate and dried over  $\text{Na}_2\text{SO}_4$ . Purification of the concentrated compound was carried out by column chromatography on silica gel using ethyl acetate and acetone as eluent. A pale yellowish foam; 87 mg, 87%. FTIR:  $\nu$  ( $\text{cm}^{-1}$ ) 3331, 2974, 2927, 1712.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta=1.43$  (s, 72H), 2.49–2.53 (m, 28H), 2.80 (t, 16H,  $J=6$  Hz), 3.13 (s, 8H), 3.33–3.40 (m, 16H), 3.93 (d, 8H,  $J=5$  Hz), 4.12 (t, 16H,  $J=5$  Hz), 5.28 (br, 8H), 8.04 (t, 4H,  $J=6$  Hz).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta=28.5, 32.1, 39.5, 41.4, 43.1, 49.6, 58.3, 63.9, 78.5, 79.4, 155.8, 168.2, 171.3, 172.2$ . MS (MALDI):  $m/z$  calcd for  $\text{C}_{106}\text{H}_{176}\text{N}_{16}\text{O}_{44}$   $[\text{M}+\text{H}]^+$ : 2379.613; found: 2378.987.

**5.6.2. Synthesis of AG Boc-G2 12.** Compound **9** (188 mg, 0.15 mmol) was dissolved in  $\text{CH}_3\text{CN}$  (1.5 ml) and PdOH/C (190 mg) was added. After stirring for 2 h under a hydrogen atmosphere, the reaction mixture was filtered. The concentrated compound was dissolved in  $\text{CH}_3\text{CN}$  (1.5 ml), and **4** (22 mg, 0.025 mmol), PyBOP (97 mg, 0.19 mmol), and triethylamine (28  $\mu\text{l}$ , 0.20 mmol) were added. After stirring overnight at room temperature under a nitrogen atmosphere, the reaction mixture was extracted with ethyl acetate and dried over  $\text{Na}_2\text{SO}_4$ . Purification of the concentrated compound was carried out by column chromatography on silica gel using ethyl acetate and acetone as eluent. A yellowish gum; 45 mg, 36%. FTIR:  $\nu$  ( $\text{cm}^{-1}$ ) 3364, 2926, 2854, 1713.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta=1.42$  (s, 144H), 2.47–2.51 (m, 60H), 2.74–2.85 (m, 64H), 3.14 (s, 8H), 3.33–3.38 (m, 32H), 3.91 (d, 8H,  $J=4$  Hz), 4.12 (t, 48H,  $J=5$  Hz), 5.26 (br, 16H), 7.96 (t, 4H,  $J=5$  Hz).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta=28.4, 31.7, 32.6, 39.5, 49.3, 49.8, 52.0, 57.9, 63.7, 78.5, 79.3, 155.7, 171.3, 172.0$ . MS (MALDI):  $m/z$  calcd for  $\text{C}_{226}\text{H}_{384}\text{N}_{32}\text{O}_{92}$   $[\text{M}+\text{H}]^+$ : 5022.628; found: 5021.661.

**5.6.3. Synthesis of AG Boc-G3 13.** Compound **10** (162 mg, 0.063 mmol) was dissolved in  $\text{CH}_3\text{CN}$  (0.63 ml) and PdOH/C (160 mg) was added. After stirring for 4 h under a hydrogen atmosphere, the reaction mixture was filtered. The concentrated compound was dissolved in  $\text{CH}_3\text{CN}$  (0.63 ml), and **4** (9 mg, 0.010 mmol), PyBOP (41 mg, 0.079 mmol), and triethylamine (12  $\mu\text{l}$ , 0.083 mmol) were added. After stirring for 1 day at room temperature under a nitrogen atmosphere, the reaction mixture was extracted with ethyl acetate and dried over  $\text{Na}_2\text{SO}_4$ . Purification of the concentrated compound was carried out by column chromatography on silica gel using ethyl acetate and acetone as eluent. A brownish gum; 24 mg, 22%. FTIR:  $\nu$  ( $\text{cm}^{-1}$ ) 3364, 2925, 1701.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta=1.43$  (s, 288H), 2.46 (t, 124H,  $J=6$  Hz), 2.72 (t, 48H,  $J=6$  Hz), 2.82 (t, 112H,  $J=6$  Hz), 3.15 (s, 8H), 3.33–3.39 (m, 64H), 3.91 (s, 8H), 4.09–4.14 (m, 112H), 5.23 (br, 32H), 7.94 (br, 4H).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta=28.4, 32.7, 39.6, 49.7, 52.0, 62.2, 63.6, 79.4, 155.9, 172.3$ .

MS (MALDI):  $m/z$  calcd for  $C_{466}H_{800}N_{64}O_{188}$   $[M+H]^+$ : 10,308.657; found: 10,303.944.

## 5.7. Synthesis of click dendrons

**5.7.1. Synthesis of CD Boc-G1 14.** Propargyl amine hydrochloride (915 mg, 10.0 mmol) was dissolved in  $CH_3CN$  (10 ml), and **1** (6.5 g, 30.0 mmol) and triethylamine (2.8 ml, 20.00 mmol) were added. After stirring for 3 days at 45 °C under a nitrogen atmosphere, the reaction mixture was extracted with ethyl acetate and dried over  $Na_2SO_4$ . Purification of the concentrated compound was carried out by column chromatography on silica gel using ethyl acetate and hexane as eluent. A pale yellowish oil; 5.3 g, quant. FTIR:  $\nu$  ( $cm^{-1}$ ) 3955, 3728, 3592, 3356, 2976, 2104, 2013, 1706.  $^1H$  NMR ( $CDCl_3$ )  $\delta$ =1.34 (s, 18H), 2.16 (t, 1H,  $J$ =2 Hz), 2.38 (t, 4H,  $J$ =7 Hz), 2.74 (t, 4H,  $J$ =7 Hz), 3.23–3.30 (m, 4H), 3.35 (d, 2H,  $J$ =2 Hz), 4.04 (t, 4H,  $J$ =5 Hz), 5.21 (br, 2H).  $^{13}C$  NMR ( $CDCl_3$ )  $\delta$ =28.7, 33.2, 39.8, 41.9, 49.2, 63.8, 74.0, 77.9, 79.5, 156.0, 172.2. MS (FAB):  $m/z$  486  $[M+H]^+$ . HRMS (FAB):  $m/z$  calcd for  $C_{23}H_{39}N_3O_8$   $[M+H]^+$ : 486.2737; found: 486.2823.

**5.7.2. Synthesis of CD Boc-G2 15.** Compound **14** (1.2 g, 2.54 mmol) was added to TFA (5.0 ml) and stirred for 2 h. The concentrated compound was lyophilized. The deprotected compound (510 mg, 0.81 mmol) was dissolved in  $CH_3CN$  (8.1 ml), and **1** (3.5 g, 16.26 mmol) and triethylamine (676  $\mu$ l, 4.87 mmol) were added. After stirring for 4 days at 45 °C under a nitrogen atmosphere, the reaction mixture was extracted with ethyl acetate and dried over  $Na_2SO_4$ . Purification of the concentrated compound was carried out by column chromatography on silica gel using ethyl acetate and hexane as eluent. A yellowish gum; 223 mg, 24%. FTIR:  $\nu$  ( $cm^{-1}$ ) 3725, 3372, 2976, 2103, 2014, 1711.  $^1H$  NMR ( $CDCl_3$ )  $\delta$ =1.40 (s, 36H), 2.19 (t, 1H,  $J$ =2 Hz), 2.43 (t, 12H,  $J$ =7 Hz), 2.69 (t, 4H,  $J$ =6 Hz), 2.79 (t, 12H,  $J$ =7 Hz), 3.31–3.37 (m, 10H), 4.06–4.12 (m, 12H), 5.14 (br, 4H).  $^{13}C$  NMR ( $CDCl_3$ )  $\delta$ =28.1, 32.6, 32.6, 39.3, 41.5, 48.6, 49.5, 51.7, 61.9, 63.4, 73.2, 77.6, 79.1, 155.4, 171.6, 171.8. MS (FAB):  $m/z$  1146.6  $[M+H]^+$ . HRMS (FAB):  $m/z$  calcd for  $C_{53}H_{91}N_7O_{20}$   $[M+H]^+$ : 1146.6319; found: 1146.6372. MS (MALDI):  $m/z$  calcd for  $C_{53}H_{91}N_7O_{20}$   $[M+H]^+$ : 1146.632; found: 1146.663.

**5.7.3. Synthesis of CD Boc-G3 16.** Compound **15** (196 mg, 0.17 mmol) was added to TFA (3.0 ml) and stirred for 2 h. The concentrated compound was lyophilized. The deprotected compound was dissolved in  $CH_3CN$  (1.6 ml), and **1** (1.4 g, 6.56 mmol) and triethylamine (318  $\mu$ l, 2.30 mmol) were added. After stirring for 7 days at 45 °C under a nitrogen atmosphere, the reaction mixture was extracted with ethyl acetate and dried over  $Na_2SO_4$ . Purification of the concentrated compound was carried out by column chromatography on silica gel using ethyl acetate and acetone as eluent. A brownish gum; 55 mg, 14%. FTIR:  $\nu$  ( $cm^{-1}$ ) 3727, 3374, 2976, 2838, 2102, 2014, 1731.  $^1H$  NMR ( $CDCl_3$ )  $\delta$ =1.42 (s, 72H), 2.22 (t, 1H,  $J$ =2 Hz), 2.45 (t, 28H,  $J$ =7 Hz), 2.70 (t, 12H,  $J$ =6 Hz), 2.81 (t, 28H,  $J$ =7 Hz), 3.33–3.39 (m, 18H), 4.08–4.14 (m, 28H), 5.15 (br, 8H).  $^{13}C$  NMR ( $CDCl_3$ )  $\delta$ =28.4, 32.7, 32.9, 32.9, 39.6, 41.8, 48.9, 49.6, 49.8, 51.9, 52.0, 62.2, 62.4, 63.7, 73.5, 78.0, 79.4, 155.7, 171.9, 172.0, 172.0. MS (MALDI):  $m/z$  calcd for  $C_{113}H_{195}N_{15}O_{44}$   $[M+H]^+$ : 2468.832; found: 2468.015.

## 5.8. Synthesis of click chemistry dendrimers

**5.8.1. Synthesis of CC Boc-G1 17.** Compounds **2** (50 mg, 0.090 mmol) and **14** (201 mg, 0.41 mmol) were dissolved in a 4:1 solvent ratio of THF/ $H_2O$  (4.1 ml), and  $CuSO_4 \cdot 5H_2O$  (21 mg, 0.082 mmol) and sodium ascorbate (33 mg, 0.17 mmol) were added. After stirring overnight, the reaction mixture was extracted with ethyl acetate and dried over  $Na_2SO_4$ . Purification of the

concentrated compound was carried out by column chromatography on silica gel using ethyl acetate and acetone as eluent. A pale yellowish foam; 184 mg, 82%. FTIR:  $\nu$  ( $cm^{-1}$ ) 3367, 2928, 1701.  $^1H$  NMR ( $CDCl_3$ )  $\delta$ =1.34 (s, 72H), 2.41–2.45 (m, 28H), 2.69 (t, 16H,  $J$ =6 Hz), 3.23–3.29 (m, 16H), 3.75 (s, 8H), 4.03 (t, 16H,  $J$ =5 Hz), 5.06 (s, 8H), 5.30 (br, 8H), 7.58 (s, 4H).  $^{13}C$  NMR ( $CDCl_3$ )  $\delta$ =28.3, 32.5, 39.4, 42.6, 47.9, 48.7, 50.9, 53.8, 63.4, 79.1, 124.1, 144.0, 155.6, 164.8, 172.0. MS (MALDI):  $m/z$  calcd for  $C_{110}H_{176}N_{24}O_{40}$   $[M+H]^+$ : 2475.712; found: 2474.737.


**5.8.2. Synthesis of CC Boc-G2 18.** Compounds **2** (25 mg, 0.046 mmol) and **15** (318 mg, 0.28 mmol) were dissolved in a 4:1 solvent ratio of THF/ $H_2O$  (2.8 ml), and  $CuSO_4 \cdot 5H_2O$  (14 mg, 0.055 mmol) and sodium ascorbate (22 mg, 0.11 mmol) were added. After stirring overnight, the reaction mixture was extracted with ethyl acetate and dried over  $Na_2SO_4$ . Purification of the concentrated compound was carried out by column chromatography on silica gel using ethyl acetate and acetone as eluent. A yellowish gum; 85 mg, 36%. FTIR:  $\nu$  ( $cm^{-1}$ ) 3371, 2976, 2110, 1731.  $^1H$  NMR ( $CDCl_3$ )  $\delta$ =1.42 (s, 144H), 2.42–2.53 (m, 60H), 2.70 (t, 16H,  $J$ =6 Hz), 2.81 (t, 48H,  $J$ =6 Hz), 3.33–3.39 (m, 32H), 3.82 (s, 8H), 4.08–4.13 (m, 48H), 5.13 (s, 8H), 5.22 (br, 16H), 7.64 (s, 4H).  $^{13}C$  NMR ( $CDCl_3$ )  $\delta$ =28.4, 32.5, 32.9, 39.6, 42.9, 48.1, 48.7, 49.8, 52.0, 62.2, 63.7, 79.4, 124.1, 155.7, 164.9, 172.1. MS (MALDI):  $m/z$  calcd for  $C_{230}H_{384}N_{40}O_{88}$   $[M+H]^+$ : 5118.727; found: 5118.790.

**5.8.3. Synthesis of CC Boc-G3 19.** Compounds **2** (8 mg, 0.014 mmol) and **16** (221 mg, 0.089 mmol) were dissolved in a 4:1 solvent ratio of THF/ $H_2O$  (890  $\mu$ l), and  $CuSO_4 \cdot 5H_2O$  (4 mg, 0.018 mmol) and sodium ascorbate (7 mg, 0.036 mmol) were added. After stirring overnight, the reaction mixture was extracted with ethyl acetate and dried over  $Na_2SO_4$ . Purification of the concentrated compound was carried out by column chromatography on silica gel using ethyl acetate and acetone as eluent. A brownish gum; 30 mg, 21%. FTIR:  $\nu$  ( $cm^{-1}$ ) 3363, 2925, 1703.  $^1H$  NMR ( $CDCl_3$ )  $\delta$ =1.43 (s, 288H), 2.42–2.54 (m, 124H), 2.71 (t, 48H,  $J$ =6 Hz), 2.82 (t, 112H,  $J$ =7 Hz), 3.34–3.40 (m, 64H), 3.83 (s, 8H), 4.09–4.15 (m, 112H), 5.14 (s, 8H), 5.24 (br, 32H), 7.64 (s, 4H).  $^{13}C$  NMR ( $CDCl_3$ )  $\delta$ =28.5, 32.7, 32.9, 39.6, 49.6, 49.8, 52.0, 62.2, 63.7, 79.4, 155.7, 172.1, 172.1. MS (MALDI):  $m/z$  calcd for  $C_{470}H_{800}N_{72}O_{184}$   $[M+H]^+$ : 10,404.756; found: 10,403.867.

## References and notes

- [1] D.A. Tomalia, H. Baker, J. Dewald, M. Hall, G. Kallos, S. Martin, J. Roeck, J. Ryder, P. Smith, *Polym. J.* 17 (1985) 117.
- [2] D.A. Tomalia, H. Baker, J. Dewald, M. Hall, G. Kallos, S. Martin, J. Roeck, J. Ryder, P. Smith, *Macromolecules* 19 (1986) 2466.
- [3] D.A. Tomalia, A.M. Naylor, W.A. Goddard, *Angew. Chem., Int. Ed. Engl.* 29 (1990) 138.
- [4] A.W. Bosman, H.M. Janssen, E.W. Meijer, *Chem. Rev.* 99 (1999) 1665.
- [5] S. Svenson, D.A. Tomalia, *Adv. Drug Delivery Rev.* 57 (2005) 2106.
- [6] A. D'Emanuele, D. Attwood, *Adv. Drug Delivery Rev.* 57 (2005) 2147.
- [7] C. Dufes, I.F. Uchegbu, A.G. Schatzlein, *Adv. Drug Delivery Rev.* 57 (2005) 2177.
- [8] R. Duncan, L. Izzo, *Adv. Drug Delivery Rev.* 57 (2005) 2215.
- [9] H. Kobayashi, M.W. Brechbiel, *Adv. Drug Delivery Rev.* 57 (2005) 2271.
- [10] C.C. Lee, J.A. Mackay, J.M.J. Fréchet, F.C. Szoka, *Nat. Biotechnol.* 23 (2005) 1517.
- [11] C.M. Paleos, D. Tsiourvas, Z. Sideratou, *Mol. Pharm.* 4 (2007) 169.
- [12] J. Haensler, F.C. Szoka, *Bioconjugate Chem.* 4 (1993) 372.
- [13] J.F. Kukowska-Latallo, A.U. Bielinska, J. Johnson, R. Spindker, D.A. Tomalia, J.R. Baker, *Proc. Natl. Acad. Sci. U.S.A.* 93 (1996) 4897.
- [14] B.H. Zinselmeyer, S.P. Mackay, A.G. Schatzlein, I.F. Uchegbu, *Pharm. Res.* 19 (2002) 960.
- [15] A.J. Hollins, M. Benboubetra, Y. Omid, B.H. Zinselmeyer, A.G. Schatzlein, I.F. Uchegbu, S. Akhtar, *Pharm. Res.* 21 (2004) 458.
- [16] O. Boussif, F. Lezoualc'h, M.A. Zanta, M.D. Mergny, D. Scherman, B. Demeneix, J. Behr, *Proc. Natl. Acad. Sci. U.S.A.* 92 (1995) 7297.
- [17] J.H. Lee, Y. Lim, J.S. Choi, Y. Lee, T. Kim, H.J. Kim, J.K. Yoon, K. Kim, J. Park, *Bioconjugate Chem.* 14 (2003) 1214.
- [18] H. Arima, F. Kihara, F. Hirayama, K. Uekama, *Bioconjugate Chem.* 12 (2001) 476.
- [19] J.S. Choi, K. Nam, J. Park, J. Kim, J. Lee, J. Park, *J. Controlled Release* 99 (2004) 445.
- [20] K. Kono, H. Akiyama, T. Takahashi, T. Takagishi, A. Harada, *Bioconjugate Chem.* 16 (2005) 208.

- [21] L. Tziveleka, A.G. Psarra, D. Tsiourvas, C.M. Paleos, J. Controlled Release 117 (2007) 137.
- [22] T. Kim, J. Baek, J.K. Yoon, J.S. Choi, K. Kim, J. Park, Bioconjugate Chem. 18 (2007) 309.
- [23] T. Kim, J. Baek, C.Z. Bai, J. Park, Biomaterials 28 (2007) 2061.
- [24] M.A. Carnahan, M.W. Grinstaff, J. Am. Chem. Soc. 123 (2001) 2905.
- [25] M.A. Carnahan, M.W. Grinstaff, Macromolecules 34 (2001) 7648.
- [26] M.A. Carnahan, M.W. Grinstaff, Macromolecules 39 (2006) 609.
- [27] E.R. Gillies, J.M.J. Fréchet, J. Am. Chem. Soc. 124 (2002) 14137.
- [28] S. Tang, L.J. Martinez, A. Sharma, M. Chai, Org. Lett. 8 (2006) 4421.
- [29] C.C. Lee, E.R. Gillies, M.E. Fox, S.J. Guillaudeu, J.M.J. Fréchet, E.E. Dy, F.C. Szoka, Proc. Natl. Acad. Sci. U.S.A. 103 (2006) 16649.
- [30] D.G. van der Poll, H.M. Kieler-Ferguson, W.C. Floyd, S.J. Guillaudeu, K. Jerger, F. C. Szoka, J.M.J. Fréchet, Bioconjugate Chem. 21 (2010) 764.
- [31] M. Pittelkow, J.B. Christensen, Org. Lett. 7 (2005) 1295.
- [32] F. Himo, T. Lovell, R. Hilgraf, V.V. Rostovtsev, L. Noodleman, K.B. Sharpless, V.V. Fokin, J. Am. Chem. Soc. 127 (2005) 210.
- [33] P. Wu, A.K. Feldman, A.K. Nugent, C.J. Hawker, A. Scheel, B. Voit, J. Pyun, J.M.J. Fréchet, K.B. Sharpless, V.V. Fokin, Angew. Chem., Int. Ed. 43 (2004) 3928.
- [34] M.J. Joralemon, R.K. O'Reilly, J.B. Matson, A.K. Nugent, C.J. Hawker, K.L. Wooley, Macromolecules 38 (2005) 5436.
- [35] J.W. Lee, B. Kim, H.J. Kim, S.C. Han, W.S. Shin, S. Jin, Macromolecules 39 (2006) 2418.
- [36] J.W. Lee, J.H. Kim, B. Kim, J.H. Kim, W.S. Shin, S. Jin, Tetrahedron 62 (2006) 9193.
- [37] J.W. Lee, B. Kim, J.H. Kim, W.S. Shin, S. Jin, J. Org. Chem. 71 (2006) 4988.
- [38] C.N. Urbani, C.A. Bell, D.E. Lonsdale, M.R. Whittaker, M.J. Monteiro, Macromolecules 40 (2007) 7056.
- [39] J. Luten, C.F. von Nostrum, S.C. De Smedt, W.E. Hennink, J. Controlled Release 126 (2008) 97.
- [40] H.J. Kim, M.S. Kwon, J.S. Choi, B.H. Kim, J.K. Yoon, K. Kim, J. Park, Bioorg. Med. Chem. 15 (2007) 1708.
- [41] T.H. Fife, R. Singh, R. Bembi, J. Org. Chem. 67 (2002) 3179.
- [42] England, W. P.; Stoddard, G. J.; Scobbo, J. J. U.S. Patent 5,310,795, 1994.
- [43] F.M. Menger, V.A. Migulin, J. Org. Chem. 64 (1999) 8916.



# Triplex-forming ability of oligonucleotides containing 1-aryl-1,2,3-triazole nucleobases linked via a two atom-length spacer



Yoshiyuki Hari\*, Motoi Nakahara, Satoshi Obika\*

Graduate School of Pharmaceutical Sciences, Osaka University, 1-6 Yamadaoka, Suita, Osaka 565-0871, Japan

## ARTICLE INFO

### Article history:

Received 1 May 2013

Revised 22 May 2013

Accepted 23 May 2013

Available online 31 May 2013

### Keywords:

Artificial nucleobases

1,3-Dipolar cycloaddition

Triazoles

Triplex DNA

Triplex-forming oligonucleotides

## ABSTRACT

Phosphoramidites containing 2-propynyloxy or 1-butyne-4-yl as nucleobase precursors were synthesized and introduced into oligonucleotides using an automated DNA synthesizer. Copper-catalyzed alkyne-azide 1,3-dipolar cycloaddition of the oligonucleotides with various azides gave the corresponding triazolylated oligonucleotides, triplex-forming ability of these synthetic oligonucleotides with double-stranded DNA targets was evaluated by UV melting experiments. It was found that nucleobases containing 2-(1-*m*-carbonylamino-phenyl-1,2,3-triazol-4-yl)ethyl units likely interacted with A of a TA base pair in a parallel triplex DNA.

© 2013 Elsevier Ltd. All rights reserved.

## 1. Introduction

A triplex-forming oligonucleotide (TFO) that can form a triplex DNA with double-stranded DNA (dsDNA) would be a promising tool for gene-targeting therapy and genetic diagnosis.<sup>1</sup> In general, triplex DNA can be sequence-selectively formed by recognition of AT and GC base pairs in dsDNA by T and C in TFO via Hoogsteen hydrogen bonds, respectively. The sequence of dsDNA targeted by TFO is limited to these purine–pyrimidine base pairs. Therefore, towards recognition of pyrimidine–purine base pairs such as CG and TA, a number of synthetic nucleotides have been developed to date.<sup>2</sup> Concerning CG base pair recognition, several promising nucleotides have already been synthesized. In contrast, there is almost no nucleotide recognizing a TA base pair with high selectivity and stability. The difficulty to find nucleotides for TA base pair recognition is considered to be caused by disturbance of hydrogen bonding interaction with the 4-carbonyl group due to the steric bulkiness of the 5-methyl group in T.

Recently, we have tried to develop nucleotides capable of recognizing a CG base pair by means of the post-elongation modification (PEM) methods such as copper-catalyzed alkyne-azide 1,3-dipolar cycloaddition<sup>3</sup> and nucleophilic substitution reaction<sup>4</sup> to construct artificial nucleobases. Since this PEM method allows us to efficiently synthesize various derivatives in a short period, the detailed molecular design of nucleobases based on the experimen-

tal results and their evaluation can be implemented. Therefore, we considered that the PEM method can be applied to investigate nucleobases for TA base pair recognition and this could bring useful information concerning molecular design for TA recognition. In this study, nucleobases with two atom-length spacers such as methyleneoxy and ethylene were designed to avoid the steric repulsion from 5-methyl group of T and with the recognition unit to bind to the opposite A. The recognition unit was synthesized by using copper-catalyzed alkyne-azide 1,3-dipolar cycloaddition as PEM method to investigate the fine structure (Fig. 1). The details of our results obtained are described.

## 2. Results and discussion

### 2.1. Synthesis

The synthesis of an oligonucleotide **1** bearing a 2-propynyloxy group was carried out according to Scheme 1. Chlorosugar **2**




Figure 1. Nucleobases designed in this study.

\* Corresponding authors. Tel./fax: +81 6 6879 8201 (Y. Hari); tel.: +81 6 6879 8200; fax: +81 6 6879 8204 (S. Obika).

E-mail addresses: [hari@phs.osaka-u.ac.jp](mailto:hari@phs.osaka-u.ac.jp) (Y. Hari), [obika@phs.osaka-u.ac.jp](mailto:obika@phs.osaka-u.ac.jp) (S. Obika).


underwent reaction with propargyl alcohol in MeCN to give desired  $\beta$ -isomer **3** in 34% yield, the stereochemistry of which was determined by NOESY correlations of deprotected **4** shown in Scheme 1. The primary alcohol of **4** was protected by DMTrCl in the presence of pyridine and phosphorylation of **5** gave phosphoramidite **6**, a building block for oligonucleotide synthesis. The oligonucleotide precursor **1** for copper-catalyzed alkyne-azide 1,3-dipolar cycloaddition was prepared on an automated DNA synthesizer using standard phosphoramidite chemistry.

The synthesis of oligonucleotide **7** with the incorporation of 1-butyn-4-yl group was shown in Scheme 2. Reduction of compound **8** prepared according to the report<sup>5</sup> with LiAlH<sub>4</sub>, tosylation followed by treatment with lithium acetylide-ethylenediamine complex was carried out in three-steps; this gave the compound **9** in 53% yield. Phosphoramidite **10** was obtained by phosphorylation of **9** and the yield was 97%.

Copper-catalyzed alkyne-azide 1,3-dipolar cycloaddition of oligonucleotides **1** and **7** with various azides was carried out according to the reaction conditions previously reported in our earlier study.<sup>3a</sup> This was performed to convert the oligonucleotides into TFOs (Scheme 3, and see experimental for the synthesis of azides). In all cases, the corresponding TFOs **11a–h** and **12a–j** were successfully obtained by reversed-phase HPLC purification after the reaction.

## 2.2. Evaluation


In triplex DNA between TFO and dsDNA, base pair-recognition ability of the nucleobases consisting of 1-aryl-1,2,3-triazol-4-yl unit and spacer unit in TFO was evaluated by UV-melting experiment. The results of TFOs **11** with methyleneoxy spacer and TFOs **12** with ethylene spacer are summarized in Tables 1 and 2, respectively. The measurement was carried out under neutral conditions and hairpin dsDNA targets containing four kinds of natural base pairs, TA, CG, GC and AT base pairs for YZ were used (For details see footnote of Table 1). Regarding TFOs **11** with methyleneoxy spacer, when substituent at 1-position of triazole was a simple phenyl group (**11a**) without any functional group for hydrogen bond formation, the range of  $T_m$  values against all four hairpin dsDNA targets was from 20 °C to 22 °C. *o*-Hydroxyphenyl derivative (**11b**) gave almost same  $T_m$  values as **11a**, while *o*-carbamoyl congener (**11c**) led to a decrease in  $T_m$  values against any base pairs likely due to the existence of bulky carbamoyl group at the




**Scheme 2.** Reagents and conditions: (i) LiAlH<sub>4</sub>, THF, 0 °C, 4 h, (ii) TsCl, pyridine, rt, 12 h, (iii) lithium acetylide-ethylenediamine complex, THF, rt, 9 h, 53% for three-steps, (iv) *i*-Pr<sub>2</sub>NP(Cl)OCH<sub>2</sub>CH<sub>2</sub>CN, *i*-Pr<sub>2</sub>NEt, CH<sub>2</sub>Cl<sub>2</sub>, rt, 5 h, 97%, (v) DNA synthesis (C = 2'-deoxy-5-methylcytidine).

*o*-position (Fig. 2). Among TFOs **11d–h** including *m*- or *p*-substituted phenyl analogs, only TFO **11e** with *m*-hydroxymethyl group and TFO **11h** with *m*-ureidophenyl one seem to stabilize the triplexes with dsDNA (YZ = TA) compared to that observed with TFO **11a** including an unsubstituted phenyl group though without stabilization selective to dsDNA (YZ = TA) (Table 1 and Fig. 2).

Among TFOs **12** in which an ethylene spacer was incorporated, unsubstituted phenyl analog (**12a**) increased the  $T_m$  value by +2 °C against dsDNA for either TA or CG base pair compared to the corresponding methyleneoxy spacer (**11a**). *o*-Substituents (**12b** and **12c**) decreased the stability of triplexes formed with all dsDNA targets (Fig. 3). This is likely due to the steric repulsion between the *o*-substituents and the target base pairs. No major changes were observed in TFOs **12d–f**. TFO **12g** containing *m*-carbamoyl group stabilized the triplex with dsDNA (YZ = TA) and the  $\Delta T_m$  value observed was +3 °C. Significant improvement was observed in TFO **12h** containing the *m*-ureido group. The  $\Delta T_m$  value of **12h** against dsDNA with TA base pair was +6 °C, which was significantly higher than that observed ( $\Delta T_m = +2$  °C to +3 °C) against the other dsDNA targets.



**Scheme 1.** Reagents and conditions: (i) propargyl alcohol, MeCN, rt, 4 h, 34%, (ii) NaOMe, MeOH, rt, 5 h, quant., (iii) DMTrCl, pyridine, rt, 6 h, 98%, (iv) *i*-Pr<sub>2</sub>NP(Cl)OCH<sub>2</sub>CH<sub>2</sub>CN, *i*-Pr<sub>2</sub>NEt, CH<sub>2</sub>Cl<sub>2</sub>, rt, 15 h, 90%, (v) DNA synthesis (C = 2'-deoxy-5-methylcytidine).



**Scheme 3.** Reagents and conditions: (i) R-N<sub>3</sub>, CuSO<sub>4</sub>, sodium ascorbate, tris[(1-benzyl-1,2,3-triazol-4-yl)methyl]amine (TBTA), DMSO-phosphate buffer (10 mM, pH 7.0), rt, 0.5–12 h, 67–91%.

**Table 1**  
T<sub>m</sub> values (°C) of triplexes between TFOs **11** including a methyleneoxy spacer and four hairpin dsDNA targets<sup>a,b</sup>

TFO	YZ			
	TA	CG	GC	AT
11a	20	22	22	20
11b	20 (±0)	23 (+1)	20 (−2)	20 (±0)
11c	17 (−3)	21 (−1)	19 (−3)	18 (−2)
11d	21 (+1)	24 (+2)	21 (−1)	20 (±0)
11e	22 (+2)	25 (+3)	23 (+1)	23 (+3)
11f	19 (−1)	22 (±0)	19 (−3)	19 (−1)
11g	19 (−1)	23 (+1)	20 (−2)	19 (−1)
11h	23 (+3)	25 (+3)	23 (+1)	21 (+1)

<sup>a</sup> Conditions: 10 mM sodium cacodylate buffer (pH 6.8), 100 mM KCl and 50 mM MgCl<sub>2</sub>. The final concentration of each oligonucleotide used was 1.89 μM. The sequence of hairpin dsDNA targets is as follows: 5'-GGCAAAAAGAYAGAGACGC-hexaethyleneglycol-GCGTCTCTZTCTTTTTGCC-3' (YZ = TA, CG, GC or AT).

<sup>b</sup> ΔT<sub>m</sub>: Difference in the T<sub>m</sub> value from that of **11a** is shown in parenthesis.

In the above analysis on substituent in the phenyl ring (Tables 1 and 2), TFO **12** with ethylene spacer showed higher ΔT<sub>m</sub> values than TFO **11** with methyleneoxy spacer (Figs. 2 and 3). In addition, a high ΔT<sub>m</sub> value of TFO **12h** to dsDNA (YZ = TA) was observed when compared with TFO **11h** with the same *m*-ureidophenyl group. These results may imply that the ethylene spacer is less flexible than methyleneoxy spacer and consequently the


**Table 2**

T<sub>m</sub> values (°C) of triplexes between TFOs **12** including a ethylene spacer and four hairpin dsDNA targets<sup>a,b</sup>


TFO	YZ			
	TA	CG	GC	AT
12a	22	24	20	21
12b	18 (−4)	22 (−2)	19 (−1)	20 (−1)
12c	18 (−4)	20 (−4)	19 (−1)	16 (−5)
12d	22 (±0)	22 (−2)	21 (+1)	21 (±0)
12e	23 (+1)	23 (−1)	20 (±0)	19 (−2)
12f	22 (±0)	23 (−1)	19 (−1)	19 (−2)
12g	25 (+3)	25 (+1)	22 (+2)	22 (+1)
12h	28 (+6)	27 (+3)	23 (+3)	23 (+2)
12i	27 (+5)	27 (+3)	23 (+3)	22 (+1)
12j	27 (+5)	26 (+2)	21 (+1)	21 (±0)

<sup>a</sup> Conditions are shown in the footnote of Table 1.

<sup>b</sup> ΔT<sub>m</sub>: Difference in the T<sub>m</sub> value from that of **12a** is shown in parenthesis.




**Figure 2.** The difference in the T<sub>m</sub> values of TFOs **11b–h** from that of TFO **11a**.



**Figure 3.** The difference in the T<sub>m</sub> values of TFOs **12b–j** from that of TFO **12a**.

aryltriazole moiety attached to ethylene spacer lies in a suitable space for TA base pair recognition.

C<sub>2</sub>-Symmetric 3,5-bisureidophenyl group (**12i**) designed with consideration of free rotation of bond between aryl moiety and triazole ring gave almost same result observed with the monoureido derivative (**12h**). Interestingly, ΔT<sub>m</sub> value of *N*-acetyl-1,2,3,4-tetrahydroquinoline (**12j**) to the TA base pair was +5 °C. The results of TFOs **12h–j** strongly suggest that the carbonyl oxygen of the common 2-(1-*m*-carbonylaminophenyl-1,2,3-triazol-4-yl)ethyl unit forms hydrogen bond-mediated recognition with A (probably with 6-amino group in A) of the TA base pair, as shown in Figure 4. In fact, under same conditions, the T<sub>m</sub> value of triplex of TFO **12h** and dsDNA (YZ = TA) was comparable to that (T<sub>m</sub> = 29 °C) of triplex containing a T-CG base triplet forming a single hydrogen bond though this was lower than that (T<sub>m</sub> = 32 °C) of triplex containing a G-TA base triplet.<sup>6</sup> The above results do not imply that TFOs **12h–j** formed a stable and selective triplex to only dsDNA (YZ = TA) among all four dsDNA targets. However, this result would provide



**Figure 4.** Plausible recognition mode of a TA base pair by 2-[1-(*m*-ureidophenyl)-1,2,3-triazol-4-yl]ethyl nucleobase in TFO **12h**. The moiety displayed in black is 2-(1-*m*-carbonylamino-phenyl-1,2,3-triazol-4-yl)ethyl unit.

useful information on the design of nucleobase for TA base pair recognition in the formation of triplex DNA.

### 3. Conclusion

Oligonucleotides bearing two type of spacers attached to acetylene unit were synthesized and by the copper-catalyzed alkyne-azide 1,3-dipolar cycloaddition of them with various azides the preparation of oligonucleotides with the corresponding substituted triazoles was achieved. The evaluation of their triplex-forming ability with dsDNA demonstrated that 2-(1-*m*-carbonylamino-phenyl-1,2,3-triazol-4-yl)ethyl unit could make a hydrogen bond to a TA base pair though the stability and sequence-selectivity of the triplex formed was not satisfactory as assessed by UV-melting experiments. Our results obtained could contribute considerably in finding of nucleobases to recognize a TA base pair at a practical level.

### 4. Experimental

#### 4.1. General

All chemicals were purchased from chemical suppliers. For column chromatography, Fuji Silysia silica gel PSQ-100B and FL-100D were used. All melting points were measured on a Yanagimoto micro melting point apparatus and are uncorrected.  $^1\text{H}$  NMR,  $^{13}\text{C}$  NMR and  $^{31}\text{P}$  NMR spectra were recorded on a JEOL ECS400 or JEOL AL300 spectrometer. IR spectra were recorded on JASCO FT/IR-200 and JASCO FT/IR-4200 spectrometers. Optical rotations were recorded on a JASCO DIP-370 instrument. Mass spectra were measured on a JEOL JMS-600 or JEOL JMS-700 mass spectrometer. MALDI-TOF mass spectra were recorded on a Bruker Daltonics Autoflex II TOF/TOF or JEOL JMS-S3000 mass spectrometer. EYELA Cute Mixer CM-1000 was used as a shaker.

#### 4.2. Prop-2-ynyl 2-deoxy-3,5-di-*O*-toluoyl- $\beta$ -D-ribofuranoside **3**

Under a nitrogen atmosphere, propargyl alcohol (360  $\mu\text{L}$ , 9.25 mmol) was added to a solution of **2** (3.0 g, 7.71 mmol) in anhydrous  $\text{CH}_3\text{CN}$  (75 mL) at  $0^\circ\text{C}$  and the mixture was stirred for 6 h at room temperature. After addition of water, the mixture was extracted with AcOEt. The organic extracts were washed with saturated  $\text{NaHCO}_3$  aq, water and brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt = 5:1) to give compound **3** (1.2 g, 39%) as a colorless oil. Compound **3**:  $[\alpha]_{\text{D}}^{26} +138.0$  (c 1.00,  $\text{CHCl}_3$ ). IR  $\nu_{\text{max}}$  (KBr) 1718, 1611, 1273, 1178, 1109, 1020  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.29 (1H, dd,  $J = 2.0, 14.5$  Hz), 2.40–2.41 (7H, m), 2.54–2.61 (1H, m), 4.28 (1H, dd,  $J = 2.5, 15.5$  Hz), 4.50–4.56 (2H, m), 4.62–4.66 (1H, m), 5.42–5.46 (1H, m), 5.51 (1H, d,  $J = 5.5$  Hz), 7.21–7.24 (4H, m),

7.90–7.95 (4H, m).  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  21.6, 21.7, 39.2, 54.2, 64.1, 74.2, 74.4, 81.5, 102.2, 122.4, 126.9, 127.0, 129.1, 129.1, 129.6, 129.8, 143.8, 143.9, 166.2, 166.4. MS (EI)  $m/z$  408 ( $\text{M}^+$ , 100); HRMS (EI)  $m/z$  Calcd for  $\text{C}_{24}\text{H}_{24}\text{O}_6$ : 408.1573. Found 408.1580.

#### 4.3. Prop-2-ynyl 2-deoxy- $\beta$ -D-ribofuranoside **4**

Under a nitrogen atmosphere, NaOMe (240 mg, 4.41 mmol) was added to a solution of **3** (600 mg, 1.47 mmol) in anhydrous MeOH (5 mL) at room temperature and the mixture was stirred for 5 h. The solvent was removed under reduced pressure and the residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt = 1:7) to give compound **4** (251 mg, quant) as a colorless oil.  $[\alpha]_{\text{D}}^{26} +216.4$  (c 1.00,  $\text{CD}_3\text{OD}$ ); IR  $\nu_{\text{max}}$  (KBr) 3284, 2928, 1086, 1034  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  1.78 (1H, ddd,  $J = 1.5, 3.5, 14.0$  Hz), 2.26 (1H, ddd,  $J = 5.0, 8.0, 14.0$  Hz), 2.70 (1H, t,  $J = 2.5$  Hz), 3.48 (1H, dd,  $J = 5.0, 12.0$  Hz), 3.57 (1H, dd,  $J = 3.5, 12.0$  Hz), 3.80–3.84 (1H, m), 4.02–4.06 (1H, m), 5.25 (1H, dd,  $J = 1.5, 5.5$  Hz);  $^{13}\text{C}$  NMR (101 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  42.2, 54.7, 63.0, 72.2, 75.4, 80.5, 87.0, 103.1; MS (FAB)  $m/z$  174 [ $\text{M}+\text{H}$ ] $^+$ ; HRMS (FAB)  $m/z$  Calcd for  $\text{C}_8\text{H}_{12}\text{O}_4$  [ $\text{M}+\text{H}$ ] $^+$ : 173.0814. Found 173.0809.

#### 4.4. Prop-2-ynyl 2-deoxy-5-*O*-(4,4'-dimethoxytrityl)- $\beta$ -D-ribofuranoside **5**

Under a nitrogen atmosphere, 4,4'-DMTrCl (283, 0.836 mmol) was added to a solution of **4** (120 mg, 0.697 mmol) in anhydrous pyridine (5 mL) at room temperature and the mixture was stirred for 6 h. After addition of water, the mixture was extracted with AcOEt. The organic extracts were washed with water and brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt = 2:1) to give compound **5** (324 mg, 98%) as a colorless oil.  $[\alpha]_{\text{D}}^{23} +86.5$  (c 1.03,  $\text{CDCl}_3$ ); IR  $\nu_{\text{max}}$  (KBr) 1607, 1509, 1444, 1301, 1251, 1177, 1084, 1034  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  2.07 (1H, d,  $J = 13.5$  Hz), 2.25–2.31 (1H, m), 2.43 (1H, t,  $J = 2.5$  Hz), 3.16 (2H, d,  $J = 4.5$  Hz), 3.77 (6H, s), 4.19–4.24 (2H, m), 4.28 (1H, dd,  $J = 1.0, 2.5$  Hz), 5.46 (1H, d,  $J = 5.0$  Hz), 6.80–6.83 (4H, m), 7.17–7.32 (7H, m), 7.40–7.42 (2H, m).  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  14.1, 41.1, 53.9, 55.1, 63.9, 73.2, 74.3, 79.3, 86.0, 86.9, 102.7, 113.0, 126.7, 127.7, 128.0, 130.0, 135.8, 135.9, 144.7, 158.4; MS (FAB)  $m/z$  697 [ $\text{M}+\text{Na}$ ] $^+$ ; HRMS (FAB)  $m/z$  Calcd for  $\text{C}_{29}\text{H}_{30}\text{NaO}_6$  [ $\text{M}+\text{Na}$ ] $^+$ : 497.1935. Found 497.1944.

#### 4.5. Prop-2-ynyl 3-*O*-[2-cyanoethoxy(diisopropylamino)-phosphino]-2-deoxy-5-*O*-(4,4'-dimethoxytrityl)- $\beta$ -D-ribofuranoside **6**

Under a nitrogen atmosphere, 2-cyanoethyl *N,N*-diisopropylchlorophosphoramidite (40  $\mu\text{L}$ , 0.177 mmol) was added to a solution of compound **5** (70 mg, 0.148 mmol) and *N,N*-diisopropylethylamine (75  $\mu\text{L}$ , 0.443 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (2 mL) at room temperature and the mixture was stirred for 15 h. After addition of water, the solvent was removed under reduced pressure and the residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt = 4:1) to give compound **6** (90 mg, 90%) as a colorless oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  1.00 (2H, d,  $J = 3.0$  Hz), 1.11–1.16 (9H, m), 1.86–1.98 (3H, m), 2.52–2.62 (1H, m), 3.08–3.14 (1H, m), 3.28–3.41 (1H, m), 3.51–3.60 (3H, m), 3.78 (6H, s), 4.12–4.34 (4H, m), 5.43 (1H, m), 6.80–6.84 (4H, m), 7.19–7.35 (7H, m), 7.42–7.46 (2H, m);  $^{31}\text{P}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  147.3, 148.2; MS (FAB)  $m/z$  697 [ $\text{M}+\text{Na}$ ] $^+$ ; HRMS (FAB)  $m/z$  Calcd for  $\text{C}_{38}\text{H}_{47}\text{N}_2\text{NaO}_7\text{P}$  [ $\text{M}+\text{Na}$ ] $^+$ : 697.3019. Found 697.3049.

#### 4.6. 1-(1-Butyn-4-yl)-1,2-dideoxy-5-O-(4,4'-dimethoxytrityl)- $\beta$ -D-ribofuranose **9**

Under a nitrogen atmosphere, LiAlH<sub>4</sub> (360 mg, 9.47 mmol) was added to a solution of **8**<sup>5</sup> (1.2 g, 2.37 mmol) in anhydrous THF (20 mL) at room temperature and the mixture was stirred for 9 h. After addition of water, the mixture was extracted with AcOEt. The organic extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt = 5:1) to give compound **5** (1.1 g) as a colorless oil. This compound was not subjected to further purification and a portion of this was used in the next step. Under a nitrogen atmosphere, *p*-TsCl (246 mg, 1.29 mmol) was added to a solution of alcohol (500 mg, 1.08 mmol) in anhydrous pyridine (20 mL) at room temperature and the mixture was stirred for 12 h. After addition of water, the mixture was extracted with AcOEt. The organic extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. Flash silica gel column chromatography (*n*-hexane/AcOEt = 3:2) of the residue was performed to obtain appropriate compound [539 mg, R<sub>f</sub> = 0.3 (*n*-hexane/AcOEt = 1:1)] as a colorless oil, 250 mg of this compound was dissolved in anhydrous THF (3 mL). Under a nitrogen atmosphere, lithium acetylide ethylenediamine complex (82 mg, 0.889 mmol) was added to the solution at room temperature and the mixture was stirred for 9 h. After addition of saturated NH<sub>4</sub>Cl aq, the mixture was extracted with AcOEt. The organic extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt = 3:2) to give compound **9** (124 mg, 53% for three-steps) as a colorless oil. [ $\alpha$ ]<sub>D</sub><sup>24</sup> +3.0 (*c* 0.31, CHCl<sub>3</sub>); IR  $\nu_{\max}$  (KBr) 2933, 1607, 1509, 1445, 1301, 1251, 1177, 1074, 1034 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  1.73–1.80 (2H, m), 1.92–1.99 (3H, m), 2.27–2.36 (2H, m), 3.06 (1H, dd, *J* = 6.0, 10.0 Hz), 3.20 (1H, dd, *J* = 5.0, 10.0 Hz), 3.77 (6H, s), 3.86–3.90 (1H, m), 4.26–4.33 (2H, m), 6.80–6.83 (4H, m), 7.17–7.44 (9H, m); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  15.3, 34.5, 40.4, 55.1, 64.4, 68.5, 74.6, 83.9, 85.6, 86.1, 113.1, 126.7, 127.6, 128.1, 130.0, 136.0, 144.8, 158.4; HRMS (MALDI-TOF) *m/z* Calcd for C<sub>30</sub>H<sub>32</sub>NaO<sub>5</sub> [M+Na]<sup>+</sup>: 495.2142. Found 495.2141.

#### 4.7. 1-(1-Butyn-4-yl)-3-O-[2-cyanoethoxy(diisopropylamino)phosphino]-1,2-dideoxy-5-O-(4,4'-dimethoxytrityl)- $\beta$ -D-ribofuranose **10**

Under a nitrogen atmosphere, 2-cyanoethyl *N,N*-diisopropylchlorophosphoramidite (51  $\mu$ L, 0.223 mmol) was added to a solution of compound **9** (90 mg, 0.190 mmol) and *N,N*-diisopropylethylamine (97  $\mu$ L, 0.571 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (2 mL) at room temperature and the mixture was stirred for 5 h. After addition of water, the solvent was removed under reduced pressure and the residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt = 1:1) to give compound **10** (124 mg, 97%) as a colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  1.17 (3H, d, *J* = 7.0 Hz), 1.13–1.18 (9H, m), 1.74–1.89 (3H, m), 1.97–2.13 (2H, m), 2.30–2.46 (3H, m), 2.44 (1H, m), 3.07–3.18 (2H, m), 3.52–3.85 (10H, m), 4.05–4.11 (1H, m), 4.26–4.32 (1H, m), 4.40–4.47 (1H, m), 6.80–6.84 (4H, m), 7.16–7.46 (9H, m); <sup>31</sup>P NMR (162 MHz, CDCl<sub>3</sub>)  $\delta$  147.5, 147.7; MS (FAB) *m/z* 673 [M+H]<sup>+</sup>; HRMS (FAB) *m/z* Calcd for C<sub>39</sub>H<sub>50</sub>N<sub>2</sub>O<sub>6</sub>P [M+H]<sup>+</sup>: 673.3407. Found 673.3434.

#### 4.8. Oligonucleotides **1** and **7**

The synthesis of **1** and **7** was performed on a 0.2- $\mu$ mol scale or 1.0- $\mu$ mol scale on an automated DNA synthesizer (Gene Design

nS-8) using the common phosphoramidite protocol. TFOs synthesized on DMTr-ON mode were cleaved from the CPG resin and all the protecting groups on TFOs were removed by treatment with 28% NH<sub>3</sub> aq at room temperature for 3 h. The obtained crude TFOs were purified on Sep-Pak<sup>®</sup> Plus C18 cartridges (Waters) followed by reversed-phase HPLC (Waters XBridge<sup>®</sup> OST C18 2.5  $\mu$ m, 10 mm  $\times$  50 mm). The composition of the TFOs was confirmed by MALDI-TOF-MS analysis. MALDI-TOF-MS data ([M-H]<sup>-</sup>) for **1** and **7**: **1**, found 4423.01 (calcd 4423.94), **7**, found 4424.31 (calcd 4423.96).

#### 4.9. Azide synthesis

Among azide reagents used for click chemistry, 2-azidobenzamide, 3-azidobenzamide, 1,1'-(5-azido-1,3-phenylene)diurea and *N*-acetyl-7-azido-1,2,3,4-tetrahydroquinoline were new compounds which were prepared according to the following procedure.

##### 4.9.1. 2-Azidobenzamide

Under a nitrogen atmosphere, SOCl<sub>2</sub> (1 mL) was added to 2-azidobenzoic acid (200 mg, 1.23 mmol) and the mixture was refluxed for 2 h. The organic layer was evaporated. 10% aqueous NH<sub>3</sub> (3 mL) was added to the residue at 0 °C and the mixture was stirred for 0.5 h. The mixture was extracted with CHCl<sub>3</sub>. The organic extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (CHCl<sub>3</sub>/MeOH = 10:1) to give desired compound (185 mg, 93%) as yellow solids. Mp 130–131 °C. IR (KBr) 3368, 3168, 2130, 2103, 1655, 1620, 1599, 1574, 1483, 1452, 1403, 1230, 1163, 1127, 1084 cm<sup>-1</sup>. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  7.22 (1H, dt, *J* = 1.5, 7.5 Hz), 7.33 (1H, dd, *J* = 1.5, 7.5 Hz), 7.50 (1H, dt, *J* = 1.5 and 7.5 Hz), 7.56 (1H, br s), 7.57 (1H, dd, *J* = 1.5 and 7.5 Hz), 7.73 (1H, br s). <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  119.7, 124.8, 128.3, 129.7, 131.4, 136.6, 167.1. MS (EI) *m/z* 162 (M<sup>+</sup>, 100); HRMS (EI) *m/z* Calcd for C<sub>7</sub>H<sub>6</sub>N<sub>4</sub>O: 162.0542. Found 162.0550.

##### 4.9.2. 3-Azidobenzamide

Under a nitrogen atmosphere, SOCl<sub>2</sub> (2 mL) was added to 3-azidobenzoic acid (120 mg, 0.88 mmol) and the mixture was refluxed for 2 h. The organic layer was evaporated. 10% aqueous NH<sub>3</sub> (3 mL) was added to the residue at 0 °C and the mixture was stirred for 0.5 h. The mixture was extracted with CHCl<sub>3</sub>. The organic extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (CHCl<sub>3</sub>/MeOH = 9:1) to give desired compound (100 mg, 83%) as brown solids. Mp 135–136 °C. IR  $\nu_{\max}$  (KBr) 3358, 3171, 2198, 2113, 1658, 1483, 1444, 1395, 1314, 1288, 1164, 1129 cm<sup>-1</sup>. <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  7.25 (1H, d, *J* = 7.5 Hz), 7.46 (1H, br s), 7.48 (1H, t, *J* = 7.5 Hz), 7.58 (1H, s), 7.68 (1H, d, *J* = 7.5 Hz), 8.06 (1H, br s). <sup>13</sup>C NMR (101 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  118.0, 121.9, 124.2, 130.0, 136.0, 139.6, 167.1. MS (EI) *m/z* 162 (M<sup>+</sup>, 100); HRMS (EI) *m/z* Calcd for C<sub>7</sub>H<sub>6</sub>N<sub>4</sub>O: 162.0542. Found 162.0546.

##### 4.9.3. 1,1'-(5-Azido-1,3-phenylene)diurea

Under a nitrogen atmosphere, SnCl<sub>2</sub> (3.2 g, 1.70 mmol) was added to a solution of 5-iodo-1,3-dinitrobenzene (500 mg, 1.70 mmol) in anhydrous EtOH (10 mL) at room temperature and the mixture was stirred for 3 h at 70 °C. Ice (10 g) was added and the pH of the solution was controlled to approximately 9 by 10% NaOH aq. After the solids were removed by filtration through Celite<sup>®</sup>, the filtrate was extracted with AcOEt. The organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. Under a nitrogen atmosphere, 0.4 N HCl aq



(10 mL) and KNCO (340 mg, 4.08 mmol) were added to the residue and the mixture was stirred for 16 h at room temperature. The solids obtained by filtration was purified by flash silica gel column chromatography (AcOEt/MeOH = 30:1) to give 1,1'-(5-iodo-1,3-phenylene)diurea (105 mg, 19% from 5-iodo-1,3-dinitrobenzene) as light brown solids. Mp 135–138 °C. IR  $\nu_{\max}$  (KBr) 3292, 3195, 1672, 1585, 1544, 1442, 1347  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (400 MHz, DMSO- $d_6$ )  $\delta$  5.92 (4H, br s), 7.78 (1H, t,  $J = 1.8$  Hz), 7.48 (2H, d,  $J = 1.8$  Hz), 8.81 (2H, br s).  $^{13}\text{C}$  NMR (101 MHz, DMSO- $d_6$ )  $\delta$  94.6, 105.6, 118.6, 142.2, 155.8. HRMS (MALDI)  $m/z$  Calcd for  $\text{C}_8\text{H}_9\text{IN}_4\text{NaO}_2$ : 342.9662. Found 342.9652.

According to the reaction conditions reported previously,<sup>7</sup>  $\text{NaN}_3$  (26 mg, 0.41 mmol), CuI (7.7 mg, 41  $\mu\text{mol}$ ), sodium ascorbate (25 mg, 0.12 mmol) and  $N,N'$ -dimethylethylenediamine (4.4  $\mu\text{L}$ , 0.81 mmol) were added to a solution of 1,1'-(5-iodo-1,3-phenylene)diurea (95 mg, 0.30 mmol) in DMSO- $\text{H}_2\text{O}$  (5:1, 10 mL) and the mixture was stirred for 19 h at room temperature. After addition of brine, the mixture was extracted with AcOEt. The organic extracts were washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (AcOEt/MeOH = 15:1) to give the desired compound (50 mg, 72%) as white powder. Mp >300 °C. IR  $\nu_{\max}$  (KBr) 3258, 2115, 1668, 1594, 1556  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  6.92 (2H, d,  $J = 1$  Hz), 7.10 (1H, t-like,  $J = 1$  Hz).  $^{13}\text{C}$  NMR (76 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  104.3, 106.6, 142.3, 142.8, 159.1. HRMS (MALDI)  $m/z$  Calcd for  $\text{C}_8\text{H}_9\text{N}_7\text{NaO}_2$ : 258.0710. Found 258.0707.

#### 4.9.4. *N*-Acetyl-7-azido-1,2,3,4-tetrahydroquinoline

Under a nitrogen atmosphere, *t*-BuONO (270  $\mu\text{L}$ , 2.28 mmol) and  $\text{TMSN}_3$  (240  $\mu\text{L}$ , 1.51 mmol) was added to a solution of *N*-acetyl-7-amino-1,2,3,4-tetrahydroquinoline<sup>8</sup> (270 mg, 1.42 mmol) at 0 °C. The mixture was stirred for 9 h at room temperature. After addition of water, the mixture was extracted with AcOEt. The organic extracts were washed with saturated  $\text{NaHCO}_3$  aq, water and brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt = 1:1) to give the compound (280 mg, 91%) as light yellow solids. Mp 53–55 °C. IR  $\nu_{\max}$  (KBr) 2944, 2109, 2050, 1656, 1606, 1576, 1498, 1454, 1406, 1352, 1300, 1263, 1231, 1209, 1136, 1019  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  1.96 (2H, quint,  $J = 7.0$  Hz), 2.25 (3H, s), 2.72 (2H, t,  $J = 7.0$  Hz), 3.76 (2H, t,  $J = 7.0$  Hz), 6.79 (1H, d,  $J = 7.0$  Hz), 6.94 (1H, br s), 7.12 (1H, d,  $J = 7.0$  Hz).  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  23.2, 23.6, 26.3, 43.2 (br s), 115.1, 115.2, 129.4, 137.5, 139.9 (br s), 169.7. MS (FAB)  $m/z$  217  $[\text{M}+\text{H}]^+$ ; HRMS (FAB)  $m/z$  Calcd for  $\text{C}_{11}\text{H}_{13}\text{N}_4\text{O}$   $[\text{M}+\text{H}]^+$ : 217.1084. Found 217.1087.

#### 4.10. Click chemistry: general procedure

A solution of azide compound (10 mM in DMSO, 3  $\mu\text{L}$ ) was added to a mixture of  $\text{CuSO}_4$  (2 mM in  $\text{H}_2\text{O}$ , 3  $\mu\text{L}$ ), TBTA (2 mM in DMSO, 6  $\mu\text{L}$ ), sodium ascorbate (10 mM in  $\text{H}_2\text{O}$ , 3  $\mu\text{L}$ ), **1** or **7** [0.9 mM in phosphate buffer (pH 7.0), 3.3  $\mu\text{L}$ ] and  $\text{H}_2\text{O}$  (8.7  $\mu\text{L}$ ) in a 1.5 mL Eppendorf tube. The mixture was shaken at room temperature using a shaker (1000 rpm) until the reaction was complete. The entire product was purified by reversed-phase HPLC [column: Waters XBridge<sup>®</sup> OST C18 2.5  $\mu\text{m}$ , 4.6 mm  $\times$  50 mm; eluent: gradient system of MeCN/0.1 M triethylammonium acetate buffer (pH 7.0); flow rate: 1.0 mL/min] to give the desired TFO **11** or **12**, the identification was confirmed by MALDI-TOF-MS analysis. Isolated yield and MALDI-TOF-MS data ( $[\text{M}-\text{H}]^-$ ) for TFOs **11a–h** and **12a–j**: **11a**, 67%. Found 4545.85 (calcd 4545.06); **11b**, 74%. Found 4560.24 (calcd 4561.06); **11c**, 91%. Found 4588.03 (calcd 4588.09); **11d**, 85%. Found 4561.66 (calcd 4561.06); **11e**, 86%. Found 4575.27 (calcd 4575.09); **11f**, 74%. Found 4574.77 (calcd

4575.09); **11g**, 69%. Found 4587.89 (calcd 4588.09); **11h**, 79%. Found 4603.52 (calcd 4603.10); **12a**, 80%. Found 4543.77 (calcd 4543.09); **12b**, 83%. Found 4559.18 (calcd 4559.09); **12c**, 82%. Found 4586.26 (calcd 4586.11); **12d**, 83%. Found 4559.51 (calcd 4559.09); **12e**, 74%. Found 4573.72 (calcd 4573.11); **12f**, 82%. Found 4573.90 (calcd 4573.11); **12g**, 71%. Found 4586.99 (calcd 4586.11); **12h**, 77%. Found 4602.02 (calcd 4601.13); **12i**, 71%. Found 4659.83 (calcd 4659.17); **12j**, 90%. Found 4640.51 (calcd 4640.20).

#### 4.11. UV melting experiments

UV melting experiments were performed on SHIMADZU UV-1650 and SHIMADZU UV-1800 spectrophotometers equipped with  $T_m$  analysis accessory. TFOs **11** and **12** and hairpin dsDNA targets were dissolved in 10 mM sodium cacodylate buffer (pH 6.8) containing 100 mM KCl and 50 mM  $\text{MgCl}_2$  to give a final concentration of each strand of 1.89  $\mu\text{M}$ . The samples were annealed in boiling water followed by slow cooling to 5 °C. The melting profiles were recorded at 260 nm from 5 °C to 90 °C at a scan rate of 0.5 °C/min. A two-point average method was used to obtain the  $T_m$  values and the final values were determined by averaging three independent measurements which were accurate to within 1 °C.

#### Acknowledgments


This work was supported in part by the Program to Disseminate Tenure Tracking System of the Ministry of Education, Culture, Sports, Science and Technology, Japan (MEXT), a Grant-in-Aid for Scientific Research on Innovative Areas (No. 22136006) from MEXT, a Grant-in-Aid for JSPS Fellows of Japan Society for the Promotion of Science (JSPS) and Mochida Memorial Foundation for Medical and Pharmaceutical Research.

#### Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.bmc.2013.05.034>.

#### References and notes

- Reviews: (a) Praseuth, D.; Guieysse, A. L.; Hélène, C. *Biochim. Biophys. Acta* **1999**, *1489*, 181; (b) Knauer, M. P.; Glazer, P. M. *Hum. Mol. Genet.* **2001**, *10*, 2243; (c) Seidman, M. M.; Glazer, P. M. *J. Clin. Invest.* **2003**, *112*, 487; (d) Kuan, J. Y.; Glazer, P. M. *Methods Mol. Biol.* **2004**, *262*, 173; (e) Mahato, R. I.; Chen, K.; Guntaka, R. V. *Expert Opin. Drug Deliv.* **2005**, *2*, 3; (f) Sasaki, S.; Nagatsugi, F. *Curr. Opin. Chem. Biol.* **2006**, *10*, 615; (g) Chin, J. Y.; Schleifman, E. B.; Glazer, P. M. *Front. Biosci.* **2007**, *12*, 4288; (h) Simon, P.; Cannata, F.; Concordet, J.-P.; Giovannangeli, C. *Biochimica* **2008**, *90*, 1109; (i) Duca, M.; Vekhoff, P.; Oussedik, K.; Halby, L.; Arimondo, P. B. *Nucleic Acids Res.* **2008**, *36*, 5123; (j) Chin, J. Y.; Glazer, P. M. *Mol. Carcinog.* **2009**, *48*, 389; (k) Mukherjee, A.; Vasquez, K. M. *Biochimica* **2011**, *93*, 1197.
- Reviews: (a) Thuong, N. T.; Hélène, C. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 666; (b) Doronina, S. O.; Behr, J.-P. *Chem. Soc. Rev.* **1997**, *26*, 63; (c) Luyten, I.; Herdewijn, P. *Eur. J. Med. Chem.* **1998**, *33*, 515; (d) Gowers, D. M.; Fox, K. R. *Nucleic Acids Res.* **1999**, *27*, 1569; (e) Fox, K. R. *Curr. Med. Chem.* **2000**, *7*, 17; (f) Robles, J.; Grandas, A.; Pedrosa, E.; Luque, F. J.; Eritija, R.; Orzco, M. *Curr. Org. Chem.* **2002**, *6*, 1333; (g) Purwanto, M. G. M.; Weisz, K. *Curr. Org. Chem.* **2003**, *7*, 427; (h) Buchini, S.; Leumann, C. J. *Curr. Opin. Chem. Biol.* **2003**, *7*, 717; (i) Malnuit, V.; Duca, M.; Benhida, R. *Org. Biomol. Chem.* **2011**, *9*, 326; (j) Hari, Y.; Obika, S.; Imanishi, T. *Eur. J. Org. Chem.* **2012**, *56*, 2875.
- (a) Nakahara, M.; Kuboyama, T.; Izawa, A.; Hari, Y.; Imanishi, T.; Obika, S. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 3316; (b) Hari, Y.; Nakahara, M.; Pang, J.; Akabane, M.; Kuboyama, T.; Obika, S. *Bioorg. Med. Chem.* **2011**, *19*, 1162.
- Hari, Y.; Akabane, M.; Hatanaka, Y.; Nakahara, M.; Obika, S. *Chem. Commun.* **2011**, 4424.
- Hovinen, J.; Salo, H. J. *Chem. Soc., Perkin Trans. 1* **1997**, 3013.
- It is also known that G forms a single hydrogen bond with TA base pair: (a) Yoon, K.; Hobbs, C. A.; Koch, J.; Sardaro, M.; Kutny, R.; Weis, A. L. *Proc. Natl. Acad. Sci. U.S.A.* **1992**, *89*, 3840; (b) Griffin, L. C.; Dervan, P. B. *Science* **1989**, *245*, 967.
- Andersen, J.; Madsen, U.; Björkling, F.; Liang, X. *Synlett* **2005**, 2209.
- Kolmakov, K.; Belov, V. N.; Wurm, C. A.; Harke, B.; Leutenegger, M.; Eggeling, C.; Hell, S. W. *Eur. J. Org. Chem.* **2010**, 3593.



# Base-pair recognition ability of hydroxyphenyl nucleobases in parallel triplex DNA



Yoshiyuki Hari\*, Satoshi Kashima, Hiroyasu Inohara, Shin Ijitsu, Takeshi Imanishi, Satoshi Obika\*

Graduate School of Pharmaceutical Sciences, Osaka University, 1-6 Yamadaoka, Suita, Osaka 565-0871, Japan

## ARTICLE INFO

### Article history:

Received 30 April 2013

Received in revised form 21 May 2013

Accepted 23 May 2013

Available online 30 May 2013

### Keywords:

Artificial nucleobases

2',4'-BNA

Parallel motif triplex

Triplex DNA

Triplex-forming oligonucleotides

## ABSTRACT

Oligonucleotides containing 2'-deoxyribonucleotides bearing hydroxyphenyl nucleobases or their 2',4'-BNA-modified analogs were synthesized, and the triplex-forming ability of their oligonucleotides with double-stranded DNA targets was evaluated by UV melting experiments. Results showed that 2'-deoxyribonucleotide bearing 2'-hydroxyphenyl nucleobase could be recognized by a dUA base pair while no affinity to a TA base pair was observed. The 4'-BNA modification led to a further increase in the binding affinity to a dUA base pair. The 4'-BNA bearing 3-hydroxyphenyl nucleobase showed moderate binding affinity to a TA base pair, but without selectivity.

© 2013 Elsevier Ltd. All rights reserved.

## 1. Introduction

Triplex DNA is formed by interaction between double-stranded DNA (dsDNA) and an oligonucleotide, called a triplex-forming oligonucleotide (TFO). Triplex formation is useful for targeting dsDNA and can contribute to the development of various nucleic acid technologies.<sup>1</sup> However, there is the sequence limitation of target dsDNA as a problem of triplex formation. Although AT or GC base pairs within dsDNA are recognized by T or C in TFO in a parallel-oriented manner to form T-AT or protonated C-GC (C<sup>+</sup>H-GC) base triplets, respectively (Fig. 1), it is difficult to target dsDNA containing pyrimidine–purine base pairs (TA or CG base pairs).




Fig. 1. Structures of T-AT and C<sup>+</sup>H-GC base triplets.

\* Corresponding authors. Tel./fax: +81 6 6879 8201 (Y.H.); tel.: +81 6 6879 8200; fax: +81 6 6879 8204 (S.O.); e-mail addresses: [hari@phs.osaka-u.ac.jp](mailto:hari@phs.osaka-u.ac.jp) (Y. Hari), [obika@phs.osaka-u.ac.jp](mailto:obika@phs.osaka-u.ac.jp) (S. Obika).

Therefore, many artificial nucleic acids have been developed for recognizing TA or CG base pairs in triplex formation.<sup>2</sup> However, most of them target CG base-pair recognition and there are not many approaches for a TA base pair.

On the basis of the structure of the T-CG base triplet,<sup>3</sup> a previous report described the design of a 2-pyridone nucleobase (P) for CG base-pair recognition and demonstrated that 2-pyridone bound to a CG base pair and that 2',4'-BNA with P (P<sup>B</sup>) effectively recognized a CG base pair (Fig. 2).<sup>4</sup> Hydrogen bond formation between the 2-carbonyl group in P and 4-amino group in C was considered as the recognition mode (Fig. 2). This finding suggests that a hydroxyphenyl nucleobase might be a promising candidate for a TA base pair and could hydrogen-bond with the 4-carbonyl group in T through the hydroxyl group, despite possible steric hindrance by the 5-methyl group in T (Fig. 2). Therefore, in this study, TFOs including 2'-deoxyribonucleotides with 2-, 3-, and 4-hydroxyphenyl nucleobases




Fig. 2. Design of nucleobase for a TA base-pair recognition.

that have hydroxyl groups located at different positions, and TFOs including 2',4'-BNA with 2- and 3-hydroxyphenyl nucleobases, were synthesized and the triplex-forming ability of their TFOs with dsDNA was evaluated using UV melting experiments.<sup>5</sup>


## 2. Results and discussion

### 2.1. Synthesis


The synthesis of phosphoramidites **1–3** composed of a 2'-deoxyribose unit and hydroxyphenyl units is shown in Scheme 1. Coupling of compound **4**<sup>6</sup> and 2-allyloxyphenyllithium followed by the Mitsunobu reaction<sup>7</sup> using *N,N,N',N'*-tetramethyl azodicarboxamide (TMAD) and *n*-Bu<sub>3</sub>P gave **5** ( $\beta$ -isomer/ $\alpha$ -isomer=ca. 1:1), which had its allyl group deprotected to afford the desired  $\beta$ -isomer **6** and its  $\alpha$ -isomer **7**. A benzoyl group was chosen as the protecting group of the phenolic hydroxyl group in oligonucleotide synthesis. Reaction of **6** with BzCl in the presence of Et<sub>3</sub>N gave **8** in 97% yield, which was converted into diol **9** by hydrogenolysis using Pd(OH)<sub>2</sub>-C and cyclohexene. Dimethoxytritylation of **9** followed by phosphitylation afforded 2-hydroxyphenyl phosphoramidite **1**, a suitable building block for oligonucleotide synthesis. For the 3-hydroxyphenyl derivative, the  $\beta$ -isomer **11** was isolated in 28% yield in two steps via reaction of **4** with 3-methoxymethoxyphenyllithium and ring-closure. The methoxymethyl group (**11**) was converted into a benzoyl group (**14**) through a deprotection–protection process. Then, the phosphoramidite **2** was synthesized via three steps (debenzylation–dimethoxytritylation–phosphitylation). The synthesis of 4-hydroxyphenyl phosphoramidite **3** was achieved by a synthetic route similar to that of 2-hydroxyphenyl derivative **1**. Reaction of **4** with 4-allyloxyphenyllithium and ring-closure reaction yielded **17** as a 1:1 anomeric mixture. Palladium-catalyzed deallylation of **17** proceeded to give **18**, which was benzoylated to afford a separable mixture of  $\beta$ -isomer **19** and  $\alpha$ -isomer **20**. Hydrogenolysis of the  $\beta$ -isomer **19** produced **21** in 82% yield, which underwent reaction

with DMTrCl in pyridine to produce **22** in 91% yield. The desired phosphoramidite **3** was obtained via phosphitylation.

The 2',4'-BNA-modified phosphoramidites **23–25** with 2- and 3-hydroxyphenyl nucleobase units were synthesized according to the synthetic route shown in Scheme 2. A previous study reported



**Scheme 2.** Reagents and conditions: (i) 2-allyloxyphenylmagnesium iodide, THF,  $-40^{\circ}\text{C}$ ; TMAD, *n*-Bu<sub>3</sub>P, CH<sub>2</sub>Cl<sub>2</sub>, rt, 57% over two steps; (ii) NaBH<sub>4</sub>, Pd(PPh<sub>3</sub>)<sub>4</sub>, THF, rt, 80% (**29**) and quant. (**35**); (iii) BzCl, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>, rt, 93% (**30**) and 98% (**36**); (iv) 20% Pd(OH)<sub>2</sub>-C, cyclohexene, EtOH, reflux, 89% (**31**) and 95% (**37**); (v) DMTrCl, pyridine, rt, 94% (**32**) and 91% (**38**); (vi) *i*-Pr<sub>2</sub>N<sub>2</sub>POCH<sub>2</sub>CH<sub>2</sub>CN, DIHT, MeCN/THF, rt, 92% (**23**), 81% (**24**), and 71% (**25**); (vii) K<sub>2</sub>CO<sub>3</sub>, MeOH, rt, 5 min, 96%; (viii) allyl bromide, K<sub>2</sub>CO<sub>3</sub>, acetone, rt, 14 h, 87%; (ix) 3-allyloxyphenylmagnesium iodide, THF, rt; TMAD, *n*-Bu<sub>3</sub>P, CH<sub>2</sub>Cl<sub>2</sub>, rt, 70% over two steps.



**Scheme 1.** Reagents and conditions: (i) 2-allyloxyphenyllithium, THF,  $-78^{\circ}\text{C}$ ; TMAD, *n*-Bu<sub>3</sub>P, CH<sub>2</sub>Cl<sub>2</sub>, rt, 38% ( $\beta$ -isomer/ $\alpha$ -isomer=ca. 1:1) over two steps; (ii) NaBH<sub>4</sub>, Pd(PPh<sub>3</sub>)<sub>4</sub>, THF, rt, 55% (**6**) and 45% (**7**), and 91% (**18**,  $\beta$ -isomer/ $\alpha$ -isomer=ca. 1:1); (iii) BzCl, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>, rt, 97% (**8**), 98% (**14**), and 91% (**19/20**=ca. 1:1); (iv) 20% Pd(OH)<sub>2</sub>-C, EtOH,  $70^{\circ}\text{C}$ , 94% (**9**), 96% (**15**), and 82% (**21**); (v) DMTrCl, pyridine, rt, 98% (**10**), 72% (**16**), and 91% (**22**); (vi) *i*-Pr<sub>2</sub>NP(Cl)OCH<sub>2</sub>CH<sub>2</sub>CN, *i*-Pr<sub>2</sub>NEt, CH<sub>2</sub>Cl<sub>2</sub>, rt, 58% (**1**), 58% (**2**), and 95% (**3**); (vii) 3-methoxymethoxyphenyllithium, THF,  $-78^{\circ}\text{C}$ ; TMAD, *n*-Bu<sub>3</sub>P, CH<sub>2</sub>Cl<sub>2</sub>, rt, 28% (**11**) and 24% (**12**) over two steps; (viii) 10% HCl aq, THF, rt, 2 days, 91%; (ix) 4-allyloxyphenyllithium, THF,  $-60^{\circ}\text{C}$ ; TMAD, *n*-Bu<sub>3</sub>P, CH<sub>2</sub>Cl<sub>2</sub>, rt; 48% ( $\beta$ -isomer/ $\alpha$ -isomer=ca. 1:1) over two steps.

that reaction of aldehyde **26** with arylmagnesium reagents proceeded with high diastereoselectivity and that Mitsunobu reaction of the resulting products led predominantly to 2',4'-BNA-modified C-nucleosides with  $\beta$ -configuration.<sup>8</sup> Thus, **26** was treated with excess 2- or 3-allyloxyphenylmagnesium bromide in THF to give coupling compounds; ring-closure using TMAD afforded the  $\beta$ -anomers **27** and **28** in 57% and 70% yield, respectively. Interestingly, in both cases, only the  $\beta$ -anomer was isolated; its  $\alpha$ -isomer was not observed. Palladium-catalyzed reaction of 2-hydroxyphenyl derivative **27** with NaBH<sub>4</sub> gave deallylated compound **29**, which was benzoylated by BzCl in the presence of Et<sub>3</sub>N to produce **30** in high yield. After debenzoylation of **30** to produce **31**, compound **32** was obtained in 94% yield by dimethoxytritylation. Phosphitylation of **32** using (*i*-Pr<sub>2</sub>N)<sub>2</sub>POCH<sub>2</sub>CH<sub>2</sub>CN and diisopropylammonium tetrazolidate (DIHT) gave benzoyl-protected phosphoramidite **23** with a 2-hydroxyphenyl unit and 2',4'-BNA modification in 92% yield. In addition, the allyl-protected congener **24** was prepared through debenzoylation of **32**, followed by allylation and phosphitylation. Analogous to synthesis of **23**, 2',4'-BNA phosphoramidite **25** with a 3-hydroxyphenyl unit was synthesized. After conversion of the allyl group of **28** into a benzoyl group (**36**) in two steps, **37** was produced by the hydrogenolysis. The desired phosphoramidite **25** was obtained from **37** via dimethoxytritylated **38**.

Next, introduction of the phosphoramidites synthesized into oligonucleotides was investigated using an automated DNA synthesizer. The use of benzoyl-protected phosphoramidites **1–3** and **25**, with the exception of the 2',4'-BNA-modified 2-hydroxy analog **23**, successfully led to the desired TFOs according to standard phosphoramidite chemistry. For **23**, two undesirable oligonucleotides having an approximately +17 mass unit difference from molecular weight of the desired TFO were obtained. This result implies that a structure constrained by the methylene linkage between the 2'-oxygen and 4'-carbon atoms might trigger a ring-opening reaction by conventional ammonia treatment as shown in Fig. 3. Eventually, the desired TFO **42** was obtained by oligonucleotide synthesis using allyl-protected phosphoramidite **24** followed by treatment with NaBH<sub>4</sub> and Pd(PPh<sub>3</sub>)<sub>4</sub> to remove an allyl group. The sequence of TFOs prepared is listed in Fig. 4.




Fig. 3. Plausible mechanism of ring-opening reaction.

## 2.2. Evaluation

Triplex-forming ability of the TFOs **39–43** with dsDNA was evaluated through UV melting experiments at 10 mM sodium cacodylate (pH 6.8) plus 140 mM KCl and 50 mM MgCl<sub>2</sub>. As targets, hairpin dsDNAs containing five types of base pairs (i.e., TA, its demethylated UA, CG, AT, and GC base pairs) at the same position were used. Results are summarized in Table 1. In general, TFOs **39–41** containing 2'-deoxy derivatives 2H, 3H, and 4H, did not sequence-specifically form stable triplexes with dsDNA (YZ=TA)

### Sequence of TFOs

**39** : 5'-TTTTTCT(2H)TCTCTCT-3'  
**40** : 5'-TTTTTCT(3H)TCTCTCT-3'  
**41** : 5'-TTTTTCT(4H)TCTCTCT-3'  
**42** : 5'-TTTTTCT(2H<sup>B</sup>)TCTCTCT-3'  
**43** : 5'-TTTTTCT(3H<sup>B</sup>)TCTCTCT-3'




Fig. 4. TFOs synthesized in this study.

Table 1

*T<sub>m</sub>* values (°C) of triplexes between TFOs and dsDNA targets<sup>a,b</sup>

TFO (X)	YZ				
	TA	UA	CG	AT	GC
<b>39</b> (2H)	23	28	27	27	31
<b>40</b> (3H)	26	27	27	28	25
<b>41</b> (4H)	24	25	25	30	27
<b>42</b> (2H <sup>B</sup> )	28	37	29	36	39
<b>43</b> (3H <sup>B</sup> )	34	35	34	33	33

<sup>a</sup> Conditions: 10 mM sodium cacodylate buffer (pH 6.8), 140 mM KCl, and 50 mM MgCl<sub>2</sub>. The final concentration of each oligonucleotide used was 1.89  $\mu$ M. C indicates 2'-deoxy-5-methylcytidine.

<sup>b</sup> Triplexes containing T-AT and P<sup>B</sup>-CG base triplets as X-YZ had *T<sub>m</sub>* values of 44 °C and 37 °C, respectively. P<sup>B</sup> indicates 2',4'-BNA bearing 2-pyridone nucleobase.

and no significant sequence-selectivity toward a certain base pair was observed because the hydroxyl group could function not only as a hydrogen donor but also as a hydrogen acceptor. However, the *T<sub>m</sub>* value of a triplex with dsDNA (YZ=UA) by TFO **39** was greater than that with dsDNA (YZ=TA) by +5 °C. In addition, TFO **39** provided nearly the same stability to TFO **41** (X=4H) in the triplex with dsDNA (YZ=TA), while TFO **39** formed a stable triplex with dsDNA (YZ=UA) compared to TFO **41**. These results suggest that a 2-hydroxyphenyl nucleobase forms a hydrogen bond with U of a UA base pair without forming any hydrogen bond with a TA base pair because the 5-methyl group in T prevents bond formation (Fig. 5). In contrast, TFO **40** (X=3H) significantly stabilized triplexes formed




Fig. 5. Proposed structures of base triplets by 2H and 3H.

with dsDNAs (YZ=TA and UA) compared to TFO **41**. The  $T_m$  values of triplexes (X-YZ=3H-TA and 3H-UA) were comparable to that of triplex (X-YZ=2H-UA). This suggests that the 3-hydroxyphenyl nucleobase recognized not only a UA base pair, but also a TA base pair by effective avoidance of the bulky 5-methyl group in T, despite the lack of stability. When TFOs **42** and **43** containing 2',4'-BNA with 2- or 3-hydroxyphenyl nucleobases were used, affinities to any dsDNA targets increased without any sequence-selectivity. TFO **42** (X=2H<sup>B</sup>) formed a stable triplex with dsDNA (YZ=UA) rather than with dsDNA (YZ=TA), which was comparable to triplexes formed with TFOs, including 2',4'-BNA (P<sup>B</sup>) with a 2-pyridone nucleobase and dsDNA (YZ=CG).<sup>9</sup> As mentioned, P<sup>B</sup> likely bonds to a CG base pair via a single hydrogen bond. Therefore, 2-hydroxyphenyl nucleobase in 2H<sup>B</sup> forms a single hydrogen bond with an UA base pair and cannot recognize a TA base pair because of the 5-methyl group. In contrast, TFO **43** (X=3H<sup>B</sup>) had  $T_m$  values of 34 °C and 35 °C against dsDNAs (YZ=UA and TA), respectively. This stability was significantly higher than that of the triplex (X-YZ=2H<sup>B</sup>-TA), although it was slightly lower than that of triplex (X-YZ=2H<sup>B</sup>-UA). These results suggest that 3-hydroxyphenyl nucleobase in 3H<sup>B</sup> should recognize both UA and TA base pairs through a single hydrogen bond (Fig. 5). Results also indicated that 3-hydroxyphenyl nucleobase can recognize a TA base pair in a parallel triplex formation, although no sequence-selectivity was observed.

### 3. Conclusion

Syntheses of 2'-deoxyribonucleotides bearing 2-, 3-, and 4-hydroxyphenyl nucleobases and 2',4'-BNA-modified nucleotides containing 2- and 3-hydroxyphenyl nucleobases were developed. Evaluation of triplex-forming ability of TFOs containing their hydroxyphenyl nucleobases with dsDNA showed 3-hydroxyphenyl nucleobase might recognize a TA base pair through a single hydrogen bond without any sequence selectivity. Recently, we succeeded in the facile synthesis of various triazole or pyrimidinone nucleobases on oligonucleotides using post-elongation modification (PEM) and found promising nucleobases for CG base-pair recognition.<sup>10</sup> Thus, the synthesis of various 3-hydroxyphenyl units using PEM could be effective for screening of nucleobases for TA base-pair recognition. Further investigation of nucleobases based on the 3-hydroxyphenyl structure is underway.

## 4. Experimental

### 4.1. General methods

All chemicals were purchased from chemical suppliers. For column chromatography, Fuji Silysia silica gel BW-300, PSQ-100B, and FL-100D were used. All melting points were measured on a Yanagimoto micro melting point apparatus and are uncorrected. <sup>1</sup>H NMR, <sup>13</sup>C NMR, and <sup>31</sup>P NMR spectra were recorded on JEOL EX270, JEOL ECS400, and JEOL GX-500 spectrometers. IR spectra were recorded on JASCO FT/IR-200 and JASCO FT/IR-4200 spectrometers. Optical rotations were recorded on a JASCO DIP-370 instrument. Mass spectra were obtained using a JEOL JMS-600 or JEOL JMS-700 mass spectrometer. MALDI-TOF mass spectra were recorded on a Bruker Daltonics Autoflex II TOF/TOF or JEOL JMS-S3000 mass spectrometer.

### 4.2. 1-Allyloxy-2-(2-deoxy-3,5-di-O-benzyl-D-ribofuranosyl)benzene (**5**)

Under a nitrogen atmosphere, a solution of 2-deoxy-3,5-di-O-benzyl-D-ribofuranose **4**<sup>6</sup> (1.47 g, 4.69 mmol) in anhydrous THF (6 mL) was added to a solution of 2-allyloxyphenyllithium

[prepared from 2-allyloxyphenyl bromide (3.98 g, 18.7 mmol) and *n*-BuLi (2.67 M in hexane, 6.9 mL, 18.4 mmol) in anhydrous THF (20 mL) at -78 °C], and the mixture was stirred at -78 °C for 16 h. After addition of satd NH<sub>4</sub>Cl aq, the mixture was extracted with Et<sub>2</sub>O. The extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=5:1) to give the appropriate compounds ( $R_f$ =ca. 0.1), which were dissolved in CH<sub>2</sub>Cl<sub>2</sub> (7 mL); TMAD (750 mg, 4.35 mmol) and *n*-Bu<sub>3</sub>P (1.1 mL, 4.35 mL) were added at room temperature. After being stirred at room temperature for 6 h, water was added and the mixture was extracted with Et<sub>2</sub>O. The extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=15:1) to give a ca. 1:1 anomeric mixture **5** (773 mg, 38% from **4**) as a colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.77 (0.5H, ddd,  $J=7.4, 10.2, 13.2$  Hz), 1.96 (0.5H, ddd,  $J=5.8, 7.8, 12.7$  Hz), 2.56 (0.5H, ddd,  $J=1.0, 5.4, 13.2$  Hz), 2.73 (0.5H, td,  $J=6.8, 12.7$  Hz), 3.56–3.71 (2H, m), 4.12–4.13 (0.5H, m), 4.20–4.24 (0.5H, m), 4.27–4.31 (0.5H, m), 4.38–4.63 (6.5H, m), 5.22–5.25 (1H, m), 5.34–5.51 (1H, m), 5.96–6.06 (1H, m), 6.78–6.81 (1H, m), 6.91–6.98 (1H, m), 7.16–7.34 (11H, m), 7.52 (0.5H, d,  $J=7.4$  Hz), 7.59 (0.5H, d,  $J=6.8$  Hz). HRMS (FAB)  $m/z$  calcd for C<sub>28</sub>H<sub>30</sub>NaO<sub>4</sub> [M+Na]<sup>+</sup>: 453.2036; found, 453.2049.

### 4.3. 2-(2-Deoxy-3,5-di-O-benzyl- $\beta$ -D-ribofuranosyl)-1-hydroxybenzene (**6**) and 2-(2-deoxy-3,5-di-O-benzyl- $\alpha$ -D-ribofuranosyl)-1-hydroxybenzene (**7**)

Under a nitrogen atmosphere, Pd(PPh<sub>3</sub>)<sub>4</sub> (290 mg, 0.253 mmol) was added to a solution of compound **5** (545 mg, 1.27 mmol) in anhydrous THF (10 mL) and the mixture was stirred until solids of Pd(PPh<sub>3</sub>)<sub>4</sub> dissolved completely. Then, NaBH<sub>4</sub> (58 mg, 1.52 mmol) was added and the mixture was stirred at room temperature for 24 h. After addition of 3% HCl aq, the mixture was filtered through a pad of Celite®. The filtrate was extracted with AcOEt. The extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=20:1 to 10:1) to give compound **6** (270 mg, 55%) as a colorless oil and compound **7** (221 mg, 45%) as a colorless oil. Compound **6**: [ $\alpha$ ]<sub>D</sub><sup>25</sup> +54.0 (*c* 1.01, CHCl<sub>3</sub>). IR  $\nu_{\max}$  (KBr) 3338, 3028, 2867, 1495, 1455, 1247, 1095, 1077 cm<sup>-1</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.20–2.33 (2H, m), 3.55 (1H, dd,  $J=2.5, 10.0$  Hz), 3.67 (1H, dd,  $J=3.0, 10.0$  Hz), 4.18–4.20 (1H, m), 4.24–4.26 (1H, m), 4.47 (1H, d,  $J=12.0$  Hz), 4.51 (1H, d,  $J=12.0$  Hz), 4.54 (1H, d,  $J=12.0$  Hz), 4.68 (1H, d,  $J=12.0$  Hz), 5.20 (1H, dd,  $J=6.0, 10.5$  Hz), 6.80 (1H, ddd,  $J=2.0, 8.0, 8.0$  Hz), 6.89 (1H, dd,  $J=2.0, 8.0$  Hz), 7.01 (1H, d,  $J=2.0, 8.0$  Hz), 7.18 (1H, ddd,  $J=2.0, 8.0, 8.0$  Hz), 7.26–7.36 (10H, m), 8.10 (1H, s). <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  39.4, 69.5, 71.3, 73.5, 81.3, 82.6, 84.7, 117.4, 119.4, 123.8, 125.9, 127.7, 127.8, 127.9, 127.9, 128.1, 128.5, 129.3, 137.3, 137.8, 155.8. HRMS (FAB)  $m/z$  calcd for C<sub>25</sub>H<sub>26</sub>NaO<sub>4</sub> [M+Na]<sup>+</sup>: 413.1723; found 413.1742. Compound **7**: [ $\alpha$ ]<sub>D</sub><sup>25</sup> +6.6 (*c* 0.86, CHCl<sub>3</sub>). IR  $\nu_{\max}$  (KBr) 3337, 3033, 2925, 2862, 1494, 1455, 1246, 1105, 1074 cm<sup>-1</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.23 (1H, ddd,  $J=4.0, 8.0, 12.5$  Hz), 2.65 (1H, ddd,  $J=7.0, 7.0, 12.5$  Hz), 3.58–3.64 (2H, m), 4.27–4.32 (1H, m), 4.37 (1H, ddd,  $J=4.5, 4.5, 4.5$  Hz), 4.49 (1H, d,  $J=12.0$  Hz), 4.50 (1H, d,  $J=12.0$  Hz), 4.55 (1H, d,  $J=12.0$  Hz), 4.57 (1H, d,  $J=12.0$  Hz), 5.23 (1H, dd,  $J=7.0, 8.0$  Hz), 6.81 (1H, ddd,  $J=2.0, 8.0, 8.0$  Hz), 6.88 (1H, dd,  $J=2.0, 8.0$  Hz), 6.96 (1H, dd,  $J=2.0, 8.0$  Hz), 7.17 (1H, ddd,  $J=2.0, 8.0, 8.0$  Hz), 7.25–7.38 (10H, m), 8.09 (1H, s). <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  39.4, 70.2, 71.6, 73.5, 79.9, 81.2, 82.9, 117.3, 119.5, 124.6, 127.5, 127.6, 127.7, 127.7, 127.8, 128.4, 129.1, 137.5, 137.5, 137.9, 155.4. HRMS (FAB)  $m/z$  calcd for C<sub>25</sub>H<sub>27</sub>O<sub>4</sub> [M+H]<sup>+</sup>: 391.1904; found 391.1895.

#### 4.4. 1-Benzoyloxy-2-(2-deoxy-3,5-di-O-benzyl- $\beta$ -D-ribofuranosyl)benzene (8)

Under a nitrogen atmosphere, BzCl (88  $\mu$ L, 0.753 mmol) was added to a solution of compound **6** (270 mg, 0.627 mmol) and Et<sub>3</sub>N (105  $\mu$ L, 0.753 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (6 mL) at room temperature, and the mixture was stirred for 2 h. After addition of water, the mixture was extracted with Et<sub>2</sub>O. The extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=10:1) to give compound **8** (332 mg, 97%) as a colorless oil. [ $\alpha$ ]<sub>D</sub><sup>24</sup> +57.1 (*c* 0.90, CHCl<sub>3</sub>). IR  $\nu_{\max}$  (KBr) 3062, 3032, 2867, 1738, 1489, 1452, 1263, 1219, 1181, 1081 cm<sup>-1</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.86 (1H, ddd, *J*=6.0, 10.5, 13.0 Hz), 2.38 (1H, ddd, *J*=1.5, 5.0, 13.0 Hz), 3.56 (1H, dd, *J*=5.0, 10.0 Hz), 3.64 (1H, dd, *J*=5.0, 10.0 Hz), 4.09–4.11 (1H, m), 4.20–4.23 (1H, m), 4.40 (2H, s), 4.57 (2H, s), 5.35 (1H, dd, *J*=5.0, 10.5 Hz), 7.14–7.36 (13H, m), 7.48 (2H, dd, *J*=8.0, 8.0 Hz), 7.63 (1H, dd, *J*=8.0, 8.0 Hz), 7.65 (1H, dd, *J*=8.0, 8.0 Hz), 8.20 (2H, d, *J*=8.0 Hz). <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  39.8, 70.9, 73.4, 75.0, 81.2, 83.5, 122.1, 126.3, 126.7, 127.5, 127.5, 127.6, 127.6, 128.2, 128.3, 128.3, 128.128.7, 129.3, 130.1, 133.7, 134.2, 137.9, 138.1, 147.7, 165.0. HRMS (FAB) *m/z* calcd for C<sub>32</sub>H<sub>31</sub>O<sub>5</sub> [M+H]<sup>+</sup>: 495.2166; found 495.2165.

#### 4.5. 1-Benzoyloxy-2-(2-deoxy- $\beta$ -D-ribofuranosyl)benzene (9)

A solution of compound **8** (191 mg, 0.386 mmol), 20% Pd(OH)<sub>2</sub>/C (120 mg), and cyclohexene (3.9 mL, 38.6 mmol) in EtOH (8 mL) was stirred at 70 °C for 1.5 h. The mixture was filtered through a pad of Celite® and the filtrate was concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (CHCl<sub>3</sub>/MeOH=60:1) to give compound **9** (114 mg, 94%) as a colorless oil. [ $\alpha$ ]<sub>D</sub><sup>22</sup> +35.0 (*c* 1.03, CHCl<sub>3</sub>). IR  $\nu_{\max}$  (KBr) 3410, 2918, 1732, 1451, 1263, 1219, 1179, 1086 cm<sup>-1</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.01 (1H, ddd, *J*=6.5, 9.0, 13.5 Hz), 2.19 (1H, ddd, *J*=2.5, 6.0, 13.5 Hz), 2.33 (2H, s), 3.58–3.69 (2H, m), 3.84 (1H, ddd, *J*=4.0, 4.0, 4.0 Hz), 4.25–4.28 (1H, m), 5.31 (1H, dd, *J*=6.0, 9.0 Hz), 7.14 (1H, dd, *J*=1.0, 8.0 Hz), 7.28 (1H, ddd, *J*=1.0, 1.0, 8.0 Hz), 7.34 (1H, ddd, *J*=1.0, 8.0, 8.0 Hz), 7.50 (1H, ddd, *J*=1.0, 8.0, 8.0 Hz), 7.53 (2H, ddd, *J*=1.0, 8.0, 8.0 Hz), 7.64 (1H, ddd, *J*=1.0, 8.0, 8.0 Hz), 8.19 (2H, dd, *J*=1.0, 8.0 Hz). <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  42.6, 63.1, 73.2, 74.9, 86.6, 122.5, 126.4, 126.5, 128.6, 128.7, 129.1, 130.1, 133.8, 148.0, 165.3. HRMS (MALDI-TOF) *m/z* calcd for C<sub>18</sub>H<sub>18</sub>NaO<sub>5</sub> [M+Na]<sup>+</sup>: 337.1046; found 337.1052.

#### 4.6. 1-Benzoyloxy-2-[2-deoxy-5-O-(4,4'-dimethoxytrityl)- $\beta$ -D-ribofuranosyl]benzene (10)

Under a nitrogen atmosphere, DMTrCl (60 mg, 0.176 mmol) was added to a solution of compound **9** (46.0 mg, 0.146 mmol) in anhydrous pyridine (2 mL) at room temperature and the mixture was stirred for 2 h. After addition of water, the mixture was extracted with Et<sub>2</sub>O. The extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=5:1 to 3:1) to give compound **10** (89 mg, 98%) as a white solid. [ $\alpha$ ]<sub>D</sub><sup>26</sup> +30.2 (*c* 1.02, CHCl<sub>3</sub>). IR  $\nu_{\max}$  (KBr) 3456, 2933, 1737, 1607, 1509, 1450, 1252, 1176, 1079 cm<sup>-1</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.69 (1H, d, *J*=4.0 Hz), 2.03 (1H, ddd, *J*=6.5, 9.0, 13.0 Hz), 2.22 (1H, ddd, *J*=3.0, 6.5, 13.0 Hz), 3.28 (1H, dd, *J*=5.0, 10.0 Hz), 3.35 (1H, dd, *J*=5.0, 10.0 Hz), 3.79 (6H, s), 3.98 (1H, ddd, *J*=5.0, 5.0, 5.0 Hz), 4.32–4.35 (1H, m), 5.35 (1H, dd, *J*=6.5, 9.0 Hz), 6.82 (4H, d, *J*=9.0 Hz), 7.15–7.36 (10H, m), 7.44–7.53 (4H, m), 7.62–7.68 (2H, m), 8.19 (2H, d, *J*=8.0 Hz). <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  42.8, 55.2, 64.3, 74.3, 74.6, 85.5, 86.2, 113.1, 122.2, 126.3, 126.7, 126.8, 127.8, 128.2, 128.2, 128.7, 129.2, 130.1, 130.2, 133.7, 134.6, 136.0, 144.8, 147.6, 158.5, 164.9. HRMS

(FAB) *m/z* calcd for C<sub>39</sub>H<sub>36</sub>NaO<sub>7</sub> [M+Na]<sup>+</sup>: 639.2353; found 639.2350.

#### 4.7. 1-Benzoyloxy-2-{3-O-[2-cyanoethoxy(diisopropylamino)phosphino]-2-deoxy-5-O-(4,4'-dimethoxytrityl)- $\beta$ -D-ribofuranosyl]benzene (1)

Under a nitrogen atmosphere, *i*-Pr<sub>2</sub>NP(Cl)OCH<sub>2</sub>CH<sub>2</sub>CN (22  $\mu$ L, 0.098 mmol) was added to a solution of compound **10** (40 mg, 0.065 mmol) and *i*-Pr<sub>2</sub>NET (57 mL, 0.327 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (1 mL) at room temperature and the mixture was stirred for 1 h. After addition of satd NaHCO<sub>3</sub> aq, the mixture was extracted with Et<sub>2</sub>O. The extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=5:1) to give compound **1** (31 mg, 58%) as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  0.90 (3H, d, *J*=6.8 Hz), 0.96 (3H, d, *J*=6.4 Hz), 1.06–1.09 (6H, m), 1.89–2.01 (1H, m), 2.20–2.43 (3H, m), 3.23–3.35 (2H, m), 3.38–3.48 (3H, m), 3.53–3.58 (1H, m), 3.79 (3H, s), 3.79 (3H, s), 4.15–4.16 (1H, m), 4.39–4.47 (1H, m), 5.29–5.36 (1H, m), 6.80–6.84 (4H, m), 7.14–7.38 (10H, m), 7.46–7.56 (4H, m), 7.64–7.74 (2H, m), 8.18–8.23 (2H, m). <sup>31</sup>P NMR (CDCl<sub>3</sub>)  $\delta$  147.3, 148.3. HRMS (FAB) *m/z* calcd for C<sub>48</sub>H<sub>53</sub>N<sub>2</sub>NaO<sub>8</sub>P [M+Na]<sup>+</sup>: 839.3432; found 839.3421.

#### 4.8. 3-(2-Deoxy-3,5-di-O-benzyl- $\beta$ -D-ribofuranosyl)-1-(methoxymethoxy)benzene (11) and 3-(2-deoxy-3,5-di-O-benzyl- $\alpha$ -D-ribofuranosyl)-1-(methoxymethoxy)benzene (12)

Under a nitrogen atmosphere, a solution of compound **4** (180 mg, 0.573 mmol) in anhydrous THF (5 mL) was added to a solution of 3-methoxymethoxyphenyllithium [prepared from 3-methoxymethoxyphenyl bromide (690 mg, 3.18 mmol) and *n*-BuLi (1.65 M in hexane, 1.9 mL, 3.18 mmol) in anhydrous THF (20 mL) at -78 °C], and the mixture was stirred at -78 °C for 2 h. After addition of satd NH<sub>4</sub>Cl aq, the mixture was extracted with AcOEt. The extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=20:1) to give compound **11** (78 mg, 28%) as a colorless oil and compound **12** (66 mg, 24%) as a colorless oil. Compound **11**: [ $\alpha$ ]<sub>D</sub><sup>23</sup> +26.2 (*c* 1.03, CHCl<sub>3</sub>). IR  $\nu_{\max}$  (KBr) 2892, 1488, 1454, 1147, 1076 cm<sup>-1</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.91 (1H, ddd, *J*=6.0, 10.5, 13.0 Hz), 2.36 (1H, ddd, *J*=1.0, 6.0, 13.0 Hz), 3.45 (3H, s), 3.58 (1H, dd, *J*=5.5, 10.5 Hz), 3.67 (1H, dd, *J*=4.5, 10.5 Hz), 4.16 (1H, m), 4.30 (1H, ddd, *J*=2.5, 4.5, 5.5 Hz), 4.56 (2H, d, *J*=12.0 Hz), 4.60 (2H, d, *J*=12.0 Hz), 5.12 (1H, dd, *J*=5.0, 10.5 Hz), 5.14 (2H, s), 6.94 (1H, ddd, *J*=1.0, 2.0, 8.0 Hz), 7.01 (1H, m), 7.06 (1H, m), 7.20–7.38 (11H, m). <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  41.1, 55.9, 71.0, 71.1, 73.4, 80.2, 81.5, 83.8, 94.4, 114.0, 115.1, 119.6, 127.6, 127.6, 127.7, 128.3, 128.4, 129.3, 138.1, 138.2, 143.4, 157.3. HRMS (FAB) *m/z* calcd for C<sub>27</sub>H<sub>30</sub>NaO<sub>5</sub> [M+Na]<sup>+</sup>: 457.1985; found 457.1985. Compound **12**: [ $\alpha$ ]<sub>D</sub><sup>23</sup> +18.3 (*c* 1.24, CHCl<sub>3</sub>). IR  $\nu_{\max}$  (KBr) 2894, 1487, 1454, 1151, 1077 cm<sup>-1</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.06 (1H, ddd, *J*=6.5, 8.0, 13.0 Hz), 2.62 (1H, ddd, *J*=6.5, 6.5, 13.0 Hz), 3.45 (3H, s), 3.62 (2H, m), 4.26 (1H, ddd, *J*=4.5, 6.6, 6.5 Hz), 4.36 (1H, ddd, *J*=4.5, 4.5, 4.5 Hz), 4.49 (2H, d, *J*=12.0 Hz), 4.59 (2H, d, *J*=12.0 Hz), 5.05 (1H, dd, *J*=6.5, 8.0 Hz), 5.16 (2H, s), 6.93 (1H, ddd, *J*=1.0, 2.0, 8.0 Hz), 7.03 (1H, m), 7.08 (1H, m), 7.22–7.37 (11H, m). <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  40.9, 56.0, 70.8, 71.7, 73.5, 80.0, 80.6, 82.9, 94.4, 113.9,

115.1, 119.5, 127.6, 127.6, 128.3, 128.4, 129.4, 138.1, 144.5. HRMS (FAB)  $m/z$  calcd for  $C_{27}H_{30}NaO_5$   $[M+Na]^+$ : 457.1985; found 457.1988.

#### 4.9. 3-(2-Deoxy-3,5-di-O-benzyl- $\beta$ -D-ribofuranosyl)-1-hydroxybenzene (13)

10% HCl aq (7 mL) was added to a solution of compound **11** (286 mg, 0.657 mmol) in THF (10 mL), and mixture was stirred at room temperature for 2 days. The mixture was extracted with  $Et_2O$  and the extracts were washed with water and brine, dried over  $Na_2SO_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/ $AcOEt$ =5:1) to give compound **13** (234 mg, 91%) as a colorless oil.  $[\alpha]_D^{23} +20.5$  (c 0.96,  $CHCl_3$ ). IR  $\nu_{max}$  (KBr) 3331, 3032, 2861, 1591, 1456, 1077  $cm^{-1}$ .  $^1H$  NMR ( $CDCl_3$ )  $\delta$  1.89 (1H, ddd,  $J=5.5, 10.0, 13.0$  Hz), 2.33 (1H, ddd,  $J=1.5, 5.0, 13.0$  Hz), 3.58 (1H, dd,  $J=5.0, 10.0$  Hz), 3.67 (1H, dd,  $J=4.5, 10.0$  Hz), 4.15 (1H, m), 4.30 (1H, ddd,  $J=2.5, 4.5, 5.0$  Hz), 4.55 (2H, s), 4.58 (2H, s), 5.08 (1H, dd,  $J=5.0, 10.0$  Hz), 5.21 (1H, s), 6.67 (1H, ddd,  $J=1.0, 2.0, 8.0$  Hz), 6.80 (1H, m), 6.86 (1H, m), 7.13 (1H, dd,  $J=8.0, 8.0$  Hz), 7.24–7.36 (10H, m).  $^{13}C$  NMR ( $CDCl_3$ )  $\delta$  41.1, 71.1, 73.4, 80.3, 81.4, 83.7, 112.9, 114.6, 118.4, 127.7, 127.7, 128.4, 128.4, 129.5, 138.0, 138.2, 143.4, 155.8. HRMS (FAB)  $m/z$  calcd for  $C_{25}H_{26}NaO_4$   $[M+Na]^+$ : 413.1723; found 413.1712.

#### 4.10. 1-Benzoyloxy-3-(2-deoxy-3,5-di-O-benzyl- $\beta$ -D-ribofuranosyl)benzene (14)

Under a nitrogen atmosphere,  $BzCl$  (90  $\mu L$ , 0.779 mmol) was added to a solution of compound **13** (234 mg, 0.599 mmol) and  $Et_3N$  (109  $\mu L$ , 0.779 mmol) in anhydrous  $CH_2Cl_2$  (6 mL) at room temperature and the mixture was stirred for 30 min. After addition of water, the mixture was extracted with  $Et_2O$ . The extracts were washed with water and brine, dried over  $Na_2SO_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/ $AcOEt$ =20:1 to 10:1) to give compound **14** (290 mg, 98%) as a colorless oil.  $[\alpha]_D^{23} +12.7$  (c 1.03,  $CHCl_3$ ). IR  $\nu_{max}$  (KBr) 2860, 1738, 1451, 1233, 1078  $cm^{-1}$ .  $^1H$  NMR ( $CDCl_3$ )  $\delta$  1.93 (1H, ddd,  $J=6.5, 10.0, 13.0$  Hz), 2.40 (1H, ddd,  $J=1.0, 5.0, 13.0$  Hz), 3.58 (1H, dd,  $J=5.0, 10.5$  Hz), 3.67 (1H, dd,  $J=4.5, 10.5$  Hz), 4.16 (1H, m), 4.31 (1H, ddd,  $J=2.0, 4.5, 5.0$  Hz), 4.56 (2H, s), 4.58 (2H, s), 5.17 (1H, dd,  $J=5.0, 10.0$  Hz), 7.12 (1H, ddd,  $J=1.0, 2.0, 8.0$  Hz), 7.22–7.39 (13H, m), 7.50 (2H, dd,  $J=8.0, 8.0$  Hz), 7.63 (1H, dd,  $J=8.0, 8.0$  Hz), 8.20 (2H, d,  $J=8.0$  Hz).  $^{13}C$  NMR ( $CDCl_3$ )  $\delta$  41.2, 71.0, 73.4, 79.9, 81.5, 83.9, 119.2, 120.8, 123.5, 127.5, 127.6, 127.7, 128.3, 128.4, 128.5, 129.3, 129.6, 130.1, 133.5, 138.0, 138.1, 143.8, 151.0, 165.1. HRMS (FAB)  $m/z$  calcd for  $C_{32}H_{30}NaO_5$   $[M+Na]^+$ : 517.1985; found 517.2000.

#### 4.11. 1-Benzoyloxy-3-(2-deoxy- $\beta$ -D-ribofuranosyl)benzene (15)

A solution of compound **14** (230 mg, 0.466 mmol), 20%  $Pd(OH)_2/C$  (150 mg), and cyclohexene (4.7 mL, 46.6 mmol) in  $EtOH$  (10 mL) was stirred at 70 °C for 1.5 h. The mixture was filtered through a pad of Celite® and the filtrate was concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/ $AcOEt$ =2:1) to give compound **15** (140 mg, 96%) as a colorless oil.  $[\alpha]_D^{25} +19.1$  (c 1.88,  $CHCl_3$ ). IR  $\nu_{max}$  (KBr) 3372, 2929, 1736, 1263  $cm^{-1}$ .  $^1H$  NMR ( $CDCl_3$ )  $\delta$  1.95 (1H, ddd,  $J=6.5, 10.0, 13.0$  Hz), 2.22 (1H, ddd,  $J=2.0, 5.0, 13.0$  Hz), 2.94 (1H, br s), 3.13 (1H, br s), 3.68 (2H, m), 3.96 (1H, m), 4.30 (1H, m), 5.16 (1H, dd,  $J=5.0, 10.0$  Hz), 7.11 (1H, ddd,  $J=1.0, 2.0, 8.0$  Hz), 7.21 (2H, m), 7.37 (1H, dd,  $J=8.0, 8.0$  Hz), 7.49 (2H, dd,  $J=8.0, 8.0$  Hz), 7.62 (1H, dd,  $J=8.0, 8.0$  Hz), 8.17 (2H, dd,  $J=2.0, 8.0$  Hz).  $^{13}C$  NMR ( $CDCl_3$ )  $\delta$  43.6, 63.2, 73.4, 79.4, 87.4, 119.2, 120.9, 123.6, 128.6, 129.3, 129.5, 130.1, 133.7,

143.4, 151.0, 163.3. HRMS (MALDI-TOF)  $m/z$  calcd for  $C_{18}H_{18}NaO_5$   $[M+Na]^+$ : 337.1046; found 337.1058.

#### 4.12. 1-Benzoyloxy-3-[2-deoxy-5-O-(4,4'-dimethoxytrityl)- $\beta$ -D-ribofuranosyl]benzene (16)

Under a nitrogen atmosphere,  $DMTrCl$  (49 mg, 0.145 mmol) was added to a solution of compound **15** (38 mg, 0.121 mmol) in anhydrous pyridine (1.5 mL) at room temperature and the mixture was stirred for 2 h. After addition of water, the mixture was extracted with  $Et_2O$ . The extracts were washed with water, dried over  $Na_2SO_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/ $AcOEt$ =5:1 to 3:1) to give compound **16** (53 mg, 72%) as a white solid.  $[\alpha]_D^{23} +23.2$  (c 1.16,  $AcOEt$ ). IR  $\nu_{max}$  (KBr) 3454, 2932, 1736, 1607, 1508, 1252, 1176, 1080  $cm^{-1}$ .  $^1H$  NMR ( $CDCl_3$ )  $\delta$  1.76 (1H, d,  $J=4.0$  Hz), 2.08 (1H, ddd,  $J=6.0, 10.0, 13.5$  Hz), 2.28 (1H, ddd,  $J=2.0, 6.0, 13.5$  Hz), 3.25 (1H, dd,  $J=4.5, 10.0$  Hz), 3.35 (1H, dd,  $J=5.0, 10.0$  Hz), 3.75 (6H, s), 4.06 (1H, ddd,  $J=4.5, 5.0, 7.0$  Hz), 4.46 (1H, m), 5.20 (1H, dd,  $J=6.0, 10.0$  Hz), 6.75–6.85 (4H, m), 7.10–7.40 (11H, m), 7.45 (2H, dd,  $J=1.0, 8.0$  Hz), 7.48 (2H, dd,  $J=8.0, 8.0$  Hz), 7.62 (1H, ddd,  $J=1.0, 8.0, 8.0$  Hz), 8.20 (2H, dd,  $J=1.5, 8.0$  Hz).  $^{13}C$  NMR ( $CDCl_3$ )  $\delta$  44.0, 55.1, 64.5, 74.6, 79.4, 86.2, 86.4, 113.1, 119.2, 120.8, 123.4, 126.7, 127.8, 128.1, 128.5, 129.3, 129.5, 130.0, 130.1, 130.1, 133.5, 135.9, 136.0, 143.9, 144.8, 151.0, 158.4, 158.4, 165.1. HRMS (FAB)  $m/z$  calcd for  $C_{39}H_{36}NaO_7$   $[M+Na]^+$ : 639.2353; found 639.2376.

#### 4.13. 1-Benzoyloxy-3-{3-O-[2-cyanoethoxy(diisopropylamino)phosphino]-2-deoxy-5-O-(4,4'-dimethoxytrityl)- $\beta$ -D-ribofuranosyl}benzene (2)

Under a nitrogen atmosphere, *i*- $Pr_2NP(Cl)OCH_2CH_2CN$  (30  $\mu L$ , 0.134 mmol) was added to a solution of compound **16** (41 mg, 0.067 mmol) and *i*- $Pr_2NEt$  (58  $\mu L$ , 0.335 mmol) in anhydrous  $CH_2Cl_2$  (1 mL) at room temperature and the mixture was stirred for 30 min. After addition of satd  $NaHCO_3$  aq, the mixture was extracted with  $Et_2O$ . The extracts were washed with water, dried over  $Na_2SO_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/ $AcOEt$ =5:1 to 4:1) to give compound **2** (32 mg, 58%) as a white solid.  $^1H$  NMR ( $CDCl_3$ )  $\delta$  1.08 (3H, d,  $J=6.8$  Hz), 1.15–1.19 (9H, m), 2.02–2.11 (1H, m), 2.34–2.47 (2H, m), 2.61 (1H, dd,  $J=6.3, 6.8$  Hz), 3.23–3.34 (2H, m), 3.53–3.87 (4H, m), 3.74 (3H, s), 3.74 (3H, s), 4.24–4.25 (1H, m), 4.49–4.53 (1H, m), 5.18–5.22 (1H, m), 6.76–6.82 (4H, m), 7.12–7.18 (2H, m), 7.22–7.27 (2H, m), 7.31–7.41 (7H, m), 7.45–7.52 (4H, m), 7.59–7.63 (1H, m), 8.12–8.15 (2H, m).  $^{31}P$  NMR ( $CDCl_3$ )  $\delta$  147.8, 148.0. HRMS (FAB)  $m/z$  calcd for  $C_{48}H_{53}N_2NaO_8P$   $[M+Na]^+$ : 839.3422; found 839.3444.

#### 4.14. 1-Allyloxy-4-(2-deoxy-3,5-di-O-benzyl-D-ribofuranosyl)benzene (17)

Under a nitrogen atmosphere, a solution of compound **4** (1.00 g, 3.18 mmol) in anhydrous THF (30 mL) was added to a solution of 4-allyloxyphenyllithium [prepared from 4-allyloxyphenyl bromide (2.70 g, 12.7 mmol) and *s*- $BuLi$  (1.05 M in hexane and cyclohexane, 12.1 mL, 12.7 mmol) in anhydrous THF (130 mL) at –78 °C], and the mixture was stirred at –60 °C for 1 h. After addition of satd  $NH_4Cl$  aq, the mixture was extracted with  $CH_2Cl_2$ . The extracts were dried over  $Na_2SO_4$  and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/ $AcOEt$ =3:2) to give appropriate compounds ( $R_f$ =ca. 0.5), which were dissolved in  $CH_2Cl_2$  (30 mL);  $TMAD$  (600 mg, 3.48 mmol) and *n*- $Bu_3P$  (0.87 mL, 3.48 mL) were added at room temperature. After being stirred at room temperature for 19 h, water was added and the mixture was extracted with  $Et_2O$ . The

extracts were washed with water and brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=5:1) to give a ca. 1:1 anomeric mixture **17** (660 mg, 48% from **4**) as a colorless oil.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.92 (0.5H, ddd,  $J=5.9, 11.0, 13.2$  Hz), 2.05 (0.5H, ddd,  $J=6.2, 8.4, 12.8$  Hz), 2.31 (0.5H, dd,  $J=5.1, 13.2$  Hz), 2.59 (0.5H, td,  $J=6.6, 12.8$  Hz), 3.55–3.69 (2H, m), 4.15–4.17 (0.5H, m), 4.23–4.36 (1.5H, m), 4.46–4.64 (6H, m), 5.00–5.09 (1H, m), 5.26–5.30 (1H, m), 5.37–5.43 (1H, m), 5.99–6.11 (1H, m), 6.85–6.89 (2H, m), 7.26–7.34 (12H, m). HRMS (FAB)  $m/z$  calcd for  $\text{C}_{29}\text{H}_{31}\text{O}_4$   $[\text{M}+\text{H}]^+$ : 431.2217; found 431.2224.

#### 4.15. 4-(2-Deoxy-3,5-di-O-benzyl-D-ribofuranosyl)-1-hydroxybenzene (18)

Under a nitrogen atmosphere,  $\text{Pd}(\text{PPh}_3)_4$  (207 mg, 0.179 mmol) was added to a solution of compound **17** (385 mg, 0.895 mmol) in anhydrous THF (10 mL) and the mixture was stirred until solids of  $\text{Pd}(\text{PPh}_3)_4$  dissolved completely. Then,  $\text{NaBH}_4$  (41 mg, 1.07 mmol) was added and the mixture was stirred at room temperature for 17 h. After addition of 3% HCl aq, the mixture was filtered through a pad of Celite®. The filtrate was extracted with  $\text{Et}_2\text{O}$ . The extracts were washed with water and brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography ( $\text{CHCl}_3$  to  $\text{CHCl}_3/\text{MeOH}=60:1$ ) to give a ca. 1:1 anomeric mixture **18** (318 mg, 91%) as a colorless oil.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.91 (0.5H, ddd,  $J=5.9, 10.8, 13.2$  Hz), 2.04 (0.5H, ddd,  $J=6.4, 8.8, 12.7$  Hz), 2.31 (0.5H, ddd,  $J=1.0, 4.9, 13.2$  Hz), 2.59 (0.5H, td,  $J=6.4, 12.7$  Hz), 3.56–3.68 (2H, m), 4.15–4.17 (0.5H, m), 4.23–4.30 (1H, m), 4.33 (0.5H, td,  $J=4.9, 9.3$  Hz), 4.47–4.62 (4H, m), 5.01 (0.5H, dd,  $J=6.4, 8.8$  Hz), 5.07 (0.5H, dd,  $J=4.9, 10.8$  Hz), 5.12 (0.5H, br s), 5.14 (0.5H, br s), 6.71–6.75 (2H, m), 7.20–7.35 (12H, m). HRMS (FAB)  $m/z$  calcd for  $\text{C}_{25}\text{H}_{26}\text{NaO}_4$   $[\text{M}+\text{Na}]^+$ : 413.1723; found 413.1742.

#### 4.16. 1-Benzoyloxy-4-(2-deoxy-3,5-di-O-benzyl- $\beta$ -D-ribofuranosyl)benzene (19) and 1-benzoyloxy-4-(2-deoxy-3,5-di-O-benzyl- $\alpha$ -D-ribofuranosyl)benzene (20)

Under a nitrogen atmosphere,  $\text{BzCl}$  (129  $\mu\text{L}$ , 1.11 mmol) was added to a solution of compound **18** (318 mg, 0.923 mmol) and  $\text{Et}_3\text{N}$  (154  $\mu\text{L}$ , 1.11 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (10 mL) at room temperature and the mixture was stirred for 1.5 h. After addition of water, the mixture was extracted with  $\text{Et}_2\text{O}$ . The extracts were washed with water and brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=20:1) to give a separable mixture of **19** and **20** (365 mg, 91%, **19/20**=ca. 1:1) as a colorless oil. Compound **19**:  $[\alpha]_D^{22} +14.7$  ( $c$  1.06,  $\text{CHCl}_3$ ). IR  $\nu_{\text{max}}$  (KBr) 2860, 1739, 1263, 1078  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.94 (1H, ddd,  $J=6.0, 10.5, 13.0$  Hz), 2.38 (1H, ddd,  $J=1.5, 5.5, 13.0$  Hz), 3.60 (1H, dd,  $J=5.0, 10.0$  Hz), 3.67 (1H, dd,  $J=5.0, 10.0$  Hz), 4.18 (1H, m), 4.28 (1H, ddd,  $J=2.5, 5.0, 5.0$  Hz), 4.58 (2H, d,  $J=14.0$  Hz), 4.60 (2H, d,  $J=14.0$  Hz), 5.18 (1H, dd,  $J=5.5, 10.5$  Hz), 7.20 (2H, ddd,  $J=1.0, 1.0, 8.0$  Hz), 7.25–7.38 (10H, m), 7.42 (2H, ddd,  $J=1.0, 1.0, 8.0$  Hz), 7.51 (2H, ddd,  $J=1.0, 8.0, 8.0$  Hz), 7.63 (1H, ddd,  $J=1.0, 1.0, 8.0$  Hz), 8.20 (2H, ddd,  $J=1.0, 1.0, 8.0$  Hz).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  41.3, 71.1, 71.1, 73.5, 80.0, 81.5, 83.9, 121.5, 127.2, 127.6, 127.7, 128.4, 128.4, 128.5, 129.5, 130.2, 133.6, 138.1, 138.2, 139.4, 150.2, 165.2. HRMS (FAB)  $m/z$  calcd for  $\text{C}_{32}\text{H}_{30}\text{NaO}_5$   $[\text{M}+\text{Na}]^+$ : 517.1985; found 517.2005. Compound **20**:  $[\alpha]_D^{24} +7.4$  ( $c$  0.87,  $\text{CHCl}_3$ ). IR  $\nu_{\text{max}}$  (KBr) 2864, 1739, 1264, 1078  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  2.08 (1H, ddd,  $J=6.0, 7.5, 13.5$  Hz), 2.65 (1H, ddd,  $J=7.0, 7.0, 13.5$  Hz), 3.59–3.67 (2H, m), 4.28 (1H, ddd,  $J=4.5, 6.5, 6.5$  Hz), 4.38 (1H, ddd,  $J=4.5, 4.5, 4.5$  Hz), 4.49 (1H, d,  $J=12.0$  Hz), 4.51 (1H, d,  $J=12.0$  Hz), 4.59 (1H, d,  $J=12.0$  Hz), 4.61 (1H, d,  $J=12.0$  Hz), 5.12 (1H, dd,  $J=7.0, 7.5$  Hz), 7.18 (2H, d,  $J=8.0$  Hz),

7.24–7.38 (10H, m), 7.46 (2H, d,  $J=8.0$  Hz), 7.51 (2H, dd,  $J=8.0, 8.0$  Hz), 7.64 (1H, dd,  $J=8.0, 8.0$  Hz), 8.20 (2H, d,  $J=8.0$  Hz).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  41.5, 70.8, 71.7, 73.5, 79.8, 80.6, 82.9, 121.5, 127.2, 127.6, 127.6, 127.7, 128.4, 128.5, 130.1, 133.5, 138.0, 138.2, 140.3, 150.1, 165.2. HRMS (FAB)  $m/z$  calcd for  $\text{C}_{32}\text{H}_{30}\text{NaO}_5$   $[\text{M}+\text{Na}]^+$ : 517.1985; found 517.1974.

#### 4.17. 1-Benzoyloxy-4-(2-deoxy- $\beta$ -D-ribofuranosyl)benzene (21)

A solution of compound **19** (170 mg, 0.343 mmol), 20%  $\text{Pd}(\text{OH})_2/\text{C}$  (100 mg), and cyclohexene (3.5 mL, 34.3 mmol) in EtOH (7 mL) was stirred at 70 °C for 2 h. The mixture was filtered through a pad of Celite® and the filtrate was concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=2:1) to give compound **21** (89 mg, 82%) as a white solid.  $[\alpha]_D^{22} +34.8$  ( $c$  1.05,  $\text{CHCl}_3$ ). IR  $\nu_{\text{max}}$  (KBr) 3390, 2932, 1737, 1508, 1270, 1200, 1079  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  2.05 (1H, ddd,  $J=6.0, 10.5, 13.0$  Hz), 2.14 (1H, br s), 2.19 (1H, br s), 2.26 (1H, ddd,  $J=1.5, 5.5, 13.0$  Hz), 3.76 (2H, m), 4.01 (1H, ddd,  $J=4.5, 4.5, 4.5$  Hz), 4.42 (1H, m), 5.20 (1H, dd,  $J=5.5, 10.5$  Hz), 7.20 (2H, ddd,  $J=1.0, 1.0, 8.0$  Hz), 7.41 (2H, ddd,  $J=1.0, 1.0, 8.0$  Hz), 7.51 (2H, ddd,  $J=1.0, 8.0, 8.0$  Hz), 7.65 (1H, dddd,  $J=1.0, 1.0, 8.0, 8.0$  Hz), 8.20 (2H, dd,  $J=1.0, 8.0$  Hz).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  44.0, 63.4, 73.7, 79.6, 87.3, 121.7, 127.2, 128.6, 129.4, 130.2, 133.7, 138.8, 150.4, 165.3. HRMS (FAB)  $m/z$  calcd for  $\text{C}_{18}\text{H}_{18}\text{NaO}_5$   $[\text{M}+\text{Na}]^+$ : 337.1046; found 337.1035.

#### 4.18. 1-Benzoyloxy-4-[2-deoxy-5-O-(4,4'-dimethoxytrityl)- $\beta$ -D-ribofuranosyl]benzene (22)

Under a nitrogen atmosphere,  $\text{DMTrCl}$  (106 mg, 0.313 mmol) was added to a solution of compound **21** (82 mg, 0.260 mmol) in anhydrous pyridine (3 mL) at room temperature and the mixture was stirred for 1 h. After addition of water, the mixture was extracted with  $\text{Et}_2\text{O}$ . The extracts were washed with water, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=3:1) to give compound **22** (145 mg, 91%) as a white solid.  $[\alpha]_D^{23} +13.8$  ( $c$  1.06,  $\text{CHCl}_3$ ). IR  $\nu_{\text{max}}$  (KBr) 3454, 3004, 2933, 2837, 1739, 1607, 1508, 1263, 1079  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.87 (1H, br s), 2.05 (1H, ddd,  $J=5.5, 10.0, 13.0$  Hz), 2.25 (1H,  $J=1.5, 5.5, 13.0$  Hz), 3.28 (1H, dd,  $J=4.5, 10.0$  Hz), 3.36 (1H, dd,  $J=5.0, 10.0$  Hz), 3.78 (6H, s), 4.07 (1H, m), 4.30 (1H, m), 5.20 (1H, dd,  $J=5.5, 10.0$  Hz), 6.82 (4H, d,  $J=9.0$  Hz), 7.15–7.55 (15 H, m), 7.63 (1H, dd,  $J=7.5, 7.5$  Hz), 8.20 (1H, d,  $J=7.5$  Hz).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  43.9, 55.2, 64.4, 74.6, 79.5, 86.2, 86.3, 113.1, 121.5, 126.8, 127.1, 127.8, 128.2, 128.6, 129.5, 130.1, 130.1, 133.6, 136.0, 139.5, 144.8, 150.2, 158.4, 165.2. HRMS (FAB)  $m/z$  calcd for  $\text{C}_{39}\text{H}_{36}\text{NaO}_7$   $[\text{M}+\text{Na}]^+$ : 639.2353; found 639.2359.

#### 4.19. 1-Benzoyloxy-4-{3-O-[2-cyanoethoxy(diisopropylamino)phosphino]-2-deoxy-5-O-(4,4'-dimethoxytrityl)- $\beta$ -D-ribofuranosyl]benzene (3)

Under a nitrogen atmosphere, *i*-Pr<sub>2</sub>NP(Cl)OCH<sub>2</sub>CH<sub>2</sub>CN (36  $\mu\text{L}$ , 0.161 mmol) was added to a solution of compound **22** (66 mg, 0.108 mmol) and *i*-Pr<sub>2</sub>NEt (96  $\mu\text{L}$ , 0.554 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (1 mL) at room temperature and the mixture was stirred for 25 min. After addition of satd  $\text{NaHCO}_3$  aq, the mixture was extracted with  $\text{Et}_2\text{O}$ . The extracts were washed with water, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=5:1) to give compound **3** (84 mg, 95%) as a white solid.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.09 (3H, d,  $J=6.8$  Hz), 1.16–1.20 (9H, m), 2.00–2.09 (1H, m), 2.32–2.49 (2H, m), 2.63 (1H, dd,  $J=6.3, 6.9$  Hz), 3.23–3.36 (2H, m), 3.53–3.89 (4H, m), 3.79 (3H, s), 3.79 (3H, s), 4.24–4.25 (1H, m), 4.51–4.55 (1H, m), 5.17–5.22 (1H, m), 6.80–6.84 (4H, m), 7.17–7.30



(5H, m), 7.35–7.38 (4H, m), 7.47–7.53 (6H, m), 7.62–7.66 (1H, m), 8.19–8.21 (2H, m).  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ )  $\delta$  147.8, 148.0. HRMS (FAB)  $m/z$  calcd for  $\text{C}_{48}\text{H}_{53}\text{N}_2\text{NaO}_8\text{P}$  [ $\text{M}+\text{Na}$ ] $^+$ : 839.3422; found 839.3443.

#### 4.20. 1-Allyloxy-2-(3,5-di-*O*-benzyl-2-*O*,4-*C*-methylene- $\beta$ -*D*-ribofuranosyl)benzene (27)

Under a nitrogen atmosphere, a solution of compound **26**<sup>8</sup> (1.95 g, 5.69 mmol) in anhydrous THF (50 mL) was added to a solution of 2-allyloxyphenylmagnesium iodide [prepared from 2-allyloxyphenyl iodide (5.92 g, 22.8 mmol) and *i*-PrMgCl·LiCl (1.3 M in THF, 17.5 mL, 22.8 mmol) in anhydrous THF (180 mL) at  $-40^\circ\text{C}$ ], and the mixture was stirred at  $-40^\circ\text{C}$  for 11 h. After addition of satd  $\text{NH}_4\text{Cl}$  aq, the mixture was extracted with AcOEt. The extracts were washed with water and brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=3:1 to 1:1) to give an appropriate compound ( $R_f$ =ca. 0.5, *n*-hexane/AcOEt=1:1), which was dissolved in anhydrous  $\text{CH}_2\text{Cl}_2$  (50 mL); TMAD (1.11 g, 6.47 mmol) and *n*-Bu<sub>3</sub>P (1.61 mL, 6.47 mmol) were added at room temperature and the mixture was stirred for 2 h. The mixture was concentrated under reduced pressure and the residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=5:1) to give compound **27** (1.48 g, 57% from **26**) as a white powder.  $[\alpha]_D^{25} +60.3$  (*c* 1.36,  $\text{CHCl}_3$ ). IR  $\nu_{\text{max}}$  (KBr) 3063, 3031, 2876, 1596, 1490, 1454, 1239, 1102, 1030  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.87, 3.88 (2H, AB,  $J=11$  Hz), 4.02 (1H, s), 4.08, 4.12 (2H, AB,  $J=8$  Hz), 4.41 (2H, s), 4.40, 4.50 (2H, AB,  $J=12$  Hz), 4.67 (2H, s), 5.26 (1H, dd,  $J=1, 11$  Hz), 5.40 (1H, dd,  $J=1, 17$  Hz), 5.35 (1H, s), 5.94–6.08 (1H, m), 6.80 (1H, d,  $J=8$  Hz), 6.94 (1H, t,  $J=8$  Hz), 7.18–7.40 (11H, m), 7.54 (1H, d,  $J=8$  Hz).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  66.5, 68.3, 71.8, 73.3, 73.5, 78.0, 79.0, 80.5, 85.4, 110.7, 117.0, 120.5, 126.7, 127.3, 127.4, 127.4, 127.5, 127.5, 128.1, 128.2, 128.2, 132.8, 137.5, 138.0, 154.2. Mass (EI)  $m/z$  458 ( $\text{M}^+$ , 2.5). Anal. Calcd for  $\text{C}_{29}\text{H}_{30}\text{O}_5$ : C, 75.96; H, 6.59; found: C, 75.94; H, 6.63.

#### 4.21. 2-(3,5-Di-*O*-benzyl-2-*O*,4-*C*-methylene- $\beta$ -*D*-ribofuranosyl)-1-hydroxybenzene (29)

Under a nitrogen atmosphere,  $\text{Pd}(\text{PPh}_3)_4$  (370 mg, 0.32 mmol) was added to a solution of compound **27** (1.46 g, 3.18 mmol) in anhydrous THF (30 mL) and the mixture was stirred until solids of  $\text{Pd}(\text{PPh}_3)_4$  dissolved completely. Then,  $\text{NaBH}_4$  (361 mg, 9.55 mmol) was added and the mixture was stirred at room temperature for 19 h. After addition of 3% HCl aq, the mixture was filtered through a pad of Celite<sup>®</sup>. The filtrate was extracted with AcOEt. The extracts were washed with water and brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=5:2) to give compound **29** (1.07 g, 80%) as a white powder.  $[\alpha]_D^{25} +14.0$  (*c* 1.19,  $\text{CHCl}_3$ ). IR  $\nu_{\text{max}}$  (KBr) 3336, 3031, 2877, 1496, 1456, 1100, 1029  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.72 (2H, s), 3.97, 4.04 (2H, AB,  $J=8$  Hz), 4.31 (2H, d,  $J=2$  Hz), 4.50, 4.59 (2H, AB,  $J=11$  Hz), 4.66 (2H, d,  $J=2$  Hz), 5.42 (1H, s), 6.80–6.91 (3H, m), 7.13–7.37 (11H, m), 8.55 (1H, s).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  64.1, 72.3, 72.8, 73.6, 77.8, 82.1, 86.5, 118.0, 119.5, 120.5, 126.2, 127.5, 127.7, 127.8, 128.3, 128.4, 128.9, 137.2, 137.3, 155.3. HRMS (MALDI-TOF)  $m/z$  calcd for  $\text{C}_{26}\text{H}_{26}\text{NaO}_5$  [ $\text{M}+\text{Na}$ ] $^+$ : 441.1678; found, 441.1676.

#### 4.22. 1-Benzoyloxy-2-(3,5-di-*O*-benzyl-2-*O*,4-*C*-methylene- $\beta$ -*D*-ribofuranosyl)benzene (30)

Under a nitrogen atmosphere, BzCl (42  $\mu\text{L}$ , 0.36 mmol) was added to a solution of compound **29** (112 mg, 0.28 mmol) and  $\text{Et}_3\text{N}$  (50  $\mu\text{L}$ , 0.36 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (2 mL) at room temperature and the mixture was stirred for 3 h. After addition of water and

concentrated under reduced pressure, the mixture was extracted with AcOEt. The extracts were washed with water and brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=7:1) to give compound **30** (130 mg, 93%) as a colorless oil.  $[\alpha]_D^{25} +52.1$  (*c* 1.14,  $\text{CHCl}_3$ ). IR  $\nu_{\text{max}}$  (KBr) 3063, 3032, 2877, 1737, 1489, 1452, 1259, 1203, 1029  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.86 (2H, s), 4.01, 4.07 (2H, AB,  $J=8$  Hz), 4.09 (1H, s), 4.34 (1H, s), 4.44, 4.53 (2H, AB,  $J=12$  Hz), 4.66, 4.67 (2H, AB,  $J=12$  Hz), 5.29 (1H, s), 7.18–7.39 (13H, m), 7.48–7.54 (2H, m), 7.61–7.67 (2H, m), 8.14–8.20 (2H, m).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  66.3, 72.1, 73.4, 73.6, 77.7, 79.5, 80.1, 85.7, 122.1, 126.1, 127.3, 127.4, 127.5, 127.6, 128.2, 128.3, 128.4, 128.7, 128.9, 133.8, 137.4, 138.0, 147.0, 164.7. Mass (EI)  $m/z$  522 ( $\text{M}^+$ , 1.3). Anal. Calcd for  $\text{C}_{33}\text{H}_{30}\text{O}_6$ : C, 75.84; H, 5.79; found: C, 75.49; H, 5.88.

#### 4.23. 1-Benzoyloxy-2-(2-*O*,4-*C*-methylene- $\beta$ -*D*-ribofuranosyl)benzene (31)

A solution of compound **30** (263 mg, 0.503 mmol), 20%  $\text{Pd}(\text{OH})_2/\text{C}$  (100 mg), and cyclohexene (5.0 mL, 50.3 mmol) in EtOH (5 mL) was refluxed for 1 h. The mixture was filtered through a pad of Celite<sup>®</sup> and the filtrate was concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=1:2 to 1:5) to give compound **31** (153 mg, 89%) as a white solid.  $[\alpha]_D^{25} +4.2$  (*c* 0.94,  $\text{CH}_3\text{OH}$ ). IR  $\nu_{\text{max}}$  (KBr) 3379, 2947, 1736, 1452, 1264, 1200, 1173, 1026  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  3.87, 3.98 (2H, AB,  $J=8$  Hz), 3.90, 3.93 (2H, AB,  $J=11$  Hz), 4.13 (1H, s), 4.13 (1H, s), 5.11 (1H, s), 7.19–7.23 (1H, m), 7.31–7.38 (2H, m), 7.54–7.60 (2H, m), 7.68–7.73 (2H, m), 8.17–8.21 (2H, m).  $^{13}\text{C}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  59.3, 71.5, 73.6, 80.9, 83.7, 88.3, 123.4, 127.2, 128.3, 129.6, 129.9, 130.2, 131.0, 132.7, 135.1, 148.5, 166.1. Mass (EI)  $m/z$  342 ( $\text{M}^+$ , 0.3). Anal. Calcd for  $\text{C}_{19}\text{H}_{18}\text{O}_6$ : C, 66.66; H, 5.30; found: C, 66.34; H, 5.33.

#### 4.24. 1-Benzoyloxy-2-[5-*O*-(4,4'-dimethoxytrityl)-2-*O*,4-*C*-methylene- $\beta$ -*D*-ribofuranosyl]benzene (32)

Under a nitrogen atmosphere,  $\text{DMTrCl}$  (85 mg, 0.25 mmol) was added to a solution of compound **31** (58 mg, 0.17 mmol) in anhydrous pyridine (1 mL) at room temperature and the mixture was stirred for 3 h. After addition of satd  $\text{NaHCO}_3$  aq, the mixture was extracted with AcOEt. The extracts were washed with water and brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=2:1) to give compound **32** (121 mg, 94%) as a white solid.  $[\alpha]_D^{25} -13.5$  (*c* 1.00,  $\text{CHCl}_3$ ). IR  $\nu_{\text{max}}$  (KBr) 3458, 3007, 2949, 2836, 1737, 1606, 1508, 1451, 1254, 1175, 1068, 1030  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (acetone-*d*<sub>6</sub>)  $\delta$  3.46, 3.54 (2H, AB,  $J=11$  Hz), 3.79 (6H, s), 3.91 (2H, s), 4.16 (1H, s), 4.35 (1H, s), 5.19 (1H, s), 6.89–6.92 (1H, m), 7.22–7.46 (10H, m), 7.57–7.66 (4H, m), 7.74–7.79 (1H, m), 7.91–7.94 (1H, m), 8.23–8.26 (2H, m).  $^{13}\text{C}$  NMR (acetone-*d*<sub>6</sub>)  $\delta$  55.6, 61.0, 71.8, 73.6, 80.8, 82.9, 86.9, 86.9, 113.9, 123.5, 127.0, 127.6, 128.4, 128.6, 129.0, 129.4, 129.8, 130.1, 130.8, 131.0, 131.0, 133.1, 134.9, 136.7, 136.9, 146.2, 148.3, 159.5, 165.3. HRMS (FAB)  $m/z$  calcd for  $\text{C}_{40}\text{H}_{36}\text{NaO}_8$  [ $\text{M}+\text{Na}$ ] $^+$ : 667.2308; found, 667.2304.

#### 4.25. 1-Benzoyloxy-2-[3-*O*-[2-cyanoethoxy(diisopropylamino)phosphino]-5-*O*-(4,4'-dimethoxytrityl)-2-*O*,4-*C*-methylene- $\beta$ -*D*-ribofuranosyl]benzene (23)

Under a nitrogen atmosphere, *i*-Pr<sub>2</sub>NP(Cl)OCH<sub>2</sub>CH<sub>2</sub>CN (46  $\mu\text{L}$ , 0.206 mmol) was added to a solution of compound **32** (33 mg, 0.051 mmol) and *i*-Pr<sub>2</sub>NEt (71  $\mu\text{L}$ , 0.408 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (1 mL) at room temperature and the mixture was stirred for 1 h. After addition of satd  $\text{NaHCO}_3$  aq, the mixture was extracted with AcOEt. The extracts were washed with satd

NaHCO<sub>3</sub> aq, water, and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=4:1) to give compound **23** (40 mg, 92%) as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.87 (3H, d, *J*=6.4 Hz), 0.94 (3H, d, *J*=6.9 Hz), 1.07–1.10 (6H, m), 2.29–2.32 (1H, m), 2.43–2.47 (1H, m), 3.39–3.56 (5H, m), 3.65–3.71 (1H, m), 3.81 (3H, s), 3.82 (3H, s), 3.86–3.90 (1H, m), 4.01–4.07 (1H, m), 4.31–4.45 (2H, m), 5.33 (0.5H, s), 5.36 (0.5H, s), 6.84–6.88 (4H, m), 7.22–7.46 (10H, m), 7.52–7.58 (4H, m), 7.65–7.69 (1H, m), 7.88–7.96 (1H, m), 8.19–8.24 (2H, m). <sup>31</sup>P NMR (acetone-*d*<sub>6</sub>) δ 146.9, 147.7. HRMS (FAB) *m/z* calcd for C<sub>49</sub>H<sub>54</sub>N<sub>2</sub>O<sub>9</sub>P [M+H]<sup>+</sup>: 845.3567; found, 845.3565.

#### 4.26. 2-[5-*O*-(4,4'-Dimethoxytrityl)-2-*O*,4-*C*-methylene-β-*D*-ribofuranosyl]-1-hydroxybenzene (**33**)

Under a nitrogen atmosphere, K<sub>2</sub>CO<sub>3</sub> (103 mg, 0.75 mmol) was added to a solution of compound **32** (120 mg, 0.19 mmol) in MeOH (5 mL) at room temperature and the mixture was stirred for 5 min. After removal of MeOH under reduced pressure and the mixture was extracted with AcOEt. The extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=3:1) to give compound **33** (96 mg, 96%) as a white solid. [α]<sub>D</sub><sup>22</sup> –2.3 (c 0.57, CHCl<sub>3</sub>). IR ν<sub>max</sub> (KBr) 3338, 3006, 2949, 1606, 1507, 1457, 1296, 1250, 1030 cm<sup>-1</sup>. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>) δ 2.76 (1H, br s), 3.32, 3.39 (2H, AB, *J*=10 Hz), 3.66 (6H, s), 3.82 (2H, s), 4.09 (1H, s), 5.10 (1H, s), 6.70–6.80 (6H, m), 6.97–7.03 (1H, m), 7.08–7.23 (3H, m), 7.28–7.33 (4H, m), 7.44–7.54 (3H, m). <sup>13</sup>C NMR (acetone-*d*<sub>6</sub>) δ 55.6, 61.0, 71.8, 73.6, 80.8, 82.9, 86.9, 86.9, 113.9, 123.5, 127.0, 127.6, 128.4, 128.6, 129.0, 129.4, 129.8, 130.1, 130.8, 131.0, 131.0, 133.1, 134.9, 136.7, 136.9, 146.2, 148.3, 159.5, 165.3. HRMS (FAB) *m/z* calcd for C<sub>33</sub>H<sub>32</sub>NaO<sub>7</sub> [M+Na]<sup>+</sup>: 563.2046; found, 563.2050.

#### 4.27. 1-Allyloxy-2-[5-*O*-(4,4'-dimethoxytrityl)-2-*O*,4-*C*-methylene-β-*D*-ribofuranosyl]benzene (**34**)

Under a nitrogen atmosphere, K<sub>2</sub>CO<sub>3</sub> (27 mg, 0.20 mmol) and allyl bromide (17 μL, 0.20 mmol) were added to a solution of compound **33** (96 mg, 0.18 mmol) in anhydrous acetone (16 mL) at room temperature and the mixture was stirred for 14 h. After addition of water, acetone was removed under reduced pressure. The mixture was extracted with AcOEt. The extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=3:1) to give compound **34** (90 mg, 87%) as a white solid. [α]<sub>D</sub><sup>22</sup> +5.1 (c 0.93, CHCl<sub>3</sub>). IR ν<sub>max</sub> (KBr) 3431, 3007, 2943, 2836, 1605, 1508, 1453, 1298, 1249, 1178, 1078, 1031 cm<sup>-1</sup>. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>) δ 3.47, 3.53 (2H, AB, *J*=10 Hz), 3.77 (6H, s), 3.96 (2H, s), 4.21 (1H, s), 4.64 (2H, s), 5.25 (1H, s), 5.26 (1H, d, *J*=10 Hz), 5.47 (1H, d, *J*=17 Hz), 6.05–6.15 (1H, m), 6.88–7.02 (6H, m), 7.20–7.35 (4H, m), 7.44–7.47 (4H, m), 7.59–7.62 (2H, m), 7.75–7.78 (1H, m). <sup>13</sup>C NMR (acetone-*d*<sub>6</sub>) δ 55.6, 61.3, 69.1, 72.1, 73.5, 79.3, 81.1, 82.5, 86.7, 86.9, 112.2, 113.9, 117.1, 121.4, 127.6, 127.9, 128.6, 129.1, 129.3, 129.3, 131.0, 131.1, 134.5, 136.8, 137.0. HRMS (FAB) *m/z* calcd for C<sub>36</sub>H<sub>36</sub>NaO<sub>7</sub> [M+Na]<sup>+</sup>: 603.2359; found 603.2357.

#### 4.28. 1-Allyloxy-2-{3-*O*-[2-cyanoethoxy(diisopropylamino)phosphino]-5-*O*-(4,4'-dimethoxytrityl)-2-*O*,4-*C*-methylene-β-*D*-ribofuranosyl}benzene (**24**)

Under a nitrogen atmosphere, (*i*-Pr<sub>2</sub>N)<sub>2</sub>POCH<sub>2</sub>CH<sub>2</sub>CN (64 μL, 0.20 mmol) was added to a solution of compound **34** (69 mg, 0.12 mmol) and DIHT (24 mg, 0.12 mmol) in anhydrous MeCN/THF (2 mL, 3:1) at room temperature. After being stirred for 4 h, the

mixture was concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=2:1) to give compound **24** (75 mg, 81%) as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.82 (3H, d, *J*=6.6 Hz), 0.90 (3H, d, *J*=6.6 Hz), 1.04–1.08 (6H, m), 2.26–2.31 (1H, m), 2.37–2.42 (1H, m), 3.36–3.55 (5H, m), 3.64–3.71 (1H, m), 3.80 (3H, s), 3.80 (3H, s), 3.88–3.94 (1H, m), 4.05–4.11 (1H, m), 4.21 (0.5H, d, *J*=7.4 Hz), 4.34 (0.5H, d, *J*=9.9 Hz), 4.42 (0.5H, s), 4.50 (0.5H, s), 4.56–4.59 (2H, m), 5.24–5.31 (1H, m), 5.37–5.40 (1.5H, m), 5.43–5.47 (0.5H, m), 5.97–6.12 (1H, m), 6.81–6.99 (6H, m), 7.21–7.34 (4H, m), 7.41–7.46 (4H, m), 7.54–7.57 (2H, m), 7.71–7.73 (0.5H, m), 7.77–7.79 (0.5H, m). <sup>31</sup>P NMR (acetone-*d*<sub>6</sub>) δ 146.9, 147.7. HRMS (FAB) *m/z* calcd for C<sub>45</sub>H<sub>54</sub>N<sub>2</sub>O<sub>8</sub>P [M+H]<sup>+</sup>: 781.3618; found, 781.3638.

#### 4.29. 1-Allyloxy-3-(3,5-di-*O*-benzyl-2-*O*,4-*C*-methylene-β-*D*-ribofuranosyl)benzene (**28**)

Under a nitrogen atmosphere, a solution of compound **26** (1.38 g, 4.03 mmol) in anhydrous THF (20 mL) was added to a solution of 3-allyloxyphenylmagnesium iodide [prepared from 2-allyloxyphenyl iodide (4.19 g, 16.1 mmol) and *i*-PrMgCl·LiCl (1.3 M in THF, 12.4 mL, 16.1 mmol) in anhydrous THF (50 mL) at room temperature], and the mixture was stirred for 15 h. After addition of satd NH<sub>4</sub>Cl aq, the mixture was extracted with AcOEt. The extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=3:1 to 1:1) to give an appropriate compound (*R*<sub>f</sub>=ca. 0.5, *n*-hexane/AcOEt=1:1), which was dissolved in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (30 mL); TMAD (616 mg, 3.58 mmol) and *n*-Bu<sub>3</sub>P (0.89 mL, 3.58 mmol) were added at room temperature and the mixture was stirred for 2 h. After addition of water, the mixture was extracted with Et<sub>2</sub>O. The extracts were washed with water, brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=5:1) to give compound **28** (1.30 g, 70% from **26**) as a colorless oil. [α]<sub>D</sub><sup>20</sup> +7.1 (c 1.45, CHCl<sub>3</sub>). IR ν<sub>max</sub> (KBr) 3030, 2878, 1600, 1449, 1281, 1105, 1031 cm<sup>-1</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 3.86 (2H, s), 4.06 (1H, s), 4.07, 4.10 (2H, AB, *J*=8 Hz), 4.22 (1H, s), 4.43, 4.49 (2H, AB, *J*=12 Hz), 4.62 (2H, s), 4.65 (2H, s), 5.15 (1H, s), 5.25 (1H, d, *J*=11 Hz), 5.36 (1H, d, *J*=17 Hz), 5.92–6.06 (1H, m), 6.79 (1H, d, *J*=8 Hz), 6.85 (1H, d, *J*=8 Hz), 6.92 (1H, s), 7.19–7.35 (11H, m). <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 66.5, 68.6, 72.0, 73.5, 73.6, 77.5, 80.9, 83.7, 85.7, 111.5, 113.7, 117.5, 127.4, 127.5, 127.5, 127.6, 128.2, 128.2, 129.3, 133.1, 137.3, 137.9, 140.7, 158.5. Mass (EI) *m/z* 458 (M<sup>+</sup>, 12.5). Anal. Calcd for C<sub>29</sub>H<sub>30</sub>O<sub>5</sub>: C, 75.96; H, 6.59; found: C, 75.66; H, 6.62.

#### 4.30. 3-(3,5-Di-*O*-benzyl-2-*O*,4-*C*-methylene-β-*D*-ribofuranosyl)-1-hydroxybenzene (**35**)

Under a nitrogen atmosphere, Pd(PPh<sub>3</sub>)<sub>4</sub> (162 mg, 0.14 mmol) was added to a solution of compound **28** (640 mg, 1.40 mmol) in anhydrous THF (150 mL) and the mixture was stirred until solids of Pd(PPh<sub>3</sub>)<sub>4</sub> dissolved completely. Then, NaBH<sub>4</sub> (159 mg, 4.19 mmol) was added and the mixture was stirred at room temperature for 17 h. After addition of 3% HCl aq, the mixture was filtered through a pad of Celite®. The filtrate was extracted with AcOEt. The extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=5:2) to give compound **35** (594 mg, quant.) as a colorless oil. [α]<sub>D</sub><sup>25</sup> +11.7 (c 1.36, CHCl<sub>3</sub>). IR ν<sub>max</sub> (KBr): 3311, 3030, 2879, 1595, 1453, 1362, 1206, 1105, 1032 cm<sup>-1</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 3.85, 3.88 (2H, AB, *J*=11 Hz), 4.08 (1H, s), 4.07, 4.11 (2H, AB, *J*=8 Hz), 4.22 (1H, s), 4.38, 4.49 (2H, AB, *J*=12 Hz), 4.66 (2H, s), 5.12 (1H, s), 5.59 (1H, s), 6.70 (1H, d, *J*=8 Hz), 6.80 (1H, d, *J*=8 Hz), 6.78 (1H, s), 7.13–7.34 (11H, m). <sup>13</sup>C NMR

(CDCl<sub>3</sub>)  $\delta$  66.3, 72.1, 73.5, 73.7, 77.4, 80.9, 83.6, 85.8, 112.2, 114.4, 117.2, 127.5, 127.6, 127.7, 128.2, 128.3, 129.6, 137.3, 137.9, 140.8, 155.9. HRMS (MALDI-TOF)  $m/z$  calcd for C<sub>26</sub>H<sub>26</sub>NaO<sub>5</sub> [M+Na]<sup>+</sup>: 441.1678; found, 441.1677.

#### 4.31. 1-Benzoyloxy-3-(3,5-di-O-benzyl-2-O,4-C-methylene- $\beta$ -D-ribofuranosyl)benzene (36)

Under a nitrogen atmosphere, BzCl (200  $\mu$ L, 1.70 mmol) was added to a solution of compound **35** (594 mg, 1.42 mmol) and Et<sub>3</sub>N (240  $\mu$ L, 1.70 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (15 mL) at room temperature and the mixture was stirred for 17 h. After addition of satd NaHCO<sub>3</sub> aq, the mixture was extracted with AcOEt. The extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=3:1) to give compound **36** (730 mg, 95%) as a colorless oil. [ $\alpha$ ]<sub>D</sub><sup>25</sup> +28.3 (c 1.49, CHCl<sub>3</sub>). IR  $\nu_{\max}$  (KBr) 3423, 3030, 2867, 1599, 1490, 1450, 1260, 1097 cm<sup>-1</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  3.85, 3.87 (2H, AB, *J*=11 Hz), 4.07 (1H, s), 4.09, 4.12 (2H, AB, *J*=8 Hz), 4.25 (1H, s), 4.46, 4.52 (2H, AB, *J*=11 Hz), 4.64 (2H, s), 5.20 (1H, s), 7.11–7.42 (14H, m), 7.48–7.54 (2H, m), 7.62–7.67 (1H, m), 8.19 (2H, d, *J*=7 Hz). <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  68.7, 69.5, 71.9, 73.7, 74.5, 75.2, 81.7, 94.8, 89.1, 113.2, 114.3, 117.5, 119.5, 127.6, 127.8, 128.2, 128.4, 129.3, 133.1, 137.3, 137.4, 142.0, 158.5. HRMS (MALDI-TOF)  $m/z$  calcd for C<sub>33</sub>H<sub>30</sub>NaO<sub>6</sub> [M+Na]<sup>+</sup>: 545.1940; found, 545.1936.

#### 4.32. 1-Benzoyloxy-3-(2-O,4-C-methylene- $\beta$ -D-ribofuranosyl)benzene (37)

A solution of compound **36** (730 mg, 1.40 mmol), 20% Pd(OH)<sub>2</sub>/C (200 mg), and cyclohexene (14 mL, 140 mmol) in EtOH (14 mL) was refluxed for 1 h. The mixture was filtered through a pad of Celite<sup>®</sup> and the filtrate was concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=1:2 to 1:5) to give compound **37** (452 mg, 95%) as a white solid. [ $\alpha$ ]<sub>D</sub><sup>26</sup> –17.4 (c 0.97, CHCl<sub>3</sub>). IR  $\nu_{\max}$  (KBr) 3381, 2947, 1733, 1591, 1487, 1447, 1264, 1146, 1030 cm<sup>-1</sup>. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.69 (1H, br s), 3.05 (1H, br s), 3.95 (2H, s), 3.97, 4.04 (2H, AB, *J*=8 Hz), 4.16 (1H, s), 4.23 (1H, s), 5.13 (1H, s), 7.08–7.18 (3H, m), 7.26 (1H, s), 7.35–7.41 (1H, m), 7.47–7.53 (2H, m), 7.61–7.66 (1H, m), 8.17 (2H, d, *J*=7 Hz). <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  59.0, 71.0, 72.3, 82.9, 83.6, 86.7, 118.6, 120.9, 122.6, 128.5, 129.1, 129.6, 130.1, 133.7, 140.5, 151.0, 165.2. Mass (FAB)  $m/z$  343 (M+H<sup>+</sup>). Anal. Calcd for C<sub>19</sub>H<sub>18</sub>O<sub>6</sub>: C, 66.66; H, 5.30; found: C, 66.55; H, 5.33.

#### 4.33. 1-Benzoyloxy-3-[5-O-(4,4'-dimethoxytrityl)-2-O,4-C-methylene- $\beta$ -D-ribofuranosyl]benzene (38)

Under a nitrogen atmosphere, DMTrCl (175 mg, 0.515 mmol) was added to a solution of compound **37** (147 mg, 0.429 mmol) in anhydrous pyridine (4 mL) at room temperature and the mixture was stirred for 12 h. After addition of satd aq NaHCO<sub>3</sub> solution, the mixture was extracted with AcOEt. The extracts were washed with water, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=3:1 to 1:1) to give compound **38** (252 mg, 91%) as a white solid. [ $\alpha$ ]<sub>D</sub><sup>22</sup> –12.8 (c 0.85, CHCl<sub>3</sub>). IR  $\nu_{\max}$  (KBr): 3478, 3006, 2950, 1736, 1606, 1508, 1447, 1254, 1178, 1072, 1032 cm<sup>-1</sup>. <sup>1</sup>H NMR (acetone-*d*<sub>6</sub>)  $\delta$  2.85 (1H, s), 3.41, 3.55 (2H, AB, *J*=11 Hz), 3.74 (3H, s), 3.75 (3H, s), 3.98, 4.00 (2H, AB, *J*=8 Hz), 4.18 (1H, s), 4.27, 4.41 (2H, AB, *J*=5 Hz), 5.15 (1H, s), 6.84–6.89 (4H, m), 7.11–7.31 (4H, m), 7.38–7.61 (11H, m), 7.68–7.75 (1H, m), 8.13–8.16 (2H, m). <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  55.2, 60.2, 71.7, 72.9, 82.8, 83.3, 85.9, 86.3, 113.1, 118.6, 120.8, 122.7, 126.7, 127.8, 128.0, 128.4, 129.3, 129.4, 129.9, 130.0, 130.1, 133.5, 135.6,

135.6, 141.0, 144.5, 151.0, 158.3, 165.0. HRMS (EI)  $m/z$  calcd for C<sub>40</sub>H<sub>36</sub>O<sub>8</sub> [M]<sup>+</sup>: 644.2410; found: 644.2410.

#### 4.34. 1-Benzoyloxy-3-[3-O-(2-cyanoethoxy(diisopropylamino)phosphino)-5-O-(4,4'-dimethoxytrityl)-2-O,4-C-methylene- $\beta$ -D-ribofuranosyl]benzene (25)

Under a nitrogen atmosphere, *i*-Pr<sub>2</sub>NP(Cl)OCH<sub>2</sub>CH<sub>2</sub>CN (45  $\mu$ L, 0.20 mmol) was added to a solution of compound **38** (32 mg, 0.05 mmol) and *i*-Pr<sub>2</sub>NEt (87  $\mu$ L, 0.50 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (1 mL) at 0 °C and the mixture was stirred at room temperature for 1 h. After addition of satd NaHCO<sub>3</sub> aq, the mixture was extracted with AcOEt. The extracts were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (*n*-hexane/AcOEt=3:1) to give compound **25** (30 mg, 71%) as a white solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  0.88 (3H, d, *J*=6.9 Hz), 0.96 (3H, d, *J*=6.8 Hz), 1.06–1.09 (6H, m), 2.30–2.33 (1H, m), 2.42–2.45 (1H, m), 3.40–3.49 (5H, m), 3.66–3.69 (1H, m), 3.75–3.76 (6H, m), 3.92–3.95 (1H, m), 4.10–4.15 (1H, m), 4.21 (0.5H, d, *J*=7.4 Hz), 4.28 (0.5H, s), 4.35 (0.5H, d, *J*=9.3 Hz), 4.39 (0.5H, s), 5.26 (1H, s), 6.80–6.83 (4H, m), 7.14–7.17 (2H, m), 7.24–7.27 (3H, m), 7.39–7.53 (10H, m), 7.60–7.64 (1H, m), 8.12–8.16 (2H, m). <sup>31</sup>P NMR (CDCl<sub>3</sub>)  $\delta$  148.2, 148.4. HRMS (FAB)  $m/z$  calcd for C<sub>49</sub>H<sub>54</sub>N<sub>2</sub>O<sub>9</sub>P [M+H]<sup>+</sup>: 845.3567; found, 845.3573.

#### 4.35. Synthesis of TFOs

The 0.2  $\mu$ mol scale synthesis of TFOs **39–43** was performed on an automated DNA synthesizer (Applied Biosystems Expedite<sup>™</sup> 8909 or Genedesign nS-8) using a standard phosphoramidite protocol. By treatment with 28% NH<sub>3</sub> aq at room temperature for 4 h, TFOs synthesized on DMTr-ON mode were cleaved from the CPG resin and all protecting groups on TFOs were removed. The crude TFOs **39–41** and **43** obtained were purified with NENSORB<sup>™</sup> PREP or Sep-Pak<sup>®</sup> Plus C18 cartridges followed by reversed-phase HPLC (ChemcoPak<sup>®</sup> CHEMCOSORB 300-5C18, 4.6 mm $\times$ 250 mm or Waters XBridge<sup>®</sup> MS C<sub>18</sub> 2.5  $\mu$ m, 10 mm $\times$ 50 mm). For TFO **42**, before reversed-phase HPLC purification, treatment with NaBH<sub>4</sub> (2.2 mg, 58.2  $\mu$ mol) and Pd(PPh<sub>3</sub>)<sub>4</sub> (2.1 mg, 1.9  $\mu$ mol) in H<sub>2</sub>O (200  $\mu$ L) at room temperature for 3 h was performed to remove an allyl group. The composition of the TFOs was confirmed by MALDI-TOF-MS analysis. MALDI-TOF-MS data ([M–H]<sup>–</sup>) for TFOs **39–43**; **39**, found 4464.44 (calcd 4463.98); **40**, found 4463.89 (calcd 4463.98); **41**, found 4464.43 (calcd 4463.98); **42**, found 4491.60 (calcd 4492.00); **43**, found 4492.34 (calcd 4492.00).

#### 4.36. UV melting experiments

UV melting experiments were performed on SHIMADZU UV-1650 and SHIMADZU UV-1800 spectrophotometers equipped with *T*<sub>m</sub> analysis accessory. The TFOs **39–43** and hairpin dsDNA targets were dissolved in 10 mM sodium cacodylate buffer (pH 6.8) containing 140 mM KCl and 50 mM MgCl<sub>2</sub> to give a final concentration of each strand of 1.89  $\mu$ M. The samples were annealed in boiling water followed by slow cooling to 5 °C. The melting profiles were recorded at 260 nm from 5 °C to 90 °C at a scan rate of 0.5 °C/min. The two-point average method was used to obtain the *T*<sub>m</sub> values and the final values were determined by averaging three independent measurements, which were accurate to within 1 °C.

#### Acknowledgements

This work was supported in part by the Program to Disseminate Tenure Tracking System of the Ministry of Education, Culture, Sports, Science and Technology, Japan (MEXT), a Grant-in-Aid for

Scientific Research on Innovative Areas (No. 22136006) from MEXT, and the Mochida Memorial Foundation for Medical and Pharmaceutical Research.

## References and notes

- Reviews: (a) Praseuth, D.; Guieysse, A. L.; Hélène, C. *Biochim. Biophys. Acta, Gene Struct. Expression* **1999**, *1489*, 181–206; (b) Seidman, M. M.; Glazer, P. M. *J. Clin. Invest.* **2003**, *112*, 487–494; (c) Chin, J. Y.; Schleifman, E. B.; Glazer, P. M. *Front. Biosci.* **2007**, *12*, 4288–4297; (d) Simon, P.; Cannata, F.; Concordet, J.-P.; Giovannangeli, C. *Biochimie* **2008**, *90*, 1109–1116; (e) Duca, M.; Vekhoff, P.; Oussedik, K.; Halby, L.; Arimondo, P. B. *Nucleic Acids Res.* **2008**, *36*, 5123–5138; (f) Chin, J. Y.; Glazer, P. M. *Mol. Carcinog.* **2009**, *48*, 389–399; (g) Mukherjee, A.; Vasquez, K. M. *Biochimie* **2011**, *93*, 1197–1208.
- Reviews: (a) Thuong, N. T.; Hélène, C. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 666–690; (b) Doronina, S. O.; Behr, J.-P. *Chem. Soc. Rev.* **1997**, *26*, 63–71; (c) Gowers, D. M.; Fox, K. R. *Nucleic Acids Res.* **1999**, *27*, 1569–1577; (d) Fox, K. R. *Curr. Med. Chem.* **2000**, *7*, 17–37; (e) Robles, J.; Grandas, A.; Pedrosa, E.; Luque, F. J.; Eritija, R.; Orzco, M. *Curr. Org. Chem.* **2002**, *6*, 1333–1368; (f) Purwanto, M. G. M.; Weisz, K. *Curr. Org. Chem.* **2003**, *7*, 427–446; (g) Buchini, S.; Leumann, C. J. *Curr. Opin. Chem. Biol.* **2003**, *7*, 717–726; (h) Malnuit, V.; Duca, M.; Benhida, R. *Org. Biomol. Chem.* **2011**, *9*, 326–336; (i) Hari, Y.; Obika, S.; Imanishi, T. *Eur. J. Org. Chem.* **2012**, 2875–2887.
- (a) Yoon, K.; Hobbs, C. A.; Koch, J.; Sardaro, M.; Kutny, R.; Weis, A. L. *Proc. Natl. Acad. Sci. U.S.A.* **1992**, *89*, 3840–3844; (b) Radhakrishnan, I.; Patel, D. J. *J. Mol. Biol.* **1994**, *241*, 600–619.
- Obika, S.; Hari, Y.; Sekiguchi, M.; Imanishi, T. *Angew. Chem., Int. Ed.* **2001**, *40*, 2079–2081.
- A part of the results was reported at the International Symposium on Nucleic Acids Chemistry: (a) Obika, S.; Hari, Y.; Inohara, H.; Imanishi, T. *Nucleic Acids Res. Suppl.* **2001**, 171–172; (b) Inohara, H.; Obika, S.; Imanishi, T. *Nucleic Acids Symp. Ser.* **2004**, 63–64.
- Wierenga, W.; Skulnick, H. I. *Carbohydr. Res.* **1981**, *90*, 41–52.
- Tsunoda, T.; Otsuka, J.; Yamamiya, Y.; Ito, S. *Chem. Lett.* **1994**, 539–542.
- (a) Obika, S.; Hari, Y.; Morio, K.; Imanishi, T. *Tetrahedron Lett.* **2000**, *41*, 215–219; (b) Hari, Y.; Obika, S.; Sakaki, M.; Morio, K.; Yamagata, Y.; Imanishi, T. *Tetrahedron* **2002**, *58*, 3051–3063.
- A significant increase in the  $T_m$  value of the triplex with dsDNA (YZ=CG) was not observed by replacement of 2H with 2H<sup>B</sup>. This may imply that the 2-hydroxyphenyl nucleobase in 2H<sup>B</sup> worked only as a hydrogen-donor by the 2',4'-BNA modification though the reason is unclear.
- (a) Hari, Y.; Nakahara, M.; Pang, J.; Akabane, M.; Kuboyama, T.; Obika, S. *Bioorg. Med. Chem.* **2011**, *19*, 1162–1166; (b) Hari, Y.; Akabane, M.; Hatanaka, Y.; Obika, S. *Chem. Commun.* **2011**, 4424–4426.