

### Ⅲ. 研究成果の刊行に関する一覧表

研究成果の刊行に関する一覧表レイアウト

書籍

著者氏名	論文タイトル名	書籍全体の編集者名	書籍名	出版社名	出版地	出版年	ページ
Shima, F., Yoshikawa, Y., Matsumoto, S., and Kataoka, T.	Discovery of small-molecule Ras inhibitors that display antitumor activity and Kataby interfering with Ras•GTP-effector interaction.	Fuyuhiko Tamano, Chaning J. Der	The Enzymes	Academic Press	Burlington	2013	1-23

別刷2

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Shima, F., Yoshikawa, Y., Ye, M., Araki, M., Matsumoto, S., Liao, J., Hu, L., Sugimoto, T., Ijiri, Y., Takeda, A., Nishiyama, Y., Sato, C., Muraoka, S., Tamura, A., Osoda, T., Tsuda, K., Miyakawa, T., Fukunishi, H., Shimada, J., Kumazaki, T., Yamamoto, M., and Kataoka, T.	<i>In silico</i> discovery of novel small-molecule Ras inhibitors that display antitumor activity by blocking the Ras-effector interaction.	Proc. Natl. Acad. Sci. USA.	110 (20)	8182-8187	2013
島 扶美, 熊坂 崇, 松本 篤幸, 吉川 陽子, 山本 雅貴, 片岡 徹	<i>ras</i> がん遺伝子産物の新規立体構造情報を利用した分子標的がん治療薬の開発	日本放射光学会誌 放射光	27 (1)	3-9	2014
Shima, F., and Kataoka, T.	Structure-based drug design of small-molecule Ras inhibitors having antitumor activity.	Spring-8 Research Frontiers		in press.	2013

別刷1

別刷3

## IV. 研究成果の刊行物・別刷

# *In silico* discovery of small-molecule Ras inhibitors that display antitumor activity by blocking the Ras–effector interaction

Fumi Shima<sup>a,1,2</sup>, Yoko Yoshikawa<sup>a,1</sup>, Min Ye<sup>a</sup>, Mitsugu Araki<sup>a,b</sup>, Shigeyuki Matsumoto<sup>a</sup>, Jingling Liao<sup>a</sup>, Lizhi Hu<sup>a</sup>, Takeshi Sugimoto<sup>a</sup>, Yuichi Ijiri<sup>a</sup>, Azusa Takeda<sup>a</sup>, Yuko Nishiyama<sup>a</sup>, Chie Sato<sup>a</sup>, Shin Muraoka<sup>a</sup>, Atsuo Tamura<sup>b</sup>, Tsutomu Osoda<sup>c</sup>, Ken-ichiro Tsuda<sup>d</sup>, Tomoya Miyakawa<sup>c</sup>, Hiroaki Fukunishi<sup>e</sup>, Jiro Shimada<sup>e</sup>, Takashi Kumasaka<sup>f</sup>, Masaki Yamamoto<sup>g</sup>, and Tohru Kataoka<sup>a,2</sup>

<sup>a</sup>Division of Molecular Biology, Department of Biochemistry and Molecular Biology, Kobe University Graduate School of Medicine, 7-5-1 Kusunoki-cho, Chuo-ku, Kobe 650-0017, Japan; <sup>b</sup>Department of Chemistry, Kobe University Graduate School of Science, 1-1 Rokkodai, Nada-ku, Kobe 657-8501, Japan; <sup>c</sup>Business Innovation Center and <sup>d</sup>Intellectual Asset Research and Development Unit and Planning Division, NEC Corporation, 1753 Shimonumabe, Nakahara-ku, Kawasaki 211-8666, Japan; <sup>e</sup>Green Innovation Research Laboratories, NEC Corporation, Miyukigaoka 34, Tsukuba 305-8501, Japan; and <sup>f</sup>Japan Synchrotron Radiation Research Institute and <sup>g</sup>RIKEN SPring-8 Center, 1-1-1 Kouto, Sayo-cho, Sayo-gun, Hyogo 679-5198, Japan

Edited by Michelle R. Arkin, University of California, San Francisco, CA, and accepted by the Editorial Board April 2, 2013 (received for review October 16, 2012)

**Mutational activation of the Ras oncogene products (H-Ras, K-Ras, and N-Ras) is frequently observed in human cancers, making them promising anticancer drug targets. Nonetheless, no effective strategy has been available for the development of Ras inhibitors, partly owing to the absence of well-defined surface pockets suitable for drug binding. Only recently, such pockets have been found in the crystal structures of a unique conformation of Ras-GTP. Here we report the successful development of small-molecule Ras inhibitors by an *in silico* screen targeting a pocket found in the crystal structure of M-Ras-GTP carrying an H-Ras-type substitution P40D. The selected compound Kobe0065 and its analog Kobe2602 exhibit inhibitory activity toward H-Ras-GTP-c-Raf-1 binding both *in vivo* and *in vitro*. They effectively inhibit both anchorage-dependent and -independent growth and induce apoptosis of H-*ras*<sup>G12V</sup>-transformed NIH 3T3 cells, which is accompanied by down-regulation of downstream molecules such as MEK/ERK, Akt, and RalA as well as an upstream molecule, Son of sevenless. Moreover, they exhibit antitumor activity on a xenograft of human colon carcinoma SW480 cells carrying the *K-ras*<sup>G12V</sup> gene by oral administration. The NMR structure of a complex of the compound with H-Ras-GTP<sup>T35S</sup>, exclusively adopting the unique conformation, confirms its insertion into one of the surface pockets and provides a molecular basis for binding inhibition toward multiple Ras-GTP-interacting molecules. This study proves the effectiveness of our strategy for structure-based drug design to target Ras-GTP, and the resulting Kobe0065-family compounds may serve as a scaffold for the development of Ras inhibitors with higher potency and specificity.**

molecular targeted therapy | small-molecule inhibitor

**R**as oncoproteins belong to the Ras family of small GTPases and function as molecular switches by cycling between GTP-bound active and GDP-bound inactive forms in intracellular signaling pathways controlling cell growth, differentiation, and apoptosis (1). Interconversion between the two forms, which mainly involves the conformational changes of two flexible regions called switch I (residues 32–38) and switch II (residues 60–75), is reciprocally catalyzed by guanine nucleotide exchange factors (GEFs) and GTPase-activating proteins (GAPs) (2). In particular, GEFs such as Son of sevenless (Sos) mediate upstream signals to enhance formation of the GTP-bound form. The oncogenic potential of Ras is activated by point mutations mainly involving the codons 12 and 61, which impair the intrinsic GTPase activity and, moreover, render Ras insensitive to the GAP action, leading to constitutive activation of downstream effectors such as Raf kinases including c-Raf-1 and B-Raf, PI3Ks, and Ral guanine nucleotide dissociation stimulator (RalGDS) family proteins (1). These

mutations are observed in about 15–20% of human cancers, and specifically in about 60–90% and 30–50% of pancreatic and colorectal carcinomas, respectively (1, 3, 4). Cancer cells carrying the *ras* oncogene are known to exhibit a phenomenon called oncogene addiction, where their survival becomes dependent on the activated oncogene function (3). Consequently, inhibition of the activated Ras function has been shown to lead not only to reversal of the transformed phenotypes but also to cell death and tumor regression (4, 5). Despite their importance as an anticancer drug target, there is no effective molecular targeted therapy for Ras at present; the once highly anticipated farnesyl transferase inhibitors, which inhibit the posttranslational lipid modification, farnesylation, of Ras necessary for membrane targeting, have failed in clinical trials (1, 6). Although farnesylthiosalicylic acid has been reported to inhibit Ras by antagonizing its interaction with the Ras-escort proteins, its antitumor activity remains unclear (7).

Although recent success in drug discovery using structure-based drug design (SBDD) for AIDS and influenza has boosted hopes for the application of SBDD to anticancer drug development, Ras have been presumed refractory to this approach because they lack apparently “druggable” pockets on their surface, as seen from their crystal structures (1). Recently, by X-ray crystallography and NMR spectroscopy we solved the tertiary structures of H-Ras, its homolog M-Ras, and their mutants in complex with a non-hydrolyzable GTP analog, guanosine 5′-(β,γ-imido)triphosphate (GppNHp), all of which corresponded to a unique conformation (8–10) undergoing dynamic equilibrium with the previously known conformation. Intriguingly, the structures possessed surface pockets that seem suitable for drug binding. In this paper, we have applied SBDD to target Ras-GTP by using the structural information on these surface pockets. We report the successful discovery of a unique class of small-molecule compounds that have

Author contributions: F.S. and T. Kataoka designed research; F.S., Y.Y., M. Ye, M.A., S. Matsumoto, J.L., L.H., T.S., Y.I., A. Takeda, Y.N., C.S., S. Muraoka, T.O., K.-i.T., T.M., H.F., and J.S. performed research; F.S., Y.Y., M.A., S. Matsumoto, S. Muraoka, A. Tamura, T.M., T. Kumasaka, and M. Yamamoto analyzed data; and F.S. and T. Kataoka wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. M.R.A. is a guest editor invited by the Editorial Board.

Data deposition: NMR, atomic coordinates, chemical shifts, and restraints have been deposited in the Protein Data Bank, [www.pdb.org](http://www.pdb.org) (PDB ID code 2lwi).

<sup>1</sup>F.S. and Y.Y. contributed equally to this work.

<sup>2</sup>To whom correspondence may be addressed. E-mail: [kataoka@people.kobe-u.ac.jp](mailto:kataoka@people.kobe-u.ac.jp) or [sfumi@med.kobe-u.ac.jp](mailto:sfumi@med.kobe-u.ac.jp).

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1217730110/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1217730110/-DCSupplemental).


potent activity to block the interactions of Ras-GTP with their multiple effector molecules and, moreover, display antitumor activity on a xenograft of human colon carcinoma cells carrying the *K-ras<sup>G12V</sup>* gene.

## Results

### Discovery of Small-Molecule Compounds Inhibiting Ras–Raf Interaction by SBDD.

Aiming to discover small-molecule compounds fitting into the surface pockets of the unique conformation of Ras-GTP, we applied the molecular mechanics Poisson–Boltzman surface area (MMPB-SA) method with an Assisted Model Building and Energy Refinement (AMBER)96 force field to carry out a computer docking screen of a virtual library containing 40,882 compounds based on the high-resolution (1.35 Å) crystal structure of M-Ras<sup>P40D</sup>.GppNHp (9). Ninety-seven candidates were selected and examined in vitro for their activity to inhibit the binding of

M-Ras<sup>P40D</sup>.GTP and H-Ras-GTP to the Ras-binding domain (RBD, amino acids 50–131) of c-Raf-1. Only one compound, named Kobe0065 (Fig. 1A), exhibited potent activity to competitively inhibit the binding of H-Ras-GTP to c-Raf-1 RBD with a  $K_i$  value of  $46 \pm 13 \mu\text{M}$  as estimated from the binding kinetics (Fig. S1). A subsequent computer-assisted similarity search of ~160,000 compounds based on the Tanimoto coefficient selected 273 candidates, among which one positive, named Kobe2602 (Fig. 1A), with a  $K_i$  value of  $149 \pm 55 \mu\text{M}$  (Fig. S1), was identified. These two compounds, added to the culture medium at 2 and 20  $\mu\text{M}$ , effectively reduced the amount of c-Raf-1 associated with H-Ras<sup>G12V</sup> in NIH 3T3 cells in a dose-dependent manner, indicating the inhibition of the cellular activity of Ras (Fig. 1B). A rough estimate of the  $\text{IC}_{50}$  value for the cellular Ras–Raf-binding inhibition was around 10  $\mu\text{M}$  (Fig. 1B), which was not much different from the  $K_i$  values for the in vitro Ras–Raf-binding inhibition considering the



**Fig. 1.** Inhibition of various downstream targets of Ras by the Kobe0065-family compounds. (A) Chemical structures of the compounds. (B) NIH 3T3 cells were transfected with pEF-BOS-HA-H-Ras<sup>G12V</sup> or an empty vector and treated with the 2 and 20  $\mu\text{M}$  compound or the vehicle (DMSO) in the presence of 2% FBS for 1 h. Cell lysate was subjected to detection of c-Raf-1 coimmunoprecipitated with an anti-H-Ras antibody (Top) and total c-Raf-1 (Middle) by Western blotting with an anti-c-Raf-1 antibody. Immunoprecipitated H-Ras<sup>G12V</sup> were detected by an anti-HA antibody (Bottom). The numbers above the lanes show the values of H-Ras-bound/total c-Raf-1 relative to that of the vehicle-treated cells. (C) Lysate was prepared from cells treated with 20  $\mu\text{M}$  Kobe0065, 20  $\mu\text{M}$  Kobe2602, or 2  $\mu\text{M}$  sorafenib as described in B and subjected to detection of phosphorylated MEK (pMEK) and ERK (pERK) by Western blotting with anti-pMEK and anti-pERK antibodies. Total amounts of MEK, ERK, and HA-tagged H-Ras<sup>G12V</sup> were detected by anti-MEK, anti-ERK, and anti-HA antibodies, respectively. The numbers above the lanes show the values of pMEK/tMEK and pERK/tERK relative to those of the vehicle-treated cells. Four independent experiments yielded essentially equivalent results. (D) Recombinant c-Raf-1 was incubated with recombinant MEK in the presence of 20  $\mu\text{M}$  Kobe0065, 20  $\mu\text{M}$  Kobe2602, or 2  $\mu\text{M}$  sorafenib, and pMEK formed was detected by Western blotting with an anti-pMEK antibody. (E) Lysate was prepared from cells treated with the indicated concentrations of the compound as described in B and subjected to the measurements of phosphorylated Akt (pAKT) by Western blotting with an anti-pAkt antibody and of RalA-GTP pulled down with GST-Sec5(1–99) immobilized on glutathione-sepharose resin by Western blotting with an anti-RalA antibody. Four independent experiments yielded essentially equivalent results.


quite low cellular concentration of Raf. A similar inhibitory effect was also observed with NIH 3T3 cells overexpressing K-Ras<sup>G12V</sup> (Fig. S2).

**Inhibitory Effects of the Kobe0065-Family Compounds on Various Ras Signaling Pathways.** We examined the effect of the compounds on the cellular activity of Raf, which is dependent on its interaction with Ras-GTP. Both Kobe0065 and Kobe2602 at 20  $\mu$ M efficiently inhibited the phosphorylation of MEK and ERK, downstream kinases of Raf in NIH 3T3 cells transiently expressing H-Ras<sup>G12V</sup>, although the effect was slightly weaker than that of 2  $\mu$ M sorafenib (11), an inhibitor of multiple protein kinases including Raf (Fig. 1C). However, they failed to inhibit the kinase activity of c-Raf-1 measured in vitro (Fig. 1D), indicating the absence of direct inhibitory activity on Raf. Furthermore, the compound-treated cells showed substantial decreases of phosphorylated Akt and RalA-GTP, downstream molecules of PI3Ks and RalGDS, respectively, in a manner dependent on the compound concentrations (Fig. 1E), suggesting that the compounds exerted inhibitory effects toward multiple Ras effectors through inactivation of Ras.

We next examined the effect of the compounds on Sos. Sos has two distinct Ras-binding sites: the GEF domain catalyzing the GDP-GTP exchange on Ras through interaction with Ras-GDP and the distal site allosterically accelerating the GEF catalytic activity through interaction with Ras-GTP (12), and thereby functions not only as a regulator but also as an effector of Ras. In vitro GDP-GTP exchange assays using mSos1 and mSos1<sup>W729E</sup>, carrying an inactivating mutation of the distal site (12), showed that Kobe0065 at 50  $\mu$ M almost completely abolished the accelerating effect of the distal site without apparently affecting the catalytic activity of the GEF domain itself (Fig. 2A and Fig. S3A), suggesting that the compounds inhibited Ras-GTP but not Ras-GDP. The IC<sub>50</sub> value of Kobe0065 was estimated to be around 20  $\mu$ M (Fig. S3B). Kobe2602 also showed the same but weaker activity with an IC<sub>50</sub> value of around 100  $\mu$ M (Fig. S3C). This finding raised a possibility that the observed in vivo inhibition of H-Ras<sup>G12V</sup> by the compounds (Fig. 1C and E) might be accounted for by the decreased Ras-GTP level owing to Sos inhibition. However, we found that this was not the case, because the cellular Ras<sup>G12V</sup>-GTP level was almost unaffected by either overexpression or siRNA-mediated knockdown of mSos1 in either NIH 3T3 cells transiently expressing H-Ras<sup>G12V</sup> or human colon carcinoma SW480 cells carrying *K-ras*<sup>G12V</sup> (Fig. 2C and D).

**Inhibitory Effects of the Kobe0065-Family Compounds on the Growth of Cancer Cell Culture and Tumor Xenograft.** We next tested the effect of Kobe0065 and Kobe2602 on anchorage-independent proliferation of H-ras<sup>G12V</sup>-transformed NIH 3T3 cells. The compounds efficiently inhibited colony formation in soft agar in a dose-dependent manner (Fig. 3A and Fig. S3A). The IC<sub>50</sub> values were around 0.5 and 1.4  $\mu$ M for Kobe0065 (Fig. 3B) and Kobe2602 (Fig. S3B), respectively, which were comparable to the value of 2.1  $\mu$ M observed for sorafenib (Fig. S3B). By contrast, the compounds failed to inhibit colony formation of NIH 3T3 cells transformed by the activated *c-raf-1* gene carrying the S259A/Y340D/Y341D mutations (Fig. 3C and Fig. S4A), whereas sorafenib exhibited potent inhibitory activity.

We then assessed the *ras* specificity of inhibition by using several cancer cell lines carrying various oncogenes. The compounds effectively inhibited the colony formation of cancer cells carrying the activated *ras* oncogenes, such as SW480 and PANC-1 (K-ras<sup>G12V</sup>), EJ-1 (H-ras<sup>G12V</sup>), HT1080 (*N-ras*<sup>Q61L</sup>), and DLD-1 and HCT116 (H-ras<sup>G13D</sup>), but showed much weaker inhibition on those without the *ras* mutation, such as A375, T-47D, LNCap, BxPC-3, MCF-7, HepG2, and HeLa (Table S1). DLD-1 and HCT116 were sensitive to the compounds even though they carried additional activating mutations in PI3K, suggesting that




**Fig. 2.** Inhibition of Sos by the Kobe0065-family compounds and effect of the Sos activity on the cellular Ras<sup>G12V</sup>-GTP level. (A) GST-H-Ras(1-166)-GDP immobilized on glutathione-sepharose resin were incubated with [ $\gamma$ -<sup>35</sup>S]GTPγS and purified 6×His-tagged mSos1(563–1,049), wild-type, or a W729E mutant at 25 °C in the presence or absence of 50  $\mu$ M Kobe0065. The radioactivity pulled down by glutathione-sepharose resin was measured. Three independent experiments yielded essentially equivalent results. (B) NIH 3T3 cells were transfected with pEF-BOS-HA-H-Ras<sup>G12V</sup> in combination with pCMV-mSos1 or siRNA against mSos1. H-Ras<sup>G12V</sup>-GTP pulled down by GST-c-Raf-1-RBD from the cell lysate was detected by an anti-HA antibody (Upper). Total amounts of HA-H-Ras<sup>G12V</sup> in the lysates was also measured (Lower). (C) SW480 cells were transfected with pCMV-mSos1. K-Ras<sup>G12V</sup>-GTP pulled down by GST-c-Raf-1-RBD from cell lysates was detected by an anti-K-Ras antibody (Upper). Total amounts of K-Ras<sup>G12V</sup> in the lysates were also measured (Lower).

the activated PI3K alone might be insufficient to sustain their anchorage independence. We next examined the effect of the compounds on anchorage-dependent proliferation (Fig. 3D). The compounds at 20  $\mu$ M almost completely inhibited the proliferation of H-ras<sup>G12V</sup>-transformed NIH 3T3 cells in the presence of 2% FBS. The IC<sub>50</sub> values were ~1.5 and 2  $\mu$ M for Kobe0065 and Kobe2602, respectively, which were a bit higher than that (0.8  $\mu$ M) for sorafenib (Fig. S4C). The compound-treated cells exhibited frequent apoptosis (Fig. 3E), suggesting a contribution of the oncogene addiction mechanism to the antiproliferative effect.

We next assessed the antitumor activity of the compounds by using a xenograft of SW480 cells in nude mice. Daily oral administration of the compounds at the dose of 80 mg/kg caused ~40–50% inhibition of the tumor growth, which was weaker than the 65% inhibition by sorafenib (Fig. 4A). By doubling the dose to 160 mg/kg, the activity of Kobe0065 became more evident. During these compound treatments the mice did not exhibit any significant body weight loss (Fig. S5). Immunostaining of the tumor sections showed that the ERK activation was substantially compromised by the compound administration (Fig. 4B). Moreover, the compound-treated tumors showed a prominent increase of apoptotic cells (Fig. S6A), suggesting a contribution of the oncogene addiction mechanism to the antitumor effect. In contrast to the case of sorafenib, an antiangiogenesis effect was not observed (Fig. S6B).

**Molecular Basis for Interaction of Ras-GTP with the Kobe0065-Family Compounds.** We used NMR spectroscopy to obtain structural information on the compound-binding interface on Ras-GTP.




**Fig. 3.** Inhibition of proliferation of H-*ras*<sup>G12V</sup>-transformed cells by the Kobe0065-family compounds. (A) H-*ras*<sup>G12V</sup>-transformed NIH 3T3 cells ( $1 \times 10^3$  cells) were inoculated in 2 mL of DMEM containing 10% FBS, 0.33% SeaPlaque agarose, and the indicated concentrations of the compound. After incubation at 37 °C for 14 d, the number of colonies >200  $\mu$ m in diameter was counted under a dissecting microscope. (B) The IC<sub>50</sub> value for Kobe0065 was estimated from the dose–response curve. (C) Effects of the 20  $\mu$ M compounds on soft agar colony formation of *c-raf-1*<sup>S259A/Y340D/Y341D</sup>-transformed NIH 3T3 cells were measured similarly as described in A. (D) H-*ras*<sup>G12V</sup>-transformed NIH 3T3 cells were cultured under a low-serum condition (2% FBS) in the presence of the 20  $\mu$ M compound. Each point represents the cell number relative to that of the 0-h treatment. The values are presented as the mean  $\pm$  SEM (A,  $n = 4$ ; B,  $n = 7$ ; and C,  $n = 3$ ). All experiments were performed in duplicate. One-way ANOVA with Dunnett’s test was used for the statistical analyses. \* $P < 0.001$ . (E) H-*ras*<sup>G12V</sup>-transformed NIH 3T3 cells cultured in the presence of the 20  $\mu$ M compound for 24 h in 2% FBS were subjected to staining with DAPI (Upper) and the TUNEL assay for detection of apoptotic cells (Lower). A representative image is shown for each group.


The NMR structure corresponding to the unique conformation of H-Ras-GppNHp was determined by using only its T35S mutant (13), because this mutation almost eliminated the slow conformational exchange process (14), which made NMR analysis of the wild-type protein impractical. Because of the low water solubility of Kobe0065 and Kobe2602, which made measurements of the NOEs impossible, we chose to use a water-soluble analog named Kobe2601 (Fig. 5A), which had also been identified by the similarity search of Kobe0065. Kobe2601 showed weak inhibitory activity toward in vitro Ras–Raf binding with a  $K_i$  value of  $773 \pm 49 \mu$ M (Fig. S1). NOEs between the benzene rings of Kobe2601 and the side chains of H-Ras<sup>T35S</sup>-GppNHp were detected and the collected data were used for calculation of the tertiary structure of the H-Ras<sup>T35S</sup>-GppNHp–Kobe2601 complex (Tables S2 and S3). The fluorobenzene moiety of Kobe2601 was located in close

proximity to the side chains of Lys5, Leu56, Met67, Gln70, Tyr71, and Thr74 of H-Ras (Fig. 5A and B). These six residues formed a hydrophobic surface pocket in the neighborhood of switch I (Fig. S7A), indicating that the fluorobenzene ring was inserted into the pocket through hydrophobic interaction. However, the dinitrobenzene moiety of Kobe2601 was located near switch II but not tightly fixed. Although direct assignment of the Kobe2601-interacting residues on wild-type H-Ras was difficult, measurement of the backbone amide <sup>1</sup>H, <sup>15</sup>N heteronuclear single quantum coherence (HSQC) spectra of H-Ras-GppNHp revealed that the resonances from Leu56, Met67, and their neighboring residues underwent significant chemical shift changes and line broadening by the addition of Kobe2601 (Fig. 5C and Fig. S8), suggesting sharing of a common binding pocket with H-Ras<sup>T35S</sup>-GppNHp.

Superimposition of the NMR structure of the H-Ras<sup>T35S</sup>-Kobe2601 complex with the crystal structures of various Ras–effector complexes (15–17) revealed that Kobe2601 overlapped with the effector-binding interfaces (Fig. S7B–E). As for c-Raf-1



**Fig. 4.** Anti-proliferative activity of the Kobe0065-family compounds on a tumor xenograft. (A) Female athymic nude mice were implanted with SW480 cells ( $5 \times 10^6$  cells) in their right flanks. When the tumor sizes reached  $52 \pm 3 \text{ mm}^3$ , the compounds were administered orally for five consecutive days per week for 17–20 d at the indicated doses and the tumor volumes were continuously monitored. The values are presented as the mean  $\pm$  SEM;  $n = 8$ –10 per group.  $P = 0.086$  ( $t$  test) for 80 mg/kg Kobe0065,  $P < 0.05$  for 160 mg/kg Kobe0065 and 80 mg/kg Kobe2602, and  $P < 0.01$  for 80 mg/kg sorafenib at day 17. One-way ANOVA with Tukey’s test was used to analyze the significance of tumor size changes compared with the vehicle-treated group. (B) Phosphorylated ERK was detected by immunohistochemistry with an anti-pERK antibody in sections of tumors, which were treated daily with the 80 mg/kg compound for 17 d. The percentage of pERK-positive cells is shown on the top of each panel. A representative image is shown for each group.



**Fig. 5.** Molecular basis for the interaction of Ras-GTP with the Kobe0065-family compounds. (A) The lowest energy solution structure of the H-Ras<sup>T355</sup>.GppNHp-Kobe2601 complex. H-Ras<sup>T355</sup>.GppNHp is shown by a surface model (switch I, yellow; switch II, green) and Kobe2601 is shown by a space-filling model (C, black; O, red; N, blue; H, gray; S, yellow; and F, orange). (B) Spatial arrangements of the residues giving NOE contacts with Kobe2601. Stick representations of the residues giving intermolecular NOEs (red), Kobe2601 (cyan), and GppNHp (magenta) are shown on the backbone structure of H-Ras<sup>T355</sup>.GppNHp. (C) The residues that exhibited chemical shift perturbation and line broadening in the presence of Kobe2601 (Fig. S5) are shown on the crystal structure of H-Ras-GppNHp (PDB ID code 5P21). Moderately perturbed residues with  $0.01 \leq \Delta\delta < 0.015$  and  $I/I_0 \leq 0.8$ , orange; strongly perturbed residues with  $\Delta\delta \geq 0.015$  and  $I/I_0 \leq 0.8$ , red; missing residues or residues exhibiting resonance overlaps, cyan. The models were generated using MOLMOL (27) and PyMOL (DeLano Scientific, LLC).

RBD (15), both the fluorene and nitrobenzene moieties of Kobe2601 were likely to cause steric hindrance with its surface residues (Fig. S7B), supporting our observation of the competitive inhibition by Kobe0065 and Kobe2602. Moreover, a major part of Kobe2601, including the thiosemicarbazide and nitrobenzene moieties, was predicted to interfere with PI3K (16) much more heavily than with c-Raf-1 RBD (Fig. S7C), which may account for the efficient inhibition of Akt phosphorylation (Fig. 1E). Likewise, Kobe2601 was predicted to interfere with the Ras-interacting domain of RalGDS (17) (Fig. S7D) and also more heavily with the distal site of hSos (11) (Fig. S7E), which was experimentally demonstrated in this work (Figs. 1E and 2A).

Because the residues forming the compound-binding pocket are well conserved among Ras family members (Fig. S9), the Kobe0065-family compounds were predicted to exhibit rather broad specificity. This was indeed the case when various small GTPases in their GppNHp-bound forms were tested for direct interaction with Kobe0065 and Kobe2602 by relaxation-edited 1D <sup>1</sup>H NMR (18) (Fig. S10). The compounds bound efficiently to M-Ras, Rap2A, and RalA but weakly to Rap1A compared with H-Ras. As for Rho family small GTPases, both Kobe0065 and Kobe2602 showed very weak, if any, binding activity toward Cdc42 and Rac1, whereas Kobe0065, but not Kobe2602, seemed to have some binding activity toward RhoA. Also, we found that both Kobe0065 and Kobe2602 bound to H-Ras-GDP as well in the 1D <sup>1</sup>H NMR analysis. This was rather unexpected considering no apparent inhibitory effect of the compounds on the intrinsic GEF catalytic activity of Sos (Fig. 2A and Fig. S3A). Interpretation of

the significance of this result on the mode of action of the Kobe0065-family compounds will require further structural information on their actual binding site on H-Ras-GDP, which is totally lacking at present.

## Discussion

Recently, Maurer et al. (19) reported discovery of small-molecule compounds that bound to K-Ras-GDP and inhibited the Sos-mediated nucleotide exchange both in vitro and in vivo. Their crystal structure analyses of the complexes of the compounds, benzimidazole (BZIM), benzamidine (BZDN), and 4,6-dichloro-2-methyl-3-aminoethyl-indole (DCAI), with K-Ras-GDP, K-Ras in complex with guanosine-5'-[γ-thio]triphosphate (GTPγS), and K-Ras in complex with guanosine-5'-[(β,γ)-methylene]triphosphate, respectively, provided a molecular basis for inhibition of the Ras-GDP-Sos interaction but not the K-Ras-GTP-effector interaction; the compounds apparently interfered with the binding of K-Ras to Sos but not any effectors. In a sharp contrast, our compounds exhibited a prominent inhibitory activity at both the biochemical and cellular levels toward H-Ras-GTP and K-Ras-GTP and effectively interfered with the Ras-effector interactions, although they also showed a sign of binding activity toward H-Ras-GDP in the 1D <sup>1</sup>H NMR analysis (Fig. S10).

Although the residues whose interaction with BZDN and DCAI detected by the HSQC analysis of H-Ras-GTP (19) showed some overlap with those identified by our NOE analysis with Kobe2601, a considerable difference existed in the location of the binding pockets and the orientation of the compounds (Fig. S11), which seemed to account for the difference in their ability to interfere with the effector interaction (Fig. S7 shows the effector interaction sites). Namely, the binding pocket for BZDN and DCAI in K-Ras-GTP is located close to Asp54, whose side chain forms a direct hydrogen bond with the NH group of BZDN, whereas Kobe2601 is too far to establish any direct interactions with Asp54. Sun et al. (20) also reported discovery of small-molecule compounds inhibiting K-Ras-GDP, which showed only the inhibition toward the Sos-mediated nucleotide exchange in vitro and shared the binding pocket and the orientation of the compounds on Ras-GDP with BZIM, BZDN, and DCAI. At present, it is not clear whether Sos inhibition is an effective strategy for suppressing the constitutively activated Ras mutants, considering the great reduction of their GTPase activity and a vast excess of free GTP over GDP in cellular concentrations. In this regard, our results showing that the Ras<sup>G12V</sup>.GTP level was almost unaffected by the mSos1 level (Fig. 2B and C) indicated that H-Ras<sup>G12V</sup> escaped from the regulation by Sos. However, Sos inhibition might be effective for some cancer types, considering that the function of wild-type Ras is required for the growth of tumors carrying the activated *ras* oncogene (21).

In conclusion, we found that the Kobe0065-family compounds bind to Ras-GTP and exhibit antiproliferative activity toward cancer cells carrying the activated *ras* oncogenes, by a strategy based on SBDD. The compounds efficiently inhibit the interaction of Ras-GTP with their multiple effectors including Raf, PI3K, and RalGDS and a regulator/effector Sos and show rather broad binding specificity toward various Ras family small GTPases, which may account for their higher potency at the cellular level compared with that of the in vitro binding inhibition. Although the inhibitory activity is not particularly potent at present with the order of  $10^{-6}$  to  $10^{-5}$  M, the Kobe0065-family compounds may serve as a lead scaffold for the development of Ras inhibitors with higher potency and specificity and low toxicity that are suitable for clinical application. For this purpose, we would propose two possible strategies for structural optimization: the addition of a functional group that gains a hydrogen-bonding or ionic interaction with the charged residues such as Asp54 to increase the avidity and the avoidance of the



thiosemicarbazide structure, which is generally considered to lead to cellular toxicity.

## Materials and Methods

**In Silico Docking Screening.** Structure-based screening was targeted at a surface pocket of M-Ras<sup>P40D</sup>GppNHp (PDB ID code 3KKP) (9) surrounded by the two switch regions and the nucleotide. The MMPB-SA method was used with an AMBER96 force field, where the solvent effect on binding free energy was introduced upon simulation (22). A virtual library containing 40,882 compounds (Namiki Shoji Co., Ltd., [www.namiki-s.co.jp](http://www.namiki-s.co.jp)) was filtered by application of "Lipinski's rule of five" for the selection of drug-like compounds, yielding 40,307 compounds to be screened. Upon docking simulation, the targeting pocket was specified by amino acid residues located within a 6.5-Å distance from the probe points, which were generated by referring to the position of Asp67 (corresponding to Asp57 of H-Ras) in M-Ras<sup>P40D</sup>GppNHp. The initial 3D Ras-compound docking structures and electric charges of the molecules in the presence or absence of water molecules around a Mg<sup>2+</sup> ion were calculated by using Sievegene in myPresto software (23, 24) and Tripos software, respectively. Candidates were selected based on the calculated docking free energy values and the Nihon Electric Company's original scoring functions.

1. Karnoub AE, Weinberg RA (2008) Ras oncogenes: Split personalities. *Nat Rev Mol Cell Biol* 9(7):517–531.
2. Vetter IR, Wittinghofer A (2001) The guanine nucleotide-binding switch in three dimensions. *Science* 294(5545):1299–1304.
3. Weinstein IB (2002) Cancer. Addiction to oncogenes—The Achilles heel of cancer. *Science* 297(5578):63–64.
4. Chin L, et al. (1999) Essential role for oncogenic Ras in tumour maintenance. *Nature* 400(6743):468–472.
5. Podsypanina K, Politi K, Beverly LJ, Varmus HE (2008) Oncogene cooperation in tumor maintenance and tumor recurrence in mouse mammary tumors induced by Myc and mutant Kras. *Proc Natl Acad Sci USA* 105(13):5242–5247.
6. James GL, Goldstein JL, Brown MS (1995) Polylysine and CVIM sequences of K-RasB dictate specificity of prenylation and confer resistance to benzodiazepine peptidomimetic in vitro. *J Biol Chem* 270(11):6221–6226.
7. Rotblat B, Ehrlich M, Haklai R, Kloog Y (2008) The Ras inhibitor farnesylthiosalicylic acid (Salirasib) disrupts the spatiotemporal localization of active Ras: A potential treatment for cancer. *Methods Enzymol* 439:467–489.
8. Ye M, et al. (2005) Crystal structure of M-Ras reveals a GTP-bound "off" state conformation of Ras family small GTPases. *J Biol Chem* 280(35):31267–31275.
9. Shima F, et al. (2010) Structural basis for conformational dynamics of GTP-bound Ras protein. *J Biol Chem* 285(29):22696–22705.
10. Muraoka S, et al. (2012) Crystal structures of the state 1 conformations of the GTP-bound H-Ras protein and its oncogenic G12V and Q61L mutants. *FEBS Lett* 586(12):1715–1718.
11. Wilhelm S, et al. (2006) Discovery and development of sorafenib: A multikinase inhibitor for treating cancer. *Nat Rev Drug Discov* 5(10):835–844.
12. Margarit SM, et al. (2003) Structural evidence for feedback activation by Ras.GTP of the Ras-specific nucleotide exchange factor SOS. *Cell* 112(5):685–695.
13. Araki M, et al. (2011) Solution structure of the state 1 conformer of GTP-bound H-Ras protein and distinct dynamic properties between the state 1 and state 2 conformers. *J Biol Chem* 286(45):39644–39653.
14. Ito Y, et al. (1997) Regional polyesterism in the GTP-bound form of the human c-Ha-Ras protein. *Biochemistry* 36(30):9109–9119.
15. Nassar N, et al. (1995) The 2.2 Å crystal structure of the Ras-binding domain of the serine/threonine kinase c-Raf1 in complex with Rap1A and a GTP analogue. *Nature* 375(6532):554–560.
16. Pacold ME, et al. (2000) Crystal structure and functional analysis of Ras binding to its effector phosphoinositide 3-kinase gamma. *Cell* 103(6):931–943.
17. Huang L, Hofer F, Martin GS, Kim SH (1998) Structural basis for the interaction of Ras with RalGDS. *Nat Struct Biol* 5(6):422–426.
18. Hajduk PJ, Olejniczak ET, Fesik SW (1997) One-dimensional relaxation- and diffusion-edited NMR methods for screening compounds that bind to macromolecules. *J Am Chem Soc* 119:12257–12261.
19. Maurer T, et al. (2012) Small-molecule ligands bind to a distinct pocket in Ras and inhibit SOS-mediated nucleotide exchange activity. *Proc Natl Acad Sci USA* 109(14):5299–5304.
20. Sun Q, et al. (2012) Discovery of small molecules that bind to K-Ras and inhibit Sos-mediated activation. *Angew Chem Int Ed Engl* 51(25):6140–6143.
21. Lim KH, Ancrile BB, Kashatus DF, Counter CM (2008) Tumour maintenance is mediated by eNOS. *Nature* 452(7187):646–649.
22. Sitkoff D, Sharp KA, Honig B (1994) Accurate calculation of hydration free energies using macroscopic solvent models. *J Phys Chem* 98:1978–1988.
23. Fukunishi Y, Mikami Y, Nakamura H (2003) The filing potential method: A method for estimating the free energy surface for protein-ligand docking. *J Phys Chem B* 107:13201–13210.
24. Fukunishi Y, Mikami Y, Nakamura H (2005) Similarities among receptor pockets and among compounds: analysis and application to in silico ligand screening. *J Mol Graph Model* 24(1):34–45.
25. Carhart RE, Smith DH, Venkataraghavan R (1985) Atom pairs as molecular features in structure-activity studies: Definition and applications. *J Chem Inf Comput Sci* 25:64–73.
26. Willett P, Barnard JM, Downs GM (1998) Chemical similarity searching. *J Chem Inf Comput Sci* 38:983–996.
27. Koradi R, Billeter M, Wuthrich K (1996) MOLMOL: A program for display and analysis of macromolecular structures. *J Mol Graph* 14(1):51–55.

# Supporting Information

Shima et al. 10.1073/pnas.1217730110

## SI Materials and Methods

**Compounds and Cells.** Kobe0065: *N*-(3-chloro-4-methylphenyl)-2-{2,6-dinitro-4-(trifluoromethyl)phenyl}hydrazinecarbothioamide, 2601: 2-(2,4-dinitrophenyl)-*N*-(4-fluorophenyl)hydrazinecarbothioamide, and 2602: 2-{2,6-dinitro-4-(trifluoromethyl)phenyl}-*N*-(4-fluorophenyl)hydrazinecarbothioamide were purchased from Namiki Shoji Co., Ltd. Sorafenib was purchased from LC Laboratories Inc. Various human cancer cell lines including SW480 were obtained from the European Collection of Cell Culture and cultured in media recommended by the supplier. H-Ras<sup>G12V</sup>- and c-Raf-1<sup>S259A/Y340D/Y341D</sup>-transformed NIH 3T3 cells were cloned from foci obtained by focus forming assays using pT24 (1) containing the BamHI fragment of the genomic H-Ras<sup>G12V</sup> gene and pH8-flag-c-raf-1<sup>S259A/Y340D/Y341D</sup> (2), respectively, and cultured in DMEM supplemented with 10% FBS, 100 IU/mL penicillin, and 100 µg/mL streptomycin.

**Biochemical Assays.** H-Ras (residues 1–166) and GST-c-Raf-1-Ras-binding domain (RBD; residues 50–131) were produced in *Escherichia coli* and purified as described previously (3, 4). For the in vitro binding inhibition assays, H-Ras(1–166), preloaded with [ $\gamma$ -<sup>35</sup>S]GTP $\gamma$ S, was incubated with GST-c-Raf-1-RBD(50–131) at 25 °C for 30 min in the presence of the compound and the amount of bound H-Ras was quantified as the radioactivity pulled down by glutathione-sepharose resin. The  $K_i$  value for the compound was calculated as described in Fig. S1. For in vivo assays, NIH 3T3 cells were transfected with pEF-BOS-HA-H-Ras<sup>G12V</sup> or pEF-BOS-HA-K-Ras<sup>G12V</sup>, cultured for 18 h at 10% (vol/vol) FBS, and then incubated in the presence of the compound at 2% (vol/vol) FBS for 1 h. Cells were lysed in 50 mM Tris-HCl (pH 7.4), 150 mM NaCl, 1% Nonidet P-40, 10% (vol/vol) glycerol, 1 mM EDTA, 1 mM DTT, phosphatase inhibitor mixture (Nacalai Tesque), and protease inhibitor mixture (Nacalai Tesque) and subjected to detection of c-Raf-1 coimmunoprecipitated with an anti-H-Ras antibody (C-20; Santa Cruz) by Western blotting with an anti-c-Raf-1 antibody (C-12; Santa Cruz), of phosphorylated MEK and ERK with anti-pMEK1/2 (p217/p221; Cell Signaling) and anti-ERK1/2 (p202/204; Cell Signaling) antibodies, and of phosphorylated Akt with an anti-pAkt antibody (Ser473; Cell Signaling) and of RalA-GTP pulled down with GST-Sec5(1–99) immobilized on glutathione-sepharose resin by an anti-RalA antibody (Becton Dickinson). HA-tagged H-Ras<sup>G12V</sup>·GTP was detected by an anti-HA antibody (Cell Signaling). In vitro assays for the kinase activity of recombinant c-Raf-1 were performed by using a Raf-1 Kinase Assay kit (Millipore). In vitro GDP–GTP exchange assays were done by incubating 600 nM GST-H-Ras(1–166)·GDP immobilized on glutathione-sepharose resin with 11 µM [ $\gamma$ -<sup>35</sup>S]GTP $\gamma$ S (1,500 cpm/pmol) at 25 °C in the presence of purified 6×His-tagged mouse Son of sevenless (mSos)1(563–1,049) (180 nM each), wild-type, or a W729E mutant (5) in buffer B [50 mM Tris-HCl (pH 7.4), 50 mM NaCl, 5 mM MgCl<sub>2</sub>, 1 mM DTT, and 20 mM imidazol]. The radioactivity remaining on the resin after an intensive washing was quantified by liquid scintillation counting. Varying concentrations of compounds were added to the reaction mixtures to observe their inhibitory effect.

**Colony Formation Assays.** Cells ( $10^3$  to  $10^4$ ) were inoculated in 2 mL of DMEM containing 10% (vol/vol) FBS, 0.33% SeaPlaque aga-

rose, and one of the compounds and overlaid onto bottom agar consisting of 4 mL of DMEM containing 10% (vol/vol) FBS, 0.6% SeaPlaque agarose, and the same concentration of the compound in a six-well culture plate. After incubation at 37 °C for 14–21 d, the number of colonies >200 µm in diameter was counted under a dissecting microscope.

**Cell Proliferation Assays.** Cells ( $2 \times 10^3$ ) were seeded in a 96-well plate and cultured in DMEM containing 2% (vol/vol) FBS in the presence of one of the compounds. Viable cell numbers were measured by formazan formation using a Cell Counting Kit 8 (Dojindo). Apoptotic cells were detected by a standard TUNEL assay using an In Situ Cell Detection kit (Roche).

**Tumor Xenografts.** Cells ( $5 \times 10^6$ ) were implanted into the right flanks of female athymic nude mice (6–8 wk old; CLEA Japan, Inc.). After tumor sizes reached ~50 mm<sup>3</sup> on average, compounds suspended in Cremophor:ethanol:water (1:1:6) were administered orally for five consecutive days per week for 17 d. Tumor volumes ( $V$ ) were calculated with the following formula:  $V = A \times B^2/2$ , where  $A$  is the largest diameter and  $B$  is the perpendicular diameter. Dissected tumors after 17-d administration of the 80 mg/kg compounds were fixed in 4% (wt/vol) paraformaldehyde and embedded in paraffin. Their sections were subjected to immunohistochemistry with an anti-ERK1/2 antibody or an anti-CD31 antibody using a HISTMOUSE-PLUS kit (ZYMED Laboratories). Apoptotic cells were detected by a TUNEL assay. Statistical significance for groups of three or more was determined by one-way ANOVA with Tukey's test for post hoc analysis.

**NMR Spectroscopy and Structure Calculation.** <sup>15</sup>N/<sup>13</sup>C-labeled H-Ras<sup>T35S</sup>(1–166) was purified and loaded with guanosine 5'-( $\beta$ ,  $\gamma$ -imido) triphosphate as described previously (4). The NMR measurements were performed at 5 °C on an Avance III-600 spectrometer (Bruker) equipped with a 5-mm TCI cryogenic probe on a mixture of 1.0 mM H-Ras<sup>T35S</sup>·GppNHp and 3.8 mM Kobe2601 in 80% <sup>2</sup>H<sub>2</sub>O and 20% DMSO-*d*<sub>6</sub>. Based on assignments of the protein resonances in the absence of DMSO at 25 °C (4), 18 intermolecular distance restraints were obtained from 3D <sup>13</sup>C-NOESY and 2D homonuclear NOESY spectra (mixing time 250 ms). For the structure calculation, the intraprotein distance restraints and backbone torsion angle restraints obtained in the absence of DMSO at 25 °C (4) were incorporated, assuming that the protein structure does not change upon compound binding. Among the 100 structures generated by CYANA 2.1 (6), the 20 structures with the lowest target functions were refined using CNS 1.2 (7) for selection of the lowest-energy complex structures (PDB ID code 2lwi and Table S3). The <sup>1</sup>H-<sup>15</sup>N HSQC spectrum of wild-type H-Ras·GppNHp were recorded on a DMX-750 spectrometer (Bruker) at 5 °C on 1.0 mM <sup>15</sup>N-labeled H-Ras(1–166)·GppNHp in the absence or presence of 3.3 mM Kobe2601 in 90% <sup>1</sup>H<sub>2</sub>O and 10% DMSO-*d*<sub>6</sub>. Direct interactions of Kobe0065 and Kobe2602 with various small GTPases of the Ras and Rho families were examined by relaxation-edited 1D NMR (8). All of the spectra were acquired with a Carr–Purcell–Meiboom–Gill spin-lock time of 400 ms.

1. Taparowsky E, et al. (1982) Activation of the T24 bladder carcinoma transforming gene is linked to a single amino acid change. *Nature* 300(5894):762–765.

2. Tamada M, Hu CD, Kariya K, Okada T, Kataoka T (1997) Membrane recruitment of Raf-1 is not the only function of Ras in Raf-1 activation. *Oncogene* 15(24):2959–2964.