

labeled pancreatic beta-cells. *Am J Physiol Endocrinol Metab* 2003;284:E177–E183.

29 Higuchi Y, Shiraki N, Yamane K et al. Synthesized basement membranes direct the differentiation of mouse embryonic stem cells into pancreatic lineages. *J Cell Sci* 2010;123:2733–2742.

30 Okita K, Ichisaka T, Yamanaka S. Generation of germline-competent induced pluripotent stem cells. *Nature* 2007;448:313–317.

31 Bosco D, Meda P, Halban PA et al. Importance of cell-matrix interactions in rat islet beta-cell secretion in vitro: Role of alpha6beta1 integrin. *Diabetes* 2000;49:233–243.

32 Takahashi K, Tanabe K, Ohnuki M et al. Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 2007;131:861–872.

33 Fusaki N, Ban H, Nishiyama A et al. Efficient induction of transgene-free human pluripotent stem cells using a vector based on Sendai virus, an RNA virus that does not integrate into the host genome. *Proc Jpn Acad, Ser B, Phys Biol Sci* 2009;85:348–362.

34 D'Amour KA, Bang AG, Eliazer S et al. Production of pancreatic hormone-expressing endocrine cells from human embryonic stem cells. *Nat Biotechnol* 2006;24:1392–1401.

35 Hitsuda T, Michiue H, Kitamatsu M et al. A protein transduction method using oligo-arginine (3R) for the delivery of transcription factors into cell nuclei. *Biomaterials* 2012;33:4665–4672.

36 Carlotti F, Zaldumbide A, Charif H et al. The 45-kDa form of Pdx-1 does not result from post-translational modifications. *Biochem Biophys Res Commun* 2008;370:225–229.

37 Oliver-Krasinski JM, Stoffers DA. On the origin of the beta cell. *Genes Dev* 2008;22:1998–2021.

38 Murtaugh LC. Pancreas and beta-cell development: From the actual to the possible. *Development* 2007;134:427–438.

39 Hammar E, Parnaud G, Bosco D et al. Extracellular matrix protects pancreatic beta-cells against apoptosis: Role of short- and long-term signaling pathways. *Diabetes* 2004;53:2034–2041.

40 Parnaud G, Bosco D, Berney T et al. Proliferation of sorted human and rat beta cells. *Diabetologia* 2008;51:91–100.

41 Bosco D, Gonelle-Gispert C, Wollheim CB et al. Increased intracellular calcium is required for spreading of rat islet beta-cells on extracellular matrix. *Diabetes* 2001;50:1039–1046.

42 Stoffers DA, Thomas MK, Habener JF. Homeodomain protein IDX-1: A master regulator of pancreas development and insulin gene expression. *Trends Endocrinol Metab* 1997;8:145–151.

43 Habener JF, Stoffers DA. A newly discovered role of transcription factors involved in pancreas development and the pathogenesis

of diabetes mellitus. *Proc Assoc Am Physicians* 1998;110:12–21.

44 Guz Y, Montminy MR, Stein R et al. Expression of murine STF-1, a putative insulin gene transcription factor, in beta cells of pancreas, duodenal epithelium and pancreatic exocrine and endocrine progenitors during ontogeny. *Development* 1995;121:11–18.

45 Naya FJ, Stellrecht CM, Tsai MJ. Tissue-specific regulation of the insulin gene by a novel basic helix-loop-helix transcription factor. *Genes Dev* 1995;9:1009–1019.

46 Schwitzgebel VM, Scheel DW, Connors JR et al. Expression of neurogenin3 reveals an islet cell precursor population in the pancreas. *Development* 2000;127:3533–3542.

47 Hang Y, Stein R. MafA and MafB activity in pancreatic β cells. *Trends Endocrinol Metab* 2011;22:364–373.

48 Matsuoka TA, Zhao L, Artner I et al. Members of the large Maf transcription family regulate insulin gene transcription in islet beta cells. *Mol Cell Biol* 2003;23:6049–6062.

49 Olbrot M, Rud J, Moss LG et al. Identification of beta-cell-specific insulin gene transcription factor RIPE3b1 as mammalian MafA. *Proc Natl Acad Sci USA* 2002;99:6737–6742.

50 Kataoka K, Han SJ, Shioda S et al. MafA is a glucose-regulated and pancreatic beta-cell-specific transcriptional activator for the insulin gene. *J Biol Chem* 2002;277:49903–49910.

51 Murtaugh LC, Melton DA. Genes, signals, and lineages in pancreas development. *Annu Rev Cell Dev Biol* 2003;19:71–89.

52 Jensen J. Gene regulatory factors in pancreatic development. *Dev Dyn* 2004;229:176–200.

53 Wessells NK, Cohen JH. Early pancreas organogenesis: Morphogenesis, tissue interactions, and mass effects. *Dev Biol* 1967;15:237–270.

54 Wessells NK, Evans J. Ultrastructural studies of early morphogenesis and cytodifferentiation in the embryonic mammalian pancreas. *Dev Biol* 1968;17:413–446.

55 Ahlgren U, Jonsson J, Edlund H. The morphogenesis of the pancreatic mesenchyme is uncoupled from that of the pancreatic epithelium in IPF1/PDX1-deficient mice. *Development* 1996;122:1409–1416.

56 Offield MF, Jetton TL, Labosky PA et al. PDX-1 is required for pancreatic outgrowth and differentiation of the rostral duodenum. *Development* 1996;122:983–995.

57 Mutoh H, Fung BP, Naya FJ et al. The basic helix-loop-helix transcription factor BETA2/NeuroD is expressed in mammalian enteroendocrine cells and activates secretin gene expression. *Proc Natl Acad Sci USA* 1997;94:3560–3564.

58 Matsuoka TA, Artner I, Henderson E et al. The MafA transcription factor appears to be responsible for tissue-specific expression of insulin. *Proc Natl Acad Sci USA* 2004;101:2930–2933.

59 Christodoulou C, Longmire TA, Shen SS et al. Mouse ES and iPS cells can form similar definitive endoderm despite differences in imprinted genes. *J Clin Invest* 2011;121:2313–2325.

60 Mei Y, Hollister-Lock J, Bogatyrev SR et al. A high throughput micro-array system of polymer surfaces for the manipulation of primary pancreatic islet cells. *Biomaterials* 2010;31:8989–8995.

61 Lefebvre VH, Otonkoski T, Ustinov J et al. Culture of adult human islet preparations with hepatocyte growth factor and 804G matrix is mitogenic for duct cells but not for beta-cells. *Diabetes* 1998;47:134–137.

62 Hayek A, Beattie GM, Cirulli V et al. Growth factor/matrix-induced proliferation of human adult beta-cells. *Diabetes* 1995;44:1458–1460.

63 Riddelle KS, Green KJ, Jones JC. Formation of hemidesmosomes in vitro by a transformed rat bladder cell line. *J Cell Biol* 1991;112:159–168.

64 Langhofer M, Hopkinson SB, Jones JC. The matrix secreted by 804G cells contains laminin-related components that participate in hemidesmosome assembly in vitro. *J Cell Sci* 1993;105:753–764.

65 Hormia M, Falk-Marzillier J, Plopper G et al. Rapid spreading and mature hemidesmosome formation in HaCaT keratinocytes induced by incubation with soluble laminin-5r. *J Invest Dermatol* 1995;105:557–561.

66 Baker SE, Hopkinson SB, Fitchmun M et al. Laminin-5 and hemidesmosomes: Role of the alpha 3 chain subunit in hemidesmosome stability and assembly. *J Cell Sci* 1996;109:2509–2520.

67 Colognato H, Yurchenco PD. Form and function: The laminin family of heterotrimers. *Dev Dyn* 2000;218:213–234.

68 Aumailley M, El Khal A, Knöss N et al. Laminin 5 processing and its integration into the ECM. *Matrix Biol* 2003;22:49–54.

69 Teitelman G, Alpert S, Polak JM et al. Precursor cells of mouse endocrine pancreas co-express insulin, glucagon and the neuronal proteins tyrosine hydroxylase and neuropeptide Y, but not pancreatic polypeptide. *Development* 1993;118:1031–1039.

70 Kunisada Y, Tsubooka-Yamazoe N, Shoji M et al. Small molecules induce efficient differentiation into insulin-producing cells from human induced pluripotent stem cells. *Stem Cell Res (Amst)* 2012;8:274–284.

71 Merino JF, Nacher V, Raurell M et al. Optimal insulin treatment in syngeneic islet transplantation. *Cell Transplant* 2000;9:11–18.

72 Cheng X, Ying L, Lu L et al. Self-renewing endodermal progenitor lines generated from human pluripotent stem cells. *Cell Stem Cell* 2012;10:371–384.



See www.StemCellsTM.com for supporting information available online.

Citations

This article has been cited by 1 HighWire-hosted articles:
<http://stemcellstm.alphamedpress.org/content/3/1/114#otherarticles>

Quantitative PCR Measurement of tRNA 2-Methylthio Modification for Assessing Type 2 Diabetes Risk

Peiyu Xie,^{1†} Fan-Yan Wei,^{1†} Shoji Hirata,² Taku Kaitsuka,¹ Tsutomu Suzuki,² Takeo Suzuki,^{2*} and Kazuhito Tomizawa^{1*}

BACKGROUND: Genetic variants in the human *CDKAL1* (CDK5 regulatory subunit associated protein 1-like 1) gene have been associated with reduced insulin secretion and type 2 diabetes (T2D). *CDKAL1* is a methylthiotransferase that catalyzes 2-methylthio (ms^2) modification of the adenine at position 37 (A37) of cytoplasmic tRNA^{Lys}-(UUU). We investigated the ms^2 -modification level of tRNA^{Lys}-(UUU) as a direct readout of *CDKAL1* enzyme activity in human samples.

METHOD: We developed a quantitative PCR (qPCR)-based method to measure ms^2 modification. tRNA^{Lys}-(UUU) was reverse-transcribed with 2 unique primers: Reverse primer r1 was designed to anneal to the middle of this tRNA, including the nucleotide at A37, and reverse primer r2 was designed to anneal to the region downstream (3') of A37. Subsequent qPCR was performed to detect the corresponding transcribed cDNAs.

RESULTS: The efficiency of reverse transcription of tRNA^{Lys}-(UUU) was ms^2 -modification dependent. The relative difference in threshold cycle number obtained with the r1 or r2 primer yielded the ms^2 -modification level in tRNA^{Lys}-(UUU) precisely as predicted by an original mathematical model. The method was capable of measuring ms^2 -modification levels in tRNA^{Lys}-(UUU) in total RNA isolated from human peripheral blood samples, revealing that the ms^2 -modification rate in tRNA^{Lys}-(UUU) was decreased in individuals carrying the *CDKAL1* genotype associated with T2D. In addition, the ms^2 -modification level was correlated with insulin secretion.

CONCLUSIONS: The results point to the critical role of ms^2 modification in T2D and to a potential clinical use

of a simple and high-throughput method for assessing T2D risk.

© 2013 American Association for Clinical Chemistry

Nucleotides in tRNAs are intensively modified after transcription. These posttranscriptional modifications are necessary for the structural integrity and decoding activity of tRNA (1–3). Defects in tRNA modifications can lead to such pathologies as type 2 diabetes (T2D)³ and mitochondrial diseases (4–6). Recent genome-wide association studies have shown single-nucleotide polymorphisms (SNPs) in an intron region of *CDKAL1* (CDK5 regulatory subunit associated protein 1-like 1)⁴ are associated with T2D (7, 8). Individuals carrying the risk allele of *CDKAL1* have decreased insulin secretion (7). We recently reported that *CDKAL1* is a tRNA-modifying enzyme that catalyzes 2-methylthio (ms^2) transformation of N⁶-threonylcarbamoyladenosine (t⁶A) to 2-methylthio-N⁶-threonylcarbamoyladenosine (ms^2 t⁶A) at position 37 of cytoplasmic tRNA^{Lys}-(UUU) (9). The ms^2 modification is required for accurate decoding of the lysine codon (4). In pancreatic β -cell-specific *Cdkal1* knockout (KO) mice, the deficiency in ms^2 modification of tRNA^{Lys}-(UUU) produces aberrant translation of the lysine codon in proinsulin, which leads to the development of T2D symptoms, including reduced insulin secretion and impaired blood glucose control (4). These results not only suggest that a decrease in the ms^2 -modification level in tRNA^{Lys}-(UUU) is critical to the development of T2D but also raise the possibility for the use of ms^2 -modification rates in assessing T2D risk.

¹ Department of Molecular Physiology, Kumamoto University, Kumamoto, Japan;

² Department of Chemistry and Biotechnology, Graduate School of Engineering, University of Tokyo, Tokyo, Japan.

* Address correspondence to these authors: Kazuhito Tomizawa at Department of Molecular Physiology, Kumamoto University, Kumamoto 860-8556, Japan. Fax 81-96-373-5052; e-mail tomikt@kumamoto-u.ac.jp. Takeo Suzuki at Department of Chemistry and Biotechnology, Graduate School of Engineering, University of Tokyo, Bunkyo-ku, Tokyo 113-8656, Japan. Fax 81-3-5841-0550; e-mail t_suzuki@chembio.t.u-tokyo.ac.jp.

[†] These authors contributed equally to this work.
Received May 28, 2013; accepted July 23, 2013.

Previously published online at DOI: 10.1373/clinchem.2013.210401

³ Nonstandard abbreviations: T2D, type 2 diabetes; SNP, single-nucleotide polymorphism; ms^2 , 2-methylthio; t⁶A, N⁶-threonylcarbamoyladenosine; ms^2 t⁶A, 2-methylthio-N⁶-threonylcarbamoyladenosine; KO, knockout (mice); qPCR, quantitative PCR; qPCR-MTR, qPCR-based method for detecting methylthiolation of tRNA; WT, wild-type; Ct, threshold cycle; Ct₁, Ct number obtained from a qPCR with a cDNA template generated by primer r1; dCt_{2/r1}, Ct₂ – Ct₁; T_m, melting temperature; CIR₃₀, corrected insulin response at 30 min; A37, adenine at position 37.

⁴ Genes: *CDKAL1*, CDK5 regulatory subunit associated protein 1-like 1; *Cdkal1*, *Mus musculus* gene CDK5 regulatory subunit associated protein 1-like 1.

We therefore aimed to measure ms^2 -modification status as a direct marker for investigating CDKAL1 enzyme activity in human samples and its association with T2D risk; however, current methods of measuring site-specific tRNA modifications, such as mass spectrometry and the primer-extension method (6, 10), are not applicable to clinical samples, for several reasons. First, these methods are time-consuming and require expertise and expensive equipment. Second, they are unable to handle multiple samples in parallel. Third, mass spectrometry in particular requires a large quantity of RNA for isolating tRNAs, although analyses use only a small amount of tRNA. Thus, there is a need to develop a method that is applicable to the measurement of ms^2 modifications in clinical samples. We found that ms^2 modification attenuates reverse transcription because it sterically hinders Watson–Crick base pairing. Taking advantage of this finding, we developed a quantitative PCR (qPCR)-based method for detecting methylthiolation of tRNA (qPCR-MtR), which overcomes the disadvantages of the current method by allowing measurement of ms^2 modifications in multiple samples in as little as 3 h.

Materials and Methods

PARTICIPANT RECRUITMENT

Blood samples were obtained from nondiabetic volunteers at Kumamoto University. Informed consent was obtained before blood was collected. The Ethics Committee of Kumamoto University approved the experiments regarding analysis of human genetic materials and collecting human blood (approval ID, Genome 159).

RNA ISOLATION

For isolation of total RNA from mouse liver, we used a guanidinium thiocyanate/phenol/chloroform method with TRIzol reagent (Life Technologies, Grand Island, NY) according to the manufacturer's protocol. One particular tRNA, $\text{tRNA}^{\text{Lys}}(\text{UUU})$, was further purified with reciprocal circulating chromatography, as described previously (4, 11). The $\text{tRNA}^{\text{Lys}}(\text{UUU})$ isolated from wild-type (WT) and *Cdkal1* KO mice were subjected to qPCR-MtR (Fig. 1A–C). Threshold cycle (Ct) values calculated with the manufacturer's software are indicated in this figure. To calculate the modification rate, we combined purified $\text{tRNA}^{\text{Lys}}(\text{UUU})$ isolated from WT mice and *Cdkal1* KO mice at the indicated ratios and carried out qPCR-MtR analyses (Fig. 1D). The calculated modification ratio was obtained as described in Fig. 1 in the Data Supplement that accompanies the online version of this article at <http://www.clinchem.org/content/vol59/issue11>. The value for the constant *E* for the efficiency of target amplification was assumed to be 1.

For absolute quantification of ms^2 modification in crude total RNA, we subjected total RNA isolated from WT and *Cdkal1* KO mice to qPCR-MtR (Fig. 2, A and B). To calculate the modification rate, we combined total RNA isolated from each mouse at the indicated ratios and subjected the combined total RNA to qPCR-MtR (Fig. 2C). Calculations are as presented in Fig. 1 in the online Data Supplement.

We isolated total RNA from peripheral blood by mixing 1.5 mL fresh peripheral blood with the erythrocyte-lysis buffer provided in the QIAamp RNA Blood Mini Kit (Qiagen). Total RNA was isolated according to the manufacturer's protocol (Fig. 3, A and B). To examine the appropriateness of the RNA-isolation methods, we recruited another 18 individuals. We added erythrocyte-lysis buffer to 3 mL blood and then aliquoted identical volumes of the mixture into 2 tubes. We isolated total RNA in one of the tubes with the QIAamp RNA Blood Mini Kit (Qiagen) and the total RNA in the other tube with TRIzol reagent. Total RNA was then subjected to qPCR-MtR amplification as described below.

For isolation of total RNA for mass spectrometry analysis, 50-mL peripheral blood samples were collected from each individual. Leukocytes were collected with the erythrocyte-lysis buffer provided in the QIAamp RNA Blood Mini Kit (Qiagen). Total RNA in leukocytes was purified with TRIzol reagent according to the manufacturer's protocol. A particular tRNA, $\text{tRNA}^{\text{Lys}}(\text{UUU})$, was further purified with reciprocal circulating chromatography.

CALCULATIONS

The mathematical model used to measure the modification rate is described in Fig. 1 in the online Data Supplement. The Ct number for a sample was obtained via qPCR with a cDNA template generated by primer r1 (Ct_{r1}) or by primer r2 (Ct_{r2}). The difference between Ct_{r1} and Ct_{r2} (hereafter dCt_{r2r1}) was calculated by subtracting Ct_{r1} from Ct_{r2} . The dCt_{r2r1} number for individual samples was used to calculate the absolute modification rate directly when the absolute modification rates of 2 samples were known, or to compare the modification rate of one sample relative to that of another.

PRIMER DESIGN

The DNA sequences of $\text{tRNA}^{\text{Lys}}(\text{UUU})$ were retrieved from a tRNA database (tRNAdb, <http://trna.bioinf.uni-leipzig.de/>) (12). The sequences of the primers used for detecting the ms^2 modification in $\text{tRNA}^{\text{Lys}}(\text{UUU})$ are as follows: forward primer, 5'-GTCGG-TAGAGCATCAGACTT-3'; reverse primer r1, 5'-CCTGGACCTCAGATTAAAA-3'; reverse primer r2, 5'-GAACAGGGACTTGAACCCTG-3'. All primer se-




Fig. 1. Absolute quantification of the ms² modification in purified tRNA^{Lys}(UUU) by qPCR-MtR analysis.

A, Flow chart of the qPCR-MtR method for measuring ms² modification in tRNA^{Lys}(UUU). B and C, qPCR-MtR of tRNA^{Lys}(UUU) purified from WT and *Cdkal1* KO mice. Shown are representative amplifications of a cDNA template generated with reverse primer r1 or reverse primer r2. Ct values calculated with the manufacturer's software are indicated. D, Purified tRNA^{Lys}(UUU) isolated from WT and *Cdkal1* KO mice were combined at the indicated ratios and subjected to qPCR-MtR. The calculated modification rate was obtained from the ddCt value and compared with the expected modification rate.

quences were screened with Primer 3 software (<http://frodo.wi.mit.edu/primer3/>) to eliminate primers with self-complementarity and extremely low melting-temperature (T_m) values.

REVERSE TRANSCRIPTION AND qPCR


The total amount of RNA isolated from cells and tissues was adjusted to 100 ng/ μ L in RNase-free water, unless otherwise indicated. To avoid genomic contamination, we digested 2 μ L (200 ng) total RNA with 5 U DNase I (Roche Applied Science) in a 20- μ L reaction at 37 °C for 20 min. We then heat-inactivated the DNase I at 75 °C for 10 min. After the DNase treatment, we mixed 2.5 μ L digested total RNA with 1 μ L 20 μ mol/L reverse primer r1 or reverse primer r2, heat-denatured the mixture at 65 °C for 10 min, and cooled it rapidly on ice for at least 5 min. With the mixture still on ice, recombinant reverse transcriptase (Transcriptor; Roche Applied Science) was added to a final concentra-

tion of 0.5 U/ μ L. Reverse transcription was performed in a total reaction volume of 10 μ L at 55 °C for 30 min and then heat-inactivated at 85 °C for 5 min. We subjected 2 μ L synthesized cDNA to qPCR with the SYBR Premix Ex Taq kit (Takara) and the ABI PRISM 7300 Real-Time PCR System (Life Technologies) according to the manufacturers' directions.

For analysis of isolated tRNA^{Lys}(UUU), we used the Transcriptor First Strand cDNA Synthesis Kit, the LightCycler 480 SYBR Green I Master, and the LightCycler 480 real-time PCR system (Roche Applied Science). cDNA was synthesized in a 20- μ L volume from 3 fmol of tRNA prepared at defined proportions of tRNA with and without the ms² modification. The qPCR was then performed with 0.75 μ L synthesized cDNA.

MASS SPECTROMETRIC ANALYSIS OF tRNA

Isolated tRNA^{Lys}(UUU) was digested with RNase A to obtain pyrimidine-ending oligonucleotides, which



were then analyzed with liquid chromatography–mass spectrometry, as described previously (4).

ANIMALS AND CELL CULTURE

Cdkal1 KO mice were generated as described previously (4). The Animal Ethics Committee of Kumamoto University approved all animal procedures (approval ID: B24–134, B24–132).


ORAL GLUCOSE TOLERANCE TEST

Human individuals underwent fasting beginning at 9 PM the night before the test day. They were then administered a 75-g load of glucose (Ajinomoto) between 9 AM and 11 AM on the test day. Peripheral blood

samples were obtained while the individuals were fasting and 30 min after glucose loading. Plasma glucose and serum insulin concentrations were measured by SRL Inc. The corrected insulin response at 30 min (CIR₃₀) was calculated as $I_{30}/[G_{30} \times (G_{30} - 70)]$, where I_{30} is the insulin concentration in microunits per milliliter at 30 min and G_{30} is the glucose concentration in milligrams per deciliter at 30 min (7).

GENOTYPING OF SNPs IN *CDKALI*

Genomic DNA was purified from 200 μ L peripheral human blood with the QIAamp DNA Blood Mini Kit (Qiagen) and adjusted to 10 ng/ μ L with distilled water. The




SNP (rs7754840) in *CDKAL1* was examined with the TaqMan SNP Genotyping Assay Kit (Life Technologies).

Results

MEASURING ms^2 MODIFICATION OF A37 IN PURIFIED $tRNA^{Lys}(UUU)$ BY qPCR-MIR

Posttranscriptional chemical modifications in tRNA are required to stabilize tRNA structure and codon–

anticodon interaction (1, 2). Interestingly, previous studies have shown that some of these modifications could attenuate primer extension, possibly owing to steric hindrance (6, 13–16). We investigated the molecular structure of the ms^2 modification of adenine at position 37 (A37) of $tRNA^{Lys}(UUU)$ in the presence of thymine during reverse transcription (Fig. 4A and 4B). In our model, which is based on a previous study of 2-methylthio- N^6 -isopentenyladenosine at position 37



in bacterial tRNA^{Phe} (17), the calculated distance between the S of the ms² group and the O at position 2 of the thymine base was 0.22 nm, whereas the calculated van der Waals radii of S and O were 0.18 nm and 0.15 nm, respectively (18). Therefore, the S–O distance is sufficiently short that steric hindrance prevents Watson–Crick pairing between ms²t⁶A and T, whereas the N⁶-threonylcarbonyl group does not affect Watson–Crick pairing with T (Fig. 4B). We then speculated that the ms² group might affect Watson–Crick pairing with thymine during reverse transcription because of the former's large van der Waals radii. To verify this speculation, we carried out reverse transcription with 2 primers specifically designed to amplify the region including position 37 in tRNA^{Lys}(UUU) (step 2, Fig. 1A). Reverse primer r1 was designed to anneal with the specific region encompassing A37, whereas reverse primer r2 was designed to anneal with a specific region downstream (3') of A37. The cDNA generated with either primer r1 or r2 was quantified by qPCR (step 3, Fig. 1A). If the ms² modification attenuates reverse transcription of fully modified tRNA^{Lys}(UUU), we would expect the amount of cDNA generated by primer r2 to be less than that generated by primer r1. In hypomodified tRNA^{Lys}(UUU), on the other hand, we would expect the amount of cDNA generated by primer r2 to be close to that generated by primer r1, in proportion to the reduction in ms² modification.

tRNA^{Lys}(UUU) with and tRNA^{Lys}(UUU) without the ms² modification were purified from liver of a WT mouse or a *Cdkal1* null mouse (4), respectively, and subjected to qPCR–MtR. As expected, use of a fully modified WT tRNA^{Lys}(UUU) as template markedly increased Ct_{r2} compared with Ct_{r1} (Fig. 1B). In contrast, use of unmodified tRNA^{Lys}(UUU) as template reduced Ct_{r2} to a level close to Ct_{r1} (Fig. 1C). These results suggest that ms² modification at A37 indeed attenuated reverse transcription when primer r2 was used, whereas reverse transcription was unaffected when primer r1 was used. Thus, Ct_{r2} reflects the extent of ms² modification, whereas Ct_{r1} reflects the total number of tRNA molecules in a tRNA sample.

ABSOLUTE QUANTIFICATION OF ms² MODIFICATION BY qPCR–MtR

By combining the conventional ddCt method and an original mathematical model, we demonstrated that the difference between Ct_{r1} and Ct_{r2} (i.e., dCt_{r2r1} = Ct_{r2} – Ct_{r1}) in a given sample can be used as an index that represents the modification rate (see Fig. 1 in the online Data Supplement). In short, the smaller the dCt_{r2r1} value, the less the ms² modification of the tRNA. In addition, the model enabled us to calculate the absolute modification rate of an unknown sample from 2 reference samples with known absolute modification rates.

To verify our calculation, we prepared tRNA^{Lys}-(UUU) with a known ms²-modification rate by combining tRNA^{Lys}(UUU) molecules from WT and *Cdkal1* KO mice at defined ratios and then performed qPCR-MtR (Fig. 1D). The dCt value decreased in proportion to the reduction in the ms²-modification rate. Furthermore, the calculated ms²-modification rate for the WT and modified tRNA mixtures (samples 2–4, Fig. 1D) accurately predicted the expected ms²-modification rate when fully ms²-modified tRNA (sample 5 in Fig. 1D, modification rate = 1; *c* in equation 14 in Fig. 1 in the online Data Supplement) and unmodified tRNA (sample 1 in Fig. 1D, modification rate = 0; *a* in equation 14 in Fig. 1 in the online Data Supplement) were used as references.

APPLICATION OF qPCR-MtR TO SAMPLES OF CRUDE TOTAL RNA
Purifying tRNA^{Lys}(UUU) is a time-consuming process requiring milligrams of total RNA. Because clinical samples are usually too small to yield this much total RNA, adapting qPCR-MtR for a small amount of crude RNA would be ideal for clinical samples. We subjected 200 ng total RNA isolated from 2 WT mice and 2 *Cdkal1* KO mice to qPCR-MtR analysis. In agreement with the results obtained with purified tRNA^{Lys}(UUU), we obtained a marked reduction in the Ct₂ value for the samples of total RNA isolated from *Cdkal1* KO mice, compared with WT mice (Fig. 2, A and B). Furthermore, the calculated ms²-modification rate obtained from the dCt_{r2r1} value accurately predicted the expected ms²-modification rate (Fig. 2C). In addition, we were able to calculate the absolute ms²-modification rate even when we used only 2 ng total RNA for qPCR-MtR (see Table 1 in the online Data Supplement). All procedures were finished in <3 h.

DECREASED ms²-MODIFICATION LEVEL OF tRNA^{Lys}(UUU) IN INDIVIDUALS CARRYING RISK *CDKAL1* SNPs

Because qPCR-MtR successfully detected ms² modification in tRNA^{Lys}(UUU), we applied this method via a column procedure and with 100 ng total RNA isolated from peripheral blood samples of nondiabetic individuals (*n* = 86). In addition, we determined relative modification levels by using the dCt_{r2r1} value, because a reference for the absolute ms²-modification rate in human tRNA^{Lys}(UUU) was not available. We compared the dCt_{r2r1} values obtained for total RNA isolated from 20 individuals homozygous for the T2D-associated risk allele (C/C) of the *CDKAL1* SNP (rs7754840) and 35 individuals heterozygous for the risk allele (G/C) with the dCt_{r2r1} values obtained for 31 individuals homozygous for the nonrisk allele (G/G). dCt_{r2r1} values were significantly lower in individuals carrying the risk allele (G/C and C/C) (Fig. 3A), suggesting that ms² modification was suppressed in individuals carrying the T2D-

associated risk allele for the *CDKAL1* SNP. The accuracy of the qPCR-MtR result was confirmed by mass spectrometric analysis. tRNA^{Lys}(UUU) was purified from 50 μg of total RNA isolated from 50 mL of blood from 2 individuals of same group carrying risk (C/C) or nonrisk (G/G) alleles of *CDKAL1* SNPs and analyzed with mass spectrometry. The amount of ms²-unmodified tRNA^{Lys}(UUU) was increased 1.47-fold (Fig. 3C) in individuals carrying the risk allele, compared with individuals carrying the nonrisk allele (see Fig. 2 in the online Data Supplement for a representative mass chromatogram). This result was compatible with the qPCR results obtained with the corresponding total-RNA samples (dCt_{nonrisk} – dCt_{risk} = 1).

tRNA is lost during the purification of total RNA when a column method is used, and that loss may interfere with measurement of the correct ms²-modification level. Therefore, we purified peripheral blood RNA from additional 18 nondiabetic individuals, both by a column method and by a guanidinium thiocyanate/phenol/chloroform method with the same samples; we then compared the dCt_{r2r1} values (see Fig. 3 in the online Data Supplement). We obtained a smaller dCt_{r2r1} value with the column method than with the guanidinium thiocyanate/phenol/chloroform method. These results suggest a partial loss of tRNA during column purification; however, the dCt_{r2r1} values obtained with the 2 methods were significantly correlated (see Fig 3A in the online Data Supplement). We also used electrophoresis to compare the qPCR products obtained with each isolation method. The intensities and sizes of the bands obtained with the 2 methods were identical (see Fig. 3B in the online Data Supplement).

The risk allele of *CDKAL1* has been associated with impaired insulin secretion (7). To investigate whether the ms²-modification level of tRNA^{Lys}(UUU) was also associated with insulin secretion, we carried out oral glucose tolerance tests with 28 nondiabetic individuals. Insulin secretion (CIR₃₀) was significantly correlated with dCt_{r2r1} (Fig. 3B), suggesting that the decreased ms²-modification level is associated with the impairment of insulin secretion.

Discussion

This study has demonstrated that the method we have developed (qPCR-MtR) can detect and quantify chemical modifications in tRNA^{Lys}(UUU), both for purified RNA and crude total RNA samples. This method is based on the unique biochemical feature of ms² modification. Previous biochemical and structural studies of synthetic poly(ms²-A) and poly(U) have demonstrated that Watson–Crick poly(ms²-A)–poly(U) pairing is not possible because of the large van der Waals

radii of the ms^2 group at the C2 position of adenine (19, 20). Given the structural similarity of uracil and thymine, our results suggest that the inhibitory effect of the ms^2 group at A37 of tRNA^{Lys}(UUU) is also due to its large van der Waals radii, which prevent Watson–Crick pairing with thymine during reverse transcription. Another possibility is that the ms^2 group causes almost complete loss of reverse transcription, rather than inhibiting it; hence, a trace amount of hypomodification remaining in fully ms^2 -modified samples would give a Ct_{r2} value apparently higher than Ct_{r1} . Indeed, signals corresponding to the m/z of the ms^2 -hypomodified molecule were detected at intensities of about 3% in the mass spectrometric analyses of the isolated tRNA^{Lys}(UUU) prepared from the WT mice (data not shown); however, this putative heterogeneity would not affect the calculation in Fig. 2C (by using 0.97 as the modification rate instead of 1). Because achieving complete purification of a single species from a heterogeneous population is almost impossible, the differences between Ct_{r2} and Ct_{r1} may be due to the combined effects of strong inhibition of reverse transcription and the intrinsic heterogeneity of the samples.

Genetic variants of the human *CDKAL1* gene have been associated with impaired insulin secretion and the development of T2D across ethnic groups (7). Because the disease's development in individuals carrying the *CDKAL1* risk allele could be influenced by environmental intervention, the *CDKAL1* genotype alone is less predictive of T2D (21). *CDKAL1* enzyme activity is regulated by both genetic variation and environmental conditions. Therefore, T2D risk would be assessed more accurately if genotype information were combined with ms^2 -modification measurement, which directly reflects *CDKAL1* activity. qPCR-MtR provides an efficient means of detecting and quantifying ms^2 modification of tRNA^{Lys}(UUU) in clinical samples. Our method successfully detected ms^2 modification in total RNA isolated from peripheral blood and showed that the ms^2 -modification rate was indeed decreased in individuals carrying risk-related genetic variants in the *CDKAL1* gene. Furthermore, the ms^2 -modification level was significantly correlated with insulin secretion. These results not only implicate the importance of ms^2 modification of tRNA^{Lys}(UUU) in T2D but also demonstrate that qPCR-MtR is fully applicable to the clinical setting. Further investigations into combining the ms^2 -modification level with genetic information and other clinical indices may lead to improvements in the accuracy of assessing T2D risk.

CDKAL1 has been implicated in diseases other than T2D, such as obesity and Crohn disease (22–24). Further investigations with qPCR-MtR may shed light on the pathologic function of ms^2 modifications in

these diseases. In addition, the lack of chemical modification, such as taurine modification in the anticodon loop of mitochondrial tRNAs, has been reported for mitochondrial diseases (5, 6). Because the onset of such mitochondrial diseases depends on the number of mitochondrial DNA copies with a mutated tRNA gene, quantifying the modification level in the corresponding tRNAs would help in diagnosis and treatment. The qPCR-MtR concept could easily be adapted for measuring the modification level in these mitochondrial tRNAs if these modifications also inhibit reverse transcription.

In this study, we compared the ms^2 -modification level in blood samples from which total RNA was isolated, both with a column method and a guanidinium thiocyanate/phenol/chloroform method. The dCt_{r2r1} value obtained by a column method was smaller than that obtained with the guanidinium thiocyanate/phenol/chloroform method, suggesting a partial loss of tRNA during column purification. Nevertheless, the dCt_{r2r1} values obtained with the 2 methods were significantly correlated, and the intensities and sizes of the bands of the qPCR products obtained with the 2 methods were identical. These results suggest that a guanidinium thiocyanate/phenol/chloroform method should be applied for absolute quantification of ms^2 modifications; however, the column method can still be used for relative quantification. With respect to possible diagnostic use, the guanidinium thiocyanate/phenol/chloroform method has the advantages of convenience and producing no organic waste.

To our knowledge, qPCR-MtR is the most convenient, fastest, and least expensive way to reliably measure ms^2 modifications of tRNA^{Lys}(UUU) in crude samples of total RNA. This method could be adapted not only to the clinical assessment of T2D risk but also to biomedical research into the functions of ms^2 modifications in other diseases.

Author Contributions: All authors confirmed they have contributed to the intellectual content of this paper and have met the following 3 requirements: (a) significant contributions to the conception and design, acquisition of data, or analysis and interpretation of data; (b) drafting or revising the article for intellectual content; and (c) final approval of the published article.

Authors' Disclosures or Potential Conflicts of Interest: Upon manuscript submission, all authors completed the author disclosure form. Disclosures and/or potential conflicts of interest:

Employment or Leadership: None declared.

Consultant or Advisory Role: None declared.

Stock Ownership: None declared.

Honoraria: None declared.

Research Funding: F.Y. Wei, Ministry of Health, Labor and Welfare (H24-012) and the Astellas Foundation for Research on Metabolic Disorders; Tsutomu Suzuki, Grant-in-Aid for Scientific Research