

- 65–74 (2010).
- 19) Barbee KA. Mechanical cell injury. *Ann. N.Y. Acad. Sci.*, **1066**, 67–84 (2005).
 - 20) Pettus EH, Christman CW, Giebel ML, Povlishock JT. Traumatically induced altered membrane-permeability—Its relationship to traumatically induced reactive axonal change. *J. Neurotrauma*, **11**, 507–522 (1994).
 - 21) Ellis EF, Mckinney JS, Willoughby KA, Liang S, Povlishock JT. A new model for rapid stretch-induced injury of cells in culture—Characterization of the model using astrocytes. *J. Neurotrauma*, **12**, 325–339 (1995).
 - 22) Geddes DM, Cargill RS 2nd, LaPlaca MC. Mechanical stretch to neurons results in a strain rate and magnitude-dependent increase in plasma membrane permeability. *J. Neurotrauma*, **20**, 1039–1049 (2003).
 - 23) Cullen DK, Vernekar VN, LaPlaca MC. Trauma-induced plasmalemma disruptions in three-dimensional neural cultures are dependent on strain modality and rate. *J. Neurotrauma*, **28**, 2219–2233 (2011).
 - 24) Liu F, Huang L. Noninvasive gene delivery to the liver by mechanical massage. *Hepatology*, **35**, 1314–1319 (2002).

