

1 **Role of Notch signaling in the maintenance of human mesenchymal stem**  
2 **cells under hypoxic conditions**

3 Hiroyuki Moriyama<sup>1,\*</sup>, Mariko Moriyama<sup>1,\*</sup>, Haruki Isshi<sup>1</sup>, Shin Ishihara<sup>1</sup>, Hanayuki Okura<sup>2</sup>, Akihiro Ichinose<sup>3</sup>,  
4 Toshiyuki Ozawa<sup>4</sup>, Akifumi Matsuyama<sup>2</sup> and Takao Hayakawa<sup>1</sup>

5 <sup>1</sup> Pharmaceutical Research and Technology Institute, Kinki University, Higashi-Osaka, Osaka, Japan.

6 <sup>2</sup> Platform of therapeutics for Rare Disease and Health Policy, National Institute of Biomedical Innovation, Kobe,  
7 Hyogo, Japan.

8 <sup>3</sup> Department of Plastic Surgery, Kobe University Hospital, Kobe, Hyogo, Japan.

9 <sup>4</sup> Department of Dermatology, Osaka City University, Graduate School of Medicine, Osaka, Japan.

10  
11 \*These two authors contributed equally to this work.

12  
13 Running title: MESENCHYMAL STEM CELLS UNDER HYPOXIA

14  
15 Address correspondence to:

16 Dr. Hiroyuki Moriyama

17 Pharmaceutical Research and Technology Institute

18 Kinki University

19 3-4-1 Kowakae

20 Higashi-Osaka

21 Osaka 577-8502

22 Japan

23 Tel&Fax: +81-6-4307-4312

24 Email: moriyama@phar.kindai.ac.jp

## 1 Abstract

2 Human adipose tissue-derived multilineage progenitor cells (hADMPCs) are attractive for cell therapy and tissue  
3 engineering because of their multipotency and ease of isolation without serial ethical issues. However, their limited  
4 *in vitro* lifespan in culture systems hinders their therapeutic application. Some somatic stem cells including  
5 hADMPCs are known to be localized in hypoxic regions; thus, hypoxia may be beneficial for *ex vivo* culture of  
6 these stem cells. These cells exhibit a high level of glycolytic metabolism in presence of high oxygen levels and  
7 further increase their glycolysis rate under hypoxia. However, the physiological role of glycolytic activation and its  
8 regulatory mechanisms are still incompletely understood. Here we show that Notch signaling is required for  
9 glycolysis regulation under hypoxic conditions. Our results demonstrate that 5% O<sub>2</sub> dramatically increased the  
10 glycolysis rate, improved the proliferation efficiency, prevented senescence, and maintained the multipotency of  
11 hADMPCs. Intriguingly, these effects were not mediated by hypoxia-inducible factor (HIF), but rather by the Notch  
12 signaling pathway. 5% O<sub>2</sub> significantly increased the level of activated Notch1 and expression of its downstream  
13 gene, *HES1*. Furthermore, 5% O<sub>2</sub> markedly increased glucose consumption and lactate production of hADMPCs,  
14 which decreased back to normoxic levels upon treatment with a  $\gamma$ -secretase inhibitor. We also found that HES1  
15 was involved in induction of GLUT3, TPI, and PGK1 in addition to reduction of TIGAR and SCO2 expression.  
16 These results clearly suggest that Notch signaling regulates glycolysis under hypoxic conditions and thus likely  
17 affects the cell lifespan via glycolysis.

## 1 Introduction

2 Human adipose tissue-derived mesenchymal stem cells (MSCs), also referred to as human adipose tissue–  
3 derived multilineage progenitor cells (hADMPCs), are multipotent stem cells that can differentiate into various  
4 types of cells, including hepatocytes [1], cardiomyoblasts [2], pancreatic cells [3], and neuronal cells [4-6]. They  
5 can be easily and safely obtained from lipoaspirate without posing serious ethical issues and can also be  
6 expanded *ex vivo* under appropriate culture conditions. Moreover, MSCs, including hADMPCs, have the ability to  
7 migrate to injured areas and secrete a wide variety of cytokines and growth factors necessary for tissue  
8 regeneration [7-11]. In addition, because of their hypoimmunogenicity and immunomodulatory effects, hADMPCs  
9 are good candidates as gene delivery vehicles for therapeutic purposes [12]. Thus, hADMPCs are attractive  
10 seeding cells for cell therapy and tissue engineering. However, similar to other somatic stem cells or primary cells,  
11 hADMPCs have limited growth potential and ultimately stop proliferation as a result of cellular senescence [13],  
12 which hinders their therapeutic application.

13 Conversely, embryonic stem cells (ESCs) and induced pluripotent stem cells (iPSCs) are immortal  
14 under standard culture conditions. Recently, several groups have reported that these cells greatly rely on glycolysis  
15 for energy production even under high-oxygen conditions [14-16]. This phenomenon is known as the Warburg  
16 effect and was originally described for cancer cells by Otto Warburg in the 1920s [17]. Although mitochondrial  
17 respiration is more efficient than glycolysis in generating ATP (net yield of 30 ATPs vs. 2 ATPs), glycolysis is able  
18 to produce ATP considerably faster than mitochondrial respiration as long as glucose supplies are adequate. Thus,

1 a metabolic shift from mitochondrial respiration to glycolysis would provide a growth advantage for actively  
2 proliferating cells. Moreover, Kondoh *et al.* demonstrated that enhanced glycolysis is also involved in cellular  
3 immortalization through reduction of intrinsic ROS production [14,18,19]. Because accumulation of intrinsic ROS  
4 levels could be a major reason for replicative senescence [20], enhancing glycolysis in cultured cells might improve  
5 the quality of the cells by suppressing premature senescence. One candidate method for induction of glycolysis is  
6 application of low-oxygen conditions to activate the transcription factor, hypoxia-inducible factor (HIF). HIF-1 is  
7 known to increase the expression of most glycolytic enzymes and the glucose transporters GLUT1 and GLUT3  
8 [20]. Thus, several studies have reported that hypoxia is beneficial for the maintenance of hESCs in a pluripotent  
9 state [21,22]. Moreover, low oxygen tension has been reported to enhance the generation of induced pluripotent  
10 stem cells both from mouse and human primary fibroblasts [23].

11           Recently, hypoxic culture conditions have also been reported to confer a growth advantage, prevent  
12 premature senescence, and maintain undifferentiated states in somatic stem cells, e.g., hematopoietic stem cells  
13 (HSCs) [24], neural stem cells [25], and bone marrow-derived MSCs [26]. These stem cells reside in their local  
14 microenvironments called the “stem cell niche”, where the oxygen tension is relatively low (in the range of 1%–  
15 9%). Thus, hypoxic culture may be beneficial to these stem cells with regard to in vitro proliferation, cell survival,  
16 and differentiation. Takubo *et al.* reported that HSCs activated Pdk through HIF1 $\alpha$  in hypoxic culture conditions,  
17 resulting in maintenance of glycolytic flow and suppression of the influx of glycolytic metabolites into mitochondria,  
18 and this glycolytic metabolic state was shown to be indispensable for the maintenance of HSCs [27]. Several  
19 studies have reported that MSCs exhibit a high level of glycolytic metabolism in the presence of high oxygen

1 levels and further increase their rate of glycolysis upon culture under hypoxia [28,29]. However, a relationship  
2 between beneficial effects of hypoxic conditions and metabolic status in addition to involvement of HIFs in the  
3 metabolic changes has not been investigated in these reports.

4 In this study, we aimed to investigate the effect of 5% oxygen on hADMPCs. Our results demonstrate  
5 that culture under 5% oxygen increased the glycolysis rate, improved the proliferation efficiency, prevented the  
6 cellular senescence, and maintained the undifferentiated status of hADMPCs. Intriguingly, these effects were not  
7 mediated by HIF, but rather by Notch signaling, an important signaling pathway required for the development of  
8 many cell types and maintenance of stem cells [30,31]. 5% oxygen activated Notch signaling, resulting the  
9 upregulation of *SLC2A3*, *TPI*, and *PGK1* in addition to the downregulation of *TIGAR* and *SCO2*, which may  
10 contribute to the increase in the glycolysis rate. These observations thus provide new regulatory mechanisms for  
11 stemness maintenance obtained under 5% oxygen conditions.

## 12 13 14 **Material and Methods**

### 15 **Adipose tissue samples**

16 Subcutaneous adipose tissue samples (10–50 g each) were resected during plastic surgery from 5 female and 2  
17 male patients (age 20–60 years) as discarded tissue. The study protocol was approved by the Review Board for  
18 Human Research of Kobe University Graduate School of Medicine Foundation for Biomedical Research and

1 Innovation, Osaka City University Graduate School of Medicine, and Kinki University Pharmaceutical Research  
2 and Technology Institute (reference number: 12-043). Each subject provided signed informed consent.

#### 4 **Cell culture**

5 hADMPCs were isolated as previously reported [32-35] and maintained in a medium containing 60% DMEM-low  
6 glucose, 40% MCDB-201 medium (Sigma Aldrich, St. Louis, MO, USA), 1× insulin-transferrin-selenium (Life  
7 Technologies, NY, USA), 1 nM dexamethasone (Sigma Aldrich), 100 mM ascorbic acid 2-phosphate (Wako,  
8 Osaka, Japan), 10 ng/mL epidermal growth factor (PeproTech, NJ, USA), and 5% fetal bovine serum. The cells  
9 were plated to a density of  $5 \times 10^3$  cells/cm<sup>2</sup> on fibronectin-coated dishes, and the medium was replaced every 2  
10 days. For hypoxic culture, cells were cultured in a gas mixture composed of 90% N<sub>2</sub>, 5% CO<sub>2</sub>, and 5% O<sub>2</sub>. For  
11 maintenance of the hypoxic gas mixture, a ProOx C21 carbon dioxide and oxygen controller and a C-Chamber  
12 (Biospherix, NY, USA) were used.

#### 14 **Senescence-associated β-galactosidase staining**

15 Cells were fixed with 2% paraformaldehyde/0.2% glutaraldehyde for 5 min at room temperature and then washed 2  
16 times with phosphate-buffered saline (PBS). The cells were then incubated overnight at 37 °C with fresh  
17 senescence-associated β-galactosidase (SA-β-Gal) chromogenic substrate solution (1 mg/mL Bluo-gal (Life

1 Technologies), 40 mM citric acid (pH 6.0), 5 mM potassium ferrocyanide, 5 mM potassium ferricyanide, 150 mM  
2 NaCl, and 2 mM MgCl<sub>2</sub>).

#### 3 4 **Measurement of reactive oxygen species production**

5 Cells were harvested and incubated with 10 μM 5-(and-6)-chloromethyl-2',7'-dichlorodihydrofluorescein diacetate,  
6 acetyl ester (CM-H<sub>2</sub>DCFDA). The amount of intracellular ROS production was proportional to the green  
7 fluorescence, as analyzed using a Guava EasyCyte 8HT flow cytometer (Millipore, MA, USA) using an argon  
8 laser at 488 nm and a 525/30 nm band pass filter, and dead cells were excluded using the Live/Dead Fixable Far  
9 Red Dead Cell Stain Kit (Life Technologies).

#### 10 11 **EdU proliferation assay**

12 For assessment of cell proliferation, hADMPCs were seeded on a fibronectin-coated 6-well plate at a density of 5 ×  
13 10<sup>3</sup> cells/cm<sup>2</sup> and cultured for 3 days. Cell proliferation was detected by incorporating of 5-ethynyl-2'-deoxyuridine  
14 (EdU) and using the Click-iT EdU Alexa Fluor 488 Flow Cytometry Assay Kit (Life Technologies). Briefly, according  
15 to the manufacturer's protocol, cells were incubated with 10 μM EdU for 2 h before fixation, permeabilized, and  
16 stained with EdU. EdU-positive cells were then analyzed using the 488 nm laser of a Guava EasyCyte 8HT flow  
17 cytometer (Millipore).

1

## 2 **Flow cytometry analysis**

3 Flow cytometry analysis was performed as previously described [35]. Briefly, hADMPCs were harvested and  
4 re-suspended in staining buffer (PBS containing 1% BSA, 2 mM EDTA, and 0.01% sodium azide) at a density of  
5  $1 \times 10^6$  cells/mL, incubated for 20 min with a fluorescein isothiocyanate (FITC)-conjugated antibody against CD49b  
6 or CD98 (BioLegend, CA, USA) or a phycoerythrin (PE)-conjugated antibody against CD10, CD13, CD29, CD44,  
7 CD49a, CD49c, CD49d, CD49e, CD51/61, CD73, CD90, CD105, CD117, SSEA4, HLA-A,B,C (BioLegend),  
8 CD133/1 (Miltenyi Biotec, CA, USA), or CD166 (Beckman Coulter, CA, USA). Non-specific staining was assessed  
9 using relevant isotype controls. Dead cells were excluded using the Live/Dead Fixable Far Red Dead Cell Stain Kit  
10 (Life Technologies). FlowJo software was used for quantitative analysis.

11

## 12 **RNA extraction, cDNA generation, and quantitative polymerase chain reaction (Q-PCR)**

13 Total RNA was extracted using the RNeasy Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's  
14 instructions. cDNA was generated from 1  $\mu$ g of total RNA using the Verso cDNA Synthesis Kit (Thermo Scientific,  
15 MA, USA) and purified using the MinElute PCR Purification Kit (Qiagen). Q-PCR analysis was conducted using  
16 the SsoFast EvaGreen supermix (Bio-Rad, CA, USA) according to the manufacturer's protocols. The relative  
17 expression value for each gene was calculated using the  $\Delta\Delta C_t$  method and the most reliable internal control gene



1 was determined using geNorm Software (<http://medgen.ugent.be/~jvdesomp/genorm/>). Details of the primers  
2 used in these experiments are available upon request.

#### 3 4 **Western blot analysis**

5 Whole cell extracts were prepared by washing cells with ice-cold phosphate-buffered saline (PBS) and lysing  
6 them with M-PER Mammalian Protein Extraction Reagent (Thermo Scientific Pierce, IL, USA) according to the  
7 manufacturer' s instructions. Nuclear and cytosolic extracts were prepared as follows. Cells were washed with  
8 ice-cold PBS and lysed with lysis buffer (50 mM Tris-HCl (pH 7.5), 0.5% Triton X-100, 137.5 mM NaCl, 10%  
9 glycerol, 5 mM EDTA, 1 mM sodium vanadate, 50 mM sodium fluoride, 10 mM sodium pyrophosphate, and  
10 protease inhibitor cocktail). Then, insoluble nuclei were isolated by centrifugation and lysed with lysis buffer  
11 containing 0.5% SDS. Equal amounts of proteins were separated by sodium dodecyl sulfate polyacrylamide gel  
12 electrophoresis (SDS-PAGE), transferred to polyvinylidene fluoride (PVDF) membranes (Immobilon-P; Millipore),  
13 and probed with antibodies against cleaved Notch1 (#2421, Cell Signaling Technology, MA, USA), HIF-1 $\alpha$   
14 (#610959, BD Bioscience, CA, USA), hypoxia inducible factor 2 $\alpha$  (MAB3472, Millipore), Akt (#9272, Cell Signaling  
15 Technology), and phospho Akt (Ser473) (#4060, Cell Signaling Technology). Horseradish peroxidase  
16 (HRP)-conjugated anti-mouse or -rabbit IgG antibody (Cell Signaling Technology) was used as a secondary  
17 antibody and immunoreactive bands were visualized using Immobilon Western Chemiluminescent HRP substrate  
18 (Millipore). The band intensity was measured using the ImageJ software.

1

## 2 **Fluorescence microscopy**

3 Phase contrast and fluorescence images were obtained using a fluorescence microscope (BZ-9000; Keyence,  
4 Osaka, Japan) using BZ Analyzer Software (Keyence).

5

## 6 **Adipogenic, osteogenic, and chondrogenic differentiation procedures**

7 For adipogenic differentiation, cells were cultured in differentiation medium (Zen-Bio, NC, USA). After 7 days, half  
8 of the medium was exchanged for adipocyte medium (Zen-Bio) and this was repeated every 3 days. Three weeks  
9 after differentiation, adipogenic differentiation was confirmed by microscopic observation of intracellular lipid  
10 droplets with the aid of Oil Red O staining. Osteogenic differentiation was induced by culturing the cells in  
11 osteocyte differentiation medium (Zen-Bio). Differentiation was examined by Alizarin Red staining. For  
12 chondrogenic differentiation,  $2 \times 10^5$  hADMSCs were centrifuged at  $400 \times g$  for 10 min. The resulting pellets were  
13 cultured in chondrogenic medium (Lonza, Basel, Switzerland) for 21 days. The pellets were fixed with 4%  
14 paraformaldehyde in PBS, embedded in OCT, frozen, and sectioned at  $8 \mu\text{m}$ . The sections were incubated with  
15 PBSMT (PBS containing 0.1% Triton X-100, and 2% skim milk) for 1 h at room temperature, and then incubated  
16 with a mouse monoclonal antibody against type II collagen (Abcam, MA, USA) for 1 h. After washing with PBS,  
17 cells were incubated with Alexa 546-conjugated anti-mouse IgG to identify chondrocytes (Life Technologies). The

1 cells were counterstained with 4'-6-diamidino-2-phenylindole (DAPI) (Life Technologies) to identify cellular nuclei.

2 The sections were also stained with 1% alcian blue (Sigma Aldrich) in 3% acetic acid, pH 2.5 for 30 min.

#### 3 4 **Determination of HK, PFK, LDH, PDH, and Cox IV activities**

5 Cells ( $2 \times 10^6$ ) were lysed, and HK, PFK, LDH, or PDH activity was measured using the Hexokinase Colorimetric

6 Activity Kit, Phosphofructokinase (PFK) Activity Colorimetric Assay Kit, Lactate Dehydrogenase (LDH) Activity

7 Assay Kit, or Pyruvate Dehydrogenase Activity Colorimetric Assay Kit (all from BioVision, Milpitas, CA, USA)

8 respectively, according to the manufacturer's instructions. To measure Cox IV activity, mitochondria were isolated

9 from  $2 \times 10^7$  cells using a Mitochondria Isolation Kit (Thermo Scientific) and lysed with buffer containing n-Dodecyl

10  $\beta$ -D-maltoside, followed by measurement with the Mitochondria Activity Assay (Cytochrome C Oxidase Activity

11 Assay) Kit (BioChain Institute, Newark, CA, USA), according to the manufacturer's instructions.

## 12 13 **Results**

### 14 **5% Oxygen hypoxic culture condition increases proliferation capacity and decreases senescence**

15 hADMPCs were cultured under 20% oxygen (normoxia; Nx) or 5% oxygen (hypoxia; Hx) and their proliferation

16 capacities were examined based on the relationship between the number of cultivation days and the population

1 doubling level (PDL). Nx-cultured hADMPs ceased proliferation at a PDL of 35–40 (between 46–70 days),  
2 whereas continuous cell proliferation beyond 45 PDL was observed when hADMPs were cultured in the Hx  
3 condition (Figure 1A). To investigate whether this increase of PDL in the Hx culture condition resulted from an  
4 increase in cell cycle progression and increase in survival rates, 5-ethynyl-2'-deoxyuridine (EdU), an alternative to  
5 5-bromo-2'-deoxyuridine (BrdU), was incorporated into the genomic DNA of the hADMPs, and the amount of  
6 incorporated EdU was quantified by flow cytometry. As shown in Figure 1B, the EdU incorporation rate was  
7 significantly higher in Hx-cultured hADMPs than in Nx-cultured hADMPs, suggesting that cell growth was  
8 increased in the Hx culture condition. In addition, measurement of DNA content in hADMPs revealed a slight but  
9 significant decrease of sub-G1 peaks, which indicates the existence of apoptotic cells with degraded DNA, when  
10 the cells were cultured in the Hx condition (Figure 1C). These data suggest that the Hx culture condition increases  
11 the proliferation capacity of hADMPs by promoting their cell growth and survival rates. We also found that  
12 Nx-cultured hADMPs were larger with a more irregular shape (Figure 1D), which suggests that the Hx culture  
13 condition prevented hADMPs from entering senescence [36]. To further investigate this phenomenon, cellular  
14 senescence was measured by staining for senescence-associated  $\beta$ -galactosidase (SA- $\beta$ -Gal), which revealed  
15 that SA- $\beta$ -Gal activity was increased in Nx-cultured hADMPs at passage 17 (Figure 1E). As it has been  
16 hypothesized that senescence results from oxidative stress [20], accumulation of reactive oxygen species (ROS)  
17 in hADMPs was detected using the non-fluorescent probe, CM-H<sub>2</sub>DCFDA. Flow cytometry analysis revealed  
18 that ROS were generated at higher levels in hADMPs when cultured in the Nx condition (Figure 1F), suggesting  
19 that reduced production of ROS in the Hx condition may prevent the cells from entering replicative senescence.

## **Hypoxic culture maintains some mesenchymal stem cell properties and increases differentiation**

We then examined the cell properties of hADMPCs under Nx and Hx conditions. Initially, cell surface antigens expressed on hADMPCs were analyzed by flow cytometry. No significant difference in expression profile between hADMPCs cultured in Nx and Hx was observed; the cells were consistently positive for CD10, CD13, CD29, CD44, CD49a, CD49b, CD49c, CD49d, CD49e, CD51/61, CD54, CD59, CD73, CD90, CD98, CD105, CD166, and HLA-A, B, C, but negative for CD34, CD45, CD117, CD133 (Figure 2 and data not shown). These data were consistent with previous reports describing the expression profiles of cell surface markers of hMSCs [37,38]. To further examine the stem cell properties of hADMPCs, their potential for differentiation into adipocyte, osteocyte, and chondrocyte lineages was analyzed at passage 8. Hx-cultured hADMPCs presented enhanced differentiation into various lineages (Figure 3A and B), indicating that the Hx culture condition improved the stem cell properties of hADMPCs.

## **Hypoxic culture condition activates Notch signaling**

To reveal the molecular mechanism by which the Hx culture condition increased the proliferative capacity and maintaining the stem cell properties of hADMPCs, we next examined Notch signaling, which is required for maintaining stem-cell features of various types of stem cells [30,31]. As expected, levels of cleaved NOTCH1, an

1 activated form of NOTCH1, were significantly increased (> 2-fold) in the Hx culture condition (Figure 4A).  
2 Quantitative PCR (Q-PCR) analysis revealed that HES1, a downstream target of Notch signaling, was  
3 upregulated in Hx-cultured hADMPCs, which also indicated that Notch signaling was activated in the Hx culture  
4 condition (Figure 4B). Administration of the  $\gamma$ -secretase inhibitor DAPT at 1  $\mu$ M, which was sufficient to inhibit the  
5 proteolytic cleavage of NOTCH1 (Figure 4A), decreased the Hx-induced expression of HES1 at both mRNA and  
6 protein levels (Figure 4B and C). These data indicate that Hx increased the expression of HES1 through activation  
7 of Notch signaling. It has been reported that Notch signaling and hypoxia-inducible factor (HIF) undergo cross talk  
8 in hypoxic cells [39-42]. Therefore, HIF-1 $\alpha$  and HIF-2 $\alpha$  protein levels in hADMPCs were analyzed by western  
9 blotting. HIF-1 $\alpha$  was stabilized when a chemical hypoxia-mimicking agent, cobalt chloride, was applied in the  
10 culture, whereas no obvious increase of HIF-1 $\alpha$  was observed in the Hx culture condition (Figure 4D). However,  
11 we did not detect any HIF-2 $\alpha$  expression even in the presence of cobalt chloride (Figure 4E). Q-PCR analysis  
12 revealed that *HIF2A* mRNA was not expressed in these cells (data not shown). From these results, we concluded  
13 that neither HIF-1 $\alpha$  nor HIF-2 $\alpha$  was involved in the Hx-induced increase in the proliferative capacity and stem cell  
14 properties of hADMPCs.

15 To identify the signaling responsible for the observed effect, we next examined the Akt, NF- $\kappa$ B, and  
16 p53 signaling pathways. It has been reported that hypoxic conditions induce the activation of Akt and NF- $\kappa$ B  
17 signaling [43,44]. In addition, hypoxic conditions have been shown to inhibit p53 activity [45], and crosstalk  
18 between these pathways and Notch signaling has also been demonstrated [42,46-48]. As shown in Figure 4F, the

1 Hx condition increased Akt phosphorylation, which was not decreased by DAPT treatment. These data  
2 demonstrate that 5% oxygen activated Akt signaling but not via Notch signaling. Similarly, the hypoxic condition  
3 induced nuclear accumulation of p65, which was inhibited by DAPT treatment (Figure 4G). These data suggest  
4 that NF- $\kappa$ B signaling is regulated by Notch signaling in hADMPCs. Furthermore, p53 was not activated under the  
5 5% oxygen condition as assessed by detection of phospho-p53 and a p53 reporter assay. However, DAPT  
6 treatment significantly increased p53 activity (Figure 4H and I).

7

#### 8 **Notch signaling is indispensable for acquisition of the advantageous properties of hADMPCs**

9 We next examined the roles of Notch signaling in the proliferative capacity and stem cell properties of hADMPCs  
10 in the Hx culture condition. To inhibit Notch signaling, DAPT was added to the medium at a final concentration of 1  
11  $\mu$ M. DAPT treatment significantly decreased the PDL when hADMPCs were cultured under either 20% or 5%  
12 oxygen (Figure 5A). Intriguingly, measurement of the DNA content in hADMPCs revealed that inhibition of Notch  
13 signaling by 1  $\mu$ M DAPT significantly attenuated the decrease in apoptotic cells in the Hx condition (Figure 5B).  
14 These data suggest that 5% oxygen increases the proliferation capacity of hADMPCs through Notch signaling by  
15 promoting their survival. To examine whether Notch signaling affects the stem cell properties of hADMPCs,  
16 differentiation into adipocyte, osteocyte, and chondrocyte lineages was analyzed at passage 8. Hx-cultured  
17 hADMPCs underwent greater differentiation into all lineages as described in Figure 3, whereas application of a  
18 Notch inhibitor significantly decreased the differentiation capacity to all lineages (Figure 5C-E). In addition,

1 SA- $\beta$ -Gal staining revealed that inhibition of Notch signaling by DAPT remarkably promoted senescence in both  
2 the Nx and Hx culture conditions, suggesting that the suppression of replicative senescence observed in the Hx  
3 condition is mediated by Notch signaling.

#### 5 **Glycolysis is enhanced in the 5% oxygen condition through Notch signaling**

6 Recent studies suggest that the metabolic shift from aerobic mitochondrial respiration to glycolysis extends the life  
7 span possibly via reduction of intrinsic ROS production [18,19]. Our results demonstrate that the 5% oxygen  
8 condition reduced ROS accumulation in hADMPs (Figure 1F). In addition, the relationship between Notch  
9 signaling and glycolysis has been recently established [49,50]. We therefore considered glycolytic flux by  
10 measuring the glucose consumption and lactate production of hADMPs in the Nx or Hx culture conditions. As  
11 shown in Figure 6A, glucose consumption and lactate production were elevated in the Hx culture condition,  
12 indicating that a metabolic shift to glycolysis occurred when hADMPs were cultured in 5% oxygen. In contrast,  
13 the Notch inhibitor DAPT markedly reduced glycolytic flux as assessed by glucose consumption and lactate  
14 production (Figure 6A). To identify the genes responsible for the glycolytic change, we performed a Q-PCR  
15 analysis. As shown in Figure 6B, *SLC2A3*, *TPI*, and *PGK1*, encoding glycolytic enzymes, were upregulated in the  
16 5% oxygen condition, whereas these genes were suppressed by DAPT treatment. Interestingly, Hes1 transduction  
17 by an adenoviral vector markedly induced the mRNA expression of the same genes (Figure 6E). In addition,  
18 *SCO2*, a positive modulator of aerobic respiration, and *TIGAR*, a negative regulator of glycolysis, were



1 transcriptionally downregulated in the 5% oxygen condition, whereas DAPT treatment partially restored the  
2 expression level (Figure 6B). Adenoviral expression of Hes1 dramatically reduced *SCO2* and *TIGAR* expression  
3 (Figure 6E), which suggests that the Notch-Hes1 signaling modulates the metabolic pathway. We also measured  
4 the activities of key enzymes in glycolysis. Hexokinase activity was not changed under hypoxic conditions;  
5 however, phosphofructokinase and lactate dehydrogenase were activated in 5% oxygen condition, which was  
6 attenuated by Notch inhibition (Figure 6C). In addition, pyruvate dehydrogenase (PDH) and cytochrome c oxidase  
7 (Complex IV) activity assays showed that mitochondrial respiration decreased under the hypoxic condition and that  
8 DAPT treatment restored it (Figure 6D). Moreover, glycolytic flux in Hes1-expressing hADMPCs was positively  
9 correlated with the expression of these glycolytic genes as assessed by glucose consumption and lactate  
10 production (Figure 6F). In contrast, HES1 knockdown by adenoviral transduction of *HES1* RNAi resulted in a  
11 significant reduction of glycolytic flux (Figure 6G), demonstrating that HES1 is involved in the regulation of  
12 glycolysis.

13

#### 14 **Glycolysis supports the proliferation of hADMPCs**

15 To determine whether aerobic glycolysis is important for the proliferation of hADMPCs, hADMPCs were treated  
16 with the glycolytic inhibitor 2-deoxy-D-glucose (2-DG) or the respiration inhibitor sodium azide ( $\text{NaN}_3$ ). We found  
17 that hADMPCs were sensitive to treatment with 2-DG even at the low concentration of 0.2 mM (Figure 7A). In  
18 contrast, treatment of hADMPCs with  $\text{NaN}_3$ , rather increased their proliferation at the concentration of 1 mM and

1 supported their proliferation even at the concentration of 5 mM (Figure 7B). These data suggest that the  
2 proliferation of hADMPCs is compromised when aerobic glycolysis is blocked.

## 3 4 5 **Discussion**

6 Recent evidence suggests that hypoxic culture conditions confer a growth advantage, prevent premature  
7 senescence, and maintain undifferentiated states in ESCs, iPSCs, and somatic stem cells. However, the  
8 molecular mechanism underlying the beneficial effects of culturing these cells at low oxygen conditions remains  
9 unclear. Our findings prompted us to hypothesize that Notch signaling in physiological hypoxic conditions (5% O<sub>2</sub>)  
10 contributes to these effects on hADMPCs by modulating glycolytic flux.

11 We found that 5% O<sub>2</sub> significantly increased the proliferation capacity, decreased apoptosis, and  
12 inhibited senescence of hADMPCs (Figure 1). Moreover, 5% O<sub>2</sub> improved the differentiation of hADMPCs without  
13 affecting the expression of their cell surface markers (Figures 2 and 3). Welford et al. reported that HIF-1 $\alpha$  delays  
14 premature senescence of mouse embryonic fibroblasts under hypoxic conditions (2% O<sub>2</sub>) [51]. Tsai et al. reported  
15 that hypoxia (1% O<sub>2</sub>) inhibits senescence and maintains mesenchymal stem cell properties through accumulation  
16 of HIF-1 $\alpha$  [26]. Hypoxia was recently reported to enhance the undifferentiated status and stem cell properties in  
17 various stem and precursor cell populations via the interaction of HIF with the Notch intracellular domain to  
18 activate Notch-responsive promoters [39]. In the current study, the effects observed in 5% O<sub>2</sub> condition were

1 independent of HIF proteins because accumulation of HIF-1 $\alpha$  and HIF-2 $\alpha$  was not observed (Figure 4). Instead,  
2 our findings suggest that 5% O<sub>2</sub> activated Notch signaling, which contributed advantageous effects of hypoxic  
3 culture on hADMPCs. A pharmacological inhibitor of Notch signaling, DAPT, abrogated the hypoxic-induced  
4 Notch activation, increased proliferation capacity and lifespan, maintenance of stem cell properties, and  
5 prevention of senescence (Figure 4 and 5). Moreover, we also found that 5% O<sub>2</sub> enhanced glucose consumption  
6 and lactate production, and these effects were also attenuated by Notch inhibition (Figure 6A) and knockdown of  
7 HES1 (Figure 6G). Previously, it has been reported that Notch signaling promotes glycolysis by activating the  
8 PI(3)K-Akt pathway [49,50]. However, our results indicate that Akt signaling was not activated by Notch signaling  
9 because DAPT did not attenuate hypoxia-induced Akt phosphorylation (Figure 4F). Although Akt is unlikely to be  
10 regulated by Notch signaling in hADMPCs, it is clear in our data that Akt signaling was activated by 5% O<sub>2</sub>.  
11 Therefore, we could not rule out the possibility that the promotion of glycolysis in the 5% O<sub>2</sub> condition was caused  
12 by Akt signaling.

13           Recent evidence suggests that Notch signaling acts as a metabolic switch [49,52]. Zhou et al.  
14 demonstrated that hairy, a basic helix-loop-helix transcriptional repressor regulated by Notch signaling, was  
15 up-regulated and genes encoding metabolic enzymes, including TCA cycle enzymes and respiratory chain  
16 complexes, were down-regulated in hypoxia-tolerant flies. Intriguingly, they also found that hairy-binding elements  
17 were present in the regulatory region of the down-regulated metabolic genes. Their work thus provides new  
18 evidence that hairy acts as a metabolic switch [52]. Landor et al. demonstrated that both hyper- and hypoactive

1 Notch signaling induced glycolysis, albeit by different mechanisms. They showed that Notch activation increased  
2 glycolysis through activation of PI3K-AKT signaling, whereas decreased Notch activity inhibited mitochondrial  
3 function in a p53-dependent manner in MCF7 breast cancer cell lines [49]. Consistent with their reports, our  
4 findings that Notch signaling promoted activity of some glycolysis enzymes and inhibited mitochondrial activity  
5 (Figure 6) also suggest that Notch signaling functioned as a metabolic switch. While our data showed that Notch  
6 inhibition by DAPT resulted in reduced glycolysis (Figure 6A-C), induction of mitochondrial function (Figure 6D)  
7 and activation of p53 (Figure 4H and I) are not consistent with the report of Landor et al. This contradiction might  
8 be explained by the expression level of endogenous Notch. Landor et al. showed that in breast cancer  
9 MDA-M-231 cells, which showed higher endogenous Notch activity, high glucose uptake, and lactate production  
10 than MCF7 breast cancer cell lines, Notch inhibition by DAPT significantly reduced glucose consumption and  
11 lactate production [49]. As shown in Figure 4A, we observed that hADMPCs in 5% O<sub>2</sub> displayed high Notch  
12 activity. Moreover, the lactate-to-glucose ratio was 1.8–1.9 in hADMPCs, suggesting that hADMPCs largely rely  
13 on glycolysis for energy production (Figure 6A). In addition, it was reported that hMSCs showed a higher glycolytic  
14 rate than primary human fibroblast [53]. It appears that hADMPCs cultured under hypoxic conditions might  
15 possess cell properties similar to MDA-M-231 cells or MCF7 cells, in which stable expression of constructs  
16 NICD1-GFP produces high Notch activity.

17 Nuclear translocation of p65 was observed in hypoxic conditions, demonstrating that NF- $\kappa$ B is a direct  
18 target of Notch signaling (Figure 4G). Intriguingly, the hypoxic culture conditions in this study upregulated several