

each finger joint. *Rheumatology* 52:523-8, 2013

7. Oku K, Amengual O, Zigon P, Horita T, Yasuda S, Atsumi T. Essential role of the p38 mitogen-activated protein kinase pathway in the tissue factor gene expression by the phosphatidylserine-dependent antiprothrombin antibody. *Rheumatol* 52: 1775-84, 2013

8. Kato M, Atsumi T, Oku K, Amengual O, Nakagawa H, Fujieda Y, Otomo K, Horita T, Yasuda S, Koike T. The involvement of CD36 in the monocyte activation by antiphospholipid antibodies. *Lupus* 22:761-71, 2013

9. Ishizu A, Tomaru U, Murai T, Yamamoto T, Atsumi T, Yoshiki T, Yumura W, Yamagata K, Yamada H, Kumagai S, Kurokawa MS, Suka M, Makino H, Ozaki S. Prediction of response to treatment by gene expression profiling of peripheral blood in patients with microscopic polyangiitis. *PLoS One* 8. e63182, 2013.

10. Nakazawa D, Shida H, Tomaru U, Yoshida M, Nishio S, Atsumi T, Ishizu A. Enhanced formation and disordered regulation of NETs in MPO-ANCA-associated microscopic polyangiitis. *J Am Soc Nephrol* (in press)

11. Jin H, Arase N, Hirayasu K, Kohyama M, Suenaga T, Saito F, Tanimura K, Matsuoka K, Ebina K, Shid K, Toyama-Sorimachi N, Yasuda S, Horita T, Hiwa R, Takasugi K, Ohmura K, Yoshikawa H, Saito T, Atsumi T, Sasazuki T, Katayama I, Lanier LL, Arase H. Autoantibodies to IgG/HLA-DR complexes are associated with rheumatoid arthritis susceptibility. *Proc Natl Acad Sci U S A* (in press)

12. Kiyohara C, Washio M, Horiuchi T, Asami T, Ide S, Atsumi T, Kobashi G, Takahashi H, Tada Y. The modifying effect of NAT2 genotype on the association between systemic lupus erythematosus and consumption of alcohol and caffeine-rich beverages. *Arthritis Care Res* (in press).

13. Takahashi H, Washio M, Kiyohara C, Tada Y, Asami T, Ide S, Atsumi T, Kobashi G, Yamamoto M, Horiuchi T. Psychological stress in a Japanese population with systemic lupus erythematosus: Finding from KYSS study. *Mod Rheumatol* (in press).

14. Fukae J, Tanimura K, Atsumi T, Koike T. Sonographic synovial vascularity of synovitis in rheumatoid arthritis. *Rheumatology* (in press).

15. Amengual O, Horita T, Binder W, Norman GL, Shums Z, Kato M, Otomo K, Fujieda Y, Oku K, Bohgaki T, Yasuda S, Atsumi T. Comparative analysis of different enzyme immunoassays for assessment of phosphatidylserine-dependent antiprothrombin antibodies. *Rheumatology International* (in press)

16. Kurita T, Yasuda S, Oba K, Odani T, Kono M, Otomo K, Fujieda Y, Oku K, Bohgaki T, Amengual O, Horita T, Atsumi T. The efficacy of tacrolimus in patients with interstitial lung diseases complicated with polymyositis or dermatomyositis. *Rheumatology (Oxford)* (in press)

H. 知的財産権の出願・登録状況
(予定を含む。)

4. 特許取得
なし
5. 実用新案登録
なし
6. その他
なし

別紙4

研究成果の刊行に関する一覧表

雑誌

| 発表者氏名 | 論文タイトル名 | 発表誌名 | 巻号 | ページ | 出版年 |
|--|---|---|-----|-----------|-------------|
| Kitaori T, Sugiura-Ogasawara M, Oku K, Papisch W, Ozaki Y, Katano K, Atsumi T. | Determination of clinically significant tests for antiphospholipid antibodies and cutoff levels for obstetric APS. | submitted | | | |
| Sugiura-Ogasawara M, Ozaki Y, Suzumori N. | Management of recurrent miscarriage. invited review J | J Obstet Gynecol Res | | | in press |
| Sugiura-Ogasawara M, Suzuki S, Ozaki Y, Katano K, Suzumori N, Kitaori T. | Frequency of recurrent spontaneous abortion and its influence on further marital relationship and illness: The Okazaki Cohort Study in Japan. | J Obstet Gynaecol Res. | 39 | 126-31 | 2013 |
| Hayashi Y, Sasaki H, Suzuki S, Nishiyama T, Kitaori T, Mizutani E, Suzumori N, Sugiura-Ogasawara M. | Genotyping analyses for polymorphisms of ANXA5 gene in patients with recurrent pregnancy loss. | Fertil Steril | 100 | 1018-1024 | 2013 |
| Nakano Y, Akechi T, Furukawa T, Sugiura-Ogasawara M. | Cognitive behavior therapy for psychological distress in patients with recurrent miscarriage. | Psychology Research and Behavior Management | 6 | 37-43 | 2013 |
| Katano K, Suzuki S, Ozaki Y, Suzumori N, Kitaori T, Sugiura-Ogasawara M. | Peripheral natural killer cell activity as a predictor of recurrent pregnancy loss: a large cohort study. | Fertil Steril | 100 | 1629-34 | 2013 |
| Sugiura-Ogasawara M, Ozaki Y, Suzumori N. | Müllerian anomalies and recurrent miscarriage. | Current Opinion in Obstetrics and Gynecology | 25 | 293-298 | 2013 |
| Sugiura-Ogasawara M, Ozaki Y, Katano K, Suzumori N, Kitaori T, Mizutani E. | Abnormal embryonic karyotype is the most frequent cause of recurrent miscarriage. | Hum Reprod | 27 | 2297-2303 | 2012 |
| Souri M, Sugiura-Ogasawara M, Saito S, Kemkes-Matthes B, Meijers JC, Ichinose A. | Increase in the plasma levels of protein Z-dependent protease inhibitor in normal pregnancies but not in non-pregnant patients with unexplained recurrent miscarriage. | Thromb Haemost | 107 | 507-512 | 2012 |
| Suzumori N, Obayashi S, Kumagai K, Goto S, Yoshida A, Sugiura-Ogasawara M. | A case of microangiopathic antiphospholipid-associated syndromes (MAPS) during pregnancy- review of the literature. | Case report in Medicine | | 827-543 | 2012 |
| Mizutani E, Suzumori N, Ozaki Y, Oseto K, Yamada-Namikawa C, Nakanishi M, Sugiura-Ogasawara M. | SYCP3 mutation may not be associated with recurrent miscarriage caused by aneuploidy. | Hum Reprod | 26 | 1259-1266 | 2011 |

| | | | | | |
|---|---|----------------------------|----|---------|----------|
| Sugiura-Ogasawara, M, Ozaki Y, Katano K, Suzumori N, Mizutani E. | Uterine Anomaly and Recurrent Pregnancy Loss. | Semin Reprod Med | 29 | 514-521 | 2011 |
| Hirshfeld-Cytron J, Sugiura-Ogasawara M, Stephenson MD.. | Management of Recurrent Pregnancy Loss Associated with a Parental Carrier of a Reciprocal Translocation: A Systematic Review. | Semin Reprod Med | 29 | 470-481 | 2011 |
| Fukae J, Tanimura K, Atsumi T, Koike T. | Sonographic synovial vascularity of synovitis in rheumatoid arthritis. | Rheumatology | | | |
| Jin H, Arase N, Hirayasu K, Kohyama M, Suenaga T, Saito F, Tanimura K, Matsuoka K, Ebina K, Shid K, Toyama-Sorimachi N, Yasuda S, Horita T, Hiwa R, Takasugi K, Ohmura K, Yoshikawa H, Saito T, Atsumi T, Sasazuki T, Katayama I, Lanier LL, Arase H. | Autoantibodies to IgG/HLA-DR complexes are associated with rheumatoid arthritis susceptibility. | Proc Natl Acad Sci U S A | | | in press |
| Kiyohara C, Washio M, Horiuchi T, Asami T, Ide S, Atsumi T, Kobashi G, Takahashi H, Tada Y. | The modifying effect of NAT2 genotype on the association between systemic lupus erythematosus and consumption of alcohol and caffeine-rich beverages. | Arthritis Care Res | | | in press |
| Takahashi H, Washio M, Kiyohara C, Tada Y, Asami T, Ide S, Atsumi T, Kobashi G, Yamamoto M, Horiuchi T. | Psychological stress in a Japanese population with systemic lupus erythematosus: Finding from KYSS study. | Mod Rheumatol | | | in press |
| Nakazawa D, Shida H, Tomaru U, Yoshida M, Nishio S, Atsumi T, Ishizu A. | Enhanced formation and disordered regulation of NETs in MPO-ANCA-associated microscopic polyangiitis. | J Am Soc Nephrol | | | in press |
| Amengual O, Horita T, Binder W, Norman GL, Shums Z, Kato M, Otomo K, Fujieda Y, Oku K, Bohgaki T, Yasuda S, Atsumi T. | Comparative analysis of different enzyme immunoassays for assessment of phosphatidylserine-dependent antiprothrombin antibodies. | Rheumatology International | | | in press |
| Kurita T, Yasuda S, Oba K, Odani T, Kono M, Otomo K, Fujieda Y, Oku K, Bohgaki T, Amengual O, Horita T, Atsumi T. | The efficacy of tacrolimus in patients with interstitial lung diseases complicated with polymyositis or dermatomyositis. | Rheumatology (Oxford) | | | in press |
| Kameda H, Kanbe K, Sato E, Ueki Y, Saito K, Nagaoka S, Hidaka T, Atsumi T, Tsukano M, Kasama T, Shiozawa S, Tanaka Y, Yamanaka H, Takeuchi T. | A merged presentation of clinical and radiographic data using probability plots in a clinical trial, the JESMR study. | Ann Rheum Dis | 72 | 310-312 | 2013 |
| Amengual O, Atsumi T, Oku K, Suzuki E, Horita T, Yasuda S, Koike T. | Phospholipid scramblase 1 expression is enhanced in patients with antiphospholipid syndrome. | Mod Rheumatol | 23 | 81-88 | 2013 |

| | | | | | |
|--|--|---------------|----|-----------|------|
| Harigai M, Takamura A, Atsumi T, Dohi M, Hirata S, Kameda H, Nagasawa H, Seto Y, Koike T, Miyasaka N. | Elevation of KL-6 serum levels in clinical trials of tumor necrosis factor inhibitors in patients with rheumatoid arthritis: a report from the Japan College of Rheumatology Ad Hoc Committee for Safety of Biological DMARDs. | Mod Rheumatol | 23 | 278-296 | 2013 |
| Takamura A, Hirata S, Nagasawa H, Kameda H, Seto Y, Atsumi T, Dohi M, Koike T, Miyasaka N, Harigai M. | A retrospective study of serum KL-6 levels during treatment with biological disease-modifying antirheumatic drugs in rheumatoid arthritis patients: a report from the Ad Hoc Committee for Safety of Biological DMARDs of the Japan College of Rheumatology. | Mod Rheumatol | 23 | 297-303 | 2013 |
| Fukaya S, Matsui Y, Tomaru U, Kawakami A, Sogo S, Bohgaki T, Atsumi T, Koike T, Kasahara M, Ishizu A. | Overexpression of TNF- α -converting enzyme in fibroblasts augments dermal fibrosis after inflammation. | Lab Invest | 93 | 72-80 | 2013 |
| Fukae J, Isobe M, Kitano A, Henmi M, Sakamoto F, Narita A, Ito T, Mitsuzaki A, Shimizu M, Tanimura K, Matsuhashi M, Kamishima T, Atsumi T, Koike T | Positive synovial vascularity in patients with low disease activity indicates smouldering inflammation leading to joint damage in rheumatoid arthritis: time-integrated joint inflammation estimated by synovial vascularity in each finger joint. | Rheumatology | 52 | 523-528 | 2013 |
| Oku K, Amengual O, Zigon P, Horita T, Yasuda S, Atsumi T. | Essential role of the p38 mitogen-activated protein kinase pathway in the tissue factor gene expression by the phosphatidylserine-dependent antiprothrombin antibody. | Rheumatol | 52 | 1775-1784 | 2013 |
| Kato M, Atsumi T, Oku K, Amengual O, Nakagawa H, Fujieda Y, Otomo K, Horita T, Yasuda S, Koike T. | The involvement of CD36 in the monocyte activation by antiphospholipid antibodies. | Lupus | 22 | 761-771 | 2013 |
| Ishizu A, Tomaru U, Murai T, Yamamoto T, Atsumi T, Yoshiki T, Yumura W, Yamagata K, Yamada H, Kumagai S, Kurokawa MS, Suka M, Makino H, Ozaki S. | Prediction of response to treatment by gene expression profiling of peripheral blood in patients with microscopic polyangiitis. | PLoS One | 8 | e63182 | 2013 |

Genotyping analyses for polymorphisms of ANXA5 gene in patients with recurrent pregnancy loss

Yuko Hayashi, M.D.,^a Hidefumi Sasaki, M.D.,^b Sadao Suzuki, M.D.,^c Takeshi Nishiyama, M.D.,^c Tamao Kitaori, M.D.,^a Eita Mizutani, M.D.,^a Nobuhiro Suzumori, M.D.,^a and Mayumi Sugiura-Ogasawara, M.D.^a

^a Department of Obstetrics and Gynecology, ^b Department of Oncology, Immunology, and Surgery, and ^c Public Health, Graduate School of Medical Sciences, Nagoya City University, Nagoya, Japan

Objective: To investigate whether polymorphisms at the promoter or 5'-untranslated region of annexin A5 gene (ANXA5) influence miscarriage.

Design: Case-control study and nested case-control study.

Setting: Hospitals.

Patient(s): A total of 264 patients with two to nine recurrent pregnancy losses (RPLs) and 195 fertile control subjects.

Intervention(s): None.

Main Outcome Measure(s): The frequency of six single-nucleotide polymorphisms (SNPs) of the ANXA5 gene in RPL patients versus control subjects, and subsequent live birth rate with and without risk alleles in RPL patients.

Result(s): The minor allele was significantly more frequent in RPL patients than in control subjects for SNP5 (rs1050606). The live birth rates of patients with and without risk alleles of SNP5 were 84.0% and 84.3%, respectively, after excluding cases with abnormal embryonic karyotype, with no significant difference.

Conclusion(s): The variations with the ANXA5 gene upstream region, especially SNP5, were confirmed to be risk factors of RPL. However, presence/absence of the ANXA5 risk allele did not have any predictive effect for subsequent pregnancy outcome. This was the first study indicating the influence of ANXA5 SNPs for pregnancy outcome. (Fertil Steril® 2013;100:1018–24. ©2013 by American Society for Reproductive Medicine.)

Key Words: Annexin A5, single-nucleotide polymorphisms, recurrent pregnancy loss, cohort

Discuss: You can discuss this article with its authors and with other ASRM members at <http://fertilityforum.com/hayashiy-genotyping-polymorphisms-recurrent-pregnancy-loss/>



Use your smartphone to scan this QR code and connect to the discussion forum for this article now.*

* Download a free QR code scanner by searching for "QR scanner" in your smartphone's app store or app marketplace.

Established causes of recurrent pregnancy loss (RPL) include antiphospholipid antibody syndrome (APS), uterine anomalies, and

abnormal chromosomes, particularly translocations, in either partner (1–3). However, according to earlier reports, in about one-half of the cases seen at

research centers, the cause remains unexplained despite conventional examinations conducted to identify the cause (4, 5). Recently, we found that an abnormal embryonic karyotype was the most frequent cause of RPL, accounting for as many as 41% of the cases (6).

APS, or acquired thrombophilia, is the only treatable cause of RPL, with combined low-dose aspirin and heparin treatment having been shown to improve the live birth rate in patients with APS (7, 8). Heritable thrombophilia has been reported to be the cause in a majority of other cases of pregnancy loss of uncertain cause

Received February 21, 2013; revised June 3, 2013; accepted June 10, 2013; published online July 11, 2013.

Y.H. has nothing to disclose. H.S. has nothing to disclose. S.S. has nothing to disclose. T.N. has nothing to disclose. T.K. has received payment for manuscript preparation from the Japan Society of Thrombosis Hemostasis. E.M. has nothing to disclose. N.S. has nothing to disclose. M.S.-O. has received payment for lectures from Bayer, Sanofi-Aventis, Eisai Co., Kissei Pharmaceutical Co., Nippon Shinyaku Co., Dainippon Sumitomo Pharma, Takeda Pharmaceutical Co., Fuji Pharma Co., and Mochida Pharmaceutical Co.; for manuscript preparation from Igaku Shoin Co. and Chunchi Shinbun; and for travel expenses from the Korean Society for Reproductive Medicine.

Supported by a grant-in-aid for Scientific Research from the Ministry of Health, Labor, and Welfare and the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

Reprint requests: Mayumi Sugiura-Ogasawara, M.D., School of Medical Sciences, Obstetrics and Gynecology, Nagoya City University, Kawasumi-1, Mizuho-ku Nagoya, Aichi 4678601, Japan (E-mail: og.mym@med.nagoya-cu.ac.jp).

Fertility and Sterility® Vol. 100, No. 4, October 2013 0015-0282/\$36.00

Copyright ©2013 American Society for Reproductive Medicine, Published by Elsevier Inc. <http://dx.doi.org/10.1016/j.fertnstert.2013.06.020>

(9, 10). However, no association has been established between hereditary thrombophilia and RPL (11).

Annexin A5 (placental anticoagulant protein, encoded by the *ANXA5* gene) normally occurs on the placental villi, and its expression appears to be decreased in the presence of antiphospholipid antibodies (aPL) (12). Annexin A5 functions as an inhibitor of coagulation by its ability to bind to the anionic phospholipids exposed on the surface of platelets (13). It is abundantly expressed in the normal placenta (14), and the encoding gene possesses a complex promoter region that is subject to intricate regulation (15). It has been shown that single-nucleotide polymorphisms (SNPs) in the promoter region of the *ANXA5* gene are significantly associated with RPL, and that women with the M2 haplotype have a 2.42-fold higher risk of pregnancy loss than noncarriers (16). More recently, it has been shown that an SNP in the 5'-untranslated region of the *ANXA5* gene (SNP5: -302T>G) serves as a major risk determinant of RPL in Japanese women (17).

In the present study, we further investigated six *ANXA5* gene SNPs in the upstream region in a cohort study of Japanese women with RPL and merged the findings with previously published reports (17) for univariate analyses. Because data on the combined effect of maternal age and genetic risk factors are still lacking, we also conducted a cohort study to examine, in relation to several factors, whether a subsequent pregnancy would result in further loss or normal delivery. This was the first cohort study to investigate *ANXA5* gene polymorphism as a molecular marker of RPL.

MATERIALS AND METHODS

Patients

We analyzed the data of 264 Japanese women with a history of unexplained RPL (defined as a history of two or more pregnancy losses) who were recruited from Nagoya City University Hospital from June 2007 to November 2012. All patients underwent systematic examination, including hysterosalpingography, chromosome analysis of both partners, determination of aPL, including lupus anticoagulant (LA), by diluted activated partial prothrombin time (aPTT), diluted Russell viper venom time (RVVT), and β 2-glycoprotein I-dependent anticardiolipin antibody (18), and blood tests for hypothyroidism and diabetes mellitus, before a subsequent pregnancy. Patients with APS, uterine anomalies, and abnormal chromosomes in either partner were excluded from the analysis (19). Patients with a history of preeclampsia or abruptio placentae also were excluded.

Subsequent pregnancies of all patients were followed until February 2013. Among the patients, 225 received no medication and 39 received anticoagulant treatment although there was no evidence. Gestational age was calculated from basal body temperature charts. Ultrasonography was performed once a week from 4 to 8 weeks of gestation. Dilation and curettage was performed on the patients diagnosed as having miscarriage. Part of the villi was cultured, and the cells were harvested after 6–22 days of cultivation to analyze the chromosome. A total of 47 aborted conceptuses could be karyotyped with the use of a standard G-banding technique.

Furthermore, 195 women with at least one child and no history of infertility or miscarriage were examined as control subjects. The control subjects were recruited from Nagoya City University Hospital and Inabe General Hospital from January to April 2012. The earlier study implicated SNP5 (minor allele frequency [MAF] 0.228) as the major susceptibility locus in RPL (17). The minor allele frequency (MAF) of SNP5 was 0.228. Based on a sample size of the 264 patients already obtained, MAF of \sim 20% in the SNP, and a multiplicative model, 195 control subjects were needed for 80% power and 5% significance.

The allele frequencies of the six SNPs of the *ANXA5* gene were compared between the patients and control subjects. The subsequent pregnancy outcomes were compared between the 264 patients with and without the risk alleles.

This study was conducted with the approval of the Research Ethics Committee of Nagoya City University Graduate School of Medical Sciences and the Ethics Committee of Inabe General Hospital. Each patient provided written consent after being provided with a full explanation about the purpose of the study and the methods to be used.

DNA Analysis

Genomic DNA was extracted from peripheral blood samples with the Midi Blood DNA Extraction kit (Qiagen). A total of six SNPs were analyzed: rs112782763 (-467G>A at promoter; SNP1), rs28717001 (-448A>C at promoter; SNP2), rs28651243 (-422T>C at 5'-untranslated exon 1; SNP3), rs113588187 (-373G>A at 5'-untranslated exon 1; SNP4), rs1050606 (-302T>G; SNP5), and rs11538099 (-1C>T at one base upstream of the initiation codon in exon2; SNP6). All genotyping was carried out using Taqman polymerase chain reaction (PCR) assays (Applied Biosystems) in 96-well arrays that included two blank wells as negative control samples according to the manufacturer's instructions. Taqman Predesigned SNP Genotyping assay and Taqman MGB probes were used. Taqman PCR and genotyping analyses were carried out on the Applied Biosystems 7500 Fast Real-Time PCR System. The reaction mixtures were amplified in 1 μ L template DNA (10 ng/ μ L), 12.5 μ L 2 \times Taqman Universal Master Mix, 0.625 μ L 20 \times primer/probe mix and 10.875 μ L double-distilled H₂O in a volume of the mixture of 25 μ L. The cycling conditions were as follows: initial denaturation at 95°C for 10 minutes, followed by 40 cycles at 95°C for 15 seconds and 58°C for 1 minute. The results were automatically analyzed on the Applied Biosystems 7500 Real-Time PCR System with the use of an allelic discrimination assay program (20).

Statistical Analysis

Departure from the Hardy-Weinberg equilibrium for the six SNPs was tested with the use of an exact test (21). Because previous studies have shown that the mainland Japanese population is genetically similar, we did not examine or perform corrections for the population substructure in our sample (22, 23).

For individual SNP association analyses, univariate logistic regression analyses were performed with dominant and

TABLE 1

Allele frequencies of six *ANXA5* single-nucleotide polymorphisms (SNPs) in patients with recurrent pregnancy loss and control subjects and the odds ratios (ORs) in the codominant and dominant models.

| | | Case | Control | OR (95% CI) | P value | P value of max-statistics | HWE | MAF |
|--------------------------------|------------|------|-----------|------------------|---------|---------------------------|-------|-------|
| SNP1 -467G>A rs112782763 | Codominant | | | | .187 | | 0.233 | 0.110 |
| | G/G | 206 | 160 | Reference | | | | |
| | G/A | 55 | 30 | 1.43 (0.87–2.33) | | | | |
| | A/A | 3 | 5 | 0.47 (0.11–1.96) | | | | |
| | Dominant | | | | .287 | .443 | | |
| | G/G | 206 | 160 | Reference | | | | |
| SNP2 -448A>C rs28717001 | Codominant | | | | .071 | | 0.086 | 0.108 |
| | A/A | 207 | 162 | Reference | | | | |
| | A/C | 54 | 27 | 1.56 (0.94–2.56) | | | | |
| | C/C | 3 | 6 | 0.39 (0.10–1.59) | | | | |
| | Dominant | | | | .211 | .266 | | |
| | A/A | 207 | 162 | Reference | | | | |
| SNP3 -442T>C rs28651243 | Codominant | | | | .086 | | 0.154 | 0.122 |
| | T/T | 205 | 160 | Reference | | | | |
| | T/C | 56 | 29 | 1.51 (0.92–2.50) | | | | |
| | C/C | 3 | 6 | 0.39 (0.10–1.59) | | | | |
| | Dominant | | | | .246 | .267 | | |
| | T/T | 205 | 160 | Reference | | | | |
| SNP4 -373G>A rs113588187 | Codominant | | | | .137 | | 0.214 | 0.107 |
| | G/G | 207 | 162 | Reference | | | | |
| | G/A | 54 | 28 | 0.89 (0.92–2.50) | | | | |
| | A/A | 3 | 5 | 0.47 (0.11–2.0) | | | | |
| | Dominant | | | | .211 | .381 | | |
| | G/G | 207 | 162 | Reference | | | | |
| SNP5 -302T>G rs1050606 | Codominant | | | | .109 | | 0.099 | 0.218 |
| | T/T | 155 | 132 | Reference | | | | |
| | T/G | 93 | 51 | 1.56 (1.03–2.33) | | | | |
| | G/G | 13 | 12 | 1.14 (0.01–2.5) | | | | |
| | Dominant | | | | .049 | .100 | | |
| | T/T | 155 | 132 | Reference | | | | |
| SNP6 1C>T rs11538099 | Codominant | | | | .044 | | 0.086 | 0.108 |
| | T/T | 206 | 163 | Reference | | | | |
| | C/C | | | | | | | |
| | T/C | 55 | 26 | 1.67 (1.01–2.78) | | | | |
| | C/T | | | | | | | |
| | C/C | 3 | 6 | 0.40 (0.10–1.61) | | | | |
| Dominant | | | | .135 | .258 | | | |
| T/T | 206 | 163 | Reference | | | | | |
| | C/C | | | | | | | |
| | T/C, C/C | 58 | 32 | 1.43 (0.89–2.33) | | | | |
| | C/T, T/T | | | | | | | |

Note: CI = confidence interval; HWE = Hardy-Weinberg equilibrium; MAF = minor allele frequency (in both patients and control subjects).

Hayashi. *ANXA5* SNPs in recurrent pregnancy loss. *Fertil Steril* 2013.

codominant models using the presence or absence of PRL as the dependent variable. The two inheritance models were tentatively chosen because earlier studies supported a dominant model for these SNPs (16, 17). We estimated the crude odds ratios (ORs) and 95% confidence intervals (CIs) with the use of the two models. However, there remains some uncertainty about the underlying genetic model. Therefore, to examine the right model of inheritance statistically as well as to avoid multiple comparisons by fitting multiple inheritance models, we used the max-statistic, which selects the largest test statistic from the dominant, recessive, and additive models.

To characterize the linkage disequilibrium (LD) pattern, we estimated the r^2 values for all pairs of SNPs. Then, the haplotype frequencies were estimated with the use of the expectation-maximization algorithm. To evaluate the association between the haplotypes and the risk of pregnancy loss, logistic regression models were used. Haplotype analyses were conducted based on the most supported model by the max-statistic in univariate analysis.

In the cohort study, univariate logistic regression analyses were performed to examine the association of subsequent miscarriage with variable SNPs. Then, multivariate logistic analyses were performed to examine the association

Results of maximizing association statistics in the present study and combination with previous data reported by Miyamura et al.

| | SNP1 | SNP2 | SNP3 | SNP4 | SNP5 | SNP6 |
|--|-------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| Present study | | | | | | |
| Dominant | 1.132 | 1.568 | 1.346 | 1.568 | 3.887 | 2.230 |
| Recessive | 1.317 | 2.177 | 2.177 | 1.317 | 0.002 | 2.177 |
| Log-additive | 0.365 | 0.400 | 0.318 | 0.581 | 2.408 | 0.705 |
| max-statistic | 1.317 | 2.177 | 2.177 | 1.568 | 3.887 | 2.230 |
| P value | .443 | .266 | .267 | .381 | .100 | .258 |
| Combination with data from Miyamura et al. | | | | | | |
| Dominant | 4.955 | 5.826 | 5.300 | 6.393 | 9.943 | 6.393 |
| Recessive | 1.954 | 2.932 | 2.932 | 1.954 | 1.541 | 2.932 |
| Log-additive | 2.696 | 2.860 | 2.565 | 3.615 | 9.058 | 3.177 |
| max-statistic | 4.955 | 5.826 | 5.300 | 6.393 | 9.943 | 6.393 |
| P value | .056 | .035 | .046 | .025 | .003 | .025 |
| OR obtained with selection of the dominant model in the combination data | | | | | | |
| OR (95% CI), P value | 1.52 (1.05–2.22), 0.029 | 1.57 (1.09–2.31), 0.18 | 1.53 (1.06–2.23), 0.24 | 1.62 (1.11–2.39), 0.13 | 1.61 (1.20–2.18), 0.02 | 1.62 (1.11–2.39), 0.13 |

Note: CI = confidence interval; OR = odds ratio; SNP = single-nucleotide polymorphism.
 Hayashi. ANXA5 SNPs in recurrent pregnancy loss. *Fertil Steril* 2013.

of subsequent miscarriage with variable SNPs, age, the number of previous miscarriages, and previous live births. The age was quadrisedged (21–31, 31–34, 34–37, and 37–45 years), and the number of previous miscarriages (two, three, four, or five to nine) and previous live births (none vs. one or two) were divided so that each group had an almost equal number, and the cells with few numbers were annexed.

Statistical analyses in cross-sectional study were conducted with R software (v. 2.13.0) (24), including the SNPassoc (25) and Haplo.stat (26) packages. Analysis in the cohort study was carried out with SAS (v. 19.0). $P < .05$ was considered to denote statistical significance.

RESULTS

The mean ages (SD) of the patients and control subjects were 33.8 (4.36) and 32.1 (6.91) years, respectively. The mean numbers of previous miscarriage were 2.74 (0.96) and 0, respectively. The mean numbers of previous live births were 0.20 (0.40) and 1.53 (0.72), respectively.

The genotype frequencies for six SNPs were found to be in Hardy-Weinberg equilibrium, suggesting neither sampling bias nor mistyping of genotyping (Table 1). The carrier frequency for the minor alleles tended to be higher in the RPL group for all six SNPs (not significant) when the dominant model was selected (SNP1: G/G vs. G/A-A/A; SNP2: A/A vs. A/C-C/C; SNP3: T/T vs. T/C-C/C; SNP4: G/G vs. G/A-A/A; SNP5: T/T vs. T/G-G/G; and SNP6: C/C vs. C/T-T/T), among which the highest significance level was obtained for SNP5 ($P = .049$; Table 1). When we selected the codominant model (SNP1: G/G vs. G/A vs. A/A; SNP2: A/A vs. A/C vs. C/C; SNP3: T/T vs. T/C vs. C/C; SNP4: G/G vs. G/A vs. A/A; SNP5: T/T vs. T/G vs. G/G; and SNP6: C/C vs. C/T vs. T/T), the highest significance level was obtained for SNP6 ($P = .044$).

Then we merged our data with the previously published Japanese data (17). Based on the results of the max-statistics, the dominant model was adopted for all six polymorphisms (Table 2). The combined data suggested that the dominant model should be selected, and statistically significant correlations were obtained for all six SNPs, with the highest level of significance for SNP5 ($P = .003$). However, the ORs were < 2.0 for all six SNPs.

We next performed LD analysis among the six ANXA5 SNPs (Fig. 1). LD evaluation revealed the existence of significant correlations among SNPs 1, 2, 3, 4, and 6. All except SNP5 exhibited a strong LD (> 0.95), as reported previously (16).

Haplotype analysis indicated the presence of three major haplotypes. G-A-T-G-T-C (patient vs. control: 75.9% vs. 80.3%) and G-A-T-G-G-C (12.1% vs. 8.7%) were the major haplotypes and accounted for $> 85\%$ of the subjects. The first four (G-A-T-G) SNPs constituted the N haplotype described previously (15). The haplotypes comprising all six minor alleles (A-C-C-A-G-T) was found to be the third most common (11.4% vs. 9.2%). The first four SNPs (A-C-C-A) constituted the M2 haplotype. All other haplotypes accounted for $< 1\%$. We subsequently performed a case-control study for SNPs 1–6 and for these three haplotypes. When we selected the dominant model, there was no significant difference of the

FIGURE 1

| | SNP1 | SNP2 | SNP3 | SNP4 | SNP5 | SNP6 |
|------|------|-------|-------|-------|-------|-------|
| SNP1 | | 0.981 | 0.980 | 0.975 | 0.484 | 0.978 |
| SNP2 | | | 0.968 | 0.954 | 0.476 | 0.995 |
| SNP3 | | | | 0.952 | 0.486 | 0.965 |
| SNP4 | | | | | 0.473 | 0.991 |
| SNP5 | | | | | | 0.476 |
| SNP6 | | | | | | |

Linkage disequilibrium of the six ANXA5 single-nucleotide polymorphisms (SNPs).

Hayashi. ANXA5 SNPs in recurrent pregnancy loss. *Fertil Steril* 2013.

frequency between patients and controls among the three haplotypes.

Of the 264 patients with RPL, 185 (70.1%) subsequently gave live births. Of the 79 aborted products, normal and abnormal karyotypes were 36.2% (n = 17) and 63.8% (n = 30), respectively.

Of the patients without treatment, 71.1% (160/225) gave live birth (Table 2). The subsequent live birth rates were 70.0% and 71.9% in the patients with and without the risk allele of SNP5, respectively, and the respective percentages were 84.0% and 84.3% when cases with an abnormal embryonic karyotype were excluded. Among the 39 patients who received anticoagulant treatment, the subsequent live birth rates were 63.2% and 65.0% in the patients with and without

TABLE 3

Subsequent live birth rate in subjects with and without the SNP5 risk allele.

| | With risk allele T/G or G/G | Without risk allele T/T |
|---|--------------------------------|----------------------------|
| Subsequent live birth rate in 225 patients without treatment | 70.0% (63/90) | 71.9% (97/135) |
| Subsequent live birth rate in 39 patients with treatment | 63.2% (12/19) | 65.0% (13/20) |
| Live birth rate after excluding cases with treatment or chemical pregnancy (n = 215) | 73.3% (63/86) | 75.2% (97/129) |
| Live birth rate after excluding cases with treatment or an abnormal embryonic karyotype (n = 190) | 84.0% (63/75) | 84.3% (97/115) |

Note: SNP = single-nucleotide polymorphism.

Hayashi. ANXA5 SNPs in recurrent pregnancy loss. *Fertil Steril* 2013.

the risk allele of SNP5, respectively. Treatment had no effect on the live birth rate in patients with the risk allele of SNP5.

According to the results of the univariate analysis using SNP5, there was no increase in the pregnancy loss rate associated with the presence of the SNP5 risk allele in the 190 patients without treatment or in chemical pregnancy or miscarriage caused by an abnormal embryonic karyotype (OR 1.027, 95% CI 0.463–2.276; P = .9487).

The results of multivariate analyses using SNP5, age, number of previous miscarriages, and presence of previous live births as variables revealed the absence of any influence of the SNP5 risk allele on the subsequent pregnancy loss rate (OR 1.187, 95% CI 0.506–2.784; P = .6941). The miscarriage rate in patients with a history of five to nine miscarriages was higher than that in the patients with a history of two miscarriages (OR 6.012, 95% CI 0.976–37.037; P = .0532).

DISCUSSION

This case-control study demonstrated a significantly higher frequency of the minor alleles in the RPL group compared with the control group for SNP5 (P = .049), but not for SNPs 1–4 and 6. Existence of an association between SNP5 and RPL was originally reported by another Japanese group (17). Miyamura et al. examined patients with a history of three or more unexplained miscarriages (3.06 ± 0.3). The other inclusion and exclusion criteria were similar to those in the present study. It was a new finding that the dominant model should be selected, because the previous study did not perform max-statistics.

Earlier studies suggested that the M2 haplotype of the ANXA5 gene promoter was significantly associated with the risk of RPL. Bogdanova et al. demonstrated for the first time that a sequence variation in the promoter region of the ANXA5 gene represented a risk factor for RPL in a German population (16, 27). Another analysis of the significance of M2 strengthened the initial findings in Italian patients with RPL (28). Reporter gene assays showed that M2 reduced the in vitro activity of the ANXA5 promoter to 37%–42% of normal (16). These data suggest that these SNPs are functional and can reduce the expression of ANXA5, thereby influencing the risk of RPL. A recent report showed that the M2 allele in heterozygous placentas resulted in a reduced expression level of the ANXA5 mRNA by an average of 42% compared with the normal allele (29, 30). The M2 haplotype within the ANXA5 gene was also reported as a new thrombophilic risk factor during pregnancy (31). However, we could not discern any significant difference by max-statistics or haplotype analysis even though our present study included the largest number of subjects in both patient and control groups.

LD evaluation revealed that all except SNP5 manifested a strong LD (>0.95), consistent with the earlier report (17), suggesting that SNP5 may be an independent risk factor for RPL. The significance of SNP5 has not been analyzed in Western patients with RPL.

It has been suggested that annexin A5 molecules form an anticoagulation shield on the apical surface of the placental syncytiotrophoblasts, that may, in pregnancy, be disrupted by antiphospholipid antibodies (32, 33). In a mouse study,

infusion of anti-annexin A5 antibodies induced coagulation at the surface of the syncytiotrophoblast layer, leading to fetal loss, which indicates that annexin A5 protects the fetomaternal interface from the coagulation of maternal blood (34). The syncytiotrophoblasts that possess annexin A5 protein at their surface are of fetal origin and have a half-maternal and half-paternal genome. Tranquilli et al. proposed that carrier status of the fetus of thrombophilia such as factor V polymorphism should be considered to be risk factor for intrauterine fetal death (35). A recent report demonstrated that paternal and maternal carriage of the annexin A5 M2 haplotype were equal risk factors for RPL (36). Thus, it is still not clear whether the annexin A5 protein at the syncytiotrophoblast surface originates from the maternal circulation or is produced by the fetal syncytiotrophoblasts. Functional analysis of SNP5 or SNP5 genotyping of the fetus or placenta would be warranted to further clarify our findings.

A number of articles regarding association between SNPs and RPL have been published (27). An association between heritable thrombophilia, such as the factor V Leiden mutation, prothrombin mutation, and protein S deficiency, and RPL has been reported (9, 10), although the association is not well established (11).

The present study confirmed *ANXA5* SNP5 as a risk factor for RPL. However, the subsequent live birth rate was 84.0% and 84.3% in patients with and without the risk allele of SNP5, after the exclusion of cases with an abnormal embryonic karyotype, with no significant difference. This was the first study indicating the influence of *ANXA5* SNP5 on further pregnancy outcomes. The genome-wide association study proved that the effect of the one of many kinds of SNPs associated with a common disease is very small when the OR is relatively small (37). The *ANXA5* risk allele was not found to be a reliable clinical predictor of subsequent pregnancy outcome. Therefore, we propose that testing for this allele is not needed, because it is without clinical benefit and is an unnecessary expense.

The randomized controlled trial by Kaandrop et al. concluded that there was no effect of combined heparin and aspirin treatment for unexplained recurrent miscarriage (38). The study did not deny that anticoagulant therapy might be effective for a limited number of patients with miscarriage of unexplained cause, because abnormal embryonic karyotype was found to be the most common cause in cases with an unexplained etiology (6). Adequate selection of patients based on analysis of several kinds of SNPs might improve the therapeutic effectiveness. Further study is important to find SNPs with clinical predictive benefit associated with RPL.

REFERENCES

- Farquharson RG, Pearson JF, John L. Lupus anticoagulant and pregnancy management. *Lancet* 1984;28:228-9.
- Sugiura-Ogasawara M, Ozaki Y, Sato T, Suzumori N, Suzumori K. Poor prognosis of recurrent aborters with either maternal or paternal reciprocal translocation. *Fertil Steril* 2004;81:367-73.
- Sugiura-Ogasawara M, Ozaki Y, Kitaori T, Kumagai K, Suzuki S. Midline uterine defect size correlated with miscarriage of euploid embryos in recurrent cases. *Fertil Steril* 2010;93:1983-8.
- Clifford K, Rai R, Watson H, Regan L. An informative protocol for the investigation of recurrent miscarriage: preliminary experience of 500 consecutive cases. *Hum Reprod* 1994;9:1328-32.
- Stephenson MD. Frequency of factors associated with habitual abortion in 197 couples. *Fertil Steril* 1996;66:24-9.
- Sugiura-Ogasawara M, Ozaki Y, Katano K, Suzumori N, Kitaori T, Mizutani E. Abnormal embryonic karyotype is the most frequent cause of recurrent miscarriage. *Hum Reprod* 2012;27:2297-303.
- Cowchock FS, Reece EA, Balaban D, Branch DW, Plouffe L. Repeated fetal losses associated with antiphospholipid antibodies: a collaborative randomized trial comparing prednisone with low-dose heparin treatment. *Am J Obstet Gynecol* 1992;166:1318-23.
- Rai R, Cohen H, Dave M, Regan L. Randomised controlled trial of aspirin and aspirin plus heparin in pregnant women with recurrent miscarriage associated with phospholipid antibodies (or antiphospholipid antibodies). *BMJ* 1997;314:253-7.
- Preston FE, Rosendaal FR, Walker ID, Briët E, Berntorp E, Conard J, et al. Increased fetal loss in women with heritable thrombophilia. *Lancet* 1996 Oct 5;348:913-6.
- Kupferminc MJ, Eldor A, Steinman N, Many A, Bar-Am A, Jaffa A, et al. Increased frequency of genetic thrombophilia in women with complications of pregnancy. *N Engl J Med* 1999;340:50-2.
- Branch DW, Gibson M, Silver RM. Clinical Practice: Recurrent miscarriage. *N Engl J Med* 2010;363:1740-7.
- Rand JH, Wu XX, Guller S, Gil J, Guha A, Scher J, et al. Reduction of annexin-V (placental anticoagulant protein-I) on placental villi of women with antiphospholipid antibodies and recurrent spontaneous abortion. *Am J Obstet Gynecol* 1994;171:1566-72.
- Thiagarajan P, Tait JF. Binding of annexin V/placental anticoagulant protein I to platelets. Evidence for phosphatidylserine exposure in the procoagulant response of activated platelets. *J Biol Chem* 1990;265:17420-3.
- Morgan RO, Bell DW, Testa JR, Fernandez MP. Genomic locations of *ANX11* and *ANX13* and the evolutionary genetics of human annexins. *Genomics* 1998;48:100-10.
- Carcedo MT, Iglesias JM, Bances P, Morgan RO, Fernandez MP. Functional analysis of the human annexin A5 gene promoter: a downstream DNA element and an upstream long terminal repeat regulate transcription. *Biochem J* 2001;356:571-9.
- Bogdanova N, Horst J, Chlystun M, Croucher PJ, Nabel A, Bohring A, et al. A common haplotype of the annexin A5 (*ANXA5*) gene promoter is associated with recurrent pregnancy loss. *Hum Mol Genet* 2007;16:573-8.
- Miyamura H, Nishizawa H, Ota S, Suzuki M, Inagaki A, Egusa H, et al. Polymorphisms in the annexin A5 gene promoter in Japanese women with recurrent pregnancy loss. *Mol Hum Reprod* 2011;17:447-52.
- Ogasawara M, Aoki K, Matsuura E, Sasa H, Yagami Y. Anti beta 2glycoprotein I antibodies and lupus anticoagulant in patients with recurrent pregnancy loss: prevalence and clinical significance. *Lupus* 1996;5:587-92.
- Miyakis S, Lockshin MD, Atsumi T, Branch DW, Brey RL, Cervera R, et al. International consensus statement of an update of the classification criteria for definite antiphospholipid syndrome (APS). *J Thromb Haemost* 2006;4:295-306.
- Endo K, Konishi A, Sasaki H, Takada M, Tanaka H, Okumura H, et al. Epidermal growth factor receptor gene mutation in nonsmall cell lung cancer using highly sensitive and fast Taqman PCR assay. *Lung Cancer* 2005;50:375-84.
- Wigginton JE, Cutler DJ, Abecasis GR. Note on exact tests of Hardy-Weinberg equilibrium. *Am J Hum Genet* 2005;76:887-93.
- Nishiyama T, Kishino H, Suzuki S, Ando R, Niimura H, Uemura H, et al. Detailed analysis of Japanese population substructure with a focus on the southwest islands of Japan. *PLoS One* 2012;7:e35000.
- Yamaguchi-Kabata Y, Shimada MK, Hayakawa Y, Minoshima S, Chakraborty R, Gojobori T, et al. Japanese population structure, based on SNP genotypes from 7003 individuals compared to other ethnic groups: effects on population-based association studies. *Am J Hum Genet* 2008;83:445-56.
- R Development Core Team. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing; 2011.

- González JR, Armengol L, Solé X, Guinó E, Mercader JM, Estivill X, et al. SNPAssoc: an R package to perform whole genome association studies. *Bioinformatics* 2007;23:644–5.
- Lake SL, Lyon H, Tantisira K, Silverman EK, Weiss ST, Laird NM, et al. Estimation and tests of haplotype-environment interaction when linkage phase is ambiguous. *Hum Hered* 2003;55:56–65.
- Bogdanova N, Markoff A. Hereditary thrombophilic risk factors for recurrent pregnancy loss. *J Community Genet* 2010;1:47–53.
- Tiscia G, Colaizzo D, Chinni E, Pisanelli D, Sciannone N, Favuzzi G, et al. Haplotype M2 in the annexin A5 (ANXA5) gene and the occurrence of obstetric complications. *Thromb Haemost* 2009;102:309–13.
- Chinni E, Tiscia GL, Colaizzo D, Vergura P, Margaglione M, Grandone E. Annexin V expression in human placenta is influenced by the carriership of the common haplotype M2. *Fertil Steril* 2009;91:940–2.
- Markoff A, Gerdes S, Feldner S, Bogdanova N, Gerke V, Grandone E. Reduced allele specific annexin A5 mRNA levels in placentas carrying the M2/ANXA5 allele. *Placenta* 2010;31:937–40.
- Grandone E, Tiscia G, Colaizzo D, Chinni E, Pisanelli D, Bafunno V, et al. Role of the M2 haplotype within the annexin A5 gene in the occurrence of pregnancy-related venous thromboembolism. *Am J Obstet Gynecol* 2010;203:461.e1–5.
32. Rand JH, Wu XX. Antibody-mediated disruption of the annexin-V antithrombotic shield: a new mechanism for thrombosis in the antiphospholipid syndrome. *Thromb Haemost* 1999;82:649–55.
33. Rand JH, Wu XX, Quinn AS, Chen PP, McCrae KR, Bovill EG, et al. Human monoclonal antiphospholipid antibodies disrupt the annexin A5 anticoagulant crystal shield on phospholipids bilayers: evidence from atomic force microscopy and functional assay. *Am J Pathol* 2003;163:1193–200.
34. Wang X, Campos B, Kaetzel MA, Dedman JR. Annexin V is critical in the maintenance of murine placental integrity. *Am J Obstet Gynecol* 1999;180:1008–16.
35. Tranquilli AL, Saccucci F, Giannubilo SR, Cecati M, Nocchi L, Lorenzi S, et al. Unexplained fetal loss: the fetal side of thrombophilia. *Fertil Steril* 2010;94:378–80.
36. Rogenhofer N, Engels L, Bogdanova N, Tuitelmann F, Markoff A, Thaler C. Paternal and maternal carriage of the annexin A5 M2 haplotype are equal risk factors for recurrent pregnancy loss: a pilot study. *Fertil Steril* 2012;98:383–8.
37. Kaiser J. Human genetics. Genetic influences on disease remain hidden. *Science* 2012;338:1016–7.
38. Kaandorp SP, Goddijn M, van der Post JA, Hutten BA, Verhoeve HR, Hamulyak K, et al. Aspirin plus heparin or aspirin alone in women with recurrent miscarriage. *N Engl J Med* 2010;362:1586–96.

Cognitive behavior therapy for psychological distress in patients with recurrent miscarriage

Yumi Nakano¹
Tatsuo Akechi²
Toshiaki A Furukawa³
Mayumi Sugiura-Ogasawara⁴

¹Department of Psychology, School of Human Sciences, Sugiyama Jogakuen University, Nisshin, Aichi, Japan; ²Department of Psychiatry and Cognitive-Behavioral Medicine, Nagoya City University, Graduate School of Medical Sciences, Nagoya, Japan; ³Department of Health Promotion and Human Behavior (Cognitive-Behavioral Medicine), Kyoto University Graduate School of Medicine, Kyoto, Japan; ⁴Department of Obstetrics and Gynecology, Nagoya City University, Graduate School of Medical Sciences, Nagoya, Japan

Objective: To examine the reduction of psychiatric symptoms using individual cognitive behavior therapy (CBT) for women who suffer from recurrent miscarriage (RM) and depression and/or anxiety.

Methods: Patients with RM and a score of five or higher for K6, a self-report screening scale for depression/anxiety, were interviewed to find information about stressful situations, thoughts, and consequent behaviors that are common and potential causes of psychological distress among RM patients. We then performed individual CBT on 14 patients with RM and depression/anxiety, referring to a list from the interviews, and examined the effects of CBT by a paired *t*-test.

Results: Fourteen women received CBT. The mean number of intervention times was 8.9 sessions (standard deviation [SD], 4.6 sessions). The average Beck Depression Inventory-Second Edition and State-Trait Anxiety Inventory-state anxiety scores, self-report screening scales for depression/anxiety, decreased from 13.6 (SD, 8.2) and 49.0 (SD, 7.1) at baseline to 5.2 (SD, 4.4) and 38.0 (SD, 10.2) posttherapy, respectively. These changes were statistically significant.

Conclusion: The current preliminary open study confirmed that individual CBT was potentially useful for women with RM and depression and/or anxiety. This finding is the first step towards creating a comprehensive psychological support system for women with RM.

Keywords: spontaneous abortion, psychological support, depression, anxiety

Introduction

Miscarriage is a very common complication in pregnancy, found in 15% of the women clinically identified as being pregnant. Recurrent miscarriage (RM), defined as two or more consecutive miscarriages, develops in about 5% of couples.¹ In recent years, RM has gained social recognition as a problem related to the decreasing birthrate.²

Causes of RM include abnormal chromosomes in either partner, existence of antiphospholipid antibodies, and uterine anomalies. Abnormal embryonic karyotype is also known as a cause, accounting for 25%–50% of miscarriages. However, more than half of RM cases remain unexplained.¹

Miscarriage causes mental distress such as depression, anxiety, anger, and grief. If miscarriage is repeated, mental distress is sustained.³ Craig et al³ and Klock et al⁴ reported that mental disorder was found in two-thirds of RM patients, and Craig et al³ also reported that their anxiety level is similar to that of psychiatric outpatients. All of these suggest that RM patients are a group of people who need appropriate mental care.

In the infertility field, many studies concerning various psychological supports have been conducted for mental distress of patients. Their outcome is divided into two major kinds, improvement of psychological status and improvement of

Correspondence: Yumi Nakano
Department of Psychology, School of Human Sciences, Sugiyama Jogakuen University, 3-2005 Takenoyama, Nisshin, Aich 4700136, Japan
Tel +81 5 6174 1452
Fax +81 5 6174 3205
Email nakanotys2012@sugiyama-u.ac.jp

pregnancy rate. Three meta-analyses lead to a general conclusion that at least one of the two is to be expected.⁵⁻⁷

A few previous studies reported that some forms of psychological support including cognitive behavior therapy (CBT) for patients with single miscarriage were useful.⁸⁻¹⁰ In the field of unexplained RM, it has been also indicated that psychological support such as counseling and increased regular checkups in the beginning pregnancy can help raise fertility rate,¹¹⁻¹³ but no further progress has been made. However, unlike for infertility patients or single-miscarriage patients, little research has been done for RM patients on the effects of psychological support with an outcome of reducing anxiety and depression in daily life.

CBT is one of the psychotherapies that is generally confirmed as being effective in reducing depression and/or anxiety.¹⁴ In this study, we preliminarily examined the possibility of reducing depression and/or anxiety of the subjects through individual CBT as the first step for constructing a total psychological support system for RM. Generally, in each CBT session, the patient and the therapist utilize specific hardships in the patient's real life to collaboratively identify and examine her perspective (thoughts, recognition) and behavior. They then devise ways to control her distress or find coping strategies for her hardship. Accordingly, having information in advance about the kind of stressful situations, thoughts, and consequent behaviors which are common and potential causes of depression and/or anxiety among RM patients helps the therapist identify a problem to be discussed and in turn smoothly conduct a session. In addition, such information could be used in the course of developing a broadly available psychological support system in the future. Therefore, we first interviewed patients about what situations often cause them distress, and then conducted preliminary individual CBT sessions referring to the previous interview results and observed the change of depression and/or anxiety.

Materials and methods

Patients

Subjects were recruited from women who visited the specialized RM outpatient care at Nagoya City University Hospital during the period of April 2008 to September 2010; recruited subjects had a history of two or more consecutive miscarriages and had no children. Two weeks after systematic RM tests and explanation about the results, patients filled in K6,^{15,16} a self-report questionnaire for depression and/or anxiety, and sent it to us by mail. Patients with a score of five or higher on the K6 were selected as subjects in this study.

With those subjects, a psychiatrist (YN) conducted Diagnostic and Statistical Manual of Mental Disorders (fourth edition, text revision) (DSM-IV-TR) diagnosis with Structured Clinical Interview for DSM-IV (SCID),¹⁷ and the psychiatrist selected those who were diagnosed with mood disorder, anxiety disorder, or adjustment disorder caused by RM-related stress, pregnancy, or childbirth. Exclusion criteria included: (1) serious physical illnesses such as autoimmune disorder, cardiac disease, or chronic respiratory disease; (2) current or past psychotic disorder, bipolar disorder, cyclothymic disorder, eating disorder, developmental disorder, learning disorder, or substance dependence according to DSM-IV-TR; (3) current depression episode, dysthymic disorder, anxiety disorder, or adjustment disorder (by DSM-IV-TR) starting before the first miscarriage; (4) being pregnant at the time of the interview or beginning of CBT; and (5) previous CBT experience. As for new pregnancy during CBT, we planned to make a go/no-go decision when it happened because pregnancy may make some subjects physically (eg, morning sickness) and psychologically (eg, anxiety about miscarriage) unstable.

In general, CBT subjects were instructed to avoid pharmacotherapy. If they were already under pharmacotherapy, they were not allowed to switch medicines during CBT, except in cases of emergency. If a cause of RM was specified, its treatment had priority over CBT.

This study was approved by the Ethics Review Committee at Nagoya City University Graduate School of Medical Sciences. The subjects received explanation in writing and submitted informed consent before entering the study.

Procedure

Before starting psychological support by CBT, a researcher in this study (YN) with over 15 years of experience in clinical psychiatry and 10 years of experience as a CBT therapist interviewed a series of ten subjects in person, for 75–90 minutes each, and collected information about mental distress concerning pregnancy and children seen in their daily situations. Another psychiatrist (TF) with over 10 years of CBT experience who has been conducting research in psychosocial factors of RM, as well as an obstetrician (MS) who specializes in RM, then examined the interviews. By checking what was said (viewpoints, recognition) and how they behaved in stressful situations, as mentioned by subjects, several features, which seemed helpful in CBT, were extracted (Table 1).

The average age of the ten subjects interviewed was 32.3 (± 3.5) years and the average number of miscarriages was

Table 1 Common thoughts and behaviors among women with recurrent miscarriage who are experiencing depression and/or anxiety

1. They believe that a woman who marries but does not have a baby is not a mature adult
2. They believe that people who have children are happy and that people who do not have children are unhappy
3. They have feelings of guilt that they have killed their own fetuses in utero
4. They are afraid of both getting pregnant and of not getting pregnant
5. They are beleaguered with anxiety about the future and about how long their present situation will last
6. They are uncertain if each action in their daily life is good or bad for their next pregnancy
7. They miss many opportunities to enjoy themselves because they avoid places where there are many children and their parents, and participating in events where many mothers will be present
8. They avoid gatherings of relatives during the summer holidays or New Year holidays, or gatherings of peers such as class reunions
9. They and their spouses are likely to overlook women's fatigue and exhaustion
10. Not many women understand that their idea and their husband's idea are not the same regarding having a baby

2.7 (±0.9). No significant difference was found between these and the average age of 33.0 (standard deviation [SD] ± 4.8) and the average number of miscarriages of 2.7 (SD ± 1.0) of a series of 305 patients who visited our hospital for thorough examination. The causes of miscarriage for eight subjects were unexplained. Two suffered from uterine deformity, but they did not need surgery in order to become pregnant or to maintain pregnancy.¹⁸ The average K6 score was 8.6 (±2.5).

Those who were interviewed to extract common thoughts and behaviors continued regular treatment at the obstetrics department. They also received routine psychiatric outpatient therapy if they needed it.

After common thoughts and behaviors were obtained as indicated in Table 1, 14 subjects who met the eligibility criteria provided informed consent and entered this study between October 2008 and September 2010 (Figure 1).




Figure 1 Patient flow chart for CBT.

Abbreviations: CBT, cognitive behavior therapy; N, number; RM, recurrent miscarriage; SCID, Structured Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders (fourth edition); K6, 6-item self-report instrument for screening for clinical depression or anxiety in adults.

The psychiatrist (YN) conducted CBT according to the individual treatment method developed by Beck¹⁹ as well as by Beck et al (Table 1).²⁰ One session lasted approximately 50 minutes. The list in Table 1 was never shown to the patient since it was made as reference to help the therapist conduct smooth and effective sessions.

The basic rule for a course of sessions was that the sessions were to be held once a week, up to 16 times in total. However, in reality, flexible decisions had to be made depending on how profoundly distressed the patient was feeling in her daily life and how much the patient's depression and/or anxiety improved. In some cases, we allowed a patient to have a session once every 2 or more weeks after the 12th session.

Measure

K6

The K6 is a 6-item self-report instrument for screening for clinical depression or anxiety in adults.¹⁵ The items cover how frequently the respondents experienced symptoms of psychological distress during the previous 30 days. The possible total scores range from 0 to 24. According to a Japanese validation study of the K6 in a general population, the positive predictive value for major depressive disorder or any anxiety disorder according to the DSM-IV-TR criteria was 0.49 for K6 scores of 10 or more.¹⁶ The reliability and validity of the Japanese version has been previously confirmed.¹⁶ The internal consistency of this instrument in the current sample was sufficient (Cronbach's $\alpha = 0.89$). In this study, women with a score of five or higher on the K6 result were included as subjects.

Beck Depression Inventory-Second Edition

The Beck Depression Inventory-Second Edition (BDI-II) is a 21-item self-report instrument for measuring the severity of depression in adolescents and adults.²¹ The total scores range from 0 through 63. The severity of depression is categorized based on the following BDI-II scores in Japanese samples: a score of 13 or less is regarded as minimal (or remission); 14 to 19 is regarded as mild; 20 to 28 is regarded as moderate; and 29 or greater is regarded as severe. The reliability and validity of the Japanese version has been previously confirmed.²² The internal consistency of this instrument in the current sample was sufficient (Cronbach's $\alpha = 0.92$).

State-Trait Anxiety Inventory-state anxiety

The State-Trait Anxiety Inventory-state (STAI-s) anxiety consists of 20 items and has a maximum score of 80.²³

State anxiety refers to the degree of anxiety at a particular point. Scores of over 50 indicate an extremely high level of anxiety (neurosis level). The reliability and validity of the Japanese version has been previously confirmed.²⁴ The internal consistency of this instrument in the current sample was sufficient (Cronbach's $\alpha = 0.91$).

BDI-II and STAI-s anxiety were adopted because they are both simple and useful scales to check the change in depression and/or anxiety severity. Following the usual CBT procedure, they were conducted right before each CBT session to assess the patient's condition.

Statistical analysis

Wilcoxon signed rank tests were conducted before and after CBT to see any statistically-significant change of BDI-II and STAI state anxiety. All analyses were performed using IBM SPSS Statistics, version 18 (IBM Corporation, Armonk, NY, USA). All statistical tests were two-tailed, and an alpha value of less than 0.05 was considered statistically significant.

Results

Fourteen patients, who met the criteria and consented, received CBT. The features of the samples are shown in Table 2. No significant difference was found between these and a series of 305 patients who visited our hospital for thorough examination in terms of the average age and number

Table 2 Sample characteristics: demographic and symptomatic data

| | Mean | SD |
|---|------|--------|
| Age (years) | 34 | 4 |
| Number of previous miscarriages | 2.7 | 1.0 |
| BDI-II before CBT | 13.6 | 8.2 |
| State anxiety in STAI before CBT | 49 | 7.1 |
| Number of CBT sessions | 8.9 | 4.6 |
| | % | Number |
| Education | | |
| High school graduate | 28.6 | 4 |
| Junior college graduate | 50.0 | 7 |
| College graduate or higher | 21.4 | 3 |
| DSM-IV diagnosis | | |
| Major depressive disorder | 35.7 | 5 |
| Adjustment disorder (with depressed mood) | 35.7 | 5 |
| Adjustment disorder (with mixed anxiety and depressed mood) | 28.6 | 4 |
| Adjustment disorder (with anxiety) | 7.1 | 1 |
| Specific phobia | 7.1 | 1 |
| Panic disorder | 7.1 | 1 |
| Posttraumatic stress disorders | 7.1 | 1 |

Abbreviations: SD, standard deviation; BDI-II, Beck Depression Inventory-Second Edition; CBT, cognitive behavior therapy; STAI, The State-Trait Anxiety Inventory; DSM-IV, Diagnostic and Statistical Manual of Mental Disorders (fourth edition).

of miscarriages. The causes of RM for nine of the patients (64.2%) were unexplained. Four patients had uterine deformity but their possibility of maintaining pregnancy was not expected to improve much by a surgery.¹³ One patient was diagnosed with hyperthyroidism but her hormone value had been stable within the normal range for over 6 months under pharmacotherapy (desiccated thyroid 75 µg).

Two patients were diagnosed with moderate- to severe-level major depression disorder according to DSM-VI-TR; their BDI-II scores exceeded 30 points at baseline. The depression and/or anxiety severity of most of the patients was around the level of adjustment disorder according to DSM-IV-TR. One patient had been in pharmacotherapy and there was no change in her medication (75 mg/day sertraline, 1 mg/day lorazepam) during CBT sessions. There was one patient who became pregnant in the 14th week of the 16 originally planned sessions. Since she complained of high anxiety about miscarriage and asked for extended treatment, a decision was made to have a session once a month until she reached a stable period, totaling 18 sessions.

Figure 2 shows the changes in BDI-II and STAI-s anxiety before and after CBT. STAI data for four patients were lost. The average BDI-II and STAI-s scores decreased from 13.6 (SD, 8.2) and 49.0 (SD, 7.1) at baseline to 5.2 (SD, 4.4) and 38.0 (SD, 10.2) at posttherapy, respectively. These changes were statistically significant (Wilcoxon signed rank test: BDI-II $n = 14$, $z = -3.2$, $P = 0.001$; STAI-s $n = 10$, $z = -2.4$, $P = 0.016$).

Discussion

This is the first attempt of psychological support with CBT for patients with RM. This study preliminarily confirmed that CBT, in concert with references to the list of common

hardships prepared in advance, can decrease depression and/or anxiety of patients with RM. It is appropriate to say that, based on the SCID diagnosis and the average BDI-II scores at baseline and posttreatment, the mild depressive state of the 14 subjects in this study recovered to normal levels. It can be also said from the average scores of STAI-s anxiety at baseline and at posttreatment that the subjects came out of their previous condition at a mildly overanxious level.

It has been already mentioned that the childbirth rate of RM patients with an unexplained cause may be raised by increasing the number of checkups and providing counseling in early pregnancy.⁸⁻¹⁰ Additionally, we have already shown that depression raises the possibility of miscarriage.²⁵ All these factors indicate that recovery from depression may lead to an increased childbirth rate. Therefore, although we were unable to examine successful birthrate this time, we should, and certainly hope to, have childbirth rate as an outcome in our future research.

However, there are several limitations in this study. First, it was an open label study with no control group. There exists a possibility that depression and/or anxiety decreased as a natural course. Second, it was left unexamined whether the positive effect could be maintained after the end of treatment, despite the fact continuing improvement of depression and/or anxiety in the life of patients is the most important outcome. Moreover, since there were only two estimating points, one at the start and another at the end of CBT sessions, the process of improvement could not be discussed. The third limitation is that the strict procedure of qualitative research, such as a qualitative and descriptive research method, was not taken in constructing Table 1. However, it was designed as reference data to conduct CBT smoothly in the first place, and in fact, Table 1 fulfilled the purpose. For example, more than half of the subjects covered items 1, 3, 5, 8, and 10 in their sessions.

We routinely hand out a form to patients to write comments freely when they visit our hospital for thorough examination. Although they often mention experiencing various kinds of psychological distress, only a few patients actually come for individual CBT. As Boivin et al²⁶ pointed out, it is likely that patients with mild psychological distress do not feel a need to depend on a specialist, while those experiencing severe distress do not reach a point where they want to meet with CBT therapists, or any other specialist engaging in psychological support, because these individuals worry about how much these services may cost and what the specialist may be like. Moreover, many patients might naturally be resistant to self-disclosure.




Figure 2 Comparison of BDI-II and STAI before and after CBT.

Abbreviations: BDI-II, Beck Depression Inventory-Second Edition; STAI, State-Trait Anxiety Inventory; CBT, cognitive behavior therapy.

Recently, there have been many attempts concerning CBT on the Internet.^{27,28} The approach using the Internet may have advantages because it can be accessed from anywhere at low cost, requires less self-disclosure, calls for less concern about compatibility with a therapist, and therefore, seems easier to start. Upon creating online CBT content in the future, the information from Table 1 will contribute considerably.

On the other hand, Wischmann²⁹ recommended that face-to-face consultations be used for complicated matters for infertility patients' mental distress. Therefore, for RM patients who are suffering from depression and/or anxiety, we plan to recommend a psychological support program based on CBT via the Web before inviting patients to engage in individual CBT. Once such a system takes off, we hope to conduct a randomized controlled trial with birthrate, as well as depression and/or anxiety, as study endpoints.

This pilot study, which preliminarily indicated a decrease in depression and/or anxiety for RM patients by individual CBT, was the first step towards creating a comprehensive psychological support system for RM.

Acknowledgments

This study was supported by the Japanese Ministry of Health, Labor, and Welfare, Grant number H20-children-general-002.

Disclosure

The authors report no conflicts of interest in this work. The authors disclosed no proprietary or commercial interest in any product mentioned or concept discussed in this article.

References

- Rai R, Regan L. Recurrent miscarriage. *Lancet*. 2006;368(9535):601–611.
- Ministry of Health Labour and Welfare J. Ministry of Health, Labour and Welfare Annual Report 2011. Tokyo, Japan, 2012. http://www.mhlw.go.jp/toukei/youran/indexyk_1_2.html.
- Craig M, Tata P, Regan L. Psychiatric morbidity among patients with recurrent miscarriage. *J Psychosom Obstet Gynaecol*. 2002;23(3):157–164.
- Klock SC, Chang G, Hiley A, Hill J. Psychological distress among women with recurrent spontaneous abortion. *Psychosomatics*. 1997;38(5):503–507.
- Boivin J. A review of psychosocial interventions in infertility. *Soc Sci Med*. 2003;57(12):2325–2341.
- de Liz TM, Strauss B. Differential efficacy of group and individual/couple psychotherapy with infertile patients. *Hum Reprod*. 2005;20(5):1324–1332.
- Hämmerli K, Znoj H, Barth J. The efficacy of psychological interventions for infertile patients: a meta-analysis examining mental health and pregnancy rate. *Hum Reprod Update*. 2009;15(3):279–295.
- Nikčević AV. Development and evaluation of a miscarriage follow-up clinic. *J Reprod Infant Psychol*. 2003;21(3):207–217.
- Séjourné N, Callahan S, Chabrol H. The utility of a psychological intervention for coping with spontaneous abortion. *J Reprod Infant Psychol*. 2010;28(3):287–296.
- Swanson KM. Effects of caring, measurement, and time on miscarriage impact and women's well-being. *Nurs Res*. 1999;48(6):288–298.
- Clifford K, Rai R, Regan L. Future pregnancy outcome in unexplained recurrent first trimester miscarriage. *Hum Reprod*. 1997;12(2):387–389.
- Liddell HS, Pattison NS, Zanderigo A. Recurrent miscarriage – outcome after supportive care in early pregnancy. *Aust N Z J Obstet Gynaecol*. 1991;31(4):320–322.
- Stray-Pedersen B, Stray-Pedersen S. Etiologic factors and subsequent reproductive performance in 195 couples with a prior history of habitual abortion. *Am J Obstet Gynecol*. 1984;148(2):140–146.
- Tolin DF. Is cognitive-behavioral therapy more effective than other therapies? A meta-analytic review. *Clin Psychol Rev*. 2010;30(6):710–720.
- Kessler RC, Andrews G, Colpe LJ, et al. Short screening scales to monitor population prevalences and trends in non-specific psychological distress. *Psychol Med*. 2002;32(6):959–976.
- Furukawa TA, Kawakami N, Saitoh M, et al. The performance of the Japanese version of the K6 and K10 in the World Mental Health Survey Japan. *Int J Methods Psychiatr Res*. 2008;17(3):152–158.
- First MB, Spitzer RL, Gibbon M, Williams JBW. User's Guide for the Structured Clinical Interview for DSM-IV Axis I Disorders: SCID-I Clinician Version. Arlington, VA: American Psychiatric Press; 1997.
- Sugiura-Ogasawara M, Ozaki Y, Kitaori T, Kumagai K, Suzuki S. Midline uterine defect size is correlated with miscarriage of euploid embryos in recurrent cases. *Fertil Steril*. 2010;93(6):1983–1988.
- Beck AT. *Cognitive Therapy of Depression*. New York, NY: Guilford Press; 1979.
- Beck AT, Emery G, Greenberg R. *Anxiety Disorders and Phobias: A Cognitive Perspective*. New York, NY: Basic Books; 1985.
- Beck AT, Steer RA, Brown GK. *Manual for the Beck Depression Inventory-II*. San Antonio, TX: The Psychological Corporation; 1996.
- Hiroe T, Kojima M, Yamamoto I, et al. Gradations of clinical severity and sensitivity to change assessed with the Beck Depression Inventory-II in Japanese patients with depression. *Psychiatry Res*. 2005;135(3):229–235.
- Spielberger CD, Gorsuch RL, Lushene R. *Manual for the State-Trait Anxiety Inventory*, Consulting Psychologists press, California, 1973.
- Iwata N, Mishima N, Okabe K, Kobayashi N, Hashiguchi E, Egashira K. Psychometric properties of the State-Trait Anxiety Inventory among Japanese clinical outpatients. *J Clin Psychol*. 2000;56(6):793–806.
- Nakano Y, Oshima M, Sugiura-Ogasawara M, Aoki K, Kitamura T, Furukawa TA. Psychosocial predictors of successful delivery after unexplained recurrent spontaneous abortions: a cohort study. *Acta Psychiatr Scand*. 2004;109(6):440–446.
- Boivin J, Scanlan LC, Walker SM. Why are infertile patients not using psychosocial counselling? *Hum Reprod*. 1999;14(5):1384–1391.
- Lewis C, Pearce J, Bisson JI. Efficacy, cost-effectiveness and acceptability of self-help interventions for anxiety disorders: systematic review. *Br J Psychiatry*. 2012;200(1):15–21.
- Reger MA, Gahm GA. A meta-analysis of the effects of internet- and computer-based cognitive-behavioral treatments for anxiety. *J Clin Psychol*. 2009;65(1):53–75.
- Wischmann T. Implications of psychosocial support in infertility – a critical appraisal. *J Psychosom Obstet Gynaecol*. 2008;29(2):83–90.

Psychology Research and Behavior Management

Dovepress

Publish your work in this journal

Psychology Research and Behavior Management is an international, peer-reviewed, open access journal focusing on the science of psychology and its application in behavior management to develop improved outcomes in the clinical, educational, sports and business arenas. Specific topics covered include: Neuroscience, memory & decision making; Behavior

modification & management; Clinical applications; Business & sports performance management; Social and developmental studies; Animal studies. The manuscript management system is completely online and includes a quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/psychology-research-and-behavior-management-journal>

Peripheral natural killer cell activity as a predictor of recurrent pregnancy loss: a large cohort study

Kinue Katano, M.D.,^a Sadao Suzuki, M.D.,^b Yasuhiko Ozaki, M.D.,^a Nobuhiro Suzumori, M.D.,^a Tamao Kitaori, M.D.,^a and Mayumi Sugiura-Ogasawara, M.D.^a

Departments of ^a Obstetrics and Gynecology and ^b Public Health, Nagoya City University, Graduate School of Medical Sciences, Nagoya, Japan

Objective: To determine the predictive value of preconceptional peripheral blood natural killer (pNK) cell activity in patients with recurrent pregnancy loss (RPL).

Design: Cohort study.

Setting: University department.

Patient(s): A total of 552 patients with a history of two to six consecutive miscarriages.

Intervention(s): None.

Main Outcome Measure(s): The predictive value of preconceptional pNK cell activity for subsequent miscarriage was analyzed using multivariable logistic regression analysis, with age, number of previous miscarriages, and presence/absence of previous live births and bed rest as covariates.

Result(s): Age and number of previous miscarriages, but not high pNK cell activity, were found to be independent risk factors for a subsequent miscarriage. No effect of bed rest and previous live birth on the likelihood of live birth was observed (odds ratios 1.28 [95% confidence interval 0.81–2.02] and 0.91 [0.52–1.59], respectively).

Conclusion(s): Elevated pNK cell activity was found to not be an independent risk factor for subsequent miscarriage. Clinicians should not measure the plasma NK activity as a systematic recurrent pregnancy loss examination, because its clinical significance is yet to be established. (Fertil Steril® 2013;100:1629–34. ©2013 by American Society for Reproductive Medicine.)

Key Words: Recurrent pregnancy loss, natural killer cell activity, predictor, cohort study

Discuss: You can discuss this article with its authors and with other ASRM members at <http://fertstertforum.com/katanok-natural-killer-cell-recurrent-pregnancy-loss/>



Use your smartphone to scan this QR code and connect to the discussion forum for this article now.*

* Download a free QR code scanner by searching for "QR scanner" in your smartphone's app store or app marketplace.

Established causes of recurrent pregnancy loss (RPL) include presence of antiphospholipid antibodies in the serum, presence of uterine anomalies, and presence of abnormal chromosomes, particularly translocations, in either

partner (1–3). According to previous reports, in approximately half of the cases seen at research centers, the cause of RPL remains unexplained despite conventional examinations conducted to identify the cause (4–6).

Cytotrophoblasts that express human leukocyte antigen G (HLA-G) come in direct contact with maternal lymphocytes. Many natural killer (NK)-like large granular lymphocytes have been detected in the human decidua of early pregnancy (7). Large numbers of NK cells appear in the mid-secretory phase. Natural killer cells have been thought to play a key role in the establishment of successful pregnancy by facilitating immunologic adaptation of the semiallogenic developing embryo. Recently, Fu et al. (8) reported that recruitment of TH17 cells and local inflammation can occur at the maternal–fetal interface during natural allogenic pregnancies, and that decidual NK cells promote immune tolerance and successful pregnancy

Received April 17, 2013; revised and accepted July 26, 2013; published online September 4, 2013.

K.K. has nothing to disclose. S.S. has nothing to disclose. Y.O. has nothing to disclose. N.S. has nothing to disclose. T.K. has received payment for manuscript preparation by the Japan Society of Thrombosis Haemostasis. M.S.-O. has received payment for lectures from Bayer, Sanofi-Aventis, Eisai Co., Kissei Pharmaceutical Co., Nippon Shinyaku Co., Dainippon Sumitomo Pharma, Takeda Pharmaceutical Company Limited, Fuji Pharma Co., and Mochida Pharmaceutical Co.; payment for manuscript preparation by Igaku shoin Co. and Chunichi Shinbun; and payment for travel expenses by the Korean Society for Reproductive Medicine.

This study was supported by a Grant-in-Aid for Nagoya City University Hospital. N.S. and M.S.-O. are supported by a grant from the Japanese Ministry of Education, Science, and Technology. T.K. and M.S.-O. are supported by a grant from the Japanese Ministry of Health, Labor and Welfare.

Reprint requests: Mayumi Sugiura-Ogasawara, M.D., Department of Obstetrics and Gynecology, Nagoya City University, Graduate School of Medical Sciences, Mizuho-ku, Nagoya 4678601, Japan (E-mail: og.mym@med.nagoya-cu.ac.jp).

Fertility and Sterility® Vol. 100, No. 6, December 2013 0015-0282/\$36.00

Copyright ©2013 American Society for Reproductive Medicine. Published by Elsevier Inc. <http://dx.doi.org/10.1016/j.fertstert.2013.07.1996>

by diminishing the recruitment of inflammatory TH17 cells via interferon- γ secreted by the CD56^{bright}CD27⁺ NK cell subset. This NK cell-mediated regulatory response is lost in patients with recurrent spontaneous abortions, resulting in a prominent TH17 response and extensive local inflammation.

We previously reported (9) that elevated preconceptional peripheral blood NK (pNK) cell activity may be predictive of subsequent miscarriage in 68 patients with RPL. The 24 women with high pNK cell activity, defined as a value equal to or exceeding the mean pNK cell activity of the 47 controls + 1 SD, had a significantly higher miscarriage rate in the subsequent pregnancy than the 44 women with normal pNK cell activity (71% vs. 20%; relative risk 3.5; 95% confidence interval [CI] 1.8–6.5). However, some studies have shown no differences in pNK cell parameters between patients with RPL and controls (10–12). Whereas 90% of pNK cells are CD56^{dim} and CD16⁺, 80% of the uterine NK cells are CD56^{bright} and CD16⁻ (13, 14). Peripheral blood NK cells are phenotypically and functionally different from uterine NK cells. Decidual leukocytes have low cytotoxic activity as compared with peripheral lymphocytes (15).

The sample size of our previous study was too small. The prognostic value of measuring pNK cell parameters remains uncertain (16). Therefore, we conducted a cohort study of 552 patients with RPL to determine whether pNK might indeed be predictive of subsequent miscarriage.

MATERIALS AND METHODS

Patients

We studied 1,127 patients with a history of two or more (2 to 12) consecutive miscarriages, in whom the study examinations could be completed and subsequent pregnancies were established between January 1996 and May 2011 in Nagoya City University Hospital. Patients with identifiable causes and patients who received any kinds of treatment were excluded from the present cohort.

The conventional examinations were completed in all patients, including hysterosalpingography, transvaginal ultrasonography, chromosomal analysis of both partners, determination of the presence/absence of antiphospholipid antibodies, including lupus anticoagulant (LA), by diluted activated partial thromboplastin time (aPTT), diluted Russel viper venom time (RVVT), and β 2 glycoprotein I-dependent anticardiolipin antibody methods (17), and blood tests for hypothyroidism and diabetes mellitus, before the subsequent pregnancy.

Antiphospholipid antibody syndrome (APS) was diagnosed according to the criteria of the International Congress on Antiphospholipid Antibodies (18). Patients with APS were treated with low-dose aspirin plus heparin (19). Diabetes mellitus, hyper- or hypothyroidism, and hyperprolactinemia in the patients were controlled with medication before conception. Patients with three or more unexplained miscarriages received paternal mononuclear cell immunization from 1996 to 1999, a biologic response modifier from 1996 to 2004 (20), and low-dose aspirin and/or heparin from 2000 to 2007.

Gestational age was calculated from basal body temperature charts. A total of 654 patients were admitted for rest, and

ultrasonography was performed twice per week from 4 to 8 weeks of gestation before 2004. The pregnancy outcome of 473 patients was followed once per week by ultrasonography without admission after 2004.

The study was conducted with the approval of the Research Ethics Committee at Nagoya City University Medical School.

Measurement

Preconceptional pNK cell activity was examined in the midsecretory phase. Peripheral blood NK cell activity was measured by a chromium-51 release cytotoxicity assay, with K562 human myeloid leukemia cells as the targets. A total of 3.7×10^3 Bq ⁵¹Cr-labeled target cells (1×10^4 per well) were seeded with 2×10^5 effector cells per well (fresh peripheral blood mononuclear cells) in triplicate, in U-bottomed 96-well plates. After 4-hour incubation at 37°C, the activity in the supernatant from each well was measured in an autogramma scintillation counter. The percentage cytotoxicity was calculated as follows: $[(\text{test cpm} - \text{spontaneous cpm}) / (\text{maximum cpm} - \text{spontaneous cpm})] \times 100$, where cpm = counts per minute.

Lupus anticoagulant was detected using fivefold diluted aPTT methods, as previously described, with brain cephalin (Automated aPTT; Organon Teknica) as the phospholipid reagent (17). The 1:1 mixing test was performed at the same time. The clotting time was measured using an Option 4 Biomerieux calculator. Lupus anticoagulant was considered positive when prolonged clotting times ($>$ mean + 3 SD of 104 healthy nonpregnant control plasma, 7.37 seconds) failed to correct when mixed 1:1 with standard plasma. Diluted Russel viper venom time for LA was performed as previously described (Gradipore). To detect β 2 glycoprotein I-dependent anticardiolipin antibody, we used a modified ELISA system (Yamasa).

Analysis

Patients with identifiable causes and patients who received any kinds of treatments were excluded from the present cohort. Biochemical pregnancy, ectopic pregnancy, and hydatidiform mole were excluded from the analysis of the subsequent pregnancy outcome.

Miscarriage rate was analyzed according to pNK cell activity, age, number of previous miscarriages, and presence/absence of previous live births and bed rest. Peripheral blood NK cells, age, and previous number of miscarriages were categorized into quartiles, because they showed normal distribution. The previous number of miscarriages was categorized into two, three, four, or five to six.

Crude logistic regression was performed to examine the predictive value of pNK cell activity for subsequent miscarriage. We also examined the influence of age, previous number of miscarriages, and presence/absence of previous live birth and bed rest on the likelihood of subsequent miscarriage.

Age is well known to influence the miscarriage rate. Age is also associated with number of previous miscarriages and number of previous live births. Thus, first, we chose