

and for exploring possible therapeutic targets for each tumor type.

ACKNOWLEDGMENTS

This study was supported by the Core Research for Evolutional Science and Technology (CREST) division “Development of Fundamental Technologies for Diagnosis and Therapy Based upon Epigenome Analysis” of the Japan Science and Technology Agency (JST), the Program for Promotion of Fundamental Studies in Health Sciences from the NiBio, a Grant in Aid for the Third Term Comprehensive 10-Year Strategy for Cancer Control from the Ministry of Health, Labor and Welfare of Japan, and Grants in Aid for Scientific Research (B) and (C) from the Japan Society for the Promotion of Science (JSPS). Tissue specimens were provided by the National Cancer Center Biobank, which is supported by the National Cancer Center Research and Development Fund, Japan.

REFERENCES

- Arai, E., Chiku, S., Mori, T., Gotoh, M., Nakagawa, T., Fujimoto, H., et al. (2012). Single-CpG-resolution methylome analysis identifies clinicopathologically aggressive CpG island methylator phenotype clear cell renal cell carcinomas. *Carcinogenesis* 33, 1487–1493. doi: 10.1093/carcin/bgs177
- Arai, E., and Kanai, Y. (2010). DNA methylation profiles in precancerous tissue and cancers: carcinogenetic risk estimation and prognostication based on DNA methylation status. *Epigenomics* 2, 467–481. doi: 10.2217/epi.10.16
- Arai, E., Kanai, Y., Ushijima, S., Fujimoto, H., Mukai, K., and Hirohashi, S. (2006). Regional DNA hypermethylation and DNA methyltransferase (DNMT) 1 protein overexpression in both renal tumors and corresponding nontumorous renal tissues. *Int. J. Cancer* 119, 288–296. doi:10.1002/ijc.21807
- Arai, E., Ushijima, S., Fujimoto, H., Hosoda, F., Shibata, T., Kondo, T., et al. (2009a). Genome-wide DNA methylation profiles in both precancerous conditions and clear cell renal cell carcinomas are correlated with malignant potential and patient outcome. *Carcinogenesis* 30, 214–221. doi: 10.1093/carcin/bgn268
- Arai, E., Ushijima, S., Gotoh, M., Ojima, H., Kosuge, T., Hosoda, F., et al. (2009b). Genome-wide DNA methylation profiles in liver tissue at the precancerous stage and in hepatocellular carcinoma. *Int. J. Cancer* 125, 2854–2862. doi: 10.1002/ijc.24708
- Bae, J. B. (2013). Perspectives of international human epigenome consortium. *Genomics Inform.* 11, 7–14. doi: 10.5808/GI.2013.11.1.7
- Djebali, S., Davis, C. A., Merkel, A., Dobin, A., Lassmann, T., Mortazavi, A., et al. (2012). Landscape of transcription in human cells. *Nature* 489, 101–108. doi: 10.1038/nature11233
- Etoh, T., Kanai, Y., Ushijima, S., Nakagawa, T., Nakanishi, Y., Sasako, M., et al. (2004). Increased DNA methyltransferase 1 (DNMT1) protein expression correlates significantly with poorer tumor differentiation and frequent DNA hypermethylation of multiple CpG islands in gastric cancers. *Am. J. Pathol.* 164, 689–699. doi: 10.1016/S0002-9440(10)63156-2
- Fabbri, M., Garzon, R., Cimmino, A., Liu, Z., Zanesi, N., Callegari, E., et al. (2007). MicroRNA-29 family reverts aberrant methylation in lung cancer by targeting DNA methyltransferases 3A and 3B. *Proc. Natl. Acad. Sci. U.S.A.* 104, 15805–15810. doi:10.1073/pnas.0707628104
- Gomes, A. Q., Nolasco, S., and Soares, H. (2013). Non-coding RNAs: multi-tasking molecules in the cell. *Int. J. Mol. Sci.* 14, 16010–16039. doi: 10.3390/ijms140816010
- Gotoh, M., Arai, E., Wakai-Ushijima, S., Hiraoka, N., Kosuge, T., Hosoda, F., et al. (2011). Diagnosis and prognostication of ductal adenocarcinomas of the pancreas based on genome-wide DNA methylation profiling by bacterial artificial chromosome array-based methylated CpG island amplification. *J. Biomed. Biotechnol.* 2011, 780836. doi: 10.1155/2011/780836
- He, L., and Hannon, G. J. (2004). MicroRNAs: small RNAs with a big role in gene regulation. *Nat. Rev. Genet.* 5, 522–531. doi: 10.1038/nrg1379
- Huang, J., Wang, Y., Guo, Y., and Sun, S. (2010). Down-regulated microRNA-152 induces aberrant DNA methylation in hepatitis B virus-related hepatocellular carcinoma by targeting DNA methyltransferase 1. *Hepatology* 52, 60–70. doi: 10.1002/hep.23660
- Jinek, M., and Doudna, J. A. (2009). A three-dimensional view of the molecular machinery of RNA interference. *Nature* 457, 405–412. doi: 10.1038/nature07755
- Kanai, Y. (2008). Alterations of DNA methylation and clinicopathological diversity of human cancers. *Pathol. Int.* 58, 544–558. doi: 10.1111/j.1440-1827.2008.02270.x
- Kanai, Y. (2010). Genome-wide DNA methylation profiles in precancerous conditions and cancers. *Cancer Sci.* 101, 36–45. doi: 10.1111/j.1349-7006.2009.01383.x
- Kanai, Y., and Hirohashi, S. (2007). Alterations of DNA methylation associated with abnormalities of DNA methyltransferases in human cancers during transition from a precancerous to a malignant state. *Carcinogenesis* 28, 2434–2442. doi: 10.1093/carcin/bgm206
- Kanai, Y., Ushijima, S., Tsuda, H., Sakamoto, M., Sugimura, T., and Hirohashi, S. (1996). Aberrant DNA methylation on chromosome 16 is an early event in hepatocarcinogenesis. *Jpn. J. Cancer Res.* 87, 1210–1217. doi: 10.1111/j.1349-7006.1996.tb03135.x
- Kim, D. H., Saetrom, P., Snøve, O. Jr., and Rossi, J. J. (2008). MicroRNA-directed transcriptional gene silencing in mammalian cells. *Proc. Natl. Acad. Sci. U.S.A.* 105, 16230–16235. doi: 10.1073/pnas.0808830105
- Lister, R., O’Malley, R. C., Tonti-Filippini, J., Gregory, B. D., Berry, C. C., Millar, A. H., et al. (2008). Highly integrated single-base resolution maps of the epigenome in *Arabidopsis*. *Cell* 133, 523–536. doi: 10.1016/j.cell.2008.03.029
- Lujambio, A., Ropero, S., Ballestar, E., Fraga, M. F., Cerrato, C., Setién, F., et al. (2007). Genetic unmasking of an epigenetically silenced microRNA in human cancer cells. *Cancer Res.* 67, 1424–1429. doi: 10.1158/0008-5472.CAN-06-4218
- Miura, F., Enomoto, Y., Dairiki, R., and Ito, T. (2012). Amplification-free whole-genome bisulfite sequencing by post-bisulfite adaptor tagging. *Nucleic Acids Res.* 40, e136. doi: 10.1093/nar/gks454
- Nakagawa, T., Kanai, Y., Ushijima, S., Kitamura, T., Kakizoe, T., and Hirohashi, S. (2005). DNA hypermethylation on multiple CpG islands associated with increased DNA methyltransferase DNMT1 protein expression during multistage urothelial carcinogenesis. *J. Urol.* 173, 1767–1771. doi: 10.1097/01.ju.0000154632.11824.4d
- Nishiyama, N., Arai, E., Chihara, Y., Fujimoto, H., Hosoda, F., Shibata, T., et al. (2010). Genome-wide DNA methylation profiles in urothelial carcinomas and urothelia at the precancerous stage. *Cancer Sci.* 101, 231–240. doi: 10.1111/j.1349-7006.2009.01330.x
- Peng, D. F., Kanai, Y., Sawada, M., Ushijima, S., Hiraoka, N., Kitazawa, S., et al. (2006). DNA methylation of multiple tumor-related genes in association with overexpression of DNA methyltransferase 1 (DNMT1) during multistage carcinogenesis of the pancreas. *Carcinogenesis* 27, 1160–1168. doi: 10.1093/carcin/bgi361
- Saito, Y., Kanai, Y., Sakamoto, M., Saito, H., Ishii, H., and Hirohashi, S. (2002). Overexpression of a splice variant of DNA methyltransferase 3b, DNMT3b4, associated with DNA hypomethylation on pericentromeric satellite regions during human hepatocarcinogenesis. *Proc. Natl. Acad. Sci. U.S.A.* 99, 10060–10065. doi: 10.1073/pnas.152121799
- Saito, Y., Liang, G., Egger, G., Friedman, J. M., Chuang, J. C., and Coetzee, G. A., et al. (2006). Specific activation of microRNA-127 with downregulation of the proto-oncogene BCL6 by chromatin-modifying drugs in human cancer cells. *Cancer Cell* 9, 435–443. doi: 10.1016/j.ccr.2006.04.020
- Saito, Y., Saito, H., Liang, G., and Friedman, J. M. (2013). Epigenetic alterations and microRNA misexpression in cancer and autoimmune diseases: a critical review. *Clin. Rev. Allergy Immunol.* doi: 10.1007/s12016-013-8401-z [Epub ahead of print].
- Saito, Y., Suzuki, H., Tsugawa, H., Nakagawa, I., Matsuzaki, J., Kanai, Y., et al. (2009). Chromatin remodeling at Alu repeats by epigenetic treatment activates silenced microRNA-512-5p with downregulation of Mcl-1 in human gastric cancer cells. *Oncogene* 28, 2738–2744. doi: 10.1038/onc.2009.140
- Sato, T., Arai, E., Kohno T., Takahashi, Y., Miyata, S., Tsuta, K., et al. (2014). Epigenetic clustering of lung adenocarcinomas based on DNA methylation profiles in adjacent lung tissue: its correlation with smoking history and chronic obstructive pulmonary disease. *Int. J. Cancer* doi: 10.1002/ijc.28684
- Sato, T., Arai, E., Kohno, T., Tsuta, K., Watanabe, S., Soejima, K., et al. (2013). DNA methylation profiles at precancerous stages associated with recurrence of lung adenocarcinoma. *PLoS ONE* 8:e59444. doi: 10.1371/journal.pone.0059444

Yoshiura, K., Kanai, Y., Ochiai, A., Shimoyama, Y., Sugimura, T., and Hirohashi, S. (1995). Silencing of the E-cadherin invasion-suppressor gene by CpG methylation in human carcinomas. *Proc. Natl. Acad. Sci. U.S.A.* 92, 7416–7419. doi: 10.1073/pnas.92.16.7416

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 14 January 2014; accepted: 24 January 2014; published online: 14 February 2014.

Citation: Kanai Y and Arai E (2014) Multilayer-omics analyses of human cancers: exploration of biomarkers and drug targets based on the activities of the International Human Epigenome Consortium. Front. Genet. 5:24. doi: 10.3389/fgene.2014.00024
This article was submitted to *Epigenomics and Epigenetics*, a section of the journal *Frontiers in Genetics*.

Copyright © 2014 Kanai and Arai. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

DNA Methylation Profiles at Precancerous Stages Associated with Recurrence of Lung Adenocarcinoma

Takashi Sato^{1,2}, Eri Arai^{1*}, Takashi Kohno³, Koji Tsuta⁴, Shun-ichi Watanabe⁵, Kenzo Soejima², Tomoko Betsuyaku², Yae Kanai¹

1 Division of Molecular Pathology, National Cancer Center Research Institute, Tokyo, Japan, **2** Division of Pulmonary Medicine, Department of Medicine, Keio University School of Medicine, Tokyo, Japan, **3** Division of Genome Biology, National Cancer Center Research Institute, Tokyo, Japan, **4** Department of Pathology and Clinical Laboratories, Pathology Division, National Cancer Center Hospital, Tokyo, Japan, **5** Department of Thoracic Oncology, Thoracic Surgery Division, National Cancer Center Hospital, Tokyo, Japan

Abstract

The aim of this study was to clarify the significance of DNA methylation alterations at precancerous stages of lung adenocarcinoma. Using single-CpG resolution Infinium array, genome-wide DNA methylation analysis was performed in 36 samples of normal lung tissue obtained from patients without any primary lung tumor, 145 samples of non-cancerous lung tissue (N) obtained from patients with lung adenocarcinomas, and 145 samples of tumorous tissue (T). Stepwise progression of DNA methylation alterations from normal lung tissue to non-cancerous lung tissue obtained from patients with lung adenocarcinomas, and then tumorous tissue samples, was observed at 3,270 CpG sites, suggesting that non-cancerous lung tissue obtained from patients with lung adenocarcinomas was at precancerous stages with DNA methylation alterations. At CpG sites of 2,083 genes, DNA methylation status in samples of non-cancerous lung tissue obtained from patients with lung adenocarcinomas was significantly correlated with recurrence after establishment of lung adenocarcinomas. Among such recurrence-related genes, 28 genes are normally unmethylated (average β -values based on Infinium assay in normal lung tissue samples was less than 0.2) and their DNA hypermethylation at precancerous stages was strengthened during progression to lung adenocarcinomas ($\Delta\beta_{T-N} > 0.1$). Among these 28 genes, we focused on 6 for which implications in transcription regulation, apoptosis or cell adhesion had been reported. DNA hypermethylation of the *ADCY5*, *EVX1*, *GFRA1*, *PDE9A*, and *TBX20* genes resulted in reduced mRNA expression in tumorous tissue samples. 5-Aza-2'-deoxycytidine treatment of lung cancer cell lines restored the mRNA expression levels of these 5 genes. Reduced mRNA expression in tumorous tissue samples was significantly correlated with tumor aggressiveness. These data suggest that DNA methylation alterations at precancerous stages determine tumor aggressiveness and outcome through silencing of specific genes.

Citation: Sato T, Arai E, Kohno T, Tsuta K, Watanabe S-i, et al. (2013) DNA Methylation Profiles at Precancerous Stages Associated with Recurrence of Lung Adenocarcinoma. PLoS ONE 8(3): e59444. doi:10.1371/journal.pone.0059444

Editor: Bernard W Futscher, The University of Arizona, United States of America

Received: September 25, 2012; **Accepted:** February 14, 2013; **Published:** March 27, 2013

Copyright: © 2013 Sato et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was supported by the Program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation (NIBio, 10-42, <http://www.nibio.go.jp/part/promote/fundamental/doc/index.html>) and partially supported by a Grant-in-Aid for the Third Term Comprehensive 10-Year Strategy for Cancer Control from the Ministry of Health, Labor and Welfare of Japan (2, <http://www.mhlw.go.jp/bunya/kenkyuujigyoku/hojokin-koubo14/09.html>), National Cancer Center Research and Development Fund (23-A-11, http://www.ncc.go.jp/jp/about/rinri/kaihatsu/files/h24_ncc_research_list.pdf), and Grants-in-Aid for Scientific Research (B, 233900 90, <http://www.jsps.go.jp/j-grantsinaid/>) and for Young Scientists (B, 23790456, <http://www.jsps.go.jp/j-grantsinaid/>) from the Japan Society for the Promotion of Science (JSPS). National Cancer Center Biobank is supported by the National Cancer Center Research and Development Fund (23-A-1, http://www.ncc.go.jp/jp/about/rinri/kaihatsu/files/h24_ncc_research_list.pdf), Japan. T. Sato is an awardee of a research resident fellowship from the Foundation for Promotion of Cancer Research in Japan (<http://www.fpcr.or.jp/index.html>). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: earai@ncc.go.jp

Introduction

Lung adenocarcinoma (LADC) is increasingly recognized as a clinicopathologically and molecularly heterogeneous disease: frequent mutations of the *EGFR*, *KRAS*, *BRAF*, *TP53*, *ERBB2*, *PIK3CA* and *MET* genes and *EML4-ALK* fusions have been reported in LADCs [1,2]. In addition, recent whole-exome sequencing has revealed frequent mutation of the *CSMD3* gene [3]. However, the molecular background responsible for the clinicopathological diversity of LADCs is not yet fully understood.

As well as genetic abnormalities, epigenetic changes in human cancers have also been described [4–7]. In LADCs, silencing of the *RASSF1A*, *CDKN2A*, *RAR β* , *MGMT*, *APC*, *DAPK*, *FHIT* and *CDH13* genes due to DNA hypermethylation around their

promoter regions has been frequently observed [8]. In addition, DNA methylation alterations are known to occur even at the early and precancerous stages of carcinogenesis in various organs [5–7,9]. For example, we have reported that DNA hypermethylation at the D17S5 locus, where the *HIC-1* tumor suppressor gene has been identified, is evident even in non-cancerous lung tissue obtained from patients with non-small cell lung cancers, and is correlated with smoking history [10]. Other researchers have also reported DNA hypermethylation of specific tumor-related genes at precancerous stages associated with cigarette smoking [8,11]. However, it has been unclear whether DNA methylation status is simply altered at precancerous stages or whether DNA methylation alterations at these stages actually result in gene expression alterations in established LADCs. Moreover, in organs other than




Figure 1. Hazard ratio (HR) obtained from the Cox regression model. Correlation between DNA methylation status (average β -values) and recurrence was examined in 145 samples of tumorous tissue (A) and 145 samples of the corresponding non-cancerous lung tissue (B) obtained from patients with lung adenocarcinomas who had undergone complete resection and had not received any adjuvant therapy after surgery. All of the examined 26,455 probes of the Infinium array are shown along the chromosomes. Red color means that higher β -values of the probes were observed in recurrence-positive patients than in recurrence-negative patients ($P < 0.001$). Blue color means that lower β -values of the probes were observed in recurrence-positive patients than in recurrence-negative patients ($P < 0.001$). doi:10.1371/journal.pone.0059444.g001

the lung, it has been suggested that DNA methylation profiles at precancerous stages may determine tumor aggressiveness and outcome [12–14]. However, the clinicopathological impact of DNA methylation alterations at precancerous stages during lung carcinogenesis has not been clarified.

Recently, genome-wide DNA methylation analysis using the single-CpG resolution Infinium array has made it possible to interrogate 27,000 highly informative CpG sites, i.e. an average of two CpG sites within the proximal promoter regions of the transcription start sites of each of 14,475 consensus coding sequences in the National Center for Biotechnology Information Database, especially 3 to 20 CpG sites for more than 200 cancer-related and imprinted genes [15]. Although a few studies of lung cancers employing the Infinium assay have been reported [16,17], they did not focus on precancerous stages. In order to clarify the significance of DNA methylation alterations at precancerous stages of lung carcinogenesis, we performed the Infinium assay in association with mRNA expression and clinicopathological analyses of 36 samples of normal lung tissue (C) obtained from patients without any primary lung tumors, 145 samples of non-cancerous lung tissue (N) from patients with LADCs, and 145 corresponding samples of tissue from the tumors (T) themselves. Although the molecular classification of LADCs based on the results of the

Infinium assay will be published elsewhere, we focused on specific genes methylated at precancerous stages in the present study.

Materials and Methods

Patients and Tissue Samples

The 145 paired samples of N and the corresponding T were obtained from patients with primary LADCs who underwent lung resection at the National Cancer Center Hospital, Japan, between December 1997 and March 2008. These patients had undergone complete resection and had not received any preoperative treatment or adjuvant therapy after surgery. Eighty-one patients were males and 64 were females with a median age of 61 years (range, 30–81 yr). Histological diagnosis and grading were based on the 2004 World Health Organization classification [18]. Recurrence was diagnosed by clinicians on the basis of physical examination and imaging modalities such as computed tomography, magnetic resonance imaging, scintigraphy or positron-emission tomography, and sometimes confirmed histopathologically by biopsy.

For comparison, 36 C samples were obtained from materials that had been surgically resected from patients without any primary lung tumor. Twenty-two of these patients were males and 14 were females, with a median age of 63 years (range, 27–83 yr).

Table 1. The 28 probes for which the average β -value in N samples (β_N) was higher in recurrence-positive patients than in recurrence-negative patients, for which the average β -value in C samples (β_C) was less than 0.2, and for which the average β -value in T samples minus that in corresponding N samples ($\Delta\beta_{T-N}$) was more than 0.1.

Target ID ^a	Chromosome	Position ^b	Gene symbol	P^c	Adjusted P^d
cg00516481	21	44,073,202	<i>PDE9A</i>	9.996×10^{-4}	1.193×10^{-2}
cg01295203	8	70,984,199	<i>PRDM14</i>	8.156×10^{-5}	3.236×10^{-3}
cg02008154	7	35,293,537	<i>TBX20</i>	7.188×10^{-4}	9.917×10^{-3}
cg02909790	6	26,271,587	<i>HIST1H3G</i>	2.896×10^{-6}	5.787×10^{-4}
cg03538436	12	117,799,370	<i>NOS1</i>	7.227×10^{-4}	9.944×10^{-3}
cg03963198	5	1,882,871	<i>IRX4</i>	5.133×10^{-4}	8.186×10^{-3}
cg06005396	19	590,541	<i>HCN2</i>	9.485×10^{-4}	1.158×10^{-2}
cg06269753	8	72,755,871	<i>MSC</i>	8.627×10^{-7}	3.426×10^{-4}
cg07651242	7	45,614,720	<i>ADCY1</i>	7.403×10^{-4}	1.010×10^{-2}
cg11612345	6	168,842,491	<i>SMOC2</i>	4.264×10^{-6}	6.753×10^{-4}
cg12087643	10	118,033,370	<i>GFRA1</i>	3.144×10^{-9}	2.489×10^{-5}
cg12265829	14	24,804,022	<i>ADCY4</i>	7.265×10^{-4}	9.984×10^{-3}
cg13262687	4	147,559,579	<i>POU4F2</i>	6.246×10^{-5}	2.795×10^{-3}
cg13449778	1	179,712,298	<i>FAM163A</i>	8.195×10^{-7}	3.426×10^{-4}
cg13878010	3	123,167,276	<i>ADCY5</i>	2.339×10^{-6}	5.220×10^{-4}
cg16254309	7	145,814,152	<i>CNTNAP2</i>	2.917×10^{-4}	6.240×10^{-3}
cg16387606	1	149,804,293	<i>HIST2H4A</i>	7.441×10^{-4}	1.011×10^{-2}
cg16604516	3	13,590,419	<i>FBLN2</i>	7.181×10^{-4}	9.916×10^{-3}
cg16652259	2	172,949,501	<i>DLX1</i>	4.175×10^{-4}	7.386×10^{-3}
cg17191178	3	157,824,217	<i>SHOX2</i>	8.872×10^{-5}	3.353×10^{-3}
cg18454685	17	48,639,239	<i>CACNA1G</i>	4.326×10^{-4}	7.487×10^{-3}
cg20286200	6	133,562,267	<i>EYA4</i>	1.541×10^{-9}	2.038×10^{-5}
cg21087137	12	75,728,469	<i>GLIPR1L1</i>	1.965×10^{-6}	4.821×10^{-4}
cg22461835	8	26,723,365	<i>ADRA1A</i>	7.297×10^{-4}	9.998×10^{-3}
cg23418591	20	57,090,317	<i>LOC149773</i>	4.075×10^{-4}	7.287×10^{-3}
cg25302419	5	11,904,015	<i>CTNND2</i>	5.149×10^{-4}	8.194×10^{-3}
cg25764191	10	105,037,215	<i>INA</i>	3.834×10^{-4}	7.026×10^{-3}
cg27626299	7	27,282,431	<i>EVX1</i>	2.164×10^{-4}	5.423×10^{-3}

^aProbe ID for the Infinium HumanMethylation27 Bead Array (Illumina).

^bNational Center for Biotechnology Information database (Genome Build 37).

^cNon-adjusted P -values and.

^dBenjamini-Hochberg-adjusted P -values for the Cox regression model used for evaluation of correlation with recurrence.

doi:10.1371/journal.pone.0059444.t001

Thirty-five had undergone lung resection for metastatic lesions of primary cancers of the colon, rectum, kidney, urinary bladder, thyroid, breast, pancreas, ampulla of Vater and salivary gland, osteosarcoma, synovial sarcoma, leiomyosarcoma, rhabdomyosarcoma, liposarcoma, dermatofibrosarcoma, and myxofibrosarcoma. The remaining one patient had undergone chest wall resection for lipoma with removal of adjacent lung tissue. Histological observation confirmed that all of the C samples showed no remarkable histological abnormality and did not contain any contaminating tumor cells that had metastasized from organs other than the lung.

Tissue specimens were provided by the National Cancer Center Biobank, Japan. This study was approved by the Ethics Committee of the National Cancer Center, Tokyo, Japan, and was performed in accordance with the Declaration of Helsinki 1975. All patients included in this study provided written informed consent.

Cell Lines

The characteristics of the four lung cancer cell lines used in this study are summarized in Table S1.

Infinium Assay

Genomic DNA was extracted using a QIAamp DNA Mini kit (Qiagen, Valencia, CA, USA) and phenol-chloroform extraction followed by dialysis [19] from all tissue samples and cell lines, respectively. Five-hundred-nanogram aliquots of DNA were subjected to bisulfite conversion using an EZ DNA Methylation-Gold Kit (Zymo Research, Irvine, CA, USA). DNA methylation status at 27,578 CpG loci was examined at single-CpG resolution using the Infinium HumanMethylation27 Bead Array (Illumina, San Diego, CA, USA). After hybridization, the specifically hybridized DNA was fluorescence-labeled by a single-base extension reaction and detected using a BeadScan reader (Illumina) in accordance with the manufacturer's protocols. The data were then assembled using GenomeStudio methylation




Figure 2. Correlation between DNA methylation levels and mRNA expression levels. DNA methylation levels (average β -values) (A) and mRNA expression levels (B) for the *ADCY5*, *CNTNAP2*, *EVX1*, *GFRA1*, *PDE9A* and *TBX20* genes in samples of non-cancerous lung tissue (N) from patients with lung adenocarcinomas and samples of the corresponding tumorous tissue (T) were determined by Infinium assay and quantitative real-time reverse transcription-PCR analysis, respectively. DNA methylation levels for all six genes were significantly higher in T samples than in N samples, and levels of expression of mRNAs for the *ADCY5*, *EVX1*, *GFRA1*, *PDE9A* and *TBX20* genes were significantly lower in T samples than in N samples, although the reduction in the expression of the *CNTNAP2* gene did not reach statistical significance. These results suggested that DNA hypermethylation of the *ADCY5*, *EVX1*, *GFRA1*, *PDE9A* and *TBX20* genes may result in reduced mRNA expression in tissue samples from the same cohort. doi:10.1371/journal.pone.0059444.g002

software (Illumina). At each CpG site, the ratio of the fluorescence signal was measured using a methylated probe relative to the sum of the methylated and unmethylated probes, i.e. the so-called β -value, which ranges from 0.00 to 1.00, reflecting the methylation level of an individual CpG site.

Quantitative Real-time Reverse Transcription (RT)-PCR Analysis

Total RNA was extracted from 132 N and 151 T samples for which additional tissue specimens were available and cell lines using TRIzol reagent (Life Technologies, Carlsbad, CA, USA) in accordance with the manufacturer's instructions. cDNA was synthesized from total RNA with random primers using SuperScript III Reverse Transcriptase (Life Technologies) and pre-amplified using TaqMan PreAmp Master Mix (Life Technologies).

To evaluate mRNA expression levels, fluorescence-labeled locked nucleic acid hydrolysis probes were selected from the Universal Probe Library collection (Roche Applied Science, Mannheim, Germany) and specific PCR primers yielding intron-spanning amplicons were designed using ProbeFinder assay design software (<https://www.roche-applied-science.com/sis/rtpcr/upl/index.jsp?id=UPO30000>). The probe ID and primer sequences are summarized in Table S2. Quantitative real-time PCR was performed using TaqMan Universal Master Mix II (Life Technologies) and the relative standard curve method in the BioMark HD System (Fluidigm, South San Francisco, CA, USA). Ct values were normalized to that of *GAPDH* in the same sample. All assays were performed in triplicate.

5-aza-2'-deoxycytidine (5-aza-dC) Treatment

A549, PC9, VMRC-LCD and EBC-1 cells were seeded at a density of 9×10^5 cells per 15-cm dish on day 0 and then allowed to attach for a 24-h period. Then, 5-aza-dC (Sigma-Aldrich, St. Louis, MO, USA) was added to a final concentration of 1 μ M. Cells were passaged at a subculture ratio of 1:2 on day 3. At 24 h after replating, 5-aza-dC was added again to the same final concentration. Since toxicity had been obvious during preliminary experiments, the final concentration of 5-aza-dC was reduced to 0.1 μ M for EBC-1 cells. Genomic DNA and total RNA were extracted from all cells on days 3 and 6.

Statistics

In the Infinium assay, all CpG sites on chromosomes X and Y were excluded, to avoid any gender-specific methylation bias. The call proportions (P -values for detection of signals above the background <0.01) for 31 probes (shown in Table S3) in all of the tissue samples examined were less than 90%. Since such a low proportion may be attributable to polymorphism at the probe CpG sites, these 31 probes were excluded from the present assay, leaving a final total of 26,455 autosomal CpG sites.

Infinium probes showing ordered differences from 36 C to 145 N, and then to the 145 T samples themselves, were examined by the cumulative logit model adjusted by sex, age and experimental batch ($P < 1 \times 10^{-14}$). Correlations between β -values in N and T samples and recurrence were assessed by the Cox

regression model adjusted by sex, age and experimental batch ($P < 0.001$). Benjamini-Hochberg correction was performed to adjust for multiple testing. Differences of β -values and mRNA expression levels between N and T samples were examined by Mann-Whitney U test. Correlations between mRNA expression levels and clinicopathological parameters were assayed by analysis of variance between groups (ANOVA) and Welch's T-test: after adjusted Bonferroni correction to adjust for multiple testing, corrected P values of <0.05 were considered to be significant. All statistical analyses were performed using programming language R.

Results

DNA Methylation Profiles Associated with Recurrence are Established at Precancerous Stages

DNA methylation levels of CpG sites of the *CNTNAP2*, *EVX1*, *GFRA1*, *PDE9A* and *TBX20* genes based on the Infinium assay were clearly verified using the quantitative pyrosequencing method (Figure S1), indicating the reliability of the Infinium assay. The cumulative logit model ($P < 1 \times 10^{-14}$) revealed ordered progression of DNA methylation alterations from C to N, and then to T samples, on 3,270 probes; DNA methylation alterations occurred even in N samples compared to C samples, and such DNA methylation alterations were inherited by, or strengthened in, T samples, indicating that Ns were at precancerous stages with DNA methylation alterations. Among the 3,270 probes, the number showing average β -values in T samples minus average β -values in C samples ($\Delta\beta_{T-C}$) of >0.1 and <-0.1 were 1,209 and 1,056, respectively. Thus, when we defined differentially methylated probes as probes showing a $\Delta\beta_{T-C}$ value of >0.1 or <-0.1 , the false positivity rate by the cumulative logit model was 4.3%.

Correlations between DNA methylation status and recurrence were examined using the Cox regression model ($P < 0.001$) for 145 patients. In T samples, DNA methylation status on 944 probes for the 916 genes was significantly correlated with recurrence: on 87 probes (red dots in Figure 1A), higher β -values were observed in recurrence-positive patients than in recurrence-negative patients, whereas lower β -values on 857 probes (blue dots in Figure 1A) were observed in recurrence-positive patients. Surprisingly, even in N samples, the DNA methylation status on 2,215 probes for the 2,083 genes was significantly correlated with recurrence: on 425 probes (red dots in Figure 1B), higher β -values were observed in recurrence-positive patients than in recurrence-negative patients, whereas lower β -values on 1,790 probes (blue dots in Figure 1B) were observed in recurrence-positive patients.

In order to identify recurrence-related genes that are normally unmethylated and for which DNA hypermethylation at precancerous stages is strengthened in the established LADCs, among the 425 probes (red dots in Figure 1B), we initially focused on 28 probes for which the average β -values in C samples (β_C) were less than 0.2 and the average β -values in T samples minus that in the corresponding N samples ($\Delta\beta_{T-N}$) were more than 0.1 (Table 1).




Figure 3. DNA methylation levels and mRNA expression levels after 5-aza-2'-deoxycytidine (5-aza-dC) treatment. DNA methylation levels (β -values) and mRNA expression levels for the *ADCY5* (A), *EVX1* (B), *GFRA1* (C), *PDE9A* (D) and *TBX20* (E) genes were determined by Infinium assay and quantitative real-time reverse transcription-PCR analysis, respectively. The error bars represent the standard deviation for triplicate quantitative real-time RT-PCR analyses. DNA methylation levels and mRNA expression levels on days 3 (AZA3) and 6 (AZA6) were compared with those of untreated cells (AZA0). After 5-aza-dC treatment, reduction of DNA methylation levels and restoration of the mRNA expression levels of *ADCY5* (A), *EVX1* (B) and *TBX20* (E) were observed in both of the cell lines used. In panel C, since reduction of the DNA methylation level was not induced by 5-aza-dC in PC9 cells, restoration of mRNA expression did not occur in these cells. Panel D shows reduction of the DNA methylation level and restoration of the mRNA expression level in PC9 cells. doi:10.1371/journal.pone.0059444.g003

Silencing of Recurrence-related Genes due to DNA Hypermethylation

Among the 28 genes listed in Table 1, we further focused on 6 genes (*ADCY5* [20,21], *CNTNAP2* [22,23], *EVX1* [24], *GFRA1* [25], *PDE9A* [26,27] and *TBX20* [28]) for which implications in transcription regulation, apoptosis or cell adhesion had been reported. Quantitative real-time RT-PCR analysis of these 6 genes was performed in 132 N and 151 T samples for which total RNA was available. mRNA expression levels for the *ADCY5*, *EVX1*, *GFRA1*, *PDE9A* and *TBX20* genes in T samples were significantly lower than those in N samples, although the reduced expression of the *CNTNAP2* gene did not reach statistical significance (Figure 2B). The DNA methylation statuses (β -values in N and T samples) of the 6 genes are also shown in Figure 2A; the data suggested that DNA hypermethylation of these genes might result in reduction of mRNA expression in tissue samples from the same cohort.

DNA methylation levels of the *ADCY5*, *EVX1*, *GFRA1*, *PDE9A* and *TBX20* genes in lung cancer cell lines A549, PC9, VMRC-LCD and EBC-1 are shown in Figure S2. To examine the effects of the DNA methylation inhibitor, the top two cell lines showing the highest DNA methylation levels (β -values) were selected for each gene. In fact, mRNA expression levels determined by quantitative real-time RT-PCR analysis of the genes (with the exception of *PDE9A*) were extremely low in the cell lines selected. 5-aza-dC treatment induced marked reduction of DNA methylation levels and restored the mRNA expression levels of *ADCY5*, *EVX1*, *GFRA1* and *TBX20* (Figure 3). With regard to *GFRA1*, since reduction of the DNA methylation level was not induced by 5-aza-dC, restoration of mRNA expression did not occur in PC9 cells. Taken together with Figures 2 and 3, the data suggested that the examined genes were silenced due to DNA hypermethylation in the lung cancers. With regard to *PDE9A*, for which the mRNA expression levels were high even in the top two cell lines showing the highest levels of DNA methylation, 5-aza-dC treatment did not further increase the mRNA expression level in the two cell lines. The PC9 cells were then additionally treated with 5-aza-dC, and restoration of *PDE9A* mRNA expression due to DNA demethylation was confirmed in the cells (Figure 3).

Clinicopathological Impact of Reduced Expression of mRNA for Recurrence-related Genes

Reduced expression of mRNA for *ADCY5*, *EVX1*, *GFRA1* and *PDE9A* in T samples was correlated with clinicopathological parameters reflecting tumor aggressiveness, such as a larger tumor diameter, higher histological grade, blood vessel invasion, pleural invasion and tumor anthracosis (Table 2), although their mRNA expression levels were not predictors of recurrence that were independent of known parameters such as pathological-TNM stage and lymph node metastasis (Table S4). With regard to the correlation with histological subtype, levels of mRNA expression for *GFRA1* and *PDE9A* were significantly higher in lepidic-type LADCs showing a less invasive growth pattern than in other histological subtypes.

Discussion

The 'field cancerization' phenomenon in the lung has become evident, being especially associated with cigarette smoking [31]. We and other groups have reported DNA methylation of specific genes or chromosomal loci in non-cancerous lung tissue obtained from lung cancer patients, or in lung tissue from cancer-free smokers [8–10]. These previous data drew our attention to DNA methylation alterations at precancerous stages of LADC. However, the impact of DNA methylation alterations at precancerous stages on the expression of specific gene and clinicopathological parameters of established cancers has remained unclear. Moreover, previous examinations focusing on precancerous stages in the lung have not involved a genome-wide approach. Although Selamat et al. and Lockwood et al. have reported Infinium assay results for 59 and 43 lung cancer samples, respectively [16,17], they did not focus on precancerous stages.

Here we have reported the results of the Infinium assay for 326 lung tissue samples including 145 N samples. Our cumulative logit model analysis revealed stepwise progression of DNA methylation alterations from C to N, and then T samples on 3,270 probes. Genome-wide analysis at single-CpG resolution confirmed that DNA methylation alterations actually occurred even at precancerous stages, although the possibility that such alterations observed in N samples had been influenced by differences in tissue composition between C and N samples cannot be completely excluded. Moreover, it was clearly shown that such DNA methylation alterations had clinicopathological impact, since many probes in N samples were significantly correlated with recurrence after establishment of LADCs (Figure 1B). DNA methylation profiles determining outcome are already established at precancerous stages. The finding that the number of probes showing DNA methylation alterations significantly associated with recurrence in N samples was larger than that in T samples may have been due to the fact that passenger DNA methylation alterations occurring during progression from the precancerous stages to established cancers may have masked any clinicopathologically significant DNA methylation profiles in T samples.

Next, we focused on 28 recurrence-related genes that are normally unmethylated and for which DNA hypermethylation in N samples was strengthened in T samples (Table 1). Among these 28 genes, we further focused on *ADCY5*, *CNTNAP2*, *EVX1*, *GFRA1*, *PDE9A* and *TBX20*, based on their previously reported implications in transcription regulation, apoptosis or cell adhesion. (a) The data in *adc5*-knockout mice indicated that *ADCY5* promotes apoptosis in cardiomyocytes [20,21]. Although large-scale screening studies of leukemias have identified *ADCY5* as one of the genes that are methylated in leukemic cells [32,33], the clinicopathological impact of DNA methylation of the *ADCY5* gene has not yet been clarified in human malignancies. (b) *CNTNAP2*, a glial adhesion molecule, binds extracellularly to contactin 2, an immunoglobulin superfamily neural recognition protein [22]. *CNTNAP2* is known to be a tumor-suppressor gene for gliomas, and is disrupted by chromosomal translocations and gene mutations [23]. Although a large-scale screening study of

Table 2. Correlation between mRNA expression levels of recurrence-related genes and clinicopathological factors.

Clinicopathological parameters	Number of tumors	ADCY5		EVX1		GFRA1		PDE9A		TBX20		
		Expression ^a	P ^b	Expression ^a	P ^b	Expression ^a	P ^b	Expression ^a	P ^b	Expression ^a	P ^b	
Tumor diameter												
<2.5 cm	43	-4.95±2.10	<u>1.63×10^{-1c}</u>	-2.80±1.82	<u>4.24×10^{-1c}</u>	-3.27±2.67	<u>1.60×10^{-3c}</u>	0.50±1.39	<u>3.40×10^{-2c}</u>	-5.73±2.68	<u>7.65×10^{-1c}</u>	
≥2.5 cm, <4 cm	61	-5.25±2.39		-2.36±2.16		-4.02±2.86		0.28±1.76		-5.76±2.78		
≥4 cm	47	-6.14±2.76		-3.04±2.02		-5.74±3.28		-0.50±1.59		-6.10±2.84		
Histological subtype ^d												
Lepidic	12	-4.19±1.61	<u>2.10×10^{-1c}</u>	-2.73±2.14	<u>1.79×10^{-1c}</u>	-1.64±2.35	<u>1.71×10^{-4c}</u>	0.93±1.07	<u>1.73×10^{-3c}</u>	-6.24±1.79	<u>1.28×10^{-1c}</u>	
Acinar	24	-6.03±2.01		-2.91±2.09		-5.41±2.72		-0.20±1.36		-5.77±2.57		
Papillary	69	-5.30±2.35		-2.29±1.99		-3.90±2.65		0.23±1.63		-5.28±2.80		
Micropapillary	15	-4.69±2.19		-2.35±2.10		-3.47±2.87		1.20±1.27		-6.89±1.79		
Solid	28	-6.34±3.24		-3.62±1.88		-6.27±3.62		-0.91±1.81		-6.37±3.11		
Invasive mucinous	3	-4.64±1.46		-3.19±1.24		-2.95±1.43		0.16±1.68		-8.21±0.14		
Histological grades												
G1	56	-4.60±1.80	<u>1.28×10^{-3c}</u>	-2.68±1.90	<u>4.57×10^{-2c}</u>	-2.62±1.93	<u>4.80×10⁻¹⁰</u>	0.55±1.41	<u>1.32×10⁻⁴</u>	-5.48±2.69	<u>9.50×10⁻²</u>	
G2	65	-5.58±2.33		-2.32±2.10		-4.59±2.71		0.25±1.57		-5.74±2.67		
G3	30	-6.73±3.20		-3.54±1.92		-7.01±3.60		-1.06±1.77		-6.81±2.94		
Lymphatic invasion												
Negative	52	-5.31±2.43	1.00 ^e	-2.76±1.71	1.00 ^e	-3.95±3.38	1.00 ^e	0.06±1.64	1.00 ^e	-5.86±2.42	<u>9.82×10^{-1e}</u>	
Positive	99	-5.52±2.50		-2.66±2.19		-4.55±2.92		0.12±1.67		-5.85±2.93		
Blood vessel invasion												
Negative	43	-4.89±2.40	<u>2.38×10^{-1e}</u>	-2.77±1.84	<u>7.68×10^{-1e}</u>	-3.13±3.45	<u>2.81×10^{-2e}</u>	0.63±1.68	<u>6.39×10^{-2e}</u>	-5.71±2.60	1.00 ^e	
Positive	108	-5.66±2.47		-2.67±2.11		-4.82±2.81		-0.11±1.60		-5.92±2.83		
Pleural invasion												
Negative	78	-5.16±2.42	<u>4.45×10^{-1c}</u>	-2.77±2.15	<u>4.60×10^{-1c}</u>	-3.65±2.83	<u>7.28×10^{-3c}</u>	0.28±1.61	<u>5.82×10^{-1c}</u>	-5.95±2.51	<u>5.38×10^{-1c}</u>	
Invasion to the visceral pleura beyond the elastic fiber	33	-5.88±2.68		-2.62±2.07		-4.99±3.54		0.03±1.93		-6.41±2.96		
Invasion to the surface of the visceral pleura	22	-4.96±2.07		-2.16±1.62		-4.01±2.66		0.18±1.29		-4.98±2.82		
Invasion to the parietal pleura	18	-6.47±2.49		-3.15±1.90		-6.56±2.71		-0.65±1.58		-5.52±3.22		
Tumor anthracosis ^f												
Negative	68	-4.97±2.57	<u>8.27×10^{-2e}</u>	-2.37±2.22	<u>2.04×10^{-1e}</u>	-3.32±2.80	<u>1.08×10^{-3e}</u>	0.70±1.46	<u>1.66×10^{-4e}</u>	-5.51±2.78	<u>1.99×10^{-1e}</u>	
Positive	82	-5.87±2.30		-2.91±1.79		-5.10±3.02		-0.39±1.66		-6.09±2.69		

^aAverage of log₂-transformed mRNA expression levels/GAPDH ± standard deviation. ^bAdjusted P-values using adjusted Bonferroni correction. ^cAnalysis of variance between groups. ^dPredominant histological subtypes of LADCs were diagnosed according to the classification devised by the International Association for the Study of Lung Cancer, the American Thoracic Society and the European Respiratory Society [29]. ^eWelch's T-test. ^fCoal dust is accumulated in active fibroblast proliferation foci, which is associated with poorer prognosis of lung adenocarcinoma patients and reflects an active cancer-stromal interaction [30]. P values of <0.05 are underlined. doi:10.1371/journal.pone.0059444.t002