

in those with multiple cancers. Although further prospective study may be necessary, it would appear that methylation of *IRF4* could be a molecular marker with which to predict the development or recurrence of gastric cancer.

Several lines of evidence have suggested that *IRF5* has tumor suppressor activity, and that in response to DNA damage *IRF5* is induced by p53 to promote cell cycle arrest and apoptosis.<sup>(20,21,42)</sup> Kulaeva *et al.*<sup>(43)</sup> showed that treating spontaneously immortal Li-Fraumeni fibroblasts with DAC induces a senescence-like state, and that *IRF5* is silenced by DNA methylation in the same cells, suggesting *IRF5* is involved in mediating cellular senescence.<sup>(22)</sup> Here we showed that DAC enhanced p53-induced *IRF5* expression, and that *IRF5* expression was also induced by p63 and p73, suggesting *IRF5* is a target of the p53 gene family. Although, on average, *IRF5* methylation was not significantly higher in primary cancers than in noncancerous tissues, several cases did show high levels of *IRF5* methylation.

We found that *IRF8* expression was down-regulated in gastric cancer cell lines; that DNA methylation was well correlated with gene silencing; and that treating cells with DAC restored *IRF8* expression. This is consistent with earlier reports showing that *IRF8* is silenced in colorectal cancer cell lines in a DNA methylation-dependent manner.<sup>(23)</sup> In contrast to the data obtained with cell lines, we did not find an increase in *IRF8* methylation in primary gastric cancers, as compared to noncancerous tissues. This is in contrast to earlier studies showing that *IRF8* is methylated in cancers of the colon, esophagus, and nasopharynx.<sup>(24,37)</sup> This discrepancy may reflect the different methods used to detect methylation: methylation-specific PCR was used in those earlier studies, whereas we used bisulfate-pyrosequencing. Alternatively, methylation of *IRF8* may be an

early event in tumorigenesis, which starts in subsets of gastric epithelial cells. Consistent with that idea, Lee *et al.* reported that *IRF8* is methylated only in some esophageal tissues from esophageal cancer patients. Further study will be necessary to clarify the significance of *IRF8* methylation in primary gastric cancers.

In conclusion, we have shown that *IRF4*, *IRF5*, and *IRF8* are epigenetically silenced in gastric cancer cells. Methylation of *IRF5* was associated with CIMP and EBV infection. Moreover, the high degree of *IRF4* methylation in gastric mucosae from cancer patients suggests that DNA methylation of *IRF4* could be a useful molecular marker for gastric cancer diagnosis and risk assessment.

#### Acknowledgments

The authors thank Dr William F. Goldman for editing the manuscript. This study was supported in part by Grants-in-Aid for Scientific Research on Priority Areas (M.T., K.I., and T.T.); the Program for Developing Supporting Systems for Upgrading Education and Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (Y.S., M.T.); Grants-in-Aid for Scientific Research (S) from the Japan Society for Promotion of Science (K.I.); a Grant-in-Aid for the Third-term Comprehensive 10-year Strategy for Cancer Control (F.I., M.T.); and Grant-in-Aid for Cancer Research from the Ministry of Health, Labor, and Welfare of Japan (M.T.).

#### Disclosure Statement

The authors have no conflict of interest.

#### References

- 1 Wright PA, Williams GT. Molecular biology and gastric carcinoma. *Gut* 1993; **34**: 145-7.
- 2 Jones PA, Baylin SB. The epigenomics of cancer. *Cell* 2007; **128**: 683-92.
- 3 Sato F, Meltzer SJ. CpG island hypermethylation in progression of esophageal and gastric cancer. *Cancer* 2006; **106**: 483-93.
- 4 Suzuki H, Tokino T, Shinomura Y, Imai K, Toyota M. DNA methylation and cancer pathways in gastrointestinal tumors. *Pharmacogenomics* 2008; **9**: 1917-28.
- 5 Ushijima T. Detection and interpretation of altered methylation patterns in cancer cells. *Nat Rev Cancer* 2005; **5**: 223-31.
- 6 Nojima M, Suzuki H, Toyota M *et al.* Frequent epigenetic inactivation of SFRP genes and constitutive activation of Wnt signaling in gastric cancer. *Oncogene* 2007; **26**: 4699-713.
- 7 Sato H, Suzuki H, Toyota M *et al.* Frequent epigenetic inactivation of DICKKOPF family genes in human gastrointestinal tumors. *Carcinogenesis* 2007; **28**: 2459-66.
- 8 Maruyama R, Akino K, Toyota M *et al.* Cytoplasmic RASSF2A is a proapoptotic mediator whose expression is epigenetically silenced in gastric cancer. *Carcinogenesis* 2008; **29**: 1312-8.
- 9 Suzuki H, Itoh F, Toyota M, Kikuchi T, Kakiuchi H, Imai K. Inactivation of the 14-3-3 sigma gene is associated with 5' CpG island hypermethylation in human cancers. *Cancer Res* 2000; **60**: 4353-7.
- 10 Akino K, Toyota M, Suzuki H *et al.* Identification of DFNA5 as a target of epigenetic inactivation in gastric cancer. *Cancer Sci* 2007; **98**: 88-95.
- 11 Issa JP. DNA methylation as a therapeutic target in cancer. *Clin Cancer Res* 2007; **13**: 1634-7.
- 12 Smits KM, Cleven AH, Weijenberg MP *et al.* Pharmacoeigenomics in colorectal cancer: a step forward in predicting prognosis and treatment response. *Pharmacogenomics* 2008; **9**: 1903-16.
- 13 Hellebrekers DM, Lentjes MH, van den Bosch SM *et al.* GATA4 and GATA5 are potential tumor suppressors and biomarkers in colorectal cancer. *Clin Cancer Res* 2009; **15**: 3990-7.
- 14 Muller HM, Oberwalder M, Fiegl H *et al.* Methylation changes in faecal DNA: a marker for colorectal cancer screening? *Lancet* 2004; **363**: 1283-5.
- 15 Maekita T, Nakazawa K, Mihara M *et al.* High levels of aberrant DNA methylation in *Helicobacter pylori*-infected gastric mucosae and its possible association with gastric cancer risk. *Clin Cancer Res* 2006; **12**: 989-95.
- 16 Nakajima T, Maekita T, Oda I *et al.* Higher methylation levels in gastric mucosae significantly correlate with higher risk of gastric cancers. *Cancer Epidemiol Biomarkers Prev* 2006; **15**: 2317-21.
- 17 Takaoka A, Tamura T, Taniguchi T. Interferon regulatory factor family of transcription factors and regulation of oncogenesis. *Cancer Sci* 2008; **99**: 467-78.
- 18 Miyamoto M, Fujita T, Kimura Y *et al.* Regulated expression of a gene encoding a nuclear factor, IRF-1, that specifically binds to IFN-beta gene regulatory elements. *Cell* 1988; **54**: 903-13.
- 19 Tanaka N, Ishihara M, Lamphier MS *et al.* Cooperation of the tumour suppressors IRF-1 and p53 in response to DNA damage. *Nature* 1996; **382**: 816-8.
- 20 Barnes BJ, Kellum MJ, Pinder KE, Frisancho JA, Pitha PM. Interferon regulatory factor 5, a novel mediator of cell cycle arrest and cell death. *Cancer Res* 2003; **63**: 6424-31.
- 21 Mori T, Anazawa Y, Iizumi M, Fukuda S, Nakamura Y, Arakawa H. Identification of the interferon regulatory factor 5 gene (IRF-5) as a direct target for p53. *Oncogene* 2002; **21**: 2914-8.
- 22 Li Q, Tang L, Roberts PC *et al.* Interferon regulatory factors IRF5 and IRF7 inhibit growth and induce senescence in immortal Li-Fraumeni fibroblasts. *Mol Cancer Res* 2008; **6**: 770-84.
- 23 Yang D, Thangaraju M, Greenelch K *et al.* Repression of IFN regulatory factor 8 by DNA methylation is a molecular determinant of apoptotic resistance and metastatic phenotype in metastatic tumor cells. *Cancer Res* 2007; **67**: 3301-9.
- 24 Lee KY, Geng H, Ng KM *et al.* Epigenetic disruption of interferon-gamma response through silencing the tumor suppressor interferon regulatory factor 8 in nasopharyngeal, esophageal and multiple other carcinomas. *Oncogene* 2008; **27**: 5267-76.
- 25 Ortmann CA, Burchert A, Holzle K *et al.* Down-regulation of interferon regulatory factor 4 gene expression in leukemic cells due to hypermethylation of CpG motifs in the promoter region. *Nucleic Acids Res* 2005; **33**: 6895-905.
- 26 Yanagihara K, Ito A, Toge T, Numoto M. Antiproliferative effects of isoflavones on human cancer cell lines established from the gastrointestinal tract. *Cancer Res* 1993; **53**: 5815-21.
- 27 Ishida S, Yamashita T, Nakaya U, Tokino T. Adenovirus-mediated transfer of p53-related genes induces apoptosis of human cancer cells. *Jpn J Cancer Res* 2000; **91**: 174-80.

- 28 Maruyama R, Aoki F, Toyota M *et al.* Comparative genome analysis identifies the vitamin D receptor gene as a direct target of p53-mediated transcriptional activation. *Cancer Res* 2006; **66**: 4574–83.
- 29 Watanabe Y, Toyota M, Kondo Y *et al.* PRDM5 identified as a target of epigenetic silencing in colorectal and gastric cancer. *Clin Cancer Res* 2007; **13**: 4786–94.
- 30 Sugai T, Habano W, Jiao YF *et al.* Analysis of allelic imbalances at multiple cancer-related chromosomal loci and microsatellite instability within the same tumor using a single tumor gland from colorectal carcinomas. *Int J Cancer* 2005; **114**: 337–45.
- 31 Kusano M, Toyota M, Suzuki H *et al.* Genetic, epigenetic, and clinicopathologic features of gastric carcinomas with the CpG island methylator phenotype and an association with Epstein-Barr virus. *Cancer* 2006; **106**: 1467–79.
- 32 Driggers PH, Ennist DL, Gleason SL *et al.* An interferon gamma-regulated protein that binds the interferon-inducible enhancer element of major histocompatibility complex class I genes. *Proc Natl Acad Sci U S A* 1990; **87**: 3743–7.
- 33 Shou J, Soriano R, Hayward SW, Cunha GR, Williams PM, Gao WQ. Expression profiling of a human cell line model of prostatic cancer reveals a direct involvement of interferon signaling in prostate tumor progression. *Proc Natl Acad Sci U S A* 2002; **99**: 2830–5.
- 34 Karpf AR, Peterson PW, Rawlins JT *et al.* Inhibition of DNA methyltransferase stimulates the expression of signal transducer and activator of transcription 1, 2, and 3 genes in colon tumor cells. *Proc Natl Acad Sci U S A* 1999; **96**: 14007–12.
- 35 Satoh A, Toyota M, Ikeda H *et al.* Epigenetic inactivation of class II transactivator (CIITA) is associated with the absence of interferon-gamma-induced HLA-DR expression in colorectal and gastric cancer cells. *Oncogene* 2004; **23**: 8876–86.
- 36 Reu FJ, Bae SI, Cherkassky L *et al.* Overcoming resistance to interferon-induced apoptosis of renal carcinoma and melanoma cells by DNA demethylation. *J Clin Oncol* 2006; **24**: 3771–9.
- 37 McGough JM, Yang D, Huang S *et al.* DNA methylation represses IFN-gamma-induced and signal transducer and activator of transcription 1-mediated IFN regulatory factor 8 activation in colon carcinoma cells. *Mol Cancer Res* 2008; **6**: 1841–51.
- 38 Tshuikina M, Jernberg-Wiklund H, Nilsson K, Oberg F. Epigenetic silencing of the interferon regulatory factor ICSBP/IRF8 in human multiple myeloma. *Exp Hematol* 2008; **36**: 1673–81.
- 39 Ahlquist DA. Next generation stool DNA testing: expanding the scope. *Gastroenterology* 2009; **136**: 2068–73.
- 40 Watanabe Y, Kim HS, Castoro RJ *et al.* Sensitive and specific detection of early gastric cancer using DNA methylation analysis of gastric washes. *Gastroenterology* 2009; **136**: 2149–58.
- 41 Takeshima H, Yamashita S, Shimazu T, Niwa T, Ushijima T. The presence of RNA polymerase II, active or stalled, predicts epigenetic fate of promoter CpG islands. *Genome Res* 2009; **19**: 1974–82.
- 42 Barnes BJ, Field AE, Pitha-Rowe PM. Virus-induced heterodimer formation between IRF-5 and IRF-7 modulates assembly of the IFNA enhanceosome in vivo and transcriptional activity of IFNA genes. *J Biol Chem* 2003; **278**: 16630–41.
- 43 Kulaeva OI, Draghici S, Tang L, Kraniak JM, Land SJ, Tainsky MA. Epigenetic silencing of multiple interferon pathway genes after cellular immortalization. *Oncogene* 2003; **22**: 4118–27.

## Supporting Information

Additional Supporting Information may be found in the online version of this article:

**Fig. S1.** Real-time PCR analysis of interferon regulatory factor (*IRF*)-4, *IRF5*, and *IRF8* expression in gastric cancer cell lines.

**Fig. S2.** Induction of interferon regulatory factor (*IRF*)-5 expression by p53 (a) and of *IRF8* expression by interferon (IFN)- $\gamma$  (b).

**Fig. S3.** 5-Aza-2'-deoxycytidine (DAC) enhances suppression of cell growth by interferon.

**Fig. S4.** Methylation analysis of interferon regulatory factor (*IRF*)-4, *IRF5*, and *IRF8* after treatment with 5-aza-2'-deoxycytidine (DAC) and/or interferon (IFN).

**Fig. S5.** Receiver–operator curve (ROC) for interferon regulatory factor (*IRF*)-4 methylation to discriminate patients with a single gastric cancer from patients with *Helicobacter pylori*-positive chronic gastritis.

**Fig. S6.** Receiver–operator curve (ROC) curve for interferon regulatory factor (*IRF*)-4 methylation to discriminate patients with a single or multiple gastric cancers from patients with *Helicobacter pylori*-positive chronic gastritis.

**Table S1.** Primers used for methylation-specific PCR (MSP) used in this study.

**Table S2.** Primer sequences used for bisulfite-pyrosequencing and bisulfite-sequencing.

**Table S3.** High levels of interferon regulatory factor (*IRF*)-4 methylation are associated with multiple gastric cancers.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

## Genomic screening for genes upregulated by demethylation revealed novel targets of epigenetic silencing in breast cancer

Tomoko Fujikane · Noriko Nishikawa · Minoru Toyota · Hiromu Suzuki · Masanori Nojima · Reo Maruyama · Masami Ashida · Mutsumi Ohe-Toyota · Masahiro Kai · Toshihiko Nishidate · Yasushi Sasaki · Tousei Ohmura · Koichi Hirata · Takashi Tokino

Received: 4 May 2009 / Accepted: 12 October 2009 / Published online: 27 October 2009  
© Springer Science+Business Media, LLC. 2009

**Abstract** Breast cancer arises through the accumulation of multiple genetic alterations and epigenetic changes such as methylation, which silences gene expression in a variety of cancers. In the present study, we applied genomic screening to identify genes upregulated by the demethylating agent 5-aza-2'-deoxycytidine (DAC) in a human breast cancer cell line (MCF7). We identified 288 genes upregulated and 29 genes downregulated more than five-fold after treatment with DAC, and gene ontology analyses revealed the genes to be involved in immune responses,

apoptosis, and cell differentiation. In addition, real-time PCR analysis of ten genes silenced in MCF7 cells confirmed that they are upregulated by DAC, while bisulfite-sequencing analysis confirmed that nine of those genes were silenced by methylation. We also found that treating MCF7 cells with DAC restored induction of DFNA5 by p53, as well as by two other p53 family genes, p63 $\gamma$  and p73 $\beta$ . Introduction of NTN4 into MCF7 cells suppressed cell growth, indicating that NTN4 has tumor suppressive activity. In primary breast cancers, we detected cancer-specific methylation of NTN4, PGP9.5, and DKK3, suggesting that methylation of these genes could be useful markers for diagnosis of breast cancer. Thus, DNA methylation appears to be a common event in breast cancer, and the genes silenced by methylation could be useful targets for both diagnosis and therapy.

Tomoko Fujikane, Noriko Nishikawa, Minoru Toyota, Hiromu Suzuki contributed equally to this work.

**Electronic supplementary material** The online version of this article (doi:10.1007/s10549-009-0600-1) contains supplementary material, which is available to authorized users.

T. Fujikane · M. Toyota · H. Suzuki · R. Maruyama · M. Ohe-Toyota · Y. Sasaki · T. Tokino  
Department of Molecular Biology, Cancer Research Institute, Sapporo Medical University, Sapporo 060-8556, Japan

T. Fujikane · N. Nishikawa · T. Nishidate · T. Ohmura · K. Hirata  
First Department of Surgery, Sapporo Medical University, Sapporo 060-8556, Japan

M. Toyota · H. Suzuki · R. Maruyama  
First Department of Internal Medicine, Sapporo Medical University, Sapporo 060-8556, Japan

M. Toyota (✉) · M. Ashida · M. Kai  
Department of Biochemistry, Sapporo Medical University, South 1, West 17, Chuo-ku, Sapporo 060-8556, Japan  
e-mail: mtoyota@sapmed.ac.jp

M. Nojima  
Department of Public Health, Sapporo Medical University, Sapporo 060-8556, Japan

**Keywords** DNA methylation · Epigenetics · Gene expression

### Introduction

Epigenetic changes such as DNA methylation and histone modification are now thought to play a significant role in tumorigenesis. Under normal physiological conditions, DNA methylation is involved in such processes as X-chromosome inactivation, genome imprinting, and suppression of repetitive sequences [1], but genome-wide hypomethylation and regional hypermethylation are also common events in tumors [2]. For example, breast cancer, which continues to be one of the most commonly occurring cancers among women, worldwide [3], is known to arise through the accumulation of multiple genetic and epigenetic DNA alterations. Given that more than 1,000 genes are silenced by



DNA methylation in other types of cancers [4], the targets of epigenetic inactivation in breast cancer have just begun to be identified. To date, analysis of candidate genes for DNA methylation in breast cancer has shown that the targets of epigenetic inactivation include cell cycle regulators such as p16 [5] and 14-3-3 sigma [6], cell adhesion molecules such as E-cadherin [7], cytokines such as HIN-1 [8], genes involved in cell signaling such as RASSF1 [9], proapoptotic genes such as TMS1 [10], genes involved in development such as HOXB13 [11], and transcription factors such as activator protein-2 $\alpha$  [12]. Genomic screening approaches using cDNA microarrays, and promoter microarrays identified several novel targets of DNA methylation [13–15]. This makes identification of novel genes epigenetically inactivated in breast cancer an important step toward a better understanding of the pathogenesis of the disease. In the present study, therefore, we applied genomic screening to identify genes silenced by DNA methylation in breast cancer and confirmed the results by quantitative methylation analysis. Our findings suggest that DNA methylation is a common event in breast cancer and that many of the genes silenced by DNA methylation could represent useful targets for both diagnosis and therapy.

## Materials and methods

### Cell lines and specimens

Five breast cancer cell lines (MCF7, MB435s, MB436, MB468, and SKBR-3) were obtained from the American Type Culture Collection (Manassas, VA) or the Japanese Collection of Research Bioresources (Tokyo, Japan). All cell lines were cultured in appropriate medium supplemented with 10% fetal bovine serum and incubated under a 5% CO<sub>2</sub> atmosphere at 37°C. In addition, 75 breast cancer specimens and 15 breast tissue samples from areas adjacent to tumors were obtained from Sapporo Medical University Hospital at surgery and stored at –80°C. In accordance with institutional guidelines, all patients gave informed consent prior to collection of the specimens. Genomic DNA was extracted using the phenol/chloroform method. Total RNA was extracted from cell lines using Trizol (Life Technologies, Inc.) according to the manufacturer's instructions.

### cDNA microarray analysis

Breast cancer cells (MCF7, MB435s, MB436, MB468, and SKBR-3) were treated with DAC for 72 h, total RNA was extracted and purified using Trizol (Invitrogen) and RNA-easy (Qiagen), after which the RNA samples were quantified using NanoDrop ND-100, the quality was assessed using an

Agilent Technologies 2100 Bioanalyzer. The RNA concentration in the samples was >100 ng/ $\mu$ l, and the RNA integrity score was 8–10, with 10 being the highest possible score. Sample amplification and labeling were performed using a Low RNA Input Fluorescent Linear Amplification kit (Agilent Technologies) according to the manufacturer's instructions. Samples (1.65  $\mu$ g) labeled with Cy3 were hybridized and processed on a 4  $\times$  44 K Whole Human Genome Oligo Microarray. Scanning was performed with an Agilent G2565BA microarray scanner using the settings recommended by Agilent Technologies. After all raw data were normalized, fold-change comparisons and gene set enrichment (BROAD Institute) and gene ontology analyses were performed using GeneSpring GX 10.0. The presence of CpG islands was examined using BLAT (<http://genome.brc.mcw.edu/cgi-bin/hgBlat>) with previously described criteria [16].

### Real-time PCR

Samples (5  $\mu$ g) of RNA were reverse-transcribed using Superscript III (Invitrogen) to prepare first strand cDNA. For semi-quantitative analysis, real-time PCR was carried out using a 7900 Sequence Detection System (Applied Biosystems). The reaction mixture contained 1  $\times$  TaqMan Universal PCR Master Mix, primers and probes for each gene and 1  $\mu$ l of cDNA. GAPDH served as an endogenous control. The Taqman probes used in this study are shown in Supplementary Table 1. Each experiment was done in triplicate.

### Methylation analysis

For bisulfite-pyrosequencing, genomic DNA was treated with sodium bisulfite as described previously [17, 18], after which pyrosequencing was performed to assess the methylation status [19]. Bisulfite-PCR primers were designed using PSQ Assay Design software (Biotage, Uppsala, Sweden), and the primers and PCR conditions used were specific for each target gene. After the PCR, the biotinylated strand was captured on streptavidin-coated beads (Amersham Bioscience), and pyrosequencing was performed using PSQ HS Gold SNP reagents and a PSQ HS 96 (Biotage, Uppsala, Sweden). For each gene, the average percentage methylation of the entire CpG island was calculated, and cases in which there was more than 10% methylation were deemed to be positive for methylation. To sequence the bisulfite PCR products, the amplified fragments were cloned into a vector using a TOPO TA cloning kit (Invitrogen), after which a cycle sequencing reaction was carried out using a BigDye terminator kit (Applied Biosystems), and the DNA was sequenced using an ABI 3100 automated sequencer (Applied Biosystems).



Primer sequences used for bisulfite-pyrosequencing and bisulfite-sequencing are shown in Supplementary Table 2.

#### Infection by adenovirus

The generation, purification, and infection procedures used with replication-deficient recombinant adenovirus containing the p53 (Ad-p53), TAp63 $\gamma$  (Ad-p63 $\gamma$ ), TAp73 $\beta$  (Ad-p73 $\beta$ ), or the bacterial lacZ gene (Ad-lacZ) were described previously [20]. The relative efficiency of adenoviral infection was determined by X-gal staining of cells infected with the Ad-lacZ (control). At an MOI of 100, 90–100% of the cells were infected (data not shown).

#### Western blot analysis

Mouse anti-FLAG mAb (M2; Sigma) was used for immunoblotting. Whole cell lysates were prepared by scraping cell monolayers into radioimmunoprecipitation assay buffer without SDS [containing 150 mmol/l NaCl, 50 mmol/l Tris-HCl (pH 7.2), 1% deoxycholic acid, 1% Triton X-100, 0.25 mmol/l EDTA (pH 8.0), protease and phosphatase inhibitors, 5  $\mu$ g/ml leupeptin, 5  $\mu$ g/ml aprotinin, 1  $\mu$ g/ml pepstatin A, 1 mmol/l phenylmethylsulfonyl fluoride, 5 mmol/l NaF, and 100  $\mu$ mol/l sodium orthovanadate], and protein concentrations were determined (Lowry reagent, Bio-Rad). Equal amounts of protein were separated by SDS-PAGE and transferred to Immobilon P membranes (Millipore).

#### Chromatin immunoprecipitation assay

Chromatin immunoprecipitation (ChIP) assays were performed as described previously using a ChIP assay kit (Upstate Biotechnologies). Briefly,  $2 \times 10^6$  cells were cross-linked using 1% formaldehyde solution for 15 min at 37°C. The cells were then lysed in 200  $\mu$ l of SDS lysis buffer and sonicated to generate 300- to 800-bp DNA fragments. Following centrifugation, the cleared supernatant was diluted tenfold with ChIP dilution buffer, after which 1/50 of the extract volume was used for PCR amplification as the input control. The remaining extract was incubated with a specific antibody for 16 h at 4°C. Immune complexes were precipitated, washed, and eluted as recommended. DNA-protein cross-links were reversed by heating for 4 h at 65°C, after which the DNA fragments were purified and dissolved in 50  $\mu$ l of Tris-EDTA. One microliter of each sample was then used as a template for PCR amplification. PCR for histone analysis was carried out as described previously [21] using the primers listed in Supplementary Table 2. PCR amplification of *DFNA5* and *MDM2* containing the putative p53RE was also carried out using primers listed in Supplementary Table 2.

#### Colony formation assays

Colony formation assays were carried out as described previously [22]. Briefly, MCF7 cells ( $1 \times 10^5$  cells) were transfected with 5  $\mu$ g of pReceiver-M11-NTN4 (EX-U1401-M11, GeneCopeia) or with empty vector using Lipofectamine 2000 according to manufacturer's instructions. Cells were then plated on 60-mm culture dishes and selected for 14 days in 0.6 mg/ml G418, after which the colonies that formed were stained with Giemsa and counted using National Institutes of Health IMAGE software.

#### Statistics

To compare methylation levels between tumors and normal tissues, *t* tests were performed for all samples, and paired *t* tests were performed for matched samples from the same patients. Receiver operating characteristic (ROC) curves were constructed based on the levels of NTN4, PGP9.5, and DKK3 methylation, and *P* values were calculated by comparing the ROC curves to a reference curve. ANOVAs with post hoc Games-Howell tests were performed to compare methylation levels at different cancer stages. A scatter plot was constructed by plotting levels of FKBP6 methylation against tumor size, and a Pearson's correlation coefficient was calculated for these values. Values of *P* < 0.05 were considered significant. All statistical calculations were performed using SPSSJ 15.0 (SPSS Japan Inc.).

## Results

#### Identification of genes upregulated by DAC

The global changes in gene expression induced by DAC in MCF7 breast cancer cells were examined using an Agilent Whole Genome microarray that covers 44,000 transcripts of human genes (Supplementary Fig. 1). As compared with mock-treated cells, 288 genes were upregulated and 29 genes were downregulated more than fivefold by DAC in these cancer cells (Supplementary Tables 3, 4). Four genes (*SFRP1*, *DKK3*, *DFNA5*, *TAC1*) were recently shown to be silenced by DNA methylation in breast cancer [15, 23–25]. To identify biological processes significantly affected by demethylation, we used gene ontology analysis to assess the function of the 288 upregulated genes. Detailed results are shown in Supplementary Table 5. Treating the cells with DAC led to significant upregulation of genes involved in immune responses, the extracellular region, and cytokine activity. We also conducted a gene set enrichment analysis using functional annotation tools (Supplementary Table 6). Among 26 selected gene sets, genes involved in cell differentiation, cell development, defense responses,



apoptosis, and signal transduction were enriched in DAC-treated cells, as compared to mock-treated cells.

#### Expression analysis of genes identified by microarray

Database analysis revealed that out of 288 genes upregulated by DAC, 155 contain CpG islands in the 5' end of the gene (Supplementary Table 3). We next selected ten genes known from earlier work to be cancer-related and to have CpG islands in their 5' ends (Fig. 1). The selected genes were DFNA5, SFRP1, DKK3, PGP9.5, and LOXL4, which were all previously shown to be silenced by DNA methylation in various types of tumors [26–30]; NTN4, which encodes a member of the netrin family involved in the negative regulation of angiogenesis [31]; TRIM50, which encodes an E3 ubiquitin ligase [32]; FKBP6, which encodes an immunophilin family protein [33]; PON1, which encodes an arylesterase and whose polymorphisms are known to be associated with prostate cancer [34]; and OSBPL3, which encodes an oxysterol-binding protein that plays a role in cell adhesion [35]. Real-time PCR analysis revealed that the expression levels of all these genes were low or negligible in MCF7 cells, whereas high levels of expression—i.e., an expression ratio against GAPDH >0.01—were detected for DFNA5, SFRP1, OSBPL3, NTN4, PGP9.5, and LOXL4 in normal breast tissue; cell lines other than MCF7 showed various levels of expression (Supplementary Fig. 2). For DKK3, FKBP6, PON1, and TRIM50, expression was low—i.e., an expression ratio against GAPDH < 0.01—in normal breast tissue, and cell lines showed various levels of expression (Supplementary Fig. 3). Treatment with DAC restored expression of these genes in cell lines in which expression was otherwise low or negligible (Supplementary Fig. 3).

#### Methylation analysis of ten genes in breast cancer cell lines

To confirm methylation-dependent gene silencing, we next used bisulfite-pyrosequencing to examine the methylation status of the ten genes. This enabled us to quantify the methylation of multiple CpG sites (Fig. 2). The primers and probes were designed to detect methylation in the region around the transcription start sites. Dense methylation of nine genes (SFRP1, DFNA5, DKK3, PGP9.5, OSBPL3, NTN4, TRIM50, FKBP6, and PON1) was detected in MCF7 cells, strongly suggesting that DNA methylation is the cause of gene silencing. Various levels of methylation were detected in four other cell lines and was also associated with gene silencing (Figs. 1, 2; Supplementary Figs. 2, 3). That methylation of LOXL4 was not detected means that LOXL4 is silenced by a mechanism other than DNA methylation.

**Fig. 1** Real-time PCR analysis of genes upregulated by DAC. The expression status of DFNA5, SFRP1, DKK3, FKBP6, LOXL4, OSBPL3, NTN4, PGP9.5, PON1, and TRIM50 was confirmed by real-time PCR. The cell lines and tissues examined are shown below the columns. Cell lines were treated for 72 h with either mock (–) or 1.0  $\mu$ M DAC. The integrity of the cDNA was assessed by comparing the CT values for the genes of interest with that of GAPDH. Columns means of three experiments, bars SE


We next performed bisulfite-sequencing analysis to obtain detailed methylation profiles of the CpG sites in the region around the transcription start site of DFNA5 gene. We examined 45 CpG sites and found that DFNA5 was densely methylated in MCF7 cells, which do not express DFNA5. By contrast, little or no methylation was detected in MDA-MB435s, MDA-MB436, MDA-MB-468, and SK-Br-3 cells, which do express DFNA5 (Fig. 3). Thus, the results obtained with bisulfite-sequencing are consistent with both the bisulfite-pyrosequencing data and the DFNA5 expression status.


#### Restoration of p53-dependent transcription of DFNA5 by demethylation

It was recently reported that DFNA5 is a target gene for p53 [36]. We therefore tested whether demethylation of DFNA5 in MCF7 cells would restore its transcriptional activation by p53 and/or by two other p53 family genes, TAp63 $\gamma$  and TAp73 $\beta$ . When cells were infected with Ad-p53, Ad-p63 $\gamma$ , or Ad-p73 $\beta$ , expression of FLAG-tagged p53 family proteins was detected (Fig. 4a). In addition, p21, a cyclin-dependent inhibitor, was induced by all three vectors (Fig. 4a). We then examined expression of DFNA5 in MCF7 cells with or without treatment with DAC. We found that treating MCF7 cells with DAC restored induction of DFNA5 by p53 family genes, especially by p63, suggesting that DFNA5 is a target of the p53 family, not specifically p53, itself (Fig. 4b). We then performed ChIP assays to determine whether p63 $\gamma$  directly interacts with the p53 response element of DFNA5 (RE-DFNA5) (Fig. 4c). PCR amplification of the ChIP products revealed that one DNA fragment containing RE-DFNA5 was present in the immunoprecipitated complex with p63 $\gamma$ . As a control, we confirmed that p63 $\gamma$  binds to the p53 response element of MDM2 in vivo. These results indicate that DFNA5 can be upregulated by p63 $\gamma$  through direct interaction with RE-DFNA5.

#### Tumor suppressive activity of NTN4

Netrins and their receptors have been shown to be involved in tumorigenesis [37]. To test whether NTN4 suppresses growth of breast cancer cells, we performed colony formation assays using MCF7 cells, which express negligible levels NTN4. We found that introduction of a plasmid





**Fig. 2** Representative results of bisulfite-pyrosequencing of DFNA5 (a) and SFRP1 (b). Bisulfite-pyrosequencing was carried out using DNA from breast cancer cell lines and primary breast cancer specimens. Examined were the regions upstream from the transcription start site (DFNA5: -85 to -97; SFRP1: -65 bp to -44 bp).

Gray columns depict regions of CpG sites, and the percentage methylation at each CpG site is shown on the top. Y-axis, signal peaks expressed as a proportion of the number of nucleotides incorporated. X-axis, the nucleotides incorporated. Cell lines and specimens are shown below the columns

containing NTN4 cDNA significantly suppressed colony growth, suggesting that NTN4 does indeed have tumor suppressive activity (Fig. 5a, b).


Comparison of methylation and clinicopathological features of patients with primary breast cancer

Of the nine aforementioned genes silenced by DNA methylation in primary breast cancer, seven showed significantly higher levels of methylation in cancerous tissues than in normal breast tissues ( $P < 0.001$  for NTN4, PGP9.5, DKK3, OSBPL3, SFRP1, DFNA5;  $P < 0.01$  for PON1, Fig. 6a; Supplementary Table 7). Methylation was

also examined in paired samples of cancerous and adjacent normal breast tissues from 15 patients. Methylation of NTN4 ( $P < 0.001$ ), PGP9.5 ( $P < 0.001$ ), DKK3 ( $P = 0.006$ ), and PON1 ( $P = 0.031$ ) was significantly higher in the tumor tissue than in the adjacent breast tissue (Fig. 6b; Supplementary Table 8). The clinical usefulness of DNA methylation in distinguishing breast cancer from noncancerous tissue was confirmed by analyzing ROC curves (Fig. 6c; Supplementary Table 9). Methylation of NTN4, DKK3, and PGP9.5 showed highly discriminative ROC curve profiles, which clearly distinguished breast cancer from normal breast tissue (NTN4:  $p < 0.001$ ; DKK3:  $P < 0.001$ ; PGP9.5:  $P < 0.001$ ). When we used 16%



**Fig. 3** Bisulfite-sequencing of DFNA5 in breast cancer cell lines. Each circle represents a CpG dinucleotide. Methylation status: *open circles* unmethylated, *filled circles* methylated. At least 9 clones were sequenced for each case. The CpG sites in the region spanning  $-265$  to  $+176$  from the transcription start site were analyzed, and are indicated by vertical bars (*top*)



methylation as a cut-off value, the sensitivity was 98.6% (95% CI: 92.6–100) and the specificity was 76.5% (95% CI: 50.1–93.2), indicating that methylation of NTN4 could be a useful molecular marker for detection of breast cancer (Supplementary Table 9). We then examined the relation between methylation status and clinicopathological factors (Supplementary Table 10) and found that methylation of FKBP6 is significantly correlated with advanced stages (Fig. 7a; Supplementary Table 11,  $P = 0.014$ ) and tumor size (Fig. 7b, Supplementary Table 12,  $P = 0.017$ ). There was no correlation between methylation and other factors including stages, histological types, number of metastasis positive lymph nodes, vascular invasion.

## Discussion


### Identification of genes silenced by DNA methylation in breast cancer

In the present study, we performed a microarray analysis to identify genes silenced by DNA methylation in breast cancer. We found that 288 genes were upregulated more than fivefold after treatment with DAC. Among those,

SFRP1, DFNA5, and DKK3 are known from earlier studies to be silenced by DNA methylation in breast cancer [23, 24, 38], while PGP9.5 is known to be methylated in a variety of cancer types [39, 40]. Our findings thus confirm that, with our microarray approach, we are able to successfully identify targets of DNA methylation in breast cancer. Gene ontology analysis revealed that genes involved in immune responses, the extracellular region and cytokine activity are significantly upregulated by DAC. Consistent to those findings, Karpf et al. [41] showed that genes regulated via interferon signaling are frequently upregulated by DAC, which suggests that upregulation of genes involved in immune responses, including those involved in antigen presentation, regulation of tumor necrosis factor and/or interferon pathways, may be a general feature of DAC treatment.


### Utility of DNA methylation for molecular diagnosis in breast cancer

Although previous studies have identified numerous targets of DNA methylation in breast cancer, the usefulness of the targeted genes for diagnosis remains unclear [13–15]. In fact, those studies confirmed DNA methylation of only a



**Fig. 4** Induction of DFNA5 by p53 family members. **a** Western blot analysis of p53, TAp63 $\gamma$ , and TAp73 $\beta$ . MCF7 cells were infected with adenoviral vector encoding the protein shown on the top and were harvested 24 h after infection. Immunoblot analysis was performed using anti-FLAG antibody. Expression of p21 was examined as a control. **b** ChIP assay. PCR was performed using ChIP products, and one DNA fragment containing the RE-DFNA5 was present in the immunoprecipitated complex with TAp63 $\gamma$ . As a control, fragments MDM2 DNA were amplified. **c** Restoration of p53-dependent DFNA5 expression by DAC. MCF7 cells were treated for 72 h with either mock or 0.2  $\mu$ M DAC followed by infection with 100 MOI of Ad-lacZ, Ad-p53, Ad-63 $\gamma$ , or Ad-p73 $\beta$  for 24 h. Expression of DFNA5 was examined by real-time PCR. Columns mean of three experiments, bars SE

limited number of samples [13, 15], or the methylation analysis was not quantitative [14]. Our findings suggest DNA methylation can be used as a biomarker to detect breast cancer. The cancer can be detected using DNA from biopsy specimens, serum or breast fluid—i.e., any tissue in which genes specifically methylated at a high frequency in cancer can be identified. In the present study, bisulfite-sequencing, a semi-quantitative methylation analysis, revealed that methylation of NTN4, PGP9.5, and DKK3 occurs in a cancer-specific manner. Previous studies have shown that PGP9.5 is silenced by DNA methylation in a variety of tumors [27, 39, 40], and cancer-specific methylation of PGP9.5 has been observed in both head/neck and hepatocellular cancers [39]. On the other hand, normal tissues in the prostate, esophagus, and stomach also show PGP9.5 methylation [39], so that whether or not methylation is cancer-specific is dependent on the cancer and tissue type. Recently, Veeck et al. [24] used methylation-specific PCR to assess the methylation of DKK3 and found that the gene is methylated in 61% of breast cancers. In the present



**Fig. 5** NTN4 suppresses breast cancer cell growth. **a** Colony formation assay. MCF7 cells were transfected with NTN4 or control vector and plated. After 2 weeks, the cells were fixed with methanol and stained with Giemsa. **b** Relative colony formation efficiencies of MCF7 cells transfected with NTN4 or control plasmid (vector). Columns mean of three experiments, bars SE

study, we similarly observed that methylation of DKK3 is significantly higher in breast cancers (17.2%) than in normal breast tissues (17.2 vs. 9.8%,  $P < 0.001$ ). Moreover, we showed for the first time that NTN4 is silenced by DNA methylation in cancer. When we used 16% methylation as a cut-off value, the sensitivity was 98.6% (95% CI: 92.6–100) and specificity was 76.5% (95% CI: 50.1–93.2), indicating that methylation of NTN4 could be a good molecular marker for detection of breast cancer.

In contrast to the genes mentioned above, methylation of FKBP6, PON1, and TRIM50 was detected even in normal breast tissues, and increases in promoter methylation reportedly correlate with age in colorectal and prostate tissues [42, 43]. In this regard, methylation of SFRP1, which has been shown to correlate with aging in colon [44], was not high in breast tissue, indicating that age-related methylation is also tissue-specific and that further studies

**Fig. 6** Methylation analysis in primary breast cancers. **a** Summary of methylation levels in normal and cancerous breast tissue: *N* normal tissue, *T* cancerous tissue. \*\*\*  $P < 0.001$ , \*\*  $P < 0.01$ . **b** Analysis of NTN4, PGP9.5, DKK3, and PON1 methylation in breast cancer and adjacent normal breast tissue. **c** ROC curve analysis of NTN4, PGP9.5, and DKK3 in primary breast cancer. The area under the ROC curve for each site conveys its utility for distinguishing normal breast from breast cancer in terms of its sensitivity and specificity