

- 14 Chen Y, Takita J, Choi YL *et al*. Oncogenic mutations of ALK kinase in neuroblastoma. *Nature* 2008; **455**: 971–4.
- 15 Friedman DL, Kadan-Lottick NS, Whitton J *et al*. Increased risk of cancer among siblings of long-term childhood cancer survivors: a report from the childhood cancer survivor study. *Cancer Epidemiol Biomarkers Prev* 2005; **14**: 1922–7.
- 16 Shojaei-Brosseau T, Chompret A, Abel A *et al*. Genetic epidemiology of neuroblastoma: a study of 426 cases at the Institut Gustave-Roussy in France. *Pediatr Blood Cancer* 2004; **42**: 99–105.
- 17 Maris JM, Kyemba SM, Rebbeck TR *et al*. Molecular genetic analysis of familial neuroblastoma. *Eur J Cancer* 1997; **33**: 1923–8.
- 18 Kushner BH, Gilbert F, Helson L. Familial neuroblastoma. Case reports, literature review, and etiologic considerations. *Cancer* 1986; **57**: 1887–93.
- 19 Knudson AG Jr, Strong LC. Mutation and cancer: neuroblastoma and pheochromocytoma. *Am J Hum Genet* 1972; **24**: 514–32.
- 20 Longo L, Panza E, Schemi F *et al*. Genetic predisposition to familial neuroblastoma: identification of two novel genomic regions at 2p and 12p. *Hum Hered* 2007; **63**: 205–11.
- 21 Maris JM, Weiss MJ, Mosse Y *et al*. Evidence for a hereditary neuroblastoma predisposition locus at chromosome 16p12-13. *Cancer Res* 2002; **62**: 6651–8.
- 22 Perri P, Longo L, Cusano R *et al*. Weak linkage at 4p16 to predisposition for human neuroblastoma. *Oncogene* 2002; **21**: 8356–60.
- 23 Amiel J, Laudier B, Attie-Bitach T *et al*. Polyalanine expansion and frameshift mutations of the paired-like homeobox gene PHOX2B in congenital central hyperventilation syndrome. *Nat Genet* 2003; **33**: 459–61.
- 24 Mosse YP, Laudenslager M, Khazi D *et al*. Germline PHOX2B mutation in hereditary neuroblastoma. *Am J Hum Genet* 2004; **75**: 727–30.
- 25 Trochet D, Bourdeaut F, Janoueix-Lerosey I *et al*. Germline mutations of the paired-like homeobox 2B (PHOX2B) gene in neuroblastoma. *Am J Hum Genet* 2004; **74**: 761–4.
- 26 Raabe EH, Laudenslager M, Winter C *et al*. Prevalence and functional consequence of PHOX2B mutations in neuroblastoma. *Oncogene* 2008; **27**: 469–76.
- 27 Caren H, Abel F, Kogner P, Martinsson T. High incidence of DNA mutations and gene amplifications of the ALK gene in advanced sporadic neuroblastoma tumours. *Biochem J* 2008; **416**: 153–9.
- 28 George RE, Attiyeh EF, Li S *et al*. Genome-wide analysis of neuroblastomas using high-density single nucleotide polymorphism arrays. *PLoS ONE* 2007; **2**: e255.
- 29 Nannya Y, Sanada M, Nakazaki K *et al*. A robust algorithm for copy number detection using high-density oligonucleotide single nucleotide polymorphism genotyping arrays. *Cancer Res* 2005; **65**: 6071–9.
- 30 Yamamoto G, Nannya Y, Kato M *et al*. Highly sensitive method for genomewide detection of allelic composition in nonpaired, primary tumor specimens by use of affymetrix single-nucleotide-polymorphism genotyping microarrays. *Am J Hum Genet* 2007; **81**: 114–26.
- 31 Osajima-Hakomori Y, Miyake I, Ohira M, Nakagawa A, Nakagawa A, Sakai R. Biological role of anaplastic lymphoma kinase in neuroblastoma. *Am J Pathol* 2005; **167**: 213–22.
- 32 Shiota M, Fujimoto J, Takenaga M *et al*. Diagnosis of t(2;5)(p23;q35)-associated Ki-1 lymphoma with immunohistochemistry. *Blood* 1994; **84**: 3648–52.
- 33 Morris SW, Kirstein MN, Valentine MB *et al*. Fusion of a kinase gene, ALK, to a nucleolar protein gene, NPM, in non-Hodgkin's lymphoma. *Science* 1994; **263**: 1281–4.
- 34 Lee HH, Norris A, Weiss JB, Frasch M. Jelly belly protein activates the receptor tyrosine kinase Alk to specify visceral muscle pioneers. *Nature* 2003; **425**: 507–12.
- 35 Stoica GE, Kuo A, Aigner A *et al*. Identification of anaplastic lymphoma kinase as a receptor for the growth factor pleiotrophin. *J Biol Chem* 2001; **276**: 16772–9.
- 36 Stoica GE, Kuo A, Powers C *et al*. Midkine binds to anaplastic lymphoma kinase (ALK) and acts as a growth factor for different cell types. *J Biol Chem* 2002; **277**: 35990–8.
- 37 Duyster J, Bai RY, Morris SW. Translocations involving anaplastic lymphoma kinase (ALK). *Oncogene* 2001; **20**: 5623–37.
- 38 Iwahara T, Fujimoto J, Wen D *et al*. Molecular characterization of ALK, a receptor tyrosine kinase expressed specifically in the nervous system. *Oncogene* 1997; **14**: 439–49.
- 39 Morris SW, Naeve C, Mathew P *et al*. ALK, the chromosome 2 gene locus altered by the t(2;5) in non-Hodgkin's lymphoma, encodes a novel neural receptor tyrosine kinase that is highly related to leukocyte tyrosine kinase (LTK). *Oncogene* 1997; **14**: 2175–88.
- 40 Pulford K, Lamant L, Morris SW *et al*. Detection of anaplastic lymphoma kinase (ALK) and nucleolar protein nucleophosmin (NPM)-ALK proteins in normal and neoplastic cells with the monoclonal antibody ALK1. *Blood* 1997; **89**: 1394–404.
- 41 Shiota M, Fujimoto J, Semba T, Satoh H, Yamamoto T, Mori S. Hyperphosphorylation of a novel 80 kDa protein-tyrosine kinase similar to Ltk in a human Ki-1 lymphoma cell line, AMS3. *Oncogene* 1994; **9**: 1567–74.
- 42 Lamant L, Pulford K, Bischof D *et al*. Expression of the ALK tyrosine kinase gene in neuroblastoma. *Am J Pathol* 2000; **156**: 1711–21.
- 43 Cools J, Wlodarska I, Somers R *et al*. Identification of novel fusion partners of ALK, the anaplastic lymphoma kinase, in anaplastic large-cell lymphoma and inflammatory myofibroblastic tumor. *Genes Chromosom Cancer* 2002; **34**: 354–62.
- 44 Colleoni GW, Bridge JA, Garicochea B, Liu J, Filippa DA, Ladanyi M. ATIC-ALK: a novel variant ALK gene fusion in anaplastic large cell lymphoma resulting from the recurrent cryptic chromosomal inversion, inv(2)(p23q35). *Am J Pathol* 2000; **156**: 781–9.
- 45 Ma Z, Cools J, Marynen P *et al*. Inv(2)(p23q35) in anaplastic large-cell lymphoma induces constitutive anaplastic lymphoma kinase (ALK) tyrosine kinase activation by fusion to ATIC, an enzyme involved in purine nucleotide biosynthesis. *Blood* 2000; **95**: 2144–9.
- 46 Trinei M, Lanfrancone L, Campo E *et al*. A new variant anaplastic lymphoma kinase (ALK)-fusion protein (ATIC-ALK) in a case of ALK-positive anaplastic large cell lymphoma. *Cancer Res* 2000; **60**: 793–8.
- 47 Touriol C, Greenland C, Lamant L *et al*. Further demonstration of the diversity of chromosomal changes involving 2p23 in ALK-positive lymphoma: 2 cases expressing ALK kinase fused to CLTCL (clathrin chain polypeptide-like). *Blood* 2000; **95**: 3204–7.
- 48 Griffin CA, Hawkins AL, Dvorak C, Henkle C, Ellingham T, Periman EJ. Recurrent involvement of 2p23 in inflammatory myofibroblastic tumors. *Cancer Res* 1999; **59**: 2776–80.
- 49 Bridge JA, Kanamori M, Ma Z *et al*. Fusion of the ALK gene to the clathrin heavy chain gene, CLTC, in inflammatory myofibroblastic tumor. *Am J Pathol* 2001; **159**: 411–5.
- 50 Ma Z, Hill DA, Collins MH *et al*. Fusion of ALK to the Ran-binding protein 2 (RANBP2) gene in inflammatory myofibroblastic tumor. *Genes Chromosom Cancer* 2003; **37**: 98–105.
- 51 Panagopoulos I, Nilsson T, Domanski HA *et al*. Fusion of the SEC31L1 and ALK genes in an inflammatory myofibroblastic tumor. *Int J Cancer* 2006; **118**: 1181–6.
- 52 Debelenko LV, Arthur DC, Pack SD, Helman LJ, Schrupp DS, Tsokos M. Identification of CARS-ALK fusion in primary and metastatic lesions of an inflammatory myofibroblastic tumor. *Lab Invest* 2003; **83**: 1255–65.
- 53 Jazii FR, Najafi Z, Malekzadeh R *et al*. Identification of squamous cell carcinoma associated proteins by proteomics and loss of beta tropomyosin expression in esophageal cancer. *World J Gastroenterol* 2006; **12**: 7104–12.
- 54 Takeuchi K, Choi YL, Togashi Y *et al*. KIF5B-ALK, a novel fusion oncokine identified by an immunohistochemistry-based diagnostic system for ALK-positive lung cancer. *Clin Cancer Res* 2009; **15**: 3143–9.
- 55 Soda M, Choi YL, Enomoto M *et al*. Identification of the transforming EML4-ALK fusion gene in non-small-cell lung cancer. *Nature* 2007; **448**: 561–6.
- 56 Fujimoto J, Shiota M, Iwahara T *et al*. Characterization of the transforming activity of p80, a hyperphosphorylated protein in a Ki-1 lymphoma cell line with chromosomal translocation t(2;5). *Proc Natl Acad Sci U S A* 1996; **93**: 4181–6.
- 57 Bischof D, Pulford K, Mason DY, Morris SW. Role of the nucleophosmin (NPM) portion of the non-Hodgkin's lymphoma-associated NPM-anaplastic lymphoma kinase fusion protein in oncogenesis. *Mol Cell Biol* 1997; **17**: 2312–25.
- 58 Kuefer MU, Look AT, Pulford K *et al*. Retrovirus-mediated gene transfer of NPM-ALK causes lymphoid malignancy in mice. *Blood* 1997; **90**: 2901–10.
- 59 Chiarle R, Gong JZ, Guasparri I *et al*. NPM-ALK transgenic mice spontaneously develop T-cell lymphomas and plasma cell tumors. *Blood* 2003; **101**: 1919–27.
- 60 Turner SD, Tooze R, MacLennan K, Alexander DR. Vav-promoter regulated oncogenic fusion protein NPM-ALK in transgenic mice causes B-cell lymphomas with hyperactive Jun kinase. *Oncogene* 2003; **22**: 7750–61.
- 61 Soda M, Takada S, Takeuchi K *et al*. A mouse model for EML4-ALK-positive lung cancer. *Proc Natl Acad Sci U S A* 2008; **105**: 19893–7.
- 62 Bai RY, Dieter P, Peschel C, Morris SW, Duyster J. Nucleophosmin-anaplastic lymphoma kinase of large-cell anaplastic lymphoma is a constitutively active tyrosine kinase that utilizes phospholipase C-gamma to mediate its mitogenicity. *Mol Cell Biol* 1998; **18**: 6951–61.
- 63 Bai RY, Ouyang T, Miething C, Morris SW, Peschel C, Duyster J. Nucleophosmin-anaplastic lymphoma kinase associated with anaplastic large-cell lymphoma activates the phosphatidylinositol 3-kinase/Akt antiapoptotic signaling pathway. *Blood* 2000; **96**: 4319–27.
- 64 Nieborowska-Skorska M, Slupianek A, Xue L *et al*. Role of signal transducer and activator of transcription 5 in nucleophosmin/anaplastic lymphoma kinase-mediated malignant transformation of lymphoid cells. *Cancer Res* 2001; **61**: 6517–23.

- 65 Slupianek A, Nieborowska-Skorska M, Hoser G *et al*. Role of phosphatidylinositol 3-kinase-Akt pathway in nucleophosmin/anaplastic lymphoma kinase-mediated lymphomagenesis. *Cancer Res* 2001; **61**: 2194–9.
- 66 Amin HM, McDonnell TJ, Ma Y *et al*. Selective inhibition of STAT3 induces apoptosis and G(1) cell cycle arrest in ALK-positive anaplastic large cell lymphoma. *Oncogene* 2004; **23**: 5426–34.
- 67 Koivunen JP, Mermel C, Zejnullahu K *et al*. EML4–ALK fusion gene and efficacy of an ALK kinase inhibitor in lung cancer. *Clin Cancer Res* 2008; **14**: 4275–83.
- 68 Hubbard SR, Mohammadi M, Schlessinger J. Autoregulatory mechanisms in protein-tyrosine kinases. *J Biol Chem* 1998; **273**: 11987–90.
- 69 Lemmon MA, Schlessinger J. Cell signaling by receptor tyrosine kinases. *Cell* 2010; **141**: 1117.
- 70 Galkin AV, Melnick JS, Kim S *et al*. Identification of NVP–TAE684, a potent, selective, and efficacious inhibitor of NPM–ALK. *Proc Natl Acad Sci U S A* 2007; **104**: 270–5.
- 71 McDermott U, Iafrate AJ, Gray NS *et al*. Genomic alterations of anaplastic lymphoma kinase may sensitize tumors to anaplastic lymphoma kinase inhibitors. *Cancer Res* 2008; **68**: 3389–95.
- 72 Christensen JG, Zou HY, Arango ME *et al*. Cytoreductive antitumor activity of PF-2341066, a novel inhibitor of anaplastic lymphoma kinase and c-Met, in experimental models of anaplastic large-cell lymphoma. *Mol Cancer Ther* 2007; **6**: 3314–22.
- 73 Zou HY, Li Q, Lee JH *et al*. An orally available small-molecule inhibitor of c-Met, PF-2341066, exhibits cytoreductive antitumor efficacy through antiproliferative and antiangiogenic mechanisms. *Cancer Res* 2007; **67**: 4408–17.
- 74 Bang Y, Kwak EL, Shaw AT *et al*. Clinical activity of the oral ALK inhibitor PF-02341066 in ALK-positive patients with non-small cell lung cancer (NSCLC). *J Clin Oncol* 2010; **28**(Suppl): abstr3.

Brief report

CBL mutation in chronic myelomonocytic leukemia secondary to familial platelet disorder with propensity to develop acute myeloid leukemia (FPD/AML)

Norio Shiba,^{1,2} Daisuke Hasegawa,³ Myoung-ja Park,¹ Chisato Murata,¹ Aiko Sato-Otsubo,⁴ Chitose Ogawa,³ Atsushi Manabe,³ Hirokazu Arakawa,² Seishi Ogawa,⁴ and Yasuhide Hayashi¹

¹Department of Hematology/Oncology, Gunma Children's Medical Center, Shibukawa, Japan; ²Department of Pediatrics, Gunma University Graduate School of Medicine, Maebashi, Japan; ³Department of Pediatrics, St Luke's International Hospital, Tokyo, Japan; and ⁴Cancer Genomics Project, Graduate School of Medicine, University of Tokyo, Tokyo, Japan

Familial platelet disorder with a propensity to develop acute myeloid leukemia (FPD/AML) is a rare autosomal dominant disease characterized by thrombocytopenia, abnormal platelet function, and a propensity to develop myelodysplastic syndrome (MDS) and AML. So far, > 20 affected families have been reported. Recently, a second *RUNX1* alteration has been reported; however, no

additional molecular abnormalities have been found so far. We identified an acquired *CBL* mutation and 11q-acquired uniparental disomy (11q-aUPD) in a patient with chronic myelomonocytic leukemia (CMML) secondary to FPD with *RUNX1* mutation but not in the same patient during refractory cytopenia. This finding suggests that alterations of the *CBL* gene and *RUNX1* gene may cooper-

ate in the pathogenesis of CMML in patients with FPD/AML. The presence of *CBL* mutations and 11q-aUPD was an important "second hit" that could be an indicator of leukemic transformation of MDS or AML in patients with FPD/AML. (*Blood*. 2012; 119(11):2612-2614)

Introduction

Familial platelet disorder with a propensity to develop acute myeloid leukemia (FPD/AML) is a rare autosomal dominant disease characterized by thrombocytopenia, abnormal platelet function, and a propensity to develop myelodysplastic syndrome (MDS) and AML.^{1,2} Since Song et al reported haploinsufficiency of the *RUNX1/CBFA2* gene,³ more than 20 affected families have been reported.⁴⁻⁸ Notably, various types of mono-allelic mutations of the *RUNX1* gene have been found in patients with AML secondary to FPD.^{3,7-9} *RUNX1*, which is a key regulator of definitive hematopoiesis and myeloid differentiation, is also commonly involved in sporadic cases of MDS and AML, by translocations in AML¹⁰ and by point mutations in AML^{11,12} and MDS.¹³ Recently, a second *RUNX1* alteration has been reported⁸; however, no additional molecular abnormalities have been found so far.

In this regard, recent reports of somatic mutations of the *CBL* proto-oncogene in myeloid neoplasms are intriguing because these *CBL* mutations have been shown to result in aberrant tyrosine kinase signaling, which would also lead to the activation of RAS signaling pathways. So far, we and others have reported that *CBL* mutations occurred in a variety of myeloid neoplasms, including de novo AML,^{14,15} MDS,^{16,17} and myeloproliferative neoplasm,^{16,17} especially in chronic myelomonocytic leukemia (CMML)^{16,17} and juvenile myelomonocytic leukemia.¹⁸ The importance of *CBL* mutations for leukemogenesis has substantially increased, which prompted us to search for possible *CBL* mutations in this pedigree.

Here, we reported that *CBL* mutation developed at the time of diagnosis of CMML, but not during refractory cytopenia, in a Japanese patient with FPD/AML harboring a *RUNX1* mutation.

Methods

RUNX1 mutation analysis

DNA and RNA were extracted from peripheral blood (PB) of the proband, her sister, and their mother after obtaining informed consent. We performed mutation analysis of the *RUNX1* gene by PCR followed by direct sequencing with the use of an ABI PRISM 310 Genetic Analyzer (Applied Biosystems). For further confirmation of deletion mutations, the PCR products were subcloned with the use of a TOPO TA Cloning Kit (Invitrogen) and then sequenced. Mutations were screened from exons 1-8 of the *RUNX1* gene.

CBL mutation analysis

Because *CBL* mutations thus far reported almost exclusively involved exons 8-9 that encode Linker/RING finger domains, we confined our mutation analysis to these exons, which were subjected to direct sequencing. Because the frequency of 11q-acquired uniparental disomy (11q-aUPD) was reported as ~85%-90% in *CBL* mutations, we also analyzed the sample with Affymetrix GeneChip 250K *Nspl*.¹⁷⁻¹⁹ Genome-wide detection of copy number abnormalities or allelic imbalances was performed with CNAG/AsCNAR Version 3.0 software (<http://www.genome.umin.jp>), which enabled sensitive detection of copy number neutral loss of heterozygosity (or aUPD).¹⁹ In addition, we examined mutations of the following genes in the proband as previously reported: *FLT3*, *KIT*, *RAS*, *JAK2*, *PTPN11*, *ASXL1*, *IDH1/2*, and *MPL*.²⁰⁻²² The study adhered to the principles of the Helsinki Declaration and was conducted under the regulations enacted by the Ethics Board of Gunma Children's Medical Center.

Results and discussion

The proband (III-2), who was the second child of nonconsanguineous parents, underwent an 8-year follow-up of mild to moderate


Submitted February 2, 2011; accepted November 24, 2011. Prepublished online as *Blood* First Edition paper, December 2, 2011; DOI 10.1182/blood-2011-02-333435.

The publication costs of this article were defrayed in part by page charge

payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2012 by The American Society of Hematology

Figure 1. The family pedigree. Squares indicate males and circles indicate females. Open symbols represent unaffected persons, half-filled symbols represent persons affected by thrombocytopenia, and closed symbols represent persons affected by FPD who developed MDS/AML. The proband (III-2) is indicated by an arrow.



thrombocytopenia ($50\text{--}80 \times 10^3/\mu\text{L}$), and at that age of 10 years, her condition was diagnosed as refractory cytopenia. Cytogenetic analysis found a normal karyotype, and FISH showed neither monosomy 7 nor trisomy 8. The proband had been closely observed without any therapy for 2 years and 9 months because she did not require transfusion and her disease remained stable; however, at the age of 12 years, leukocytosis and monocytosis developed and she became dependent on platelet transfusions. Finally, the disease evolved to CMML, and allogeneic bone marrow (BM) transplantation from an unrelated donor was performed. During the entire course, the number of blast cells in PB was constantly $< 2\%$, and no additional symptoms were observed, such as hepatosplenomegaly. Her elder sister (III-1) was also followed for 10 years with mild thrombocytopenia; however, the morphologic findings of PB or BM were not compatible with myeloproliferative neoplasms.¹⁷ Because her platelet count has been gradually decreasing, allogeneic BM transplantation is being considered. Although her father (II-4) developed MDS at the age of 41 and died 2 years later, her paternal aunt (II-2) developed MDS at the age of 49 and has remained in complete remission for 11 years after successful allogeneic cord blood transplantation. Her paternal grandfather (I-1) and uncle (II-1) also had a history of thrombocytopenia (Figure 1). Direct sequencing analysis of *RUNX1* found a one-base deletion of adenine at position 2364 within exon 7, resulting in a frameshift mutation that corresponded to AML1b transcript in the proband and her sister (Figure 2A). This resulted in a frameshift after amino acid change G262GfsX21. This mutation was not detected in their mother. All these data suggested that her paternal grandfather (I-1), uncle (II-1), aunt (II-2), and her father (II-4) were considered to have FPD/AML, carrying the same *RUNX1* mutation.

Although no *CBL* mutations were found in the proband sample of refractory cytopenia before development of CMML, homozygous mutation of the *CBL*, which was located in the splice acceptor site of intron 8 (Figure 2B), was identified in the proband sample in the CMML. We also found 11q-aUPD (Figure 2C) in the proband sample, confirming a strong association of *CBL* mutations with 11q-aUPD, as previously described¹⁶⁻¹⁸; however, no mutations of any other genes, including *FLT3*, *KIT*, *RAS*, *JAK2*, *PTPN11*, *ASXL1*, *IDH1/2*, and *MPL*, were found and no additional somatic *RUNX1* alterations. No *CBL* mutations were found in her sister's sample at this time.

Inherited *RUNX1* mutations were clustered in the N-terminal region in exons 3-5, which affect the runt homology domain. Mutations in the C-terminal region, detected in the present

pedigree, have been reported less frequently so far and are considered to affect the transactivation domain (Figure 2D).




Figure 2. Mutation analysis of *RUNX1* and *CBL* genes in the pedigree. (A) Direct sequencing analysis of affected patients (III-1, III-2) and an unaffected family member (II-5) is shown. Arrow indicates a one-base deletion of adenine. (B) Mutated *CBL* is shown in the proband. (C) Identification of acquired uniparental disomy of 11q in the proband. Total copy number (tCN; red plot) is shown above the cytoband, and the results of allele-specific copy number analysis with anonymous references (ASCNAR) plots are shown below the cytoband. Larger allele is presented by a red line, and the smaller allele is presented by a blue line. Allele-specific analysis showed 11q-aUPD (blue line), which contained the *CBL* region (arrow). (D) Schematic representation of wild-type and mutated *RUNX1*. The affected *RUNX1* is truncated at the C terminus of the transactivation domain (TAD). Part of TAD is lacking in this proband (red line).

It has been postulated that disruption of the *RUNX1* gene is not sufficient to cause AML, as previously reported with monoallelic and biallelic inactivation of *Runx1* in mice^{23,24} and in mice carrying the knocked-in *Runx1-Eto* chimeric gene. These data indicate that a second-hit mutation in addition to the dysfunction of *RUNX1* is required for the development of AML. Minelli et al postulated that the mutations seen in FPD cases have a mutation effect that induces additional genetic abnormalities and promotes progression to hematologic malignancies.²⁵

Marked associations between chromosome translocation and gene mutations have been reported: *KIT* mutation in core binding leukemia, t(8;21)/*AML1-ETO* and inv(16)(p13q22)/*CBFβ-MYH11*, *FLT3-ITD* in leukemia with t(15;17)/*PML-RARα*, or with t(6;9)/*DEK-CAN*. We consider that it is important to find an association to administer clinically relevant treatment. In addition to the germline *RUNX1* mutation, we identified an acquired *CBL* mutation in the proband and assumed it to be a second hit mutation by which FPD evolved into CMML. To our knowledge, this is the first patient with FPD/AML in whom *CBL* mutation has developed. This finding suggests that alterations of the *CBL* gene and *RUNX1* could cooperate in the pathogenesis of CMML or AML in patients with FPD/AML. The presence of 11q-aUPD provided evidence that loss of the wild-type copy of *CBL* with duplication of the mutant copy was an important second hit that could be an indicator of leukemic transformation in patients with FPD/AML.

References

- Downton SB, Beardsley D, Jamison D, Blattner S, Li FP. Studies of a familial platelet disorder. *Blood*. 1985;65(3):557-563.
- Ho CY, Otterud B, Legare RD, et al. Linkage of a familial platelet disorder with a propensity to develop myeloid malignancies to human chromosome 21q22.1-22.2. *Blood*. 1996;87(12):5218-5224.
- Song WJ, Sullivan MG, Legare RD, et al. Haploinsufficiency of *CBFA2* causes familial thrombocytopenia with propensity to develop acute myelogenous leukaemia. *Nat Genet*. 1999;23(2):166-175.
- Buijs A, Poddighe P, van Wijk R, et al. A novel *CBFA2* single-nucleotide mutation in familial platelet disorder with propensity to develop myeloid malignancies. *Blood*. 2001;98(9):2856-2858.
- Michaud J, Wu F, Osato M, et al. In vitro analyses of known and novel *RUNX1/AML1* mutations in dominant familial platelet disorder with predisposition to acute myelogenous leukemia: implications for mechanisms of pathogenesis. *Blood*. 2002;99(4):1364-1372.
- Owen CJ, Toze CL, Koosch A, et al. Five new pedigrees with inherited *RUNX1* mutations causing familial platelet disorder with propensity to myeloid malignancy. *Blood*. 2008;112(12):4639-4645.
- Kirito K, Sakoe K, Shinoda D, Takiyama Y, Kaushansky K, Komatsu N. A novel *RUNX1* mutation in familial platelet disorder with propensity to develop myeloid malignancies. *Haematologica*. 2008;93(1):155-156.
- Preudhomme C, Renneville A, Bourdon V, et al. High frequency of *RUNX1* biallelic alteration in acute myeloid leukemia secondary to familial platelet disorder. *Blood*. 2009;113(22):5583-5587.
- Heller PG, Glembofsky AC, Gandhi MJ, et al. Low *Mpl* receptor expression in a pedigree with familial platelet disorder with predisposition to acute myelogenous leukemia and a novel *AML1* mutation. *Blood*. 2005;105(12):4664-4670.
- Miyoshi H, Shimizu K, Kozu T, Maseki N, Kaneko Y, Ohki M. t(8;21) breakpoints on chromosome 21 in acute myeloid leukemia are clustered within a limited region of a single gene, *AML1*. *Proc Natl Acad Sci U S A*. 1991;88(23):10431-10434.
- Osato M, Asou N, Abdalla E, et al. Biallelic and heterozygous point mutations in the runt domain of the *AML1/PEBP2alphaB* gene associated with myeloblastic leukemias. *Blood*. 1999;93(6):1817-1824.
- Taketani T, Taki T, Takita J, et al. *AML1/RUNX1* mutations are infrequent, but related to *AML-M0*, acquired trisomy 21, and leukemic transformation in pediatric hematologic malignancies. *Genes Chromosomes Cancer*. 2003;38(1):1-7.
- Harada H, Harada Y, Niimi H, et al. High incidence of somatic mutations in the *AML1/RUNX1* gene in myelodysplastic syndrome and low blast percentage myeloid leukemia with myelodysplasia. *Blood*. 2004;103(6):2316-2324.
- Sargin B, Choudhary C, Crosetto N, et al. *FLT3*-dependent transformation by inactivating *c-Cbl* mutations in AML. *Blood*. 2007;110(3):1004-1012.
- Caligiuri MA, Briesewitz R, Yu J, et al. Novel *c-CBL* and *CBL-b* ubiquitin ligase mutations in human acute myeloid leukemia. *Blood*. 2007;110(3):1022-1024.
- Dunbar AJ, Gondek LP, O'Keefe CL, et al. 250K single nucleotide polymorphism array karyotyping identifies acquired uniparental disomy and homozygous mutations, including novel missense substitutions of *c-Cbl*, in myeloid malignancies. *Cancer Res*. 2008;68(24):10349-10357.
- Sanada M, Suzuki T, Shih LY, et al. Gain-of-function of mutated *C-CBL* tumour suppressor in myeloid neoplasms. *Nature*. 2009;460(7257):904-908.
- Shiba N, Kato M, Park MJ, et al. *CBL* mutations in juvenile myelomonocytic leukemia and pediatric myelodysplastic syndrome. *Leukemia*. 2010;24(5):1090-1092.
- Nannya Y, Sanada M, Nakazaki K, et al. A robust algorithm for copy number detection using high-density oligonucleotide single nucleotide polymorphism genotyping arrays. *Cancer Res*. 2005;65(14):6071-6079.
- Pardanani AD, Levine RL, Lasho T, et al. *MPL*S15 mutations in myeloproliferative and other myeloid disorders: a study of 1182 patients. *Blood*. 2006;108(10):3472-3476.
- Rocquain J, Carbuca N, Trouplin V, et al. Combined mutations of *ASXL1*, *CBL*, *FLT3*, *IDH1*, *IDH2*, *JAK2*, *KRAS*, *NPM1*, *NRAS*, *RUNX1*, *TET2* and *WT1* genes in myelodysplastic syndromes and acute myeloid leukemias. *BMC Cancer*. 2010;10:401.
- Tartaglia M, Niemeyer CM, Fragale A, et al. Somatic mutations in *PTPN11* in juvenile myelomonocytic leukemia, myelodysplastic syndromes and acute myeloid leukemia. *Nat Genet*. 2003;34(2):148-150.
- Sun W, Downing JR. Haploinsufficiency of *AML1* results in a decrease in the number of LTR-HSCs while simultaneously inducing an increase in more mature progenitors. *Blood*. 2004;104(12):3565-3572.
- Ichikawa M, Asai T, Saito T, et al. *AML-1* is required for megakaryocytic maturation and lymphocytic differentiation, but not for maintenance of hematopoietic stem cells in adult hematopoiesis. *Nat Med*. 2004;10(3):299-304.
- Minelli A, Maserati E, Rossi G, et al. Familial platelet disorder with propensity to acute myelogenous leukemia: genetic heterogeneity and progression to leukemia via acquisition of clonal chromosome anomalies. *Genes Chromosomes Cancer*. 2004;40(3):165-171.

Acknowledgments

The authors thank Miss Sayaka Takeuchi for her excellent technical assistance.

This work was supported by a grant for cancer research and a grant for research on children and families from the Ministry of Health, Labor, and Welfare of Japan; a Grant-in-Aid for Scientific Research (B, C) and Exploratory Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan; and by a research grant for Gunma Prefectural Hospitals.

Authorship

Contribution: Y.H. and C.O. designed the study; A.M., C.O., and D.H. provided critical reagents and samples; N.S., M.P., A.S.-O., and C.M. performed the experiments; H.A. and S.O. supervised the work; N.S. and M.P. analyzed the results; N.S. and D.H. constructed the figures; N.S. and Y.H. wrote the paper; and all the authors critically reviewed and revised the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Yasuhide Hayashi, Department of Hematology/Oncology, Gunma Children's Medical Center, 779, Shimohakoda, Hokeno, Shibukawa, Gunma, 377-8577, Japan; e-mail: hayashiy-tyk@umin.ac.jp.

CBL mutations in infant acute lymphoblastic leukaemia

Infant acute lymphoblastic leukaemia (ALL) is relatively rare, occurring in approximately 2.5–5% of cases of childhood ALL (Biondi *et al*, 2000). Infant ALLs are much more likely to present with high leucocyte counts, hepatosplenomegaly and overt central nervous system (CNS) diseases (Taki *et al*, 1996). T cell phenotype is much less common in infants, while myeloid antigen co-expression and the absence of CD10 expression are more frequent in infants than in older children with ALL. When molecular techniques [such as fluorescence *in situ* hybridization (FISH) or Southern blot analysis] are used in addition to karyotype, *MLL* gene rearrangements (*MLL*-R) are found in 70–80% of infant ALL compared with only 2–4% of older children with ALL (Taki *et al*, 1996; Biondi *et al*, 2000). Thus, infant ALL appears to be biologically distinct from the disease in older children (more than 1 year old).

In this regard, recent reports of somatic mutations of the *CBL* proto-oncogene in myeloid neoplasms are intriguing, because these *CBL* mutations were shown to result in aberrant tyrosine kinase signalling, which also leads to activation of RAS signalling pathways. So far, we and others have reported that *CBL* mutations occur in a variety of myeloid neoplasms, including *de novo* acute myeloid leukaemia (AML) (Caligiuri *et al*, 2007), myelodysplastic syndrome (MDS), and myeloproliferative neoplasm, especially in chronic myelomonocytic leukaemia (CMML) (Sanada *et al*, 2009), and juvenile myelomonocytic leukaemia (JMML) (Shiba *et al*, 2010). The importance of *CBL* mutations regarding leukaemogenesis is substantially increased. Recently, we found *CBL* mutation in therapy-related AML with *MLL*-R (Shiba *et al*, 2011). Interestingly, the *MLL*-*CBL* fusion gene has been reported in a *de novo* AML case (Fu *et al*, 2003), and this prompted us to search for possible *CBL* mutations in infant ALL with *MLL*-R.

Because *CBL* mutations thus far reported were almost all clustered within exons 8–9 that encode Linker/RING finger domains (Caligiuri *et al*, 2007; Sanada *et al*, 2009; Shiba *et al*, 2010), we confined our mutation analysis to these exons, in which polymerase chain reaction-amplified exons 8–9 were subjected to direct sequencing using an ABI PRISM 310 Genetic Analyser (Applied Biosystems, Branchburg, NJ, USA). The study adhered to the principles of the Helsinki Declaration, and was conducted under the regulations enacted by the Ethics Board of Gunma Children's Medical Centre.

CBL gene analysis was performed in 41 infant ALL patients in which *MLL*-R was found in 33 patients (80.5%), including 15 patients with t(4;11)(q21;q23), 4 with t(9;11)(p22;q23) and 5 with t(11;19)(q23;p13.3). Median age at diagnosis was 4.7


months (range, 0–12 months). We also performed *CBL* gene mutation analysis in 28 B cell precursor (BCP)-ALL patients (age range, 1–14 years).

Heterozygous mutations of the *CBL* gene were identified in 2 (4.9%) of 41 infant ALL patients, but not in older children with BCP-ALL. These were located in exon 8 (Fig 1). One patient was a 3-month-old female with t(4;11)(q21;q23) and the other patient was a 6-month-old male with t(11;19)(q23;p13.3). They were registered and treated on two Japanese infant leukaemia protocols, MLL96 and MLL98 respectively (Isoyama *et al*, 2002; Kosaka *et al*, 2004). Although strong association between *CBL* mutations and 11q-acquired uniparental disomy (aUPD) has been reported (Sanada *et al*, 2009), we did not perform the single nucleotide polymorphism array analysis due to lack of DNA.

MLL-R are more frequent in younger infants; up to 90% of infant ALL less than 6 months old at diagnosis have detectable *MLL*-R compared with 30–50% of infant ALL aged 6–12 months (Taki *et al*, 1996). *MLL*-R ALL has a characteristic gene expression profile that significantly differs from that of non-*MLL*-R BCP-ALL and of AML, confirming that *MLL*-R ALL is a biologically unique leukaemia subtype.

Patient 7. 1120 A>G (M374V)

ACTGTGAGATGGGCTCC



Patient 21. 1169 A>G (D390G)

GATAAGGATGTAAGA




Fig 1. Identification of *CBL* mutations. Heterozygous mutations of the *CBL* gene were identified in Patients 7 and 21.

Thus, the distinctive presenting features and clinical behaviour of infant ALL appear to be primarily due to the high frequency of *MLL-R* in this age group. However, outcome data comparing infant and non-infant patients with *MLL-R* suggest that there may be other factors which impact the prognosis of infant ALL. Both of the patients with *CBL* mutations were diagnosed before 6 months of age. In our previous report, all of three cases with *CBL* mutation developed JMML before 4 months of age (Shiba *et al*, 2010). These data suggested that *CBL* mutation may have a strong association with very early onset disease. *CBL* mutations have been reported as germline mutations in JMML (Niemeyer *et al*, 2010). Unfortunately, we could not investigate whether the mutations in our cases were germline mutations or not, because somatic cells were not available.

CBL mutations have been found in approximately 5% of 2000 samples from patients with myeloid neoplasms, including AML transformed from MDS. Gene aberrations in addition to *MLL-R* have rarely been reported in infant ALL. No reports of ALL with *CBL* mutations have so far been reported, suggesting that the pathogenesis of infant ALL is different from paediatric or adult ALL. To our knowledge, this is the first report of infant ALL patients with 11q23 translocation/*MLL-R* and *CBL* mutations. The present study suggests that alterations of *CBL* gene and *MLL-R* may cooperatively play a pathogenic role in the development of infant ALL with *MLL-R*.

Acknowledgements

We thank Mrs. Chisato Murata for her excellent technical assistance. This work was supported by a grant for Cancer Research, and a grant for Research on Children and Families from the Ministry of Health, Labour, and Welfare of Japan, a Grant-in-Aid for Scientific Research (B, C) and Exploratory Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan and by a Research grant for Gunma Prefectural Hospitals.

References

- Biondi, A., Cimino, G., Pieters, R. & Pui, C.H. (2000) Biological and therapeutic aspects of infant leukemia. *Blood*, **96**, 24–33.
- Caligiuri, M.A., Briesewitz, R., Yu, J., Wang, L., Wei, M., Amoczky, K.J., Marburger, T.B., Wen, J., Perrotti, D., Bloomfield, C.D. & Whitman, S.P. (2007) Novel c-*CBL* and *CBL-b* ubiquitin ligase mutations in human acute myeloid leukemia. *Blood*, **110**, 1022–1024.
- Fu, J.F., Hsu, J.J., Tang, T.C. & Shih, L.Y. (2003) Identification of *CBL*, a protooncogene at 11q23.3, as a novel *MLL* fusion partner in a patient with *de novo* acute myeloid leukemia. *Genes Chromosomes Cancer*, **37**, 214–219.
- Isoyama, K., Eguchi, M., Hibi, S., Kinukawa, N., Ohkawa, H., Kawasaki, H., Kosaka, Y., Oda, T.,

- Oda, M., Okamura, T., Nishimura, S., Hayashi, Y., Mori, T., Imaizumi, M., Mizutani, S., Tsukimoto, I., Kamada, N. & Ishii, E. (2002) Risk-directed treatment of infant acute lymphoblastic leukaemia based on early assessment of *MLL* gene status: results of the Japan Infant Leukaemia Study (MLL96). *British Journal of Haematology*, **118**, 999–1010.
- Kosaka, Y., Koh, K., Kinukawa, N., Wakazono, Y., Isoyama, K., Oda, T., Hayashi, Y., Ohta, S., Moritake, H., Oda, M., Nagatoshi, Y., Kigasawa, H., Ishida, Y., Ohara, A., Hanada, R., Sako, M., Sato, T., Mizutani, S., Horibe, K. & Ishii, E. (2004) Infant acute lymphoblastic leukemia with *MLL* gene rearrangements: outcome following intensive chemotherapy and hematopoietic stem cell transplantation. *Blood*, **104**, 3527–3534.

- Niemeyer, C.M., Kang, M.W., Shin, D.H., Furlan, I., Erlacher, M., Bunin, N.J., Bunda, S., Finklestein, J.Z., Sakamoto, K.M., Gorr, T.A., Mehta, P., Schmid, I., Kropshofer, G., Corbacioglu, S., Lang, P.J., Klein, C., Schlegel, P.G., Heinzmann, A., Schneider, M., Starý, J., van, den, Heuvel-Eibrink, M.M., Hasle, H., Locatelli, F., Sakai, D., Archambeault, S., Chen, L., Russell, R.C., Sybingco, S.S., Ohh, M., Braun, B.S., Flotho, C. & Loh, M.L. (2010) Germline *CBL* mutations cause developmental abnormalities and predispose to juvenile myelomonocytic leukemia. *Nature Genetics*, **42**, 794–800.
- Sanada, M., Suzuki, T., Shih, L.Y., Otsu, M., Kato, M., Yamazaki, S., Tamura, A., Honda, H., Sakata-Yanagimoto, M., Kumano, K., Oda, H., Yamagata, T., Takita, J., Gotoh, N., Nakazaki, K., Kawamata, N., Onodera, M., Nobuyoshi, M.,

Author's contributions

TT and YH designed the study. JT, MH, TK, MS and EI provided critical reagents and samples. NS and MP performed the experiments. EI, HA and SO supervised the work. NS and MP analysed the results. NS, TT, and YH wrote the paper and all the authors critically reviewed and revised it.

Conflict of interest

The authors declare no conflict of interest.

Norio Shiba^{1,2}
 Myoung-Ja Park¹
 Tomohiko Taki³
 Junko Takita⁴
 Mitsuteru Hiwatari⁴
 Takashi Kanazawa²
 Manabu Sotomatsu¹
 Eiichi Ishii⁵
 Hirokazu Arakawa²
 Seishi Ogawa⁶
 Yasuhide Hayashi¹

¹Department of Haematology/Oncology, Gunma Children's Medical Centre, ²Department of Paediatrics, Gunma University Graduate School of Medicine, Gunma, ³Department of Molecular Diagnostics and Therapeutics, Kyoto Prefectural University of Medicine Graduate School of Medical Science, Kyoto, ⁴Department of Paediatrics, Graduate School of Medicine, University of Tokyo, Tokyo, ⁵Department of Paediatrics, Ehime University Graduate School of Medicine, Ehime, and ⁶Cancer Genomics Project, Graduate School of Medicine, University of Tokyo, Tokyo, Japan.

E-mail: hayashiy-ty@umin.ac.jp

Keywords: infant, ALL, *MLL*, *CBL*.

First published online 11 October 2011

doi: 10.1111/j.1365-2141.2011.08900.x

Hayashi, Y., Harada, H., Kurokawa, M., Chiba, S., Mori, H., Ozawa, K., Omine, M., Hirai, H., Nakauchi, H., Koeffler, H.P. & Ogawa, S. (2009) Gain-of-function of mutated C-CBL tumour suppressor in myeloid neoplasms. *Nature*, **460**, 904–908.

Shiba, N., Kato, M., Park, M.J., Sanada, M., Ito, E., Fukushima, K., Sako, M., Arakawa, H., Ogawa,

S. & Hayashi, Y. (2010) CBL mutations in juvenile myelomonocytic leukemia and pediatric myelodysplastic syndrome. *Leukemia*, **24**, 1090–1092.

Shiba, N., Taki, T., Park, M.J., Nagasawa, M., Kanazawa, T., Takita, J., Ohnishi, H., Sotomatsu, M., Arakawa, H. & Hayashi, Y. (2011) CBL

mutation in childhood therapy-related leukemia. *Leukemia*, **25**, 1356–1358.

Taki, T., Ida, K., Bessho, F., Hanada, R., Kikuchi, A., Yamamoto, K., Sako, M., Tsuchida, M., Seto, M., Ueda, R. & Hayashi, Y. (1996) Frequency and clinical significance of the MLL gene rearrangements in infant acute leukemia. *Leukemia*, **10**, 1303–1307.

Association of an increased frequency of CD14⁺HLA-DR^{lo/neg} monocytes with decreased time to progression in chronic lymphocytic leukaemia (CLL)

Clinically important immune dysregulation is an early feature of leukaemia/small lymphocytic lymphoma (CLL) that often precedes other clinical manifestations of this disease. Defects in the adaptive immune system in patients with CLL results in quantitative and qualitative abnormalities in antibody production, and profound changes in T and Natural Killer (NK) cell numbers, ratios, and function (Palmer *et al*, 2008; Gonzalez-Rodriguez *et al*, 2010). Monocytes and macrophages, critical for adaptive and innate immune responses, also have an important role in the function of the CLL cell microenvironment, and this relationship continues to be explored

(Caligaris-Cappio, 2011). However, little is known about the effects of CLL on monocyte/macrophage physiology and whether alterations in monocytes have any clinical role in CLL. We were especially interested in the possible role of immunosuppressive CD14⁺ monocytes with reduced HLA-DR expression in CLL as we have observed this phenomena in glioblastoma (Gustafson *et al*, 2010), non-Hodgkin lymphoma (Lin *et al*, 2011) and prostate cancer (Vuk-Pavlovic *et al*, 2010).

To identify potential monocyte alterations by CLL in patients, we performed flow cytometric analysis of peripheral




Fig 1. CD14⁺HLA-DR^{lo/neg} Monocytes in CLL. Blood from 29 CLL patients and 15 healthy volunteers were used for immune phenotyping. There was no age difference (median of 59 vs. 58 years, respectively; $P = 0.2896$). Patients were eligible for the clinical trial (ClinicalTrials.gov NCT00562328) if diagnosed with previously untreated high-risk CLL using standard criteria and did not meet guidelines for conventional treatment. Blood was collected before initiation of treatment and 6, 9, and 12 months after completion of treatment in patients who had a sustained response. The percentage of CD14⁺ cells with a loss of HLA-DR staining was determined and compared between CLL patients and healthy volunteers (HV; upper left). CLL patients with a sustained response to treatment had a decrease in the frequency of CD14⁺HLA-DR^{lo/neg} monocytes 12 months after completion of treatment compared to measurement prior to treatment (upper right). Kaplan-Meier survival curve comparing CLL patients with elevated ratios (>2.5 standard deviations) of CD14⁺HLA-DR^{lo/neg} monocytes when compared to healthy volunteers (dashed line) or with ratios similar to those seen in healthy volunteers (solid line; bottom panel). HR, Hazard Ratio; 95% CI, 95% confidence interval.

LETTERS TO THE EDITOR

Novel splicing-factor mutations in juvenile myelomonocytic leukemia

Leukemia (2012) 26, 1879–1881; doi:10.1038/leu.2012.45

Myelodysplastic syndromes (MDS) and myelodysplastic/myeloproliferative neoplasms (MDS/MPN) are heterogeneous groups of chronic myeloid neoplasms characterized by clonal hematopoiesis, varying degrees of cytopenia or myeloproliferative features with evidence of myelodysplasia and a propensity to acute myeloid leukemia (AML).¹ In recent years, a number of novel gene mutations, involving *TET2*, *ASXL1*, *DNMT3A*, *EZH2*, *IDH1/2*, and *c-CBL*, have been identified in adult cases of chronic myeloid neoplasms, which have contributed to our understanding of disease pathogenesis.^{2–7} However, these mutations are rare in pediatric cases, with the exception of germline or somatic *c-CBL* mutations found in 10–15% of chronic myelomonocytic leukemia (CMML) and juvenile myelomonocytic leukemia (JMML),⁸ highlighting the distinct pathogenesis of adult and pediatric neoplasms.⁹

Recently, we reported high frequencies of mutations, involving the RNA splicing machinery, that are largely specific to myeloid neoplasms, showing evidence of myeloid dysplasia in adult.¹⁰ Affecting a total of eight components of the RNA splicing machinery (*U2AF35*, *U2AF65*, *SF3A1*, *SF3B1*, *SRSF2*, *ZRSR2*, *SF1* and *PRPF40B*) commonly involved in the 3' splice-site (3'SS) recognition, these pathway mutations are now implicated in the pathogenesis of myelodysplasia.¹⁰ To investigate the role of the splicing-pathway mutations in the pathogenesis of pediatric myeloid malignancies, we have examined 165 pediatric cases with AML, MDS, chronic myeloid leukemia (CML) and JMML for

mutations in the four major splicing factors, *U2AF35*, *ZRSR2*, *SRSF2*, and *SF3B1*, commonly mutated in adult cases.

Bone marrow or peripheral blood tumor specimens were obtained from 165 pediatric patients with various myeloid malignancies, including *de novo* AML ($n=93$), MDS ($n=28$), CML ($n=17$) and JMML ($n=27$), and the genomic DNA (gDNA) was subjected to mutation analysis (Supplementary Table 1). The status of the RAS pathway mutations for the current JMML series has been reported previously (Supplementary Table 2).^{11,12} Nineteen leukemia cell lines derived from AML (YNH-1, ML-1, KASUMI-3, KG-1, HL60, inv-3, SN-1, NB4 and HEL), acute monocytic leukemia (THP-1, SCC-3, J-111, CTS, P31/FUJ, MOLM-13, IMS/MI and KOCL-48) and acute megakaryoblastic leukemia (CMS and CMY) were also analyzed for mutations. Peripheral blood gDNA from 60 healthy adult volunteers was used as controls. Informed consent was obtained from the patients and/or their parents and from the healthy volunteers. We previously showed that for *U2AF35*, *SRSF2* and *SF3B1*, most of the mutations in adult cases were observed in exons 2 and 7, exon 1, and exons 14 and 15, respectively.¹⁰ Therefore, we confirmed mutation screening to these 'hot-spot' exons. In contrast, all the coding exons were examined for *ZRSR2*, because no mutational hot spots have been detected. Briefly, the relevant exons were amplified using PCR and mutations were examined by Sanger sequencing, as previously described.¹⁰ The Fisher's exact test was used to evaluate the statistical significance of frequencies of mutations for *U2AF35*, *SF3B1*, *ZRSR2* or *SRSF2* in adult cases and pediatric cases. This study was approved by the Ethics Committee of the University of Tokyo (Approval number 948-7).




Figure 1. Novel *U2AF35* and *SRSF2* mutations detected in JMML cases. (a) Left panel: sequence chromatogram of a heterozygous mutation at R156 in N-terminal zinc-finger motifs of *U2AF35* detected in a JMML case (JMML 4) is shown. Mutated nucleotides are indicated by arrows. Right panel: illustration of functional domains and mutations of *U2AF35*. Red arrow heads indicate hot-spot mutations at S34 and Q157 detected in the adult cases.¹⁰ Blue arrow head indicates the missense mutation at R156. (b) Left panel: sequence chromatogram of a 6-bp in-frame deletion (c.518-523delAAGTCC) in *SRSF2* detected in JMML 17 is shown. Mutated nucleotides are indicated by arrows. Right panel: illustration of functional domains and mutations of *SRSF2*. Red arrow head indicates hot-spot mutation at P95 frequently detected in the adult cases.¹⁰ Blue arrow head indicates a 6-bp in-frame deletion leading to deletion of S170 and K171.

No mutations were identified in the 28 cases with pediatric MDS, which included 13 cases with refractory anemia with excess blasts, 5 with refractory cytopenia of childhood, 2 with Down syndrome-related MDS, 2 with Fanconi anemia-related MDS, 2 with secondary MDS and 4 with unclassified MDS. Similarly, no mutations were detected in 93 cases with *de novo* AML or in 17 with CML, as well as 19 leukemia-derived cell lines. Our previous study in adult patients showed the frequency of mutations in *U2AF35*, *SF3B1*, *ZRSR2* or *SRSF2* to be 60/155 cases with MDS without increased ring sideroblasts and 8/151 *de novo* AML patients, emphasizing the rarity of these mutations in pediatric MDS ($P < 5.0 \times 10^{-6}$) and AML ($P < 0.02$) compared with adult cases. We found mutations in two JMML cases, JMML 4 and JMML 17. JMML 4 carried a heterozygous *U2AF35* mutation (R156M), whereas JMML 17 had a 6-bp in-frame deletion (c.518-523delAAGTCC) in *SRSF2* that resulted in deletion of amino acids S170 and K171 (Figure 1). Both nucleotide changes found in *U2AF35* and *SRSF2* were neither identified in the 60 healthy volunteers nor registered in the dbSNP database (<http://www.ncbi.nlm.nih.gov/projects/SNP/>) or in the 1000 genomes project, indicating that they represent novel spliceosome mutations in pediatric cases.

U2AF35 is the small subunit of the U2 auxiliary factor (*U2AF*), which binds an AG dinucleotide at the 3' splice site, and has an essential role in RNA splicing.¹³ With the exception of a single A26V mutation found in a case of refractory cytopenia with multilineage dysplasia, all the *U2AF35* mutations reported in adult myeloid malignancies involved one of the two hot spots within the two zinc-finger domains, S34 and Q157, which are highly conserved across species, suggesting the gain-of-function mutations.¹⁰ In JMML 4, the R156M *U2AF35* mutation affects a conserved amino acid adjacent to Q157, suggesting it may also be a gain-of-function mutation, leading to aberrant pre-mRNA splicing possibly in a dominant fashion.

SRSF2, better known as SC35, is a member of the serine/arginine-rich (SR) family of proteins.¹⁴ *SRSF2* binds to a splicing-enhancer element in pre-mRNA and has a crucial role not only in constitutive and alternative pre-mRNA splicing but also in transcription elongation and genomic stability.¹⁴ All mutations thus far identified in adult cases exclusively involved P95 within the intervening sequence between the N-terminal RNA-binding domain and the C-terminal RS domain.¹⁰ This region interacts with other SR proteins, again suggesting that the P95 mutation may result in gain-of-function.¹⁰ This proline residue is thought to determine the relative orientation of the two flanking domains of *SRSF2*, and a substitution at this position could compromise critical interactions with other splicing factors necessary for RNA splicing to take place. In contrast, the newly identified 6-bp in-frame deletion in JMML 17 results in two conserved amino acids, S170 and K171, within the RS domain. Although it may affect protein-protein interactions, the functional significance of this deletion remains elusive.

JMML is a unique form of pediatric MDS/MPN characterized by activation of the RAS/mitogen-activated protein kinase signaling pathway; in 90% of cases, there are germ line and/or somatic mutations of *NF1*, *NRAS*, *KRAS*, *PTPN11* and *CBL*.⁸ Although JMML shares some clinical and molecular features with CMML, its spectrum of gene mutations suggests that it is a neoplasm distinct from CMML.¹⁵ This was also confirmed by the current results that the splicing-pathway mutations are rare in JMML, whereas they are extremely frequent (~60%) in CMML.¹⁰ Although the two JMML cases carrying the splicing-pathway mutations had no known RAS-pathway mutations, both the pathway mutations frequently coexisted in CMML.⁸

To summarize, no mutations of *SF3B1*, *U2AF35*, *ZRSR2* or *SRSF2* are found in pediatric MDS and AML. In our study, except for *ZRSR2*, mutations were examined focusing on the reported hot spots in adult studies, raising a possibility that we may have missed some mutations occurring in other regions. However,

these hot spots represent evolutionally conserved amino acids and have functional relevance, it is unlikely that the distribution of hot spots in children significantly differs from adult cases and as such, we could safely conclude that mutations of *SF3B1*, *U2AF35*, *ZRSR2* and *SRSF2* are rare in myeloid neoplasms in children. Finally, mutations of *U2AF35* and *SRSF2* may have some role in the pathogenesis of JMML, although further evaluations are required.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

This work was supported by Research on Measures for Intractable Diseases, Health and Labor Sciences Research Grants, Ministry of Health, Labor and Welfare, by Research on Health Sciences focusing on Drug Innovation, and the Japan Health Sciences Foundation. We would like to thank M Matsumura, M Yin, N Hoshino and S Saito for their excellent technical assistance.

J Takita^{1,2}, K Yoshida³, M Sanada³, R Nishimura¹, J Okubo¹,
A Motomura¹, M Hiwatari¹, K Oki¹, T Igarashi¹,
Y Hayashi⁴ and S Ogawa³

¹Department of Pediatrics, Graduate School of Medicine,
The University of Tokyo, Tokyo, Japan;

²Department of Cell Therapy and Transplantation Medicine,
Graduate School of Medicine, The University of Tokyo, Tokyo, Japan;

³Cancer Genomics Project, Graduate School of Medicine,
The University of Tokyo, Tokyo, Japan and

⁴Gunma Children's Medical Center, Gunma, Japan
E-mail: sogawa-tyk@umin.ac.jp

REFERENCES

- Garcia-Manero G. Myelodysplastic syndromes: 2011 update on diagnosis, risk-stratification, and management. *Am J Hematol* 2011; **86**: 490–498.
- Delhommeau F, Dupont S, Della Valle V, James C, Trannoy S, Masse A *et al*. Mutation in TET2 in myeloid cancers. *N Engl J Med* 2009; **360**: 2289–2301.
- Thol F, Friesen I, Damm F, Yun H, Weissinger EM, Krauter J *et al*. Prognostic significance of ASXL1 mutations in patients with myelodysplastic syndromes. *J Clin Oncol* 2011; **29**: 2499–2506.
- Ley TJ, Ding L, Walter MJ, McLellan MD, Lamprecht T, Larson DE *et al*. DNMT3A mutations in acute myeloid leukemia. *N Engl J Med* 2010; **363**: 2424–2433.
- Nikolovski G, Langemeijer SM, Kuiper RP, Knops R, Massop M, Tonnissen ER *et al*. Somatic mutations of the histone methyltransferase gene EZH2 in myelodysplastic syndromes. *Nature Genet* 2010; **42**: 665–667.
- Green A, Beer P. Somatic mutations of IDH1 and IDH2 in the leukemic transformation of myeloproliferative neoplasms. *N Engl J Med* 2010; **362**: 369–370.
- Sanada M, Suzuki T, Shih LY, Otsu M, Kato M, Yamazaki S *et al*. Gain-of-function of mutated C-CBL tumour suppressor in myeloid neoplasms. *Nature* 2009; **460**: 904–908.
- Perez B, Kosmider O, Cassinat B, Renneville A, Lachenaud J, Kaltenbach S *et al*. Genetic typing of CBL, ASXL1, RUNX1, TET2 and JAK2 in juvenile myelomonocytic leukaemia reveals a genetic profile distinct from chronic myelomonocytic leukaemia. *Br J Haematol* 2010; **151**: 460–468.
- Oki K, Takita J, Hiwatari M, Nishimura R, Sanada M, Okubo J *et al*. IDH1 and IDH2 mutations are rare in pediatric myeloid malignancies. *Leukemia* 2011; **25**: 382–384.
- Yoshida K, Sanada M, Shiraishi Y, Nowak D, Nagata Y, Yamamoto R *et al*. Frequent pathway mutations of splicing machinery in myelodysplasia. *Nature* 2011; **478**: 64–69.
- Chen Y, Takita J, Hiwatari M, Igarashi T, Hanada R, Kikuchi A *et al*. Mutations of the PTPN11 and RAS genes in rhabdomyosarcoma and pediatric hematological malignancies. *Genes Chromosomes Cancer* 2006; **45**: 583–591.
- Shiba N, Kato M, Park MJ, Sanada M, Ito E, Fukushima K *et al*. CBL mutations in juvenile myelomonocytic leukemia and pediatric myelodysplastic syndrome. *Leukemia* 2010; **24**: 1090–1092.