

PROTEIN DEGRADATION IN CANCER ANOREXIA—CACHEXIA

In adults, the muscle mass remains fairly constant in the absence of stimuli such as exercise, such that protein synthesis and degradation remain in balance (37). In cachexia, muscle atrophy occurs, which results from a depression in protein synthesis, an increase in protein degradation or a combination of both (37) (Fig. 2). In recent years, it has become evident that specific regulating molecules are upregulated (e.g. members of the ubiquitin–proteasome system, myostatin and apoptosis-inducing factors), whereas other factors (e.g. insulin-like growth factor 1, IGF-1) are downregulated in cachexia muscle wasting (31). A major barrier to the effective management of skeletal muscle wasting is the inadequate understanding of its underlying biological mechanisms (31). The most evident metabolic explanation for muscle decline is an imbalance between protein catabolism and anabolism (31).




Figure 2. An abbreviated schematic diagram of skeletal muscle in cancer anorexia–cachexia. Source: (31) and (69) with modification. Arrows indicate the activation of the process, -| indicates the inhibition of the process. The balance of skeletal muscle has been shifted towards protein breakdown, finally leading to the weight loss, weakness and fatigue that characterize cancer anorexia–cachexia. Muscle RING-finger protein-1 (MuRF1). Atrogin-1/muscle atrophy F-box (MAFbx). Nuclear factor kappa B (NF-κB). Insulin-like growth factor 1 (IGF-1). forkhead box O (FoxO). phosphatidylinositol-3-kinase (PI3K). mTOR = mammalian target of rapamycin. p70S6K = p70 S6 kinase.

In addition to an increase in catabolism, a reduction in anabolism has been shown to occur in cancer-related cachexia (31). Skeletal muscle wasting in cancer cachexia can be mediated by multiple factors derived from tumor and host cells (38).

At least four major proteolytic pathways (lysosomal, Ca²⁺-dependent, caspase-dependent and ubiquitin–proteasome-dependent) operate in skeletal muscle and may be altered during muscle cachexia (31). Aside from these four distinct pathways, the autophagic/lysosomal pathway must also be considered (31). In this pathway, portions of the cytoplasm and cell organelles are sequestered into autophagosomes, which subsequently fuse with lysosomes, where the proteins are digested (39). When dissecting the molecular regulation of the ubiquitin–proteasome-dependent system (UPS) and autophagy, it became evident that forkhead box O (FoxO) transcription factors play a central role (31). FoxO transcription factors, which are normally phosphorylated and inactivated by phosphatidylinositol 3-kinase (PI3K)-Akt/PKB, translocate into the cell nucleus and induce the transcription of the skeletal muscle-specific E3 ubiquitin ligases, muscle RING-finger protein-1 (MuRF1) and atrogin-1/muscle atrophy F-box (MAFbx) (40), as well as autophagy-related genes such as LC3 and Bnip3 (41). Upstream of PI3K-Akt, several factors including reactive oxygen species (ROS), TNF-α, tumor-released proteolysis-inducing factor (PIF), peroxisome proliferator-activated receptor gamma coactivator 1 alpha (PGC-1α) and IGF-1 have been shown to influence this regulatory system (40,42–44). In contrast, protein anabolic factors such as IGF-1 counteract muscle atrophy (31). Aside from inhibiting autophagy and the UPS, IGF-1 activates protein synthesis via the Akt-mammalian target of rapamycin (mTOR)–p70 S6 kinase (p70S6K) signaling pathway (45,46).

The UPS is a major intracellular system that regulates skeletal muscle wasting in response to tumor factors and inflammatory cytokines (38). In cancer cachexia, the decrease in skeletal muscle protein synthesis is partly related to the increased serum levels of PIF (31). Intravenous administration of PIF to normal mice produced a rapid decrease in body weight that was accompanied by increased mRNA levels of ubiquitin in the gastrocnemius muscle (43). There were also increased protein levels of the 20 S proteasome core and the 19 S regulatory subunit, suggesting activation of the ATP–ubiquitin-dependent proteolytic pathway (31). Recent evidence suggests that PIF decreases protein synthesis by inhibiting protein translation initiation through phosphorylation of the eukaryotic initiation factor 2 (eIF2-α) (47).

Myostatin is an extracellular cytokine that is mostly expressed in skeletal muscles and is known to play a crucial role in the negative regulation of muscle mass (48). Upon binding to the activin type IIB receptor, myostatin can initiate several different signaling cascades, resulting in decreased muscle growth and differentiation (48). Muscle size is regulated via a complex interplay of myostatin signaling with the IGF-1/PI3K/Akt pathway, which is responsible for increased protein synthesis in muscle (48). Therefore, the regulation of muscle weight is a process in which myostatin plays a central

role, but the mechanism of its action and the role of the signaling cascades involved are not fully understood (48). Myostatin upregulation was observed in the pathogenesis of muscle wasting during cancer cachexia (48).

One of the main positive regulators of muscle growth is IGF-1 (48). Under normal conditions, IGF-1 signaling seems to be dominant and blocks the myostatin pathway (49). However, an inhibition of IGF-1 was observed when myostatin was overexpressed (50,51). IGF-1 can prevent TGF- α family-mediated apoptosis (52), and it was shown that in the absence of IGF-1, the level of apoptosis in C2C12 cells treated with myostatin increased (48). The mechanism by which IGF-1 regulates myostatin signaling includes the inhibition of transcription factors responsible for the induction of atrogenes via phosphorylation through the PI3K/Akt pathway (48). Akt plays a significant role in different metabolic processes in the cell, particularly in the hypertrophic response to insulin and IGF-1 (53,54). Akt is the crossing point between the IGF-1/myostatin pathways (48). It is likely that under conditions of muscle wasting, myostatin can reverse the Akt/mTOR pathway, which is normally responsible for protein synthesis, to inhibit protein synthesis via FoxO, GSK-3 β or other unknown patterns, leading to the loss of muscle mass (48). Another factor that may contribute to decreased anabolism is angiotensin II (31). In an animal model of continuously administered angiotensin II, markedly reduced plasma IGF I levels occurred (55). Compared with a sham treatment, angiotensin II-infused hypertensive rats lost 18–26% of their body weight within 1 week, an effect that was completely reversible by losartan, an AT1 receptor antagonist (55).

There is a wealth of evidence in the current literature that oxidative stress is associated with chronic diseases, and it is assumed that an increase in ROS directs muscle cells into a catabolic state that leads to muscle wasting (31,56,57). In cachexia, ROS are regarded as crucial players for muscle protein catabolism via their stimulation of the UPS (31). Reaction products are measured as indirect markers of oxidative stress (31). In cachexia, malondialdehyde is regarded as one such indirect marker (31). In addition, experimental cancer cachexia appears to be mediated by increased nitrosative stress secondary to increased NO formation. Indeed, protein tyrosine nitration is markedly increased in the muscles of tumor-bearing rats with advanced cachexia (58). In cachexia, the increase in protein tyrosine nitration in ROS is due to significantly lower activities of the antioxidant enzymes superoxide dismutase and glutathione peroxidase (59).

Experimental data suggest that local IGF-1 may act as a regenerative agent, promoting the recruitment of stem cells to sites of muscle injury (60). Because IGF-1 is reduced in experimental models of cachexia (61), it is reasonable to assume that under conditions of cachexia, the function of satellite cells is impaired (31). Other factors controlling the differentiation of satellite cells into functional fibers include nuclear factor kappa B (NF- κ B) and myostatin (31). Data are available that demonstrate a beneficial effect of myostatin inhibition in cancer cachexia (62), but negative study results have also been

reported (63). With respect to apoptosis, several reports demonstrated an increase in apoptosis or apoptosis-related proteins in skeletal muscle after the induction of cachexia (31). The skeletal muscle of cachectic tumor-bearing animals reveals the activation of DNA fragmentation, a hallmark of apoptosis (64). In addition to DNA fragmentation, a significant up-regulation of caspase-1, -3, -6, -8, and -9 activity was also documented in the gastrocnemius muscles of tumor-bearing mice (65).

There is a relative deficiency or resistance to anabolic hormones in cachectic states. Up to 50% of men with metastatic cancer can present with low concentrations of testosterone prior to chemotherapy (66). A reduction in testosterone might lead to reduced bone mass, muscle strength and sexual function in both men and women (67,68). Low concentrations of testosterone and other anabolic hormones are major contributors to cachexia-related wasting of skeletal muscle (69). However, with respect to a correlation between body composition (including muscle mass) and the concentration of anabolic hormones, conflicting results have been reported in the current literature (31). Some studies found a correlation (66,70), whereas others reported no association (71).

TRANSLATIONAL ASPECTS OF HERBAL MEDICINE PARTICULARLY FOR CANCER ANOREXIA–CACHEXIA

Many effective chemotherapeutic agents for cancer are burdened by toxicities that can reduce patient's quality of life or hinder their effective use. 5HT3 receptor antagonists, dexamethasone and aprepitant significantly improved chemotherapy-induced nausea and vomiting. Empirical antibiotics significantly improved neutropenic fever. A multinational survey found that 35.9% of cancer patients were either past or present users of CAM. Herbal medicines were by far the most commonly used group of treatments, escalating in use from 5.3% before the diagnosis of cancer to 13.9% after the diagnosis of cancer (72). Herbal medicines are believed by the general public to be safe, cause less side-effects and less likely to cause dependency (73).

Herbal medicine in palliative treatment of cancer is a fast-emerging area. Palliative care in cancer treatment aims not only for disease control but for addressing the patient's physical and psychosocial symptoms, and improving the QOL. The mechanism involves an enhancement of signaling by ghrelin which was discovered in 1999 as an appetite-stimulating peptide from the stomach (74–77). Currently, ghrelin agonists and antagonists are being developed and tested for the treatment of anorexia/cachexia and obesity, respectively.

RIKKUNSHITO AS AN ACTIVATOR OF GHRELIN SIGNALING

It is known that plasma ghrelin levels are elevated under cachectic conditions caused by a variety of underlying disorders

(78–82). Although this phenomenon has been called ‘ghrelin resistance’, these elevations may be a compensatory response that reflects the negative energy balance state (83). Several randomized, double-blind placebo-controlled trials have demonstrated the efficacy and safety of ghrelin or growth hormone (GH) secretagogue (GHS) in patients with cancer-associated cachexia (84–86). Therefore, evaluation of the role of ghrelin in the pathogenesis and treatment of such cachectic conditions is warranted (74).

In our study, it was demonstrated that plasma acyl ghrelin concentrations in tumor-bearing rats were higher than those in free-fed normal rats, but lower than those in pair-fed normal rats, and had an inverse relationship with plasma leptin concentrations (87). These results indicate that changes in ghrelin and leptin secretion in pair-fed animals represent a compensatory mechanism in a persistent catabolic state and that these responses are attenuated in tumor-bearing rats (87). Peripheral ghrelin administration stimulates food intake in melanoma cell-bearing mice and cancer patients (85) in the short term as well as in lean, healthy men and women (88). There were similar therapeutic effects of ghrelin on anorexia and gastrointestinal (GI) dysmotility in cachectic animal models, suggesting that high plasma concentrations of ghrelin may overcome resistance to the appetite-stimulating effects of the endogenous peptide in the short term (87). Oral administration of *rikkunshito* increases plasma acyl ghrelin levels in humans, mice, rats (89,90) and dogs (87). *Rikkunshito* stimulates ghrelin secretion through 5-hydroxytryptamine (5-HT) 2b/2c receptor antagonism, and its active flavonoid ingredients such as hesperidin that antagonize 5-HT2b/2c receptor binding have been identified (89).

The central 5-HT system has been implicated in the processes of meal satiation and satiety (87). 5-HT reuptake inhibitors such as fenfluramine and 5-HT2c agonists attenuate food intake and weight gain in rodents and humans (91–93), with the involvement of potentiated MC signaling and decreased ghrelin secretion (87). 5-HT also inhibits NPY/AgRP neurons by activating the 5-HT1bR, leading to decreased orexigenic signaling and inhibitory drive onto POMC cells (87). However, the previous study has demonstrated that the 5-HT2cR has a major role in the regulation of physiological fasted and fed motor activities in addition to feeding through changes in endogenous ghrelin (90). In this study, the decreases in food intake and GI motor activities in tumor-bearing rats were recovered after administration of either a 5-HT2cR antagonist or ghrelin (87). The 5-HT concentration in the hypothalamus is increased in humans and animals with cancer (94,95); in addition, NPY and dopamine concentrations decrease simultaneously, while 5-HT concentration increases in the PVN at the onset of anorexia in tumor-bearing rats (96). These findings suggest that 5-HT2cR activation in tumor-bearing rats induces anorexia in part via decreased ghrelin secretion (87).

The hypothermia in tumor-bearing rats may be due to a state of negative energy balance or a decrease in the threshold for the activation of thermogenesis, which is involved in starvation-induced hypothermia (97). IL-1 β and leptin (98)

decrease the expression of ghrelin mRNA in the stomach, whereas IL-6 produced in various cells, including adipocytes, regulates leptin production (99). These findings suggest that cytokines have an important role in energy balance through the persistent activation of the leptin system and the inhibition of the ghrelin-NPY/agouti-related peptide orexigenic network in tumor-bearing rats (87). In addition to NPY and agouti-related peptide, the level of POMC mRNA was also decreased in the hypothalamus of the tumor-bearing rats (87). Synaptic input organization and mRNA expression of POMC neurons have been shown to be increased in adrenalectomized animals and restored by corticosterone replacement (100). Thus, activity of hypothalamic POMC neurons may be affected by changes in circulating levels of corticosterone and a state of negative energy balance (87).

Hypothalamic 5-HT and CRF activities are stimulated by proinflammatory cytokines in the circulation and the hypothalamus (101,102). A CRF receptor antagonist attenuated cancer anorexia–cachexia, and administration of the 5-HT2cR antagonist or *rikkunshito* reduced hypothalamic CRF levels and anxiety-related behaviors in tumor-bearing rats (87). The improvement in anxiety by *rikkunshito* may lead to a higher quality of life in cancer patients (87). Some studies suggest that ghrelin induce anxiety, whereas others suggest that the elevated ghrelin helps animals cope with stress by producing anxiolytic-like response (103). Future studies are needed to sort out the effect of ghrelin on anxiety-like behavior as in the case of NPY (76). Importantly, a hypothalamic 5-HT-CRF receptor pathway that regulates ghrelin secretion has a major role in cancer anorexia–cachexia (87).

It has been previously shown that a central 5-HT2cR pathway regulates ghrelin secretion without downstream activation of melanocortin 3/4 receptors (90). The 5-HT2cR is expressed in many brain regions and its expression is restricted to the central nervous system (87). Dual-neurohistochemical labeling has revealed that approximately one-half of PVN CRF-containing neurons co-express 5-HT2cR mRNA (104). In this study, 5-HT activated single CRF neurons isolated from the PVN, and the activities of the CRF neurons were blocked by simultaneous administration of *rikkunshito* (87). Moreover, intracerebroventricular administration of CRF decreased plasma acyl ghrelin in fasted rats (87). These findings suggest that CRF neurons are involved in 5-HT-regulated ghrelin secretion (87).

The GH secretagogue receptor (GHS-R) is reportedly expressed in vagal afferent neurons, and the gastric vagus nerve system is involved in the effect of ghrelin on food intake and GI motor activities (105,106). It was demonstrated that ghrelin decreased the afferent activity of the gastric vagus nerve (87). Gastric ghrelin signaling via vagal afferents stimulated the efferent activities of both the gastric and the celiac branches of the vagus nerve and suppressed the activity of the sympathetic nerve (87). Peripheral administration of a higher dose of ghrelin increased the discharge rate of the vagal efferent nerve, probably in part through the GHS-R in the ARC of the hypothalamus (87). It has also been shown that *rikkunshito*

activated the efferent vagus nerve, which may be mediated by both the vagal afferent nerve and the direct central action (87). In addition, ghrelin-induced cellular signaling in GHS-R-expressing cells was enhanced by pretreatment with *rikkunshito* and its active components, such as atractylodin, which stimulate ghrelin/GHS-R binding activity (87). Similar potentiating effects of *rikkunshito* were observed in rat ARC NPY neurons (87). These findings suggest that the physiological functions of endogenous ghrelin are enhanced by the dual actions of *rikkunshito*, which involve the stimulation of ghrelin secretion and the activation of GHS-R activity, possibly due to allosteric changes in the receptor (87). This potentiation of the ghrelin effect by *rikkunshito* on NPY neurons could be orexigenic because the activity of ghrelin-responsive NPY neurons is coupled to feeding (107,108). As mentioned in the past section, ghrelin strongly stimulates GH secretion in humans (109–112), which regulates IGF-1 levels, and increases muscle strength (113,114). Moreover, ghrelin induces the anti-inflammatory cytokine (115,116), while suppresses the production of proinflammatory cytokines (115,117–119), and inhibits the activation of NF-κB which may regulate skeletal muscle proteasome expression and protein degradation (83,116,118). Consequently, potentiation of ghrelin receptor signaling with *rikkunshito* can be valuable in the treatment of anorexia and muscle

wasting which characterize cancer anorexia–cachexia syndrome (87).

The adverse effect of (D-Lys3)-GHRP-6 on survival in tumor-bearing rats has been indicated in this study, suggesting that the potentiation of ghrelin signaling is critical to the attenuation of anorexia–cachexia and the prolongation of survival in subjects with cancer (87). *Rikkunshito* and its active component, atractylodin, prolonged survival in these animals, and this effect was enhanced by the concomitant administration of cisplatin (87). Cancer patients receiving chemotherapy or radiation therapy may experience nausea, vomiting, taste changes, stomatitis and diarrhea, which could contribute to weight loss and decreased survival (87). Therefore, cancer anorexia–cachexia syndrome is a major obstacle in cancer chemotherapy (8). The use of *rikkunshito* in tumor-bearing rats was effective not only against anorexia–cachexia, but also for promoting survival, particularly in combination with chemotherapy (87). However, daily administration of a 5-HT_{2c} receptor antagonist failed to prolong survival, suggesting that a sensitizing effect on the GHS-R may be essential for ameliorating ghrelin resistance in anorexia–cachexia in the long term (87). Pancreatic cancer patients generally respond poorly to chemotherapy, resulting in a higher frequency of anorexia–cachexia (87). These results suggest that *rikkunshito* may be useful in clinical practice for cachectic




Figure 3. The dual actions of *rikkunshito* on ghrelin signaling and their therapeutic benefits on cancer anorexia–cachexia. Source: (87) with modification. Hypothalamic CRF neurons are activated by cytokines through serotonin (5-HT) and the 5-HT_{2c} receptor (5-HT_{2c}R), which show a functional divergence. The existence of a novel 5-HT-CRF neuronal pathway inhibits ghrelin secretion and has a pathogenetic role in cancer anorexia–cachexia. *Rikkunshito* and its active component, hesperidin, a principal component of *Aurantii Nobilis Pericarpium*, stimulate ghrelin secretion from the stomach by interrupting this 5-HT-CRF pathway via 5-HT_{2c}R antagonism. Another active component, atractylodin, a principal component of *Atractylodis Lanceae Rhizoma*, potentiates the action of ghrelin, presumably by allosterically sensitizing the GHS-R on the vagal afferent terminals of the stomach or the NPY neurons of the hypothalamic arcuate nucleus (ARC). Thus, both the release of ghrelin and the potentiation of ghrelin/GHS-R signaling are important for mitigating ghrelin insufficiency and resistance, which are characteristics of cancer anorexia–cachexia. Consequently, potentiation of ghrelin receptor signaling with *rikkunshito* may be an attractive treatment for anorexia and muscle wasting and may prolong survival in patients with cancer anorexia–cachexia.

cancer patients via its dual action on ghrelin secretion and receptor sensitization (Fig. 3) (87). Prospective randomized trials are warranted.

A recent study regarding the underlying mechanisms of *rikkunshito* has shown that *rikkunshito* and its component 10-gingerol may inhibit the degradation of acyl-ghrelin by inhibiting the circulating ghrelin degrading enzyme (120). Another study has shown that administration of *rikkunshito* reversed the decrease in hypothalamic ghrelin secretion and food intake 24 h after cisplatin treatment (121). Cisplatin-induced anorexia is mediated through reduced hypothalamic ghrelin secretion (121). Cerebral serotonin 2C receptor activation partially induces decrease in hypothalamic ghrelin secretion, and *rikkunshito* suppresses cisplatin-induced anorexia by enhancing this secretion (121).

Altogether, the synergism of activity of the herbs demonstrated in these studies highlights the importance of adopting traditional approaches in the utilization of traditional medicines.

CONCLUSIONS

The best way of treating cancer cachexia is to cure cancer, but unfortunately this remains an infrequent achievement among adults with advanced solid tumors (8,122). Appetite, body weight and survival are endpoints for cancer anorexia–cachexia management. Therefore, the treatment goal for cachexia should be the reversal of the loss of body weight and muscle mass with a variety of pharmacological agents (Fig. 4) (8). The European Palliative Care Research Collaboration has developed evidence-based recommendations on classification and treatment of cachexia in advanced cancer patients as part of its clinical guideline work (1). These treatment guidelines focus on patients with advanced cancer likely to suffer from refractory cachexia. Many of these patients are receiving palliative care, and life expectancy often is short. Only little cachexia research has been done on this patient group, and the treatment guidelines had to consider whether research results from other disease stages are applicable for these patients with advanced and incurable disease and for refractory cachexia (1).

Herbal medicine could substantially influence cancer therapy as adjuvant treatment. Although herbal medicines have not been fully accepted by mainstream medicine because of the complex nature of the formulae, the stringent quality control of Japanese herbal (Kampo) medicine and reproducibility of preclinical findings, together with few adverse events, have made herbal medicines more and more attractive for the management of intractable diseases such as cancer. In recent years, studies on the evaluation of the therapeutic and toxic activities of herbal medicinal products became available and popular (123). The advances in modern biotechnology have led to the discovery of many new active constituents (123). Although the working mechanisms of some of the herbs are unclear and remain to be elucidated, they are worth further studying as newly potential therapy agents for cancer




Figure 4. The potential modalities of pharmacological intervention of cancer anorexia–cachexia syndrome. Source: (8) with modification. Agents were classified as those established (first-line) or those unproven/investigational (second-line), depending on their site or mechanism of actions. ①, inhibitors of production/release of cytokines and other factors; ②, gastroprokinetic agents with or without antiemetic effect; ③, blockers of the Cori cycle; ④ ⑤, blockers of fat and muscle tissue wasting; ⑥, appetite stimulants with or without antiemetic effect; and ⑦, antianxiety/depressant drugs. These agents should be selected on an individual basis according to the cause of cachexia or the state of the patient. *The precise actions of statins on skeletal muscle still remain controversial. First-line treatments, glucocorticoids ① ⑥, progesterones ① ⑥; second-line treatments, cannabinoids ⑥, cyproheptadine ⑥, branched-chain amino acids ⑤ ⑥, metoclopramide ② ⑥, eicosapentaenoic acid ① ④ ⑤, 5'-deoxy-5-fluorouridine ①, melatonin ①, thalidomide ①, β_2 -adrenoceptor agonists ⑤, non-steroidal anti-inflammatory drugs ① ⑥, others, anabolic steroids ⑤, pentoxifylline ①, hydrazine sulfate ③, statin ① ⑤*, angiotensin-converting-enzyme inhibitor ⑤, selective androgen receptor modulator ⑤, ghrelin agonists ① ② ③ ④ ⑤ ⑥.

treatment (123). The multicomponent herbal medicines capable of targeting multiple sites could be useful for future drug discovery. Mechanistic studies and identification of active compounds could lead to new discoveries in biological and biomedical sciences.

Funding

This work was supported by Grants-in-Aid for Scientific Research (B) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT), the Ministry of Health Labor and Welfare (MHLW) and the Japan Society for the

Promotion of Science (JSPS) Strategic Young Researcher Overseas Visits Program for Accelerating Brain Circulation.

Conflict of interest statement

None declared.

References

- Radbruch L, Elsner F, Trottenberg P, Strasser F, Fearon K. Clinical practice guidelines on cancer cachexia in advanced cancer patients. Aachen, Department of Palliative Medicine/European Palliative Care Research Collaborative 2010. http://www.eprc.org/publication_listfiles.php?id=mWdBCM15eXVlcNFk7Gnq.
- Inui A. Cancer anorexia-cachexia syndrome: are neuropeptides the key? *Cancer Res* 1999;59:4493-501.
- Tisdale M. Biology of cachexia. *J Natl Cancer Inst* 1997;89:1763-73.
- Larkin M. Thwarting the dwindling progression of cachexia. *Lancet* 1998;1336.
- Kotler D. Cachexia. *Ann Intern Med* 2000;133:622-34.
- Davis M. Cachexia and anorexia: cancer's covert killer. *Support Care Cancer* 2000;8:180-7.
- Fearon KC, Barber MD, Falconer JS, McMillan DC, Ross JA, Preston T. Pancreatic cancer as a model: inflammatory mediators, acute-phase response, and cancer cachexia. *World J Surg* 1999;23:584-8.
- Inui A. Cancer anorexia-cachexia syndrome: current issues in research and management. *CA Cancer J Clin* 2002;52:72-91.
- Friedman JM, Halaas JL. Leptin and the regulation of body weight in mammals. *Nature* 1998;395:763-70.
- Flier JS, Maratos-Flier E. Obesity and the hypothalamus: novel peptides for new pathways. *Cell* 1998;92:437-40.
- Schwartz MW, Figlewicz DP, Baskin DG, Woods SC, Porte D. Insulin in the brain: a hormonal regulator of energy balance. *Endocr Rev* 1992;13:387-414.
- Inui A. Feeding and body-weight regulation by hypothalamic neuropeptides—mediation of the actions of leptin. *Trends Neurosci* 1999;22:62-7.
- Yoshida N, Maejima Y, Sedbazar U, et al. Stressor-responsive central nesfatin-1 activates corticotropin-releasing hormone, noradrenaline and serotonin neurons and evokes hypothalamic-pituitary-adrenal axis. *Aging (Albany NY)* 2010;2:775-84.
- Plata-Salamán CR. Immunoregulators in the nervous system. *Neurosci Biobehav Rev* 1991;15:185-215.
- Plata-Salamán CR. Anorexia during acute and chronic disease. *Nutrition* 1996;12:69-78.
- Moldawer LL, Copeland EM. Proinflammatory cytokines, nutritional support, and the cachexia syndrome: interactions and therapeutic options. *Cancer* 1997;79:1828-39.
- Haslett PA. Anticytokine approaches to the treatment of anorexia and cachexia. *Semin Oncol* 1998;25:53-7.
- Mantovani G, Maccio A, Lai P, Massa E. Cytokine activity in cancer-related anorexia/cachexia: role of megestrol acetate and medroxyprogesterone acetate. *Semin Oncol* 1998;25:45-52.
- Donohoe CL, Ryan AM, Reynolds JV. Cancer cachexia: mechanisms and clinical implications. *Gastroenterol Res Pract* 2011;2011:601434.
- Sternberg EM. Neural-immune interactions in health and disease. *J Clin Invest* 1997;100:2641-7.
- Licinio J, Wong ML. American Society for Clinical Investigation. Pathways and mechanisms for cytokine signaling of the central nervous system. *J Clin Invest* 1997;100:2941-7.
- Hopkins SJ, Rothwell NJ. Cytokines and the nervous system. I: expression and recognition. *Trends Neurosci* 1995;18:83-8.
- Rothwell NJ, Hopkins SJ. Cytokines and the nervous system II: actions and mechanisms of action. *Trends Neurosci* 1995;18:130-6.
- Moldawer LL, Rogy MA, Lowry SF. The role of cytokines in cancer cachexia. *JPEN J Parenter Enteral Nutr* 1992;16:43S-9S.
- Noguchi Y, Yoshikawa T, Matsumoto A, Svaninger GS, Gelin J. Are cytokines possible mediators of cancer cachexia? *Surg Today* 1996;26:467-75.
- Matthys P, Billiau A. Cytokines and cachexia. *Nutrition* 1997;13:763-70.
- Gelin J, Moldawer L, Lönnroth C, Sherry B. Role of endogenous tumor necrosis factor α and interleukin 1 for experimental tumor growth and the development of cancer cachexia. *Cancer Res* 1991;51:415-21.
- Torelli G, Meguid M. Use of recombinant human soluble TNF receptor in anorectic tumor-bearing rats. *Am J Physiol Regul Integr Comp Physiol* 1999;277:R850-5.
- McCarthy HD, Crowder RE, Dryden S, Williams G. Megestrol acetate stimulates food and water intake in the rat: effects on regional hypothalamic neuropeptide Y concentrations. *Eur J Pharmacol* 1994;265:99-102.
- Scarlett JM, Jobst EE, Enriori PJ, et al. Regulation of central melanocortin signaling by interleukin-1 β . *Endocrinology* 2007;148:4217-25.
- Lenk K, Schuler G, Adams V. Skeletal muscle wasting in cachexia and sarcopenia: molecular pathophysiology and impact of exercise training. *J Cachex Sarcopenia Muscle* 2010;1:9-21.
- Williams A, Sun X, Fischer J. The expression of genes in the ubiquitin-proteasome proteolytic pathway is increased in skeletal muscle from patients with cancer. *Surgery* 1999;126:744-9.
- Llovera M, García-Martínez C, López-Soriano J, et al. Role of TNF receptor 1 in protein turnover during cancer cachexia using gene knockout mice. *Mol Cell Endocrinol* 1998;142:183-9.
- Ungureanu-Longrois D, Balligand J. Myocardial contractile dysfunction in the systematic inflammatory response syndrome: role of a cytokine-inducible nitric oxide synthase in cardiac myocytes. *J Mol Cell Cardiol* 1995;27:155-67.
- Sherry B, Gelin J, Fong Y, Marano M, Wei H. Anticachectin/tumor necrosis factor- α antibodies attenuate development of cachexia in tumor models. *FASEB J* 1989;3:1956-62.
- Trikha M, Corringham R, Klein B. Targeted anti-interleukin-6 monoclonal antibody therapy for cancer. *Clin Cancer Res* 2003;9:4653-65.
- Tisdale MJ. Mechanisms of cancer cachexia. *Physiol Rev* 2009;89:381-410.
- Guttridge D. Molecular mechanisms of muscle wasting in cancer cachexia. *Nutr Support Cancer* 2006;1-13.
- Lum JJ, DeBerardinis RJ, Thompson CB. Autophagy in metazoans: cell survival in the land of plenty. *Nat Rev Mol Cell Biol* 2005;6:439-48.
- Sandri M, Sandri C, Gilbert A, et al. Foxo transcription factors induce the atrophy-related ubiquitin ligase atrogin-1 and cause skeletal muscle atrophy. *Cell* 2004;117:399-412.
- Mammucari C, Milan G, Romanello V, et al. FoxO3 controls autophagy in skeletal muscle in vivo. *Cell Metab* 2007;6:458-71.
- Dodd SL, Gagnon BJ, Senf SM, Hain BA, Judge AR. Ros-mediated activation of NF- κ B and Foxo during muscle disuse. *Muscle Nerve* 2010;41:110-3.
- Lorite MJ, Smith HJ, Arnold JA, Morris A, Thompson MG, Tisdale MJ. Activation of ATP-ubiquitin-dependent proteolysis in skeletal muscle in vivo and murine myoblasts in vitro by a proteolysis-inducing factor (PIF). *Br J Cancer* 2001;85:297-302.
- Sandri M, Lin J, Handschin C, et al. PGC-1 α protects skeletal muscle from atrophy by suppressing FoxO3 action and atrophy-specific gene transcription. *Proc Natl Acad Sci USA* 2006;103:16260-5.
- Glass DJ. Signalling pathways that mediate skeletal muscle hypertrophy and atrophy. *Nat Cell Biol* 2003;5:87-90.
- Guttridge DC. Signaling pathways weigh in on decisions to make or break skeletal muscle. *Curr Opin Clin Nutr Metab Care* 2004;7:443-50.
- Eley HL, Russell ST, Tisdale MJ. Effect of branched-chain amino acids on muscle atrophy in cancer cachexia. *Biochem J* 2007;407:113-20.
- Elkina Y, Haehling S, Anker SD, Springer J. The role of myostatin in muscle wasting: an overview. *J Cachex Sarcopenia Muscle* 2011;2:143-51.
- Trendelenburg AU, Meyer A, Rohner D, Boyle J, Hatakeyama S, Glass DJ. Myostatin reduces Akt/TORC1/p70S6K signaling, inhibiting myoblast differentiation and myotube size. *Am J Physiol Cell Physiol* 2009;296:C1258-70.
- Amirouche A, Durieux A-C, Banzet S, et al. Down-regulation of Akt/mammalian target of rapamycin signaling pathway in response to myostatin overexpression in skeletal muscle. *Endocrinology* 2009;150:286-94.

51. Morissette MR, Cook SA, Buranasombati C, Rosenberg MA, Rosenzweig A. Myostatin inhibits IGF-1-induced myotube hypertrophy through Akt. *Am J Physiol Cell Physiol* 2009;297:C1124–32.
52. Chen RH, Su YH, Chuang RL, Chang TY. Suppression of transforming growth factor-beta-induced apoptosis through a phosphatidylinositol 3-kinase/Akt-dependent pathway. *Oncogene* 1998;17:1959–68.
53. Bodine SC, Stitt TN, Gonzalez M, et al. Akt/mTOR pathway is a crucial regulator of skeletal muscle hypertrophy and can prevent muscle atrophy in vivo. *Nat Cell Biol* 2001;3:1014–9.
54. Zdychová J, Komers R. Emerging role of Akt kinase/protein kinase B signaling in pathophysiology of diabetes and its complications. *Physiol Res* 2005;54:1–16.
55. Brink M, Wellen J, Delafontaine P. Angiotensin II causes weight loss and decreases circulating insulin-like growth factor I in rats through a pressor-independent mechanism. *J Clin Invest* 1996;97:2509–16.
56. Buck M, Chojkier M. Muscle wasting and dedifferentiation induced by oxidative stress in a murine model of cachexia is prevented by inhibitors of nitric oxide synthesis and antioxidants. *EMBO J* 1996;15:1753–65.
57. Laviano A, Meguid MM, Preziosa I, Rossi Fanelli F. Oxidative stress and wasting in cancer. *Curr Opin Clin Nutr Metab Care* 2007;10:449–56.
58. Barreiro E, la Puente de B, Busquets S. Both oxidative and nitrosative stress are associated with muscle wasting in tumour-bearing rats. *FEBS Lett* 2005;579:1646–52.
59. Mantovani G, Macciò A, Madeddu C, et al. Antioxidant agents are effective in inducing lymphocyte progression through cell cycle in advanced cancer patients: assessment of the most important laboratory indexes of cachexia and oxidative stress. *J Mol Med* 2003;81:664–73.
60. Musarò A, Giacinti C, Borsellino G, et al. Stem cell-mediated muscle regeneration is enhanced by local isoform of insulin-like growth factor 1. *Proc Natl Acad Sci USA* 2004;101:1206–10.
61. Costelli P, Muscaritoli M, Bossola M, et al. IGF-1 is downregulated in experimental cancer cachexia. *Am J Physiol Regul Integr Comp Physiol* 2006;291:R674–83.
62. Benny Klimek ME, Aydogdu T, Link MJ, Pons M, Koniaris LG, Zimmers TA. Acute inhibition of myostatin-family proteins preserves skeletal muscle in mouse models of cancer cachexia. *Biochem Biophys Res Commun* 2010;391:1548–54.
63. Bonetto A, Penna F, Minero VG, et al. Deacetylase inhibitors modulate the myostatin/follistatin axis without improving cachexia in tumor-bearing mice. *Curr Cancer Drug Targets* 2009;9:608–16.
64. van Royen M, Carbó N, Busquets S, et al. DNA fragmentation occurs in skeletal muscle during tumor growth: A link with cancer cachexia? *Biochem Biophys Res Commun* 2000;270:533–7.
65. Belizário JE, Lorite MJ, Tisdale MJ. Cleavage of caspases-1, -3, -6, -8 and -9 substrates by proteases in skeletal muscles from mice undergoing cancer cachexia. *Br J Cancer* 2001;84:1135–40.
66. Chlebowski RT, Heber D. Hypogonadism in male patients with metastatic cancer prior to chemotherapy. *Cancer Res* 1982;42:2495–8.
67. Zitzmann M. Hormone substitution in male hypogonadism. *Mol Cell Endocrinol* 2000;161:73–88.
68. Lobo RA. Androgens in postmenopausal women: production, possible role, and replacement options. *Obstet Gynecol Surv* 2001;56:361–76.
69. Evans WJ, Morley JE, Argilés J, et al. Cachexia: a new definition. *Clin Nutr* 2008;27:793–9.
70. Hein L, Barsh G, Pratt R, Dzau V. Behavioural and cardiovascular effects of disrupting the angiotensin II type-2 receptor gene in mice. *Nature* 1995;377:744–7.
71. Garcia J, Li H, Mann D, Epner D, Hayes T. Hypogonadism in male patients with cancer. *Cancer* 2006;106:2583–91.
72. Molassiotis A, Fernandez-Ortega P, Pud D, et al. Use of complementary and alternative medicine in cancer patients: a European survey. *Ann Oncol* 2005;16:655–63.
73. Lynch N, Berry D. Differences in perceived risks and benefits of herbal, over-the-counter conventional, and prescribed conventional, medicines, and the implications of this for the safe and effective use of herbal products. *Complement Ther Med* 2007;15:84–91.
74. Inui A. Nature Publishing Group. Ghrelin: an orexigenic and somatotrophic signal from the stomach. *Nat Rev Neurosci* 2001;2:551–60.
75. Inui A. Neuropeptide gene polymorphisms and human behavioural disorders. *Nat Rev Drug Discov* 2003;2:986–98.
76. Kamiji MM, Inui A. Neuropeptide Y receptor selective ligands in the treatment of obesity. *Endocr Rev* 2007;28:664–84.
77. Chen C-Y, Asakawa A, Fujimiya M, Lee S-D, Inui A. Ghrelin gene products and the regulation of food intake and gut motility. *Pharmacol Rev* 2009;61:430–81.
78. Nagaya N, Uematsu M, Kojima M, et al. Elevated circulating level of ghrelin in cachexia associated with chronic heart failure: relationships between ghrelin and anabolic/catabolic factors. *Circulation* 2001;104:2034–8.
79. Itoh T, Nagaya N, Yoshikawa M, et al. Elevated plasma ghrelin level in underweight patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2004;170:879–82.
80. Garcia J, Garcia-Touza M, Hijazi R. Active ghrelin levels and active to total ghrelin ratio in cancer-induced cachexia. *J Clin Endocrinol Metab* 2005;90:2920–6.
81. Shimizu Y, Nagaya N, Isobe T, et al. Increased plasma ghrelin level in lung cancer cachexia. *Clin Cancer Res* 2003;9:774–8.
82. Kerem M, Ferahkose Z, Yilmaz UT, et al. Adipokines and ghrelin in gastric cancer cachexia. *World J Gastroenterol* 2008;14:3633–41.
83. Akamizu T, Kangawa K. Ghrelin for cachexia. *J Cachex Sarcopenia Muscle* 2010;1:169–76.
84. Garcia J, Boccia R, Graham C. A phase II randomized, placebo-controlled, double-blind study of the efficacy and safety of RC-1291 (RC) for the treatment of cancer cachexia. *J Clin Oncol* 2007;25:9133.
85. Neary NM, Small CJ, Wren AM, et al. Ghrelin increases energy intake in cancer patients with impaired appetite: acute, randomized, placebo-controlled trial. *J Clin Endocrinol Metab* 2004;89:2832–6.
86. Strasser F, Lutz TA, Maeder MT, et al. Safety, tolerability and pharmacokinetics of intravenous ghrelin for cancer-related anorexia/cachexia: a randomised, placebo-controlled, double-blind, double-cross-over study. *Br J Cancer* 2008;98:300–8.
87. Fujitsuka N, Asakawa A, Uezono Y, et al. Nature Publishing Group. Potentiation of ghrelin signaling attenuates cancer anorexia-cachexia and prolongs survival. *Translational Psychiatry* 2011;1:e23-DOI. 10.1038/tp.2011.25.
88. Laferrère B, Abraham C, Russell CD, Bowers CY. Growth hormone releasing peptide-2 (GHRP-2), like ghrelin, increases food intake in healthy men. *J Clin Endocrinol Metab* 2005;90:611–4.
89. Takeda H, Sadakane C, Hattori T, et al. Rikkunshito, an herbal medicine, suppresses cisplatin-induced anorexia in rats via 5-HT2 receptor antagonism. *Gastroenterology* 2008;134:2004–13.
90. Fujitsuka N, Asakawa A, Hayashi M, et al. Selective serotonin reuptake inhibitors modify physiological gastrointestinal motor activities via 5-HT2c receptor and acyl ghrelin. *Biological Psychiatry* 2009;65:748–59.
91. Halford J, Harrold J, Boyland E, Lawton C. Serotonergic drugs: effects on appetite expression and use for the treatment of obesity. *Drugs* 2007;67:27–55.
92. Vickers SP, Dourish CT, Kennett GA. Evidence that hypophagia induced by d-fenfluramine and d-norfenfluramine in the rat is mediated by 5-HT2C receptors. *Neuropharmacology* 2001;41:200–9.
93. Vickers SP, Easton N, Webster LJ, et al. Oral administration of the 5-HT2C receptor agonist, mCPP, reduces body weight gain in rats over 28 days as a result of maintained hypophagia. *Psychopharmacology (Berl)* 2003;167:274–80.
94. Wang W, Danielsson A, Svanberg E, Lundholm K. Lack of effects by tricyclic antidepressant and serotonin inhibitors on anorexia in MCG 101 tumor-bearing mice with eicosanoid-related cachexia. *Nutrition* 2003;19:47–53.
95. Makarenko IG, Meguid MM, Gatto L, et al. Hypothalamic 5-HT1B-receptor changes in anorectic tumor bearing rats. *Neurosci Lett* 2005;376:71–5.
96. Makarenko IG, Meguid MM, Gatto L, Chen C, Ugrumov MV. Decreased NPY innervation of the hypothalamic nuclei in rats with cancer anorexia. *Brain Res* 2003;961:100–8.
97. Sakurada S, Shido O, Sugimoto N, Hiratsuka Y, Yoda T, Kanosue K. Autonomic and behavioural thermoregulation in starved rats. *J Physiol (Lond.)* 2000;526:417–24.
98. Zhao Z, Sakata I, Okubo Y, Koike K, Kangawa K, Sakai T. Gastric leptin, but not estrogen and somatostatin, contributes to the elevation of ghrelin mRNA expression level in fasted rats. *J Endocrinol* 2008;196:529–38.
99. Turrin N, Ilyin S, Gayle D. Interleukin-1beta system in anorectic catabolic tumor-bearing rats. *Curr Opin Clin Metab Care* 2004;7:419–26.

100. Gyengesi E, Liu Z-W, D'Agostino G, et al. Corticosterone regulates synaptic input organization of POMC and NPY/AgRP neurons in adult mice. *Endocrinology* 2010;151:5395–402.
101. Chen C-Y, Fujimiya M, Laviano A, Chang F-Y, Lin H-C, Lee S-D. Modulation of ingestive behavior and gastrointestinal motility by ghrelin in diabetic animals and humans. *J Chin Med Assoc* 2010;73:225–9.
102. Shintani F, Kanba S, Nakaki T. Interleukin-1 beta augments release of norepinephrine, dopamine, and serotonin in the rat anterior hypothalamus. *J Neurosci* 1993;13:3574–81.
103. Chuang J. Ghrelin's roles in stress, mood, and anxiety regulation. *Int J Pept* 2010;2010:1–5.
104. Heisler LK, Pronchuk N, Nonogaki K, et al. Serotonin activates the hypothalamic–pituitary–adrenal axis via serotonin 2C receptor stimulation. *J Neurosci* 2007;27:6956–64.
105. Date Y, Murakami N, Toshinai K, et al. The role of the gastric afferent vagal nerve in ghrelin-induced feeding and growth hormone secretion in rats. *Gastroenterology* 2002;123:1120–8.
106. Fujino K, Inui A, Asakawa A, Kihara N, Fujimura M, Fujimiya M. Ghrelin induces fasted motor activity of the gastrointestinal tract in conscious fed rats. *J Physiol (Lond.)* 2003;550:227–40.
107. Nakata M, Maekawa F, Fujiwara K, Maejima Y. Leptin suppresses ghrelin-induced activation of neuropeptide Y neurons in the arcuate nucleus via phosphatidylinositol 3-kinase-and phosphodiesterase 3-mediated pathway. *Endocrinology* 2007;148:2251–63.
108. Muroya S, Yada T, Shioda S, Takigawa M. Glucose-sensitive neurons in the rat arcuate nucleus contain neuropeptide. *Y Neurosci Lett* 1999;264:113–6.
109. Takaya K, Ariyasu H, Kanamoto N, et al. Ghrelin strongly stimulates growth hormone release in humans. *J Clin Endocrinol Metab* 2000;85:4908–11.
110. Arvat E, Maccario M, Di Vito L, et al. Endocrine activities of ghrelin, a natural growth hormone secretagogue (GHS), in humans: comparison and interactions with hexarelin, a nonnatural peptidyl GHS, and GH-releasing hormone. *J Clin Endocrinol Metab* 2001;86:1169–74.
111. Hataya Y, Akamizu T, Takaya K, et al. A low dose of ghrelin stimulates growth hormone (GH) release synergistically with GH-releasing hormone in humans. *J Clin Endocrinol Metab* 2001;86:4552.
112. Akamizu T, Takaya K, Irako T, et al. Pharmacokinetics, safety, and endocrine and appetite effects of ghrelin administration in young healthy subjects. *Eur J Endocrinol* 2004;150:447–55.
113. Gibney J, Healy ML, Sonksen PH. The growth hormone/insulin-like growth factor-I axis in exercise and sport. *Endocr Rev* 2007;28:603–24.
114. Velloso CP. Regulation of muscle mass by growth hormone and IGF-I. *Br J Pharmacol* 2008;154:557–68.
115. Gonzalez-Rey E, Chorny A, Delgado M. Therapeutic action of ghrelin in a mouse model of colitis. *Gastroenterology* 2006;130:1707–20.
116. Waseem T, Duxbury M, Ito H, Ashley SW, Robinson MK. Exogenous ghrelin modulates release of pro-inflammatory and anti-inflammatory cytokines in LPS-stimulated macrophages through distinct signaling pathways. *Surgery* 2008;143:334–42.
117. Dixit VD, Schaffer EM, Pyle RS, et al. Ghrelin inhibits leptin- and activation-induced proinflammatory cytokine expression by human monocytes and T cells. *J Clin Invest* 2004;114:57–66.
118. Wu R, Dong W, Zhou M, et al. Ghrelin attenuates sepsis-induced acute lung injury and mortality in rats. *Am J Respir Crit Care Med* 2007;176:805–13.
119. Theil M-M, Miyake S, Mizuno M, et al. Suppression of experimental autoimmune encephalomyelitis by ghrelin. *J Immunol* 2009;183:2859–66.
120. Sadakane C, Muto S, Nakagawa K, et al. 10-Gingerol, a component of rikkunshito, improves cisplatin-induced anorexia by inhibiting acylated ghrelin degradation. *Biochem Biophys Res Commun* 2011;412:506–11.
121. Yakabi K, Sadakane C, Noguchi M, et al. Reduced ghrelin secretion in the hypothalamus of rats due to cisplatin-induced anorexia. *Endocrinology* 2010;151:3773–82. <http://www.ncbi.nlm.nih.gov/pubmed/20534732>.
122. Barber MD, Ross JA, Fearon KC. Cancer cachexia. *Surg Oncol* 1999;8:133–41.
123. Huang C, Lin S, Liao P, Young S. The immunopharmaceutical effects and mechanisms of herb medicine. *Cell Mol Immunol* 2008;5:23–31.

Cancer cachexia—pathophysiology and management

Hajime Suzuki · Akihiro Asakawa ·
Haruka Amitani · Norifumi Nakamura ·
Akio Inui

Received: 15 February 2013 / Accepted: 20 February 2013 / Published online: 20 March 2013
© Springer Japan 2013

Abstract About half of all cancer patients show a syndrome of cachexia, characterized by anorexia and loss of adipose tissue and skeletal muscle mass. Cachexia can have a profound impact on quality of life, symptom burden, and a patient's sense of dignity. It is a very serious complication, as weight loss during cancer treatment is associated with more chemotherapy-related side effects, fewer completed cycles of chemotherapy, and decreased survival rates. Numerous cytokines have been postulated to play a role in the etiology of cancer cachexia. Cytokines can elicit effects that mimic leptin signaling and suppress orexigenic ghrelin and neuropeptide Y (NPY) signaling, inducing sustained anorexia and cachexia not accompanied by the usual compensatory response. Furthermore, cytokines have been implicated in the induction of cancer-related muscle wasting. Cytokine-induced skeletal muscle wasting is probably a multifactorial process, which involves a protein synthesis inhibition, an increase in protein degradation, or a combination of both. The best treatment of the cachectic syndrome is a multifactorial approach. Many drugs including appetite stimulants, thalidomide, cytokine inhibitors, steroids, nonsteroidal anti-inflammatory drugs, branched-chain amino acids, eicosapentaenoic acid, and antiserotonergic drugs have been proposed and used in

clinical trials, while others are still under investigation using experimental animals. There is a growing awareness of the positive impact of supportive care measures and development of promising novel pharmaceutical agents for cachexia. While there has been great progress in understanding the underlying biological mechanisms of cachexia, health care providers must also recognize the psychosocial and biomedical impact cachexia can have.

Keywords Cachexia · Anorexia · Cytokine · Skeletal muscle · Palliative care

Pathophysiology

Hormones and mediators

Leptin is a protein hormone that sends afferent signals from the periphery to the brain that regulates adipose tissue mass [1–3]. The level of leptin is positively correlated with body fat mass, and dynamic changes in plasma leptin concentrations in either direction can activate the efferent energy regulation pathways [1, 4]. Leptin reduces appetite and increases energy expenditure and evidently elicits these effects via the central nervous system [1, 4]. This is achieved by hypothalamic neuropeptides downstream of leptin that regulate food intake and energy expenditure. Starvation or a loss of body fat can lead to a decrease in leptin, which in turn leads to a state of positive energy balance; conversely, food intake exceeds energy expenditure. This compensatory response is mediated by the increased production of ghrelin, neuropeptide Y (NPY), and other appetite-stimulating neuropeptides, and decreased activity of anorexigenic neuropeptides such as corticotropin-releasing factor (CRF) and melanocortin (Fig. 1a).

H. Suzuki · A. Asakawa · H. Amitani · A. Inui (✉)
Department of Psychosomatic Internal Medicine,
Kagoshima University Graduate School of Medical
and Dental Sciences, 8-35-1 Sakuragaoka,
Kagoshima 890-8520, Japan
e-mail: inui@m.kufm.kagoshima-u.ac.jp

H. Suzuki · N. Nakamura
Department of Oral and Maxillofacial Surgery,
Kagoshima University Graduate School of Medical
and Dental Sciences, Kagoshima 890-8520, Japan

Thus, if a disease process such as cancer was to produce factors that induce or mimic the hypothalamic effect of excess negative feedback signaling from leptin, the expected outcome would be sustained anorexia (lack of appetite) and cachexia (muscle wasting and uncontrolled weight loss), without the usual compensatory response [5]. In fact, in tumor-bearing states, cachectic factors such as cytokines can elicit effects on energy homeostasis that mimic leptin and suppress orexigenic ghrelin and NPY signaling. Consequently, the increases and decreases in hypothalamic actions caused by these mediators induce anorexia and unopposed weight loss (Fig. 1b).

Serotonin (5-HT) may also play a role in the development of cancer-induced anorexia. This is because increased levels of plasma and brain tryptophan, the precursor of 5-HT, and interleukin (IL)-1 may underlie the increased serotonergic activity seen in the cancer cachexia. In addition, cisplatin-induced anorexia has become problematic in clinical settings. Cisplatin is a widely used and effective anti-cancer chemotherapy drug, however, the undesirable gastrointestinal side effects associated with it, such as nausea, vomiting, and anorexia, markedly decrease patients' quality of life, rendering continuation of chemotherapy difficult [6]. Cisplatin-induced gastrointestinal tract




Fig. 1 A simplified model of the hypothalamic neuro peptide circuitry in response to starvation (a) and cancer cachexia (b). Full line arrows indicate the activation of the process, and broken line arrows indicate the inhibition of the process. Under normal conditions, energy intake is determined by the hypothalamic integration of peripheral signals conveying inputs on adiposity status, digestive processes, and metabolic profile. Some of these signals such as adipocyte-derived leptin inhibit energy intake, while other signals such as stomach-derived ghrelin stimulate energy intake. In the hypothalamus, the arcuate nucleus (ARC) receives information from the periphery and integrates these inputs to modulate food intake via second-order neurons. According to the information conveyed to the brain, peripheral signals may differentially activate or inhibit POMC/CART and NPY/AgRP neurons. When an energy deficit (e.g., starvation) is signaled, orexigenic NPY/AgRP neurons are activated and anorexigenic POMC/CART neurons are inhibited, resulting in increased energy intake. When an energy excess is signaled, NPY/

AgRP neurons are inhibited and POMC/CART neurons are activated. During cancer, cachectic factors such as cytokines elicit effects on energy homeostasis that mimic leptin in some respects and suppress orexigenic Ghrelin-NPY/AgRP signaling. Increased brain cytokine expression disrupts hypothalamic neurochemistry, particularly in the ARC where cytokines activate POMC/CART neurons, while inactivate NPY/AgRP neurons. The anorexia and unopposed weight loss in cachexia could be accomplished through persistent inhibition of the NPY orexigenic network and stimulation of anorexigenic neuropeptides, although the hypothalamic pathways participating in this response remain to be determined. *AgRP* Agouti-related peptide, *MCH* melanin-concentrating hormone, *CART* cocaine- and amphetamine-related transcript, *NPY* neuropeptide Y, *POMC* pro-opiomelanocortin, *CRH* corticotropin-releasing hormone, *MC4R* melanocortin-4 receptor, *PVN* paraventricular nucleus. *LHA* lateral hypothalamic area. Source: (5) with modification