

- [33] Masuda, E., and Maeda, H. (1997) Host-mediated antitumor activity induced by neocarzinostatin and its polymer-conjugated derivative in tumor-bearing mice. In *Neocarzinostatin: The Past, Present, and Future of an Anticancer Drug*. (eds. Maeda, H., Edo, K., and Ishida, N.) pp 187-204, Springer, Tokyo.
- [34] Oda, T., Morinaga, T., and Maeda, H. (1986) Stimulation of macrophage by polyanions and its conjugated proteins and effect on cell membrane. *Proc. Soc. Exp. Biol. Med.* 181, 9-17.
- [35] Kobayashi, A., Oda, T., and Maeda, H. (1988) Protein binding of macromolecular anticancer agent SMANCS: Characterization of poly(styrene-co-maleic acid) derivatives as an albumin binding ligand. *J. Bioactive Compatible Polymers*, 3, 319-333.
- [36] Maeda, H. (2001) SMANCS and polymer-conjugated macromolecular drugs: Advantages in cancer chemotherapy. *Adv. Drug Deliv. Rev.* 46, 169-185.
- [37] Iwai, K., Maeda, H., and Konno, T. (1984) Use of oily contrast medium for selective drug targeting to tumor: Enhanced therapeutic effect and X-ray image. *Cancer Res.* 44, 2115-2121.
- [38] Konno, T., Maeda, H., Yokoyama, I. et al (1982) Use of a lipid lymphographic agent, lipiodol, as a carrier of high molecular weight antitumor agent, SMANCS, for hepatocellular carcinoma. *Cancer and Chemotherapy* 9, 2005-2015 (in Japanese).
- [39] Konno, T., Maeda, H., Iwai, K., et al (1983) Effect of arterial administration of high-molecular-weight anticancer agent SMANCS with lipid lymphographic agent on hepatoma: A preliminary report. *Eur. J. Cancer Clin. Oncol.* 19, 1053-1065.
- [40] Konno, T., Maeda, H., Iwai, K., et al (1984) Selective targeting of anti-cancer drug and simultaneous image enhancement in solid tumors by arterially administered lipid contrast medium. *Cancer* 54, 2367-2374.
- [41] Maki, S., Konno, T., and Maeda, H. (1985) Image enhancement in computerized tomography for sensitive diagnosis of liver cancer and semiquantitation of tumor selective drug targeting with oily contrast medium. *Cancer* 56, 751-757.
- [42] Matsumoto, K., Yamamoto, T., Kamata, R., and Maeda, H. (1984) Pathogenesis of serratial infection: Activation of the Hageman factor-prekallikrein cascade by serratial protease. *J. Biochem.* 96, 739-749.
- [43] Kamata, R., Yamamoto, T., Matsumoto, K., and Maeda, H. (1985) A serratial protease causes vascular permeability reaction by activation

- of the Hageman factor-dependent pathway in guinea pigs. *Infect. Immun.* 48, 747-753.
- [44] Molla, A., Yamamoto, T., Akaike, T., Miyoshi, S., and Maeda, H. (1989) Activation of Hageman factor and prekallikrein and generation of kinin by various microbial proteinases. *J. Biol. Chem.* 264, 10589-10594.
- [45] Maeda, H. (1996) Role of microbial proteases in pathogenesis. *Microbiol. Immunol.* 40, 685-699.
- [46] Maeda, H. (2002) Microbial proteinases and pathogenesis of infection. *Wiley Encyclopedia of Molecular Medicine* (Creighton, T. E., Ed.) pp 2663-2668, Volume 4, John Wiley & Sons, New York.
- [47] Akaike, T., Molla, A., Ando, M., Araki, S., and Maeda, H. (1989) Molecular mechanism of complex infection by bacteria and virus analyzed by a model using serratia protease and influenza virus in mice. *J. Virol.* 63, 2252-2259.
- [48] Maruo, K., Akaike, T., Inada, Y., Ohkubo, I., Ono, T., and Maeda, H. (1993) Effect of microbial and mite proteases on low and high molecular weight kininogens. *J. Biol. Chem.* 268, 17711-17715.
- [49] Akaike, T., Maeda, H., Maruo, K., Sakata, Y., and Sato, K. (1994) Potentiation of infectivity and pathogenesis of influenza A virus by a house dust mite protease. *J. Infect. Dis.* 170, 1023-1026.
- [50] Frizz, H., and Travis, J. (2004) Hiroshi Maeda—40 years of research. Reflections on the occasion of his retirement and 65<sup>th</sup> birthday. *Biol. Chem.* 385, 987-988.
- [51] Oda, T., Akaike, T., Hamamoto, T., Suzuki, F., Hirano, T., and Maeda, H. (1989) Oxygen radicals in influenza-induced pathogenesis and treatment with pyran polymer-conjugated SOD. *Science* 244, 974-976.
- [52] Akaike, T., Ando, M., Oda *et al* (1990) Dependence on  $O_2^-$  generation by xanthine oxidase of pathogenesis of influenza virus infection in mice. *J. Clin. Invest.* 85, 739-745.
- [53] Maeda, H., and Akaike, T. (1991) Oxygen free radicals as pathogenic molecules in viral diseases. *Proc. Soc. Exp. Biol. Med.* 198, 721-727.
- [54] Maeda, H. (2000) Paradigm shift in microbial pathogenesis: An alternative to the Koch-Pasteur paradigm in the new millennium. Abstr. for the 13<sup>th</sup> International Congress of The International Organization for Mycoplasmology, Fukuoka, Japan, 35.
- [55] Akaike, T., Noguchi, Y., Ijiri, S. *et al* (1996) Pathogenesis of influenza virus-induced pneumonia: Involvement of both nitric oxide and oxygen radicals. *Proc. Natl. Acad. Sci. U.S.A.* 93, 2448-2453.

- [56] Akaike, T., Fujii, S., Kato, A. *et al* (2000) Viral mutation accelerated by nitric oxide production during infection *in vivo*. *FASEB J.* 14, 1447-1454.
- [57] Sawa, T., Akaike, T., Ichimori, K. *et al* (2003) Superoxide generation mediated by 8-nitroguanosine, a highly redox-active nucleic acid derivative. *Biochem. Biophys. Res. Comm.*, 311, 300-306.
- [58] Maeda, H. and Akaike, T. (1998) Nitric oxide and oxygen radicals in infection, inflammation, and cancer. *Biochemistry (Moscow)*, 63, (No.7) 1007-1017.
- [59] Kuwahara, H., Kariu, T., Fan, J., and Maeda, H. (2009) Generation of drug-resistant mutants of *Helicobacter pylori* in the presence of peroxynitrite, a derivative of nitric oxide, at pathophysiological concentration. *Microbiol. Immunol.* 52, 1-7.
- [60] Kuwahara, H., Kanazawa, A., Wakamatsu, D., Morimura, S., Kida, K., Akaike, T., and Maeda, H. (2004) Antioxidative and antimutagenic activities of 4-vinyl-2, 6-dimethoxyphenol (canolol) isolated from canola oil. *J. Agric. Fd. Chem.*, 52, 4380-4387
- [61] Cao, X., Tsukamoto, T., Seki, T. *et al* (2008) 4-Vinyl-2,6-dimethoxyphenol (canolol) suppresses oxidative stress and gastric carcinogenesis in *Helicobacter pylori*-infected carcinogen-treated Mongolian gerbils. *Int. J. Cancer* 122, 1445-1454.
- [62] J. Yoshitake, T. Akaike, T. Akuta, F. Tamura, T. Ogura, H. Esumi and H. Maeda (2004) Nitric oxide as an endogenous mutagen for Sendai virus without antiviral activity. *J. Virol.*, 78, 8709-8719.
- [63] Sawa, T., Zaki, M., H., Okamoto, T. *et al* (2007) Protein S-guanylation by the biological signal 8-nitroguanosine 3',5'-cyclic monophosphate. *Nature Chem. Biol.* 3, 727-735.
- [64] Maeda, H., Matsumura, Y., and Kato, H. (1988) Purification and identification of [hydroxyprolyl<sup>3</sup>]bradykinin in ascetic fluid from a patient with gastric cancer. *J. Biol. Chem.* 263, 16051-16054.
- [65] Matsumura, Y., and Maeda, H. (1986) A new concept for macromolecular therapeutics in cancer chemotherapy: Mechanism of tumoritropic accumulation of proteins and the antitumor agent smancs. *Cancer Res.* 46, 6387-6392.
- [66] Nagamitsu, A., Greish, K., and Maeda, H. (2009) Elevating blood pressure as a strategy to increase tumor targeted delivery of macromolecular drug SMANCS: Cases of advanced solid tumors. *Jpn. J. Clin. Oncol.*, 39, 756-766.
- [67] Li, C. J., Miyamoto, Y., Kojima, Y., and Maeda, H. (1993) Augmentation of tumor delivery of macromolecular drugs with

- reduced bone marrow delivery by elevating blood pressure. *Br. J. Cancer* 67, 975-980.
- [68] Noguchi, Y., Wu, J., Duncan, R., Strohalm J., Ulbrich K., Akaike T., Maeda H. (1998) Early phase tumor accumulation of macromolecules: A great difference in clearance rate between tumor and normal tissues. *Jpn. J. Cancer Res.* 89, 307-314.
- [69] Seymour, L. W., Miyamoto, Y., Maeda, H., Brereton, M., Strohalm, J., Ulbrich, K., and Duncan, R. (1995) Influence of molecular weight on passive tumour accumulation of a soluble macromolecular drug carrier. *Eur. J. Cancer* 31, 766-770.
- [70] Maeda, H., Seymour, L. W., and Miyamoto, Y. (1992) Conjugates of anticancer agents and polymers: Advantages of macromolecular therapeutics in vivo. *Bioconj. Chem.* 3, 351-362.
- [71] Maeda, H., Wu, J., Sawa, T., Matsumura, Y., and Hori, K. (2000) Tumor vascular permeability and the EPR effect in macromolecular therapeutics. *J. Control. Release* 65, 271-284.
- [72] Skinner, S. A., Tutton, P. J. M., and O'Brien, P. E. (1990) Microvascular architecture of experimental colon tumors in the rat. *Cancer Res.* 50, 2411-2417.
- [73] Matsumura, Y., Kimura, M., Yamamoto, T., Maeda, H. (1988) Involvement of the kinin-generating cascade in enhanced vascular permeability in tumor tissue. *Jpn. J. Cancer Res.* 79, 1327-1334.
- [74] Matsumura, Y., Maruo, K., Kimura, M., Yamamoto, T., Konno, T., and Maeda, H. (1991) Kinin-generating cascade in advanced cancer patients and in vitro study. *Jpn. J. Cancer Res.* 82, 732-741.
- [75] Akaike, T., Yoshida, M., Miyamoto, Y., Sato, K., Kohno, M., Sasamoto, K., Miyazaki, K., Ueda, S., and Maeda, H. (1993) Antagonistic action of imidazolineoxyl N-oxides against endothelium-derived relaxing factor/NO through a radical reaction. *Biochemistry*, 32, 827-832.
- [76] Maeda, H., Noguchi, Y., Sato, K., Akaike, T. (1994) Enhanced vascular permeability in solid tumor is mediated by nitric oxide and inhibited by both new nitric oxide scavenger and nitric oxide synthase inhibitor. *Jpn. J. Cancer Res.* 85, 331-334.
- [77] Maeda, H., Wu, J., Okamoto, T., Maruo, K., and Akaike, T. (1999) Kallikrein-kinin in infection and cancer. *Immunopharmacology*, 43, 115-128.
- [78] Wu, J., Akaike, T., and Maeda, H. (1998) Modulation of enhanced vascular permeability in tumors by a bradykinin antagonist, a

- cyclooxygenase inhibitor, and a nitric oxide scavenger. *Cancer Res.* 58, 159-165.
- [79] Wu, J., Akaike, T., Hayashida, K., Okamoto, T., Okuyama, A., and Maeda, H. (2001) Enhanced vascular permeability in solid tumor involving peroxynitrite and matrix metalloproteinase. *Jpn. J. Cancer Res.* 92, 439-451.
- [80] Senger, D. R., Galli, S. J., Dvorak, A. M., Perruzzi, C. A., Harvey, V. S., Dvorak, H. F. (1983) Tumor cells secrete a vascular permeability factor that promotes accumulation of ascites fluid. *Science*, 219, 983-985.
- [81] Kimura N, Taniguchi S, Aoki K, Baba T. (1980) Selective localization and growth of *Bifidobacterium bifidum* in mouse tumors following intravenous administration. *Cancer Res.* 40. 2060-2068.
- [82] Zhao, M., Yang, M., Ma, H., Li, X., Tan X, Li S, Yang Z and Hoffman R M. (2006) Targeted therapy with a *salmonella* typhimurium leucine-arginine auxotroph cures orthotopic human breast tumors in nude mice. *Cancer Res.* 66, 7647-7652.
- [83] Maeda, H., Sawa, T. and Konno, T. (2001) Mechanism of tumor-targeted delivery of macromolecular drugs, including the EPR effect in solid tumor and clinical overview of the prototype polymeric drug SMANCS. *J. Cont. Release*, 74, 47-61.
- [84] Maeda, H., Bharate, G. Y., Daruwalla, J. (2009) Polymeric drugs and nanomedicines for efficient tumor targeted drug delivery based on EPR-effect. *Eur. J. Pharm. Biopharm.* 71, 409-419.
- [85] Seki, T., Fang, J., Maeda, H. (2009) Tumor targeted macromolecular drug delivery based on the enhanced permeability and retention effect in solid tumor. In *Pharm. Perspective of Cancers Therapeutics* (eds. Y. Lu and R.I. Mahato), pp 94-120, AAPS Springer Press, New York.
- [86] Iyer, A.K., Greish, K., Fang, J., Murakami, R. and Maeda, H. (2007) High-loading nanosized micelles of copoly (styrene-maleic acid) – zinc protoporphyrin for targeted delivery of a potent heme oxygenase inhibitor. *Biomaterials*, 28, 1871-1881.
- [87] Greish, K., Fang, J., Inuzuka, T., Nagamitsu, A. and Maeda, H. (2003) Macromolecular anticancer therapeutics for effective solid tumor targeting: Advantages and prospects. *Clin. Pharmacokinetics*, 42, 1089-1105.
- [88] Maeda, H. (2001) The enhanced permeability and retention (EPR) effect in tumor vasculature: The key role of tumor-selective macromolecular drug targeting. In *Advances in Enzyme Regulation* (eds. Weber, G.), 41, pp.189-207, Oxford, UK, Elsevier Science Ltd.

- [89] Hori, K., Suzuki, M., Tanda, S., Saito, D., Shinozaki, M., and Zhang, Q. H. (1991) Fluctuations in tumor blood flow under normotension and the effect of angiotensin II-induced hypertension. *Jpn. J. Cancer Res.* 82, 1309-1316.
- [90] Suzuki, M., Hori, K., Abe I., Saito, S., and Saito, H. (1981) A new approach to cancer chemotherapy: a selective enhancement of tumor blood flow with angiotensin II. *J. Natl. Cancer Inst.* 67, 663-669.
- [91] Suzuki, M., Hori, K., Abe, I., Saito, S., Satoh, H. (1984) Functional characterization of the microcirculation in tumor. *Cancer Metastasis Review.* 3, 115-126.
- [92] Maeda, H., Seki, T., Fang, J. (2008) Enhanced tumor delivery of macromolecular antitumor-drugs by topical application of nitroglycerin on superficial tumors. Abst. 67<sup>th</sup> Annual Meeting of the Jpn. Cancer Assoc. No. P-518, p.213.
- [93] Seki, T., Fang, J., and Maeda, H. (2009) Enhanced delivery of macromolecular antitumor drugs to tumors by nitroglycerin application. *Cancer Sci.* 100, 2426-2430.
- [94] Yasuda, H., Yamaya, M., Nakayama, K. et al. (2006) Randomized phase II trial comparing nitroglycerin plus vinorelbine and cisplatin with vinorelbine and cisplatin alone in previously untreated stage IIIB/IV non-small cell lung cancer, *J. Clin. Oncol.* 24, 688-694.
- [95] Yasuda, H., Nakayama, K., Watanabe, M. et al. (2006) Nitroglycerin treatment may increase response to docetaxel and carboplatin regimen via inhibitions of hypoxia-inducible factor-1 pathway and P-glycoprotein in patients with lung adenocarcinoma, *Clin. Cancer Res.* 12, 6748-6757.
- [96] Yasuda, H., Yanagihara, K., Nakayama, K. et al. (2009) Therapeutic applications of nitric oxide for malignant tumor in animal models and human studies, in: B. Bonavida (Ed.), Nitric Oxide and Cancer, Springer Science, New York.
- [97] Jordan, B.F., Misson, P.D., Demeure, R. et al. (2000) Changes in tumor oxygenation/perfusion induced by the NO donor, isosorbide dinitrate, in comparison with carbogen: monitoring by EPR and MRI, *Int. J. Radiat. Oncol. Biol. Phys.* 48, 565-570.
- [98] J.B. Mitchell, D.A. Wink, W. DeGraff, et al., Hypoxic mammalian cell radiosensitization by nitric oxide, *Cancer Res.* 53 (1993) 5845-5848.

- [99] Iyer, A.K., Khaled, G., Fang, J. and Maeda, H. (2006) Exploiting the enhanced permeability and retention effect for tumor targeting. *Drug Discovery Today*, 11, 812-818.
- [100] Nagamitsu, A., Inuzuka, T., Kawasuji, M., Maeda, H. (2003) Recent advances in SMANCS/Lipiodol therapy – enhanced targeting and delivery effect using vascular nodulator, *Drug Delivery System* (in Japanese) 18, 438-447.
- [101] Nagamitsu, A., Khaled, G., Maeda, H. (2010) Therapy of unresectable lung cancer by SMANCS/Lipiodol in combination with isosorbite dinitrate (ISDN) given arterially. Proceeding 69<sup>th</sup> Ann. Meeting of Jpn. Cancer Assoc. p.102, No.206. Sept. 22-24.
- [102] Greish, K., Nagamitsu, A., Fang, F., and Maeda, H. (2005) Copoly (styrene-maleic acid)-pirarubicin micelles: High tumor-targeting efficacy with low toxicity. *Bioconj. Chem.* 16, 230-236.
- [103] Greish, K., Sawa, T., Fang, J., Akaike, T., and Maeda, H. (2004) SMA- doxorubicin, a new polymeric micellar drug for effective anticancer targeting. *J. Controlled Release* 97, 219-230.
- [104] Iyer, A. K., Greish, K., Seki, T., Okazaki, S., Fang, J., Takeshita, K., and Maeda, H. (2007) Polymeric micelles of zinc protoporphyrin for tumor targeted delivery based on EPR effect and singlet oxygen generation. *J. Drug Target.* 15, 496-506.
- [105] Daruwalla, J., Greish, K., Wilson, C. M., Muralidharan, V., Iyer, A., Maeda, H., Christophi, C. (2009) Styrene maleic acid-pirarubicin disrupts tumor microcirculation and enhances the permeability of colorectal Liver Metastases, *J. Vascu. Res.*, 46, 218-228.
- [106] Daruwalla, J., Nikfarjam, M., Greish, K. Malcontenti-WilsonI, C., Muralidharan, Christophi, V. C. and Maeda, H. (2010) In vitro and in vivo evaluation of tumor targeting SMA-pirarubicin micelles: Survival improvement and inhibition of liver metastases. *Cancer Science*, 101, 1866-1874.
- [107] Sahoo, S. K., Sawa, T., Fang, J., Tanaka, S., Miyamoto, Y., Akaike, T., and Maeda, H. (2002) Pegylated zinc protoporphyrin: A water-soluble heme oxygenase inhibitor with tumor-targeting capacity. *Bioconj. Chem.* 13, 1031-1038.
- [108] Fang, J., Sawa, T., Akaike, T., Akuta, T., Sahoo, S. K., Griesh, K., Hamada, A., Maeda, H. (2003) In vivo antitumor activity of pegylated zinc protoporphyrin: Targeted inhibition of heme oxygenase in solid tumor. *Cancer Res.* 63, 3567-3574.
- [109] Doi, K., Akaike, T., Fujii, S., Tanaka, S., Ikebe, N., Beppu, T., Shibahara, S., Ogawa, M. and Maeda, H. (1999) Induction of haem

- oxygenase-1 by nitric oxide and ischaemia in experimental solid tumours and implications for tumour growth. *Br. J. Cancer*, 80, 1945-1954.
- [110] Fang, J., Sawa,T., Akaike,T., Greish, K. and Maeda, H. (2004) Enhanced chemotherapeutic effect against tumor cells by a heme oxygenase inhibitor, pegylated zinc protoporphyrin in combination with anticancer agents. *Int. J. Cancer* 109, 1-8.
- [111] Fang, J., Akaike, T., and Maeda, H. (2004) Antiapoptotic role of heme oxygenase and its potential as an anticancer target. *Apoptosis*, 9, 27-35.
- [112] Regehly, M., Greish, K., Rancan, F., Maeda, H., Böhm, F. and Röder, B. (2007) Water-soluble polymer conjugated of ZnPP for photodynamic tumour therapy. *Bioconj. Chem.*, 18, 494-499.
- [113] Mayerhofer, M., Gleixner, K. V., Hörmann, G., Sonneck, K., Aichberger, K. J., Ott, R. G., Greish, K., Nakamura, H., Derdak, S., Samorapoompitchit, P., Pickl, W. F., Sexl, V., Esterbauer, H., Sillaber, C., Maeda, H., and Valent, P. (2008) Targeting of heat shock protein 32 (Hsp32)/heme oxygenase-1 (HO-1) in leukemic cells in chronic myeloid leukemia: A novel approach to overcome resistance against imatinib. *Blood* 111, 2200-2210.
- [114] Hadzijusufovic, E., Rebuzzi, L., Gleixner, K. V., Ferenc, V., Kondo, R., Gruze, A., Kneidinger, M., Krauth, M.-T., Mayerhofer, M., Samorapoompitchit, P., Greish, K., Pickl, W. F., Maeda, H., Willmann, M., and Valent, P. (2008) Targeting of heat-shock protein 32/heme oxygenase-1 in canine mastocytoma cells is associated with reduced growth and induction of apoptosis. *Exp. Hematol.* 36, 1461-1470.
- [115] Nakamura, H., Fang, J., Bharate, G., Tsukigawa, K., Maeda, H. Intracellular uptake and behavior of two types zinc protoporphyrin (ZnPP) micelles, SMA-ZnPP and PEG-ZnPP as anticancer agents; Unique intracellular disintegration of SMA micelles (in press, 2011).
- [116] Maeda, H., Yamamoto, N. and Yamashita, A. (1976) Fate and distribution of [<sup>14</sup>C] succinyl neocarzinostatin in rats. *Europ. J. Cancer*, 12, 865-870.
- [117] Maeda, H., Kimura, M., Sasaki, I., Hirose, Y. and Konno, T. (1992) Toxicity of bilirubin and detoxification by PEG-bilirubin oxidase conjugate: A new tactic for treatment of jaundice. In *Poly (Ethylene Glycol) Chemistry: Biotechnical and Biomedical Applications* (ed. J. M. Harris) pp.153-169, Plenum Press, New York.
- [118] Kimura, M., Matsumura, Y., Konno, T., Miyauchi, Y. and Maeda, H. (1990) Enzymatic removal of bilirubin toxicity by bilirubin oxidase in

- vitro* and excretion of degradation products *in vivo*. *Proc. Soc. Exp. Biol. Med.*, 195, 64-69.
- [119] Maeda, H. and Konno, T. (1997) Metamorphosis of neocarzinostatin to SMANCS –Chemistry, pharmacology and clinical effect of the first prototype anticancer polymer therapeutic– In *Neocarzinostatin: The Past, Present, and Future of an Anticancer Drug*. (eds. H. Maeda, K. Edo and N. Ishida) pp. 227-267, Springer-Verlag, Tokyo.
- [120] Maeda, H., Akaike, T., Sakata, Y. and Maruo, K. (1993) Role of bradykinin in microbial infection: Enhancement of septicemia by microbial proteinases and kinin. In *Agents and Actions Supplements* 42, Proteases, Proteases Inhibitor and Protease-Derived Peptides: Importance in Human Pathophysiology and Therapeutics. (eds. J.C.Chironis and J.E. Repine) pp.159-165, Birkhauser Verlag AG, Switz.
- [121] Maeda, H., Edo, K. and Ishida, N. eds. (1997) *Neocarzinostatin: The Past, Present, and Future of an Anticancer Drug*. Springer-Verlag, Tokyo, pp. 1-289
- [122] Fojo, T., and Grady, C. (2009) How much is life worth: Cetuximab, non-small cell lung cancer, and the \$440 billion question. *J. Natl. Cancer Inst.* 101, 1044-1048.
- [123] Hind, D., Pilgrim, H., and Ward, S. (2007) Questions about adjuvant trastuzumab still remain. *Lancet* 369, 3-5.
- [124] Editorial (2008) Welcome clinical leadership at NICE. *Lancet* 372, 601.
- [125] Sjöblom, T., Jones, S., Wood, L. D. et al (2006) The consensus coding sequences of human breast and colorectal cancers. *Science* 314, 268-274.
- [126] Shah, S. P., Morin, R. D., Khattra, J., et al and Aparicio, S. (2009) Mutational evolution in a lobular breast tumour profiled at single nucleotide resolution. *Nature* 461, 809-813
- [127] Maeda, H. (2011) Tumor-selective delivery of macromolecular drugs via the EPR effect: Background and future prospects, *Bioconj. Chem.* 21, 797-802.

## INDEX

### A

- aclarubicin, 47  
activation of influenza virus, 21  
activation of iNOS, 35  
activation of kallikrein-kinin cascade, 21  
Akaike, Takaaki, 21, 57  
albumin, 13, 20, 21, 30, 31, 46  
albumin-binding properties, 13  
amino acid sequence, 8, 52  
angina pectoris, 45  
angiotensin II (AT), 2, 31, 37, 40, 41, 42  
angiotensin II (AT) induced high blood pressure, 41, 42  
antigenicity, 13  
antioxidant activity, 27  
antitumor effect of NCS, 11  
antitumor protein, 5  
ARCO, 10  
Asakawa Award, 22  
AT-II, 31, 40, 41  
AT-II-induced hypertension, 31  
AUC (area under the concentration-time curve), 33, 51  
augment the EPR, 53  
augmentation of the EPR, 41  
augmentation of EPR effect, 46

### B

- bacterial proteases, 1, 19, 21, 22  
Bharate, G., 49  
bilirubin, 47, 48  
biliverdin, 47  
bradykinin, 1, 20, 21, 29, 35, 36  
bradykinin generation, 21

### C

- cancer, 2, 18, 26, 39, 42, 43, 46, 47  
cancer chemotherapy, 46, 47  
cancer treatment, 14  
canolol, 27  
carbon monoxide (CO), 36, 47  
carcinogenesis, 1, 26, 27  
CCRF, 7, 8  
Chemical and Engineering News, 10, 52  
chemical modification, 3, 9, 23, 29  
chemical structure of SMANCS, 11  
Children's Cancer Research Foundation (CCRF), 7  
Children's Hospital Boston, 7, 34  
Christophi, C., 35, 57  
chronic myelogenous leukemia cells, 49

- cisplatin, 48  
 citation frequency, 37  
<sup>14</sup>C-labeled Lipiodol, 16, 17  
 cleavage of hemagglutinin, 21  
 clinical development of NCS, 11  
 clinical effects, 52  
 CO (carbon monoxide), 36, 48  
 collagenase (MMPs), 36  
 Commemorative Gold Medal Award, 22  
 compartment models, 12, 65  
 computed tomography (CT), 15  
 Controlled Release Society, 38  
 CT images, 16  
 CT scan images, 31, 41  
 cytochrome-P450 reductase, 24

**D**

- Dana-Farber Cancer Institute, 7  
 Daruwalla, J., 34, 57  
 Davis, California, 3, 30  
 degradation of DNA, 11  
 Department of Surgery, 13, 15, 37  
 details of the EPR Effect, 29  
 difficult-to-treat cases, 41  
 discovery of the EPR, 2, 13, 29, 32, 43  
 disintegration, 34, 49  
 DIVEMA (divinyl ether and maleic anhydride copolymer), 23, 32  
 DNA synthesis, 11  
 doxorubicin, 47  
 drawbacks of NCS, 13  
 Duncan, Ruth, 10, 32, 33, 57  
 Dvorak, H., 36

**E**

- E. K. Frey-E. Werle Foundation, 22  
 effect of SOD, 25  
 elevating blood pressure, 40  
 encapsulated drugs, 49  
 endocytotic intracellular uptake, 49  
 endogenous free, 1, 22, 23, 55

- endogenous oxidants, 48  
 endyene chromophore, 52  
 enhanced permeability and retention (EPR) effect, 12, 30  
 enhanced vascular permeability, 29  
 enhancement of EPR effect, 39, 46  
 EPR effect, 2, 12, 13, 14, 29, 30, 31, 32, 33, 35, 36, 37, 39, 40, 41, 43, 45, 46, 53  
 EPR effect and molecular weight, 33  
 EPR effect, augmentation of, 41, 46  
 EPR effect-based drugs, 53  
 EPR effect citation, 37  
 EPR effect, enhancement of, 40, 39  
 Evans blue, 20, 21, 29, 31  
 Evans blue-albumin, 21, 31, 46  
 extravasation, 20, 21, 29, 30, 31, 35, 36

**F**

- Factor XII, 20  
 factors facilitating the EPR effect, 35  
 Fang, Jun, 36, 45, 47, 48, 57  
 Farber, Sidney, 7, 8  
 Feeney, E. Robert, 3, 5, 51  
 femoral artery, 14, 15  
 fluorescein isothiocyanate (FITC), 6  
 Folkman, Judah, 34  
 free radicals, 1, 22, 23, 53, 57  
 Fritz, Hans, 22, 57  
 Fulbright, 3

**G**

- gallbladder cancer, 41, 42  
 Gandy, B. Judith, 57  
 genetic divergence, 53  
 Glaser, Charles, 8  
 Greish, Khaled, 40, 47, 48, 57

**H**

- Hageman factor, 20

Hakuaikei hospital, 40  
Harvard Medical School, 7, 34, 40  
heat shock protein, 36, 47  
*Helicobacter pylori*, 27  
hemagglutinin of influenza virus, 21  
heme oxygenase-1 (HO-1), 47  
hepatic artery, 14, 16, 17, 31, 42  
hepatocellular carcinoma (HCC), 16, 39,  
  42, 52  
hepatoma, 15, 16, 31, 32, 52  
heterogeneity of EPR effect, 39, 45  
Hinuma, Yorio, 8  
Hirano, Takashi, 23  
HO-1, 36, 47, 49  
HO-1 inhibitor, 36, 49  
honorary citizen, 32  
honorary mayor, 32  
HPMA (hydroxypropyl methacrylate  
  copolymer), 32  
HPMA-copolymer, 33  
Hsp-32, 38  
hydrophobic, 10, 13  
hypertensive state, 40, 41  
hypovascular tumors, 39  
hypoxia-inducible factor (HIF-1 $\alpha$ ), 39,  
  40, 46

**I**

immunogenicity, 13, 51  
immunoglobulin, 30  
immunosuppression, 13  
inactivation, 51, 52  
inducible form of NOS, 35  
inflammation, 1, 19, 26, 27, 35, 36, 53  
influenza virus, 1, 21, 22, 23  
influenza virus infection, 1, 21, 22, 23, 25,  
  27  
iNOS, 26, 35  
Institute of Macromolecular Chemistry,  
  32  
intracellular uptake, 9, 49  
involvement of proteases, 22  
ISDN; isosorbital dinitrate, 46

Ishida, Nakao, 5, 7, 11, 52  
Iwai, Ken, 17  
Iyer, A. K., 48

**J**

Japan NO Meeting, 55  
Japan Society of Drug Delivery System,  
  38, 55, 59  
Japanese Society for Bacteriology, 22,  
  55  
Journal of Drug Targeting, 43, 55  
Journal of Medicinal Chemistry, 10

**K**

kallikrein-kinin cascade, 1, 20, 21, 22, 35  
Kamata, Ryuji, 19  
Kanamaru, Ryunosuke, 11, 57  
Kawauchi, Hiroshi, 6  
Kimura, Masami, 35, 57  
kinin, 1, 20, 21, 22, 29, 35  
kinin generation, 22, 29  
Kobe University, 55  
Koch, Robert, postulate of, 1, 23  
Konno, Toshimitsu, 13, 14, 15, 16, 17,  
  32, 57  
Kumagai, Katsu, 6, 11  
Kumamoto, 8, 9, 32, 40, 52, 55, 57  
Kumamoto University, 9, 55  
Kumamoto University Hospital, 13  
Kumamoto University Medical School, 8,  
  9, 21  
Kuraray Company, 10  
Kuromizu, Kenji, 8  
Kuwahara, Hideo, 27

**L**

laparotomy, 14, 15  
Li, Chang, 40  
Li, Chao Hao, 8  
Linus Pauling, 23

Lipiodol, 14, 15, 16, 17, 52  
 liquid ammonia, 8  
 low vascular density, 39  
 low vascular density tumor, 40  
 lung cancer, 46  
 lymphatic clearance, 17, 37  
 lymphatic metastasis, 9  
 lymphotropic drug, 9

## M

macromolecules, 2, 9, 17, 30, 37, 40, 46  
 macrophages, 13  
 Maki, Shojiro, 32, 57  
 Maruo, Keishi, 21, 57  
 massive metastatic liver cancer, 43  
 massive renal cell carcinoma, 42  
 Matsumoto, Koki, 19, 20, 57  
 Matsumura, Yasuhiro, 29, 30, 31, 35, 57  
 Meares, Claude F., 30  
 mechanism of mutation, 27  
 metastatic liver cancers, 39, 42, 43  
 micelle formation, 47  
 micronodules, 34  
 minimum inhibitory concentration, 11  
 molecular target drugs, 52, 53  
 molecular weight and AUC, 33  
 Molla, Akhteruzzaman, 20, 57  
 Moncada, Salvador, 55, 57  
*Mongolian gerbil*, 27  
 Müller-Esterl, Werner, 22  
 mutant virus, 26  
 mutants of virus and bacteria, 24  
 mutation, 26, 27, 53  
 myocardial infarction, 45

## N

Nagai Award, 38, 59  
 Nagai Innovation Award for Outstanding Achievement, 38  
 Nagamitsu, Akinori, 40  
 Nakamura, Hideaki, 49

National Cancer Center Hospital East, 29  
 National Cancer Institute, 5  
 National Institutes of Health (NIH), 6  
 natural killer (NK) cells, 13  
 NCS, 1, 5, 6, 8, 9, 10, 11, 12, 13, 23, 30  
 Neocarzinostatin (NCS), 1, 5, 52  
 nitration of G, 24  
 nitric oxide, 2, 23, 36, 55  
 nitric oxide synthase, 23  
 nitrite ion ( $\text{NO}_2^-$ ), 45  
 nitroglycerin, 2, 45, 46  
 NO, 23, 24, 26, 47, 48  
 NO releasing agents, 47  
 NOS, 24, 26

## O

O'Brien, Paul, 34  
 Oda, Tatsuya, 23  
 Okamura, Ryoichi, 19  
 oncogene *bcl-abl*, 49  
 one-compartment model, 12,  
 Ono, Yasushi, 11  
 ONOO<sup>-</sup>, 24, 26, 36  
 ophthalmology, 19  
 Osaka University, 55  
 ovalbumin, 30  
 ovarian cancer, 44  
 ovomucoid, 3, 30

## P

pancreatic cancer, 42, 43, 44  
 pegylated proteins, 14, 52  
 PEGylated ZnPP, 47  
 PEG-ZnPP, 47, 48  
 peroxynitrite, 24, 26, 36  
 pharmacokinetics, 9, 11, 12, 32, 48, 51  
 pharmacokinetics of NCS, 11, 12  
 pin point delivery, 34  
 pinpoint targeting, 2  
 pirarubicin, 34, 47, 48  
 plasma concentration, 51

- pO<sub>2</sub>, 45, 46  
poly(styrene-co-maleic anhydride) (SMA), 10  
polymer leakage, 34  
polymer-conjugated superoxide dismutase (SOD), 22  
Postulates of Robert Koch, 1, 23  
protease, 1, 3, 19, 21, 22, 29, 51, 55  
protein drugs, 51, 52, 53  
*Pseudomonas aeruginosa*, 19  
pyran copolymer, 23  
pyran-conjugated SOD, 23, 25

**R**

- Raeder, Beate, 49  
reactive nitrogen species (RNS), 1, 19  
reactive oxygen species (ROS), 1, 19, 48, 52  
Regelson, William, 32  
renal cell carcinoma, 39, 42, 44  
renal excretion, 12  
Ringsdorf, Helmut, 32, 38, 43, 57  
ROS, 1, 19, 23, 26, 27, 48  
ROS and RNS in infection, 26  
Royal Pharmaceutical Society, 43

**S**

- Sahoo, S. K., 49  
*Salmonella*, 27  
San Antonio, 32  
scanning electron micrographs of vascular casts, 34  
scavenger of NO, 35  
Seki, Takahiro, 45  
Seldinger's method, 15  
Sendai virus, 5, 26  
*Serratia marcescens*, 19  
serratil 56K protease, 21  
Seymour, Len, 32, 57  
SMA, 1, 10, 13, 30, 34, 47, 48, 49, 51  
SMA micelles, 47, 48, 49

- SMANCS, 1, 2, 10, 13, 16, 30, 31, 32, 40, 41, 43, 44, 46, 51, 52, 53  
SMANCS application, 16  
SMANCS/Lipiodol, 14, 15, 16, 17, 31, 40, 46  
SMANCS/Lipiodol infusion, 46  
SMANCS/Lipiodol injection, 31  
SMANCS, structure of, 11  
SMANCS synthesis, 10, 37  
SMA-pirarubicin micelles, 34, 50, 70  
SMA-THP (pirarubicin), 49  
SMA-ZnPP, 47, 49  
Society of Cancer Prevention of Japan, 55  
SOD, 1, 22, 23, 25  
solid tumor tissues, 30  
solid tumors, 2, 20, 30, 32, 35, 37, 39, 53  
State of Oklahoma, 32  
stomach cancer, 42, 43  
structure-activity relationships, 9  
super oxide, 1, 22, 23, 25  
super oxide dismutase [SOD], 1, 22, 23  
Suzuki, Maro, 40

**T**

- Takaki, Asami, 57  
Takeshita, Jiro, 10  
Taniguchi, Naoyuki, 55  
Tatematsu, Shoei, 27  
taxol, 47  
the B-type staining (peripheral ring shape), 43  
therapeutic effect of SMANCS (SX)/Lipiodol (LP), 42  
THP, 47, 48, 49  
time dependent increase of tumor uptake, 33  
tissue distribution, 9, 51  
tissue oxygen tension (pO<sub>2</sub>), 45, 46  
Toda, Noboru, 57  
Tohoku University, 3, 5, 11  
Tohoku University Medical School, 5, 11

transferrin, 30  
tumor necrosis factor (TNF)- $\alpha$ , 36  
tumor nodules, 34  
tumor uptake of macromolecules, 30  
tumor-selective drug delivery, 2, 16, 34  
two-compartment model, 12

vesicorenal recirculation, 12  
viral infection, 1, 22, 23, 49  
virus infectivity, 21, 22  
vivo half-life, 23, 51  
VPF/VEGF, 36  
VX-2, 14, 16

**U**

ubiquitous characteristics, 53  
Ulbrich, Karel, 32  
University of California, 3, 8, 30  
University of Melbourne, 34  
University of Vienna, 49  
urinary bladder, 12  
urinary excretion, 12, 51, 52

**W**

Walker 256 tumor, 6

**V**


Valent, Peter, 59  
vascular density, 37, 39, 40  
vascular endothelial cell growth factor (VEGF), 36  
vascular extravasation, 20  
VEGF, 36, 40, 46

**Y**

Yamanouchi Pharmaceutical Company, 11, 52  
Yokoyama, Ikuzo, 15, 16  
Yokoyama, Mitsuhiro, 55

**Z**

zinc protoporphyrin (ZnPP), 50, 71  
ZnPP, 47, 48, 49



*Hiroshi Maeda*

# RECOLLECTIONS OF 45 YEARS IN RESEARCH

From Protein Chemistry  
to Polymeric Drugs  
to the EPR Effect  
in Cancer Therapy

[www.novapublishers.com](http://www.novapublishers.com)

ISBN 978-1-61761-101-8



9 781617 611018