

## ORIGINAL ARTICLE

# Treatment of pancreatic fibrosis with siRNA against a collagen-specific chaperone in vitamin A-coupled liposomes

Hirotoishi Ishiwatari,<sup>1</sup> Yasushi Sato,<sup>1</sup> Kazuyuki Murase,<sup>1</sup> Akihiro Yoneda,<sup>2</sup> Ryosuke Fujita,<sup>2</sup> Hiroki Nishita,<sup>2</sup> Naoko Kubo Birukawa,<sup>2</sup> Tsuyoshi Hayashi,<sup>1</sup> Tsutomu Sato,<sup>1</sup> Koji Miyanishi,<sup>1</sup> Rishu Takimoto,<sup>1</sup> Masayoshi Kobune,<sup>1</sup> Shigenori Ota,<sup>3</sup> Yasutoshi Kimura,<sup>3</sup> Koichi Hirata,<sup>3</sup> Junji Kato,<sup>1</sup> Yoshiro Niitsu<sup>2</sup>

► Additional supplementary files are published online only. To view these files please visit the journal online (<http://dx.doi.org/10.1136/gutjnl-2011-301746>).

<sup>1</sup>Fourth Department of Internal Medicine, Sapporo Medical University, Sapporo, Japan  
<sup>2</sup>Department of Molecular Target Exploration, Sapporo Medical University, Sapporo, Japan  
<sup>3</sup>First Department of Surgery, Sapporo Medical University, Sapporo, Japan

## Correspondence to

Professor Yoshiro Niitsu, Department of Molecular Target Exploration, Sapporo Medical University School of Medicine, South 1, West 17, Chuo-ku, Sapporo 060-8543, Japan; [niitsu@sapmed.ac.jp](mailto:niitsu@sapmed.ac.jp)

HI, YS and KM contributed equally to this work.

Revised 27 September 2012  
 Accepted 12 October 2012

## ABSTRACT

**Background and objective** Fibrosis associated with chronic pancreatitis is an irreversible lesion that can disrupt pancreatic exocrine and endocrine function. Currently, there are no approved treatments for this disease. We previously showed that siRNA against collagen-specific chaperone protein gp46, encapsulated in vitamin A-coupled liposomes (VA-lip-siRNAgp46), resolved fibrosis in a model of liver cirrhosis. This treatment was investigated for pancreatic fibrosis induced by dibutyltin dichloride (DBTC) and cerulein in rats.

**Methods** Specific uptake of VA-lip-siRNAgp46, conjugated with 6'-carboxyfluorescein (FAM) by activated pancreatic stellate cells (aPSCs), was analysed by fluorescence activated cell sorting (FACS). Intracellular distribution of VA-lip-siRNAgp46-FAM was examined by fluorescent microscopy. Suppression of gp46 expression by VA-lip-siRNAgp46 was assessed by immunoblotting. Collagen synthesis in aPSCs was assayed by dye-binding. Specific delivery of VA-lip-siRNAgp46 to aPSCs in DBTC rats was verified following intravenous VA-lip-siRNA-FAM and <sup>3</sup>H-VA-lip-siRNAgp46. The effect of VA-lip-siRNA on pancreatic histology in DBTC- and cerulein-treated rats was determined by Azan-Mallory staining and hydroxyproline content.

**Results** FACS analysis revealed specific uptake of VA-lip-siRNAgp46-FAM through the retinol binding protein receptor by aPSCs in vitro. Immunoblotting and collagen assay verified knockdown of gp46 and suppression of collagen secretion, respectively, by aPSCs after transduction of VA-lip-siRNAgp46. Specific delivery of VA-lip-siRNAgp46 to aPSCs in fibrotic areas in DBTC rats was confirmed by fluorescence and radioactivity 24 h after the final injection. 10 systemic VA-lip-siRNAgp46 treatments resolved pancreatic fibrosis, and suppressed tissue hydroxyproline levels in DBTC- and cerulein-treated rats.

**Conclusion** These data suggest the therapeutic potential of the present approach for reversing pancreatic fibrosis.

## INTRODUCTION

Chronic pancreatitis is characterised by inflammation and replacement of parenchymal cells with fibrotic tissue leading to functional alterations, such as debilitating exocrine and occasional endocrine insufficiency. The accumulation of fibrotic tissue results from sustained activation of pancreatic stellate cells (PSCs), which proliferate and

## Significance of this study

### What is already known on this subject?

- Pancreatic fibrosis is an irreversible lesion that can disrupt pancreatic function; there are no approved treatments.
- The accumulation of fibrotic tissue results from sustained activation of pancreatic stellate cells (PSCs).
- The characteristics of PSCs resemble those of hepatic stellate cells (HSCs).
- We have previously succeeded in resolving liver fibrosis in cirrhotic rat models by targeting HSCs with vitamin A coupled liposomes which carried siRNA against the collagen specific chaperone protein gp46.

### What are the new findings?

- Rat activated PSCs take up vitamin A-coupled liposomes (VA-lip-siRNAgp46) in a retinol binding protein-mediated fashion, similarly to activated HSCs.
- Transduction of siRNAgp46 in activated PSCs caused suppression of collagen secretion.
- Activated PSCs specifically took up siRNAgp46 encapsulated in vitamin A liposomes in the pancreas of dibutyltin dichloride (DBTC)-treated rats.
- VA-lip-siRNAgp46 treatments resolved pancreatic fibrosis, and suppressed tissue hydroxyproline levels in DBTC- and cerulein-treated rats.
- This is the first demonstration of successful targeting of antifibrotic drug to both cells and molecule which are responsible for pancreatic fibrosis.

### How might it impact on clinical practice in the foreseeable future?

- Results suggest the therapeutic potential of the present approach for reversing pancreatic fibrosis.

secrete collagen in response to stimulation by cytokines, growth factors, and reactive oxygen species from inflammatory cells and damaged pancreatic tissue.<sup>1 2</sup> Various approaches to suppress the

activation of PSCs and to eradicate activated PSCs (aPSCs) have been explored for the prevention and treatment of fibrosis associated with chronic pancreatitis.<sup>3–5</sup> However, so far no clinically applicable agents have been developed, mainly because of the inability for specific drug delivery to aPSCs.

We previously demonstrated complete resolution of liver cirrhosis in rat models using vitamin A-coupled liposomes to specifically deliver siRNA against the collagen-specific chaperone, gp46 (VA-lip-siRNA<sub>gp46</sub>), to hepatic stellate cells (HSCs) via the circulating receptor for retinol binding protein (RBP).<sup>6</sup> The characteristics of PSCs, including collagen synthesis, storage of vitamin A, expression of gp46, etc, resemble those of HSCs.<sup>7</sup> Therefore, in the present study, we examined whether our previous therapeutic approach for liver cirrhosis could be successfully applied for the treatment of pancreatic fibrosis.

## MATERIALS AND METHODS

### Preparation of siRNA<sub>gp46</sub> and its conjugate with 6'-carboxyfluorescein

A formulation of siRNA directed against gp46, a rat homologue of human HSP47, was purchased from Hokkaido System Science (Sapporo, Japan). The sense and anti-sense strands of siRNAs have been described in detail previously.<sup>6</sup> For fluorescence activated cell sorting (FACS) analyses and in vivo tracing of gp46siRNA, gp46 siRNA with 6'-carboxyfluorescein (6-FAM)-coupled to the 5' end of the sense strand was used.

A formulation of siRNA directed against HSP47 (GenBank accession no. 50454) was purchased from Hokkaido System Science. The sense and anti-sense strands of siRNAs were: HSP47, 5'-ggacagggccucac aacuaTT-3' (sense); 5'-uaguuguagagg ccuguccTT-3' (antisense).

### Preparation of vitamin A-coupled liposomes carrying siRNA<sub>gp46</sub>

Vitamin A-coupled liposomes carrying siRNA<sub>gp46</sub> (VA-lip-siRNA<sub>gp46</sub>) were prepared as described previously.<sup>6</sup>

### Animals

Male Lewis rats (Charles River, Tokyo, Japan), weighing 150–200 g, and male Wistar rats (Charles River), weighing 250–300 g, were used for dibutyltin dichloride (DBTC) and cerulein experiments, respectively. All animal procedures were approved by the Sapporo Medical College Institutional Animal Care and Use Committee.

### Isolation and cultivation of rat PSCs

Rat PSCs were isolated by density gradient centrifugation, as detailed previously.<sup>8</sup> All experiments were performed with culture-activated cells (aPSCs, passage 1–3).

### Isolation and cultivation of human PSCs

Human pancreases were obtained during surgery for chronic pancreatitis. All these patients were seen at Sapporo Medical University. Informed consent in writing was obtained from each patient. Human PSCs were isolated by outgrowth, using explant techniques from the pancreas as described previously.<sup>1</sup> In this study, experiments were performed on activated  $\alpha$ -smooth muscle actin (SMA)-positive cells between the first and third serial passages using lines.

### FACS analysis of VA-lip-siRNA<sub>gp46</sub>-FAM

Rat aPSCs were cultivated with VA-lip-siRNA<sub>gp46</sub>-FAM (50 nM of siRNA) for 30 min. For the blocking assay,  $1 \times 10^4$  cells were treated with mouse anti-RBP antibody (10  $\mu$ g/ml, BD

Pharmingen, San Diego, California, USA) for 30 min before adding VA-lip-siRNA<sub>gp46</sub>-FAM. The mean fluorescence intensity of VA-lip-siRNA<sub>gp46</sub>-FAM-treated cells was assessed on a FACScalibur with CellQuest software (Becton Dickinson, San Jose, California, USA).

### Intracellular distribution analysis of VA-lip-siRNA<sub>gp46</sub>-FAM

Distribution of VA-lip-siRNA<sub>gp46</sub>-FAM in aPSCs after transduction was analysed as described previously.<sup>6</sup>

### Western blot analysis

Protein extracts of cells and pancreas specimens were resolved over 4/20 sodium dodecylsulphate–polyacrylamide gels, transferred onto nitrocellulose membranes, probed with antibodies against HSP47 (gp46) (Stressgen Biotechnologies, Victoria, BC, Canada) or  $\beta$ -actin (Cell Signaling, Beverly, Massachusetts, USA), and then probed with peroxidase-coupled antibodies as the secondary antibody (Oncogene Research Product, Boston, Massachusetts, USA). Lastly, the cells were visualised with ECL (Amersham Life Science, Arlington Heights, Illinois, USA). Western blots were quantified using ImageJ 1.43u (NIH, Bethesda, Maryland, USA).

### Quantification of collagen production

Collagen production by rat aPSCs was measured according to the method described previously for aHSCs, except for cell cultivation period after transduction; 24 h for aHSCs and 72 h for aPSCs.<sup>6,9</sup>

### Quantitative RT-PCR

Total RNA was isolated using RNeasy mini kits (Qiagen, Hilden, Germany). Total RNA (1  $\mu$ g) was used for reverse transcription with SuperScript II (Invitrogen, Carlsbad, California, USA) plus RNaseOUT (Invitrogen) using random primers (Invitrogen) according to the manufacturer's instructions. All TaqMan primers mixed with probes (GAPDH, Rn 99999916\_s1; MMP2, Rn 02532334\_s1; COL1A1, Rn00801649\_g1; TIMP-1, Rn 00587558\_m1; Gp46, Rn 00367777\_m1; TGF $\beta$ , Rn01475963\_m1) were purchased from Applied Biosystems (Foster City, California, USA). The TaqMan reactions were performed using 7300 Fast Real Time PCR System (Applied Biosystems). The results were expressed as the ratio of the number of copies of the product gene to the number of copies of the housekeeping gene (GAPDH) from the same RNA (respective cDNA) sample and PCR run.

### Induction of pancreatic and hepatic fibrosis by DBTC

Among several models of pancreatic fibrosis,<sup>10–18</sup> we chose the DBTC model in which the common bile duct is obstructed by a plug formed with necrotic biliopancreatic ductal epithelium, because the procedure is relatively simple to perform as compared with other models and irreversible pancreatitis can be induced by a single injection of DBTC. DBTC (Sigma, St Louis, Missouri, USA) was first dissolved in ethanol (1 part) and then mixed with glycerol (2 parts) and dimethyl sulfoxide (2 parts).<sup>10</sup> For administration of DBTC, we selected the right jugular vein route to avoid any possible damage of the tail vein caused by the tail vein route for subsequent application of VA-lip-siRNA<sub>gp46</sub>.<sup>10</sup> In preliminary experiments, we found the lethality rates of dosages 8.0, 7.0 and 5.0 mg/kg body weight to be 4/4 (100%), 7/16 (44%) and 5/27 (19%), respectively. Thus, we selected the dosage of 5.0 mg/kg for the main experiments (see online supplementary figure S1).

Fibrosis, as revealed by Azan-Mallory staining, was evident in pancreatic and hepatic specimens on days 29 and 43 of DBTC injection (see online supplementary figure S1A–C).

#### Induction of pancreatic fibrosis by cerulein

Male Wistar rats received two intraperitoneal injections of 50 µg/kg cerulein (Sigma) 1 h apart every week for 6 weeks, as described by Ishibashi (see online supplementary figure S2A,B).<sup>18</sup>

#### Collagenase activity in pancreas homogenates

Collagenase activity (collagen type I) in pancreas homogenates was measured as described previously.<sup>19</sup>

#### In vivo localisation of VA-lip-siRNA random-FAM in rat pancreas

From day 43 of DBTC administration, rats were injected intravenously with 1 µl/g body weight of VA-lip-siRNA random-FAM or Lip-siRNA random-FAM (0.75 mg/kg siRNA) three times on alternating days. At 24 h after the last injection, the rats were sacrificed by saline perfusion. Pancreatic tissue was immediately embedded in OCT compound (Sakura Finetechnical, Tokyo, Japan) medium and cryogenically sectioned. Multicoloured fluorescent staining of sections and their analysis were carried out as described previously.<sup>6</sup>

#### Tissue distribution of radiolabelled VA-lip-siRNA random

<sup>3</sup>H-VA-lip-siRNA random (200 µCi), prepared as described previously, was administered via the tail vein under normal pressure in either DBTC-treated rats (day 43) or normal rats. After 24 h, the rats were sacrificed under anaesthesia, and radioactivity of each tissue was assayed as described previously.<sup>6</sup>

#### Treatment of DBTC rats and cerulein rats with VA-lip-siRNA gp46 and measurement of hydroxyproline content

Three groups of rats (n=10 per group) were used for histological evaluations. From day 43 of DBTC or cerulein administration, Lip-siRNA gp46, VA-lip-siRNA gp46 (0.75 mg/kg siRNA) or phosphate buffered saline (PBS; three times a week every other day) were injected for a total of 10 times via the tail vein under normal pressure in a volume of 1 µl/g body weight.<sup>6</sup> Hydroxyproline content in the pancreas was measured as previously described.<sup>20</sup>

#### Immunohistochemical staining for α-SMA

Pancreas was fixed with 10% paraformaldehyde. Then, immunohistochemical staining for α-SMA was performed by the dextran polymer method using monoclonal anti α-SMA antibody (1:1000, Sigma) and an Envision Kit (Dako), followed by colouring with 3, 3'-diaminobenzidine (DAB) and nuclear staining with Gill's haematoxylin solution. To accurately quantitate areas stained with α-SMA, slides from six randomly selected low-power fields (×100) per pancreas section from each rat were viewed by microscopy (Axioplan 2; Carl Zeiss) and the percentage of areas stained with α-SMA was quantified as previously described.<sup>6</sup>

#### In vitro apoptosis assay

Rat and human aPSCs under going apoptosis were stained with an in situ Cell Death Detection kit (Roche) according to the manufacturer's protocol. Slides were washed with PBS, and exposed to Prolong Gold Antifade Reagent with 4', 6-diamidino-2-phenylindole (DAPI) (Molecular Probes) to stain nuclei. The number of terminal deoxynucleotidyl transferase-mediated deoxyuridine triphosphate nick-end labelling (TUNEL)-positive

cells (green) in aPSCs were counted in 10 random high-power fields (×800) using fluorescent microscopy (Keyence, BZ-8000) for each sample.

#### Double staining of pancreas specimen for TUNEL and α-SMA

Double staining for TUNEL and α-SMA was undertaken and the number of TUNEL-positive apoptotic cells within the fibrotic bands was determined using a method modified from that described by Iredale *et al.*<sup>19</sup> Briefly, the specimen was first stained with an in situ Cell Death Detection kit (Roche) according to the manufacturer's protocol, followed by immunostaining for α-SMA (Sigma), with an alkaline phosphatase-conjugated anti-mouse second antibody (KPL, Gaithersburg, Maryland, USA) using a Vector Red alkaline phosphatase substrate kit 1 (Vector Lab, Peterborough, UK). The number of TUNEL-positive cells (brown) in α-SMA-positive areas (red), but not those in parenchymal area, were counted in 10 random high-power fields (×800).

#### Ethics approval

This research follows the tenets and regulations of the Declaration of Helsinki and has been approved by the Animal Care and the Institutional Review Board at Sapporo Medical University.

#### Statistics

Results are presented as mean±SD for each sample. Multiple comparisons between control groups and other groups were performed by Dunnett's test.

## RESULTS

#### Specific uptake of RBP bound VA-lip-siRNA gp46 by rat aPSCs


Rat aPSCs, which were stained positive for α-SMA (figure 1A), were incubated with VA-lip-siRNA gp46-FAM or Lip-siRNA gp46-FAM in the presence of 10% fetal calf serum, and were observed under a fluorescence microscope. In VA-lip-siRNA gp46-FAM treated aPSCs, fluorescence appeared as a fine granular pattern in the cytoplasm at 30 min and as denser granular patterns in the perinuclear region at 2 h (figure 1B). In contrast, in Lip-siRNA gp46-FAM treated aPSCs, no green fluorescence was seen at 30 min and perinuclear fluorescence at 2 h was very faint (figure 1B). The fluorescence intensity of VA-lip-siRNA gp46-FAM aPSCs as revealed by FACS was clearly suppressed by RBP antibody to nearly the same level as Lip-siRNA gp46-FAM aPSCs (figure 1C).

#### Suppression of gp46 expression and collagen secretion of rat aPSCs by VA-lip-siRNA gp46

Treatment of aPSCs with VA-lip-siRNA gp46 brought about dose-dependent suppression of gp46 with almost complete suppression at 50 nM, which lasted at least 72 h (figure 1D,E) while treatment with VA-lip-siRNA random or Lip-siRNA gp46 did not cause any suppression. In the culture plate of VA-lip-siRNA gp46-treated aPSCs, significantly less collagen than that of VA-lip-siRNA random-treated or non-treated aPSCs was found (figure 1F).


#### Delivery of VA-lip-siRNA random-FAM to aPSCs in vivo

The specific delivery of VA-lip-siRNA random-FAM to aPSCs in the fibrotic pancreas was examined by fluorescent emission 24 h after three injections (figure 2A). Specimens were prepared from head (figure 2B,D) and body portions (figure 2C,E) of the pancreas. In both portions, fluorescence of VA-lip-siRNA random-FAM (green) was identified predominantly in the



**Figure 1** Retinol binding protein receptor-dependent uptake of vitamin A-coupled liposomes (VA-lip-siRNAgp46)-carboxyfluorescein (FAM) by activated pancreatic stellate cells (aPSCs) and suppressive effect of siRNAgp46 on gp46 expression and collagen synthesis. (A) Representative fluorescent images of rat aPSCs. Phase-contrast micrograph image of aPSCs (left panel) and immunofluorescent staining of aPSCs with Cy3-conjugated anti- $\alpha$ -SMA antibody (red) and nuclei counterstained with DAPI (blue) (right panel). Bars=100  $\mu$ m. (B) Representative fluorescent images of the intracellular distribution of FAM-labelled siRNA. Rat aPSCs were treated with VA-lip-siRNAgp46-FAM or Lip-siRNAgp46-FAM. At 30 min, the medium was replaced with fresh medium. At the time indicated, cells were fixed and analysed by fluorescence microscopy to determine the relative intracellular distribution of siRNA-FAM (green). Bars=100  $\mu$ m. One of five representative images is shown. (C) Representative fluorescence activated cell sorting patterns of rat PSCs treated with vitamin A-free liposomes carrying siRNAgp46-FAM (Lip-siRNAgp46-FAM) or VA-lip-siRNAgp46-FAM with or without anti-RBP antibody. Mean fluorescence intensity is indicated. Five independent experiments were carried out in each set of aPSCs and the results were essentially the same. (D) Western blotting was used to analyse the expression of gp46 and  $\beta$ -actin

**Figure 2** Specific delivery of vitamin A-coupled liposomes (VA-lip-siRNA) random-carboxyfluorescein (FAM) to activated pancreatic stellate cells (aPSCs) in dibutyltin dichloride (DBTC)-treated rats. (A) Schedule of VA-lip-siRNA random-FAM injection in rats with DBTC-induced pancreatic fibrosis. Samples (B, D; pancreas head, C, E; pancreas body) were obtained from DBTC-treated rats that received three injections of VA-lip-siRNA random-FAM or Lip-siRNA random-FAM (siRNA doses of 0.75 mg/kg, 3 times every other day) ( $n=6$  per group). Representative fluorescent images of  $\alpha$ -SMA visualised by Cy3-conjugated anti- $\alpha$ -SMA antibody (red), nuclei counterstained with DAPI (blue) and siRNA random-FAM (green) in pancreas specimens obtained 24 h after the last injection. Pictures were taken at original magnification ( $\times 200$ ). Fluorescence of siRNA random-FAM (green) was identified in pancreas, predominantly in the region that stained positive for  $\alpha$ -SMA giving rise to areas with a merged yellow colour.



spotty region that stained positive only for  $\alpha$ -SMA (aPSCs) (red), giving rise to a merged yellow colour (figure 2B,C). By contrast, in rats treated with Lip-siRNA random-FAM, the yellow area in  $\alpha$ -SMA positive regions was negligible (figure 2D,E).

The yellow area in six randomly selected high-power ( $\times 200$ ) fields from each specimen occupied  $32.1 \pm 10.5\%$  of the area stained for  $\alpha$ -SMA. By contrast, the yellow area in  $\alpha$ -SMA-positive regions was negligible ( $2.0 \pm 1.5\%$ ) in a rat injected with lip-siRNA gp46-FAM.

To further elucidate the organ distribution,  $^3\text{H}$ -labelled VA-lip-siRNA random was administered to DBTC-treated rats with pancreatitis (day 43) and normal rats (table 1). Radioactivity in the pancreas of DBTC-treated rats was significantly higher (by approximately fivefold) than that of normal rats. In the liver, radioactivity in the DBTC group was also higher (approximately fourfold) than that of normal rats. Other organs, including the lungs, spleen and intestines, showed no increase in radioactivity in DBTC-treated rats, relative to normal rats.

#### Duration of gp46 suppression in aPSCs in the pancreas of DBTC rats treated with VA-lip-siRNA gp46

To examine the duration of effect of siRNA gp46 on gp46 expression, DBTC rats ( $n=15$ ) were injected intravenously with VA-lip-siRNA gp46 on day 43 (see online supplementary figure S3A). The expression of gp46 in the pancreas of rats sacrificed at day 43, 44, 45, 46 and 47 ( $n=3$  per group), was analysed by western blotting. The results for each rat per group were essentially the same. Representative western blot bands from one of three rats at each time-point and densitometric values are shown in online supplementary figure S3B,C. Suppression of gp46 expression started 24 h after initiated of treatment and lasted for at least 3 days.

#### Resolution of pancreatic fibrosis by VA-lip-siRNA gp46 in DBTC-treated rats

After 10 treatments with VA-lip-siRNA gp46 (figure 3A), an apparent reduction and statistically significant reduction analysed by computerised imaging of fibrosis represented by

(normalisation control) in aPSCs transfected with Lip-siRNA gp46, VA-lip-siRNA random or VA-lip-siRNA gp46. Dose-dependent inhibition of gp46 expression by siRNA gp46 was observed. (E) Duration of suppressive effect of siRNA gp46 on gp46 expression in aPSCs. Rat aPSCs were treated with VA-lip-siRNA gp46 or VA-lip-siRNA random. At 30 min, the medium was replaced with fresh medium. At the time indicated, the expression of gp46 and  $\beta$ -actin (normalisation control) was analysed by western blotting. Similar results were obtained in three independent experiments. (F) Collagen deposition on tissue culture plates was assayed by the dye-binding method 72 h after transduction in rat aPSCs treated with VA-lip-siRNA gp46 or with VA-lip-siRNA random. Data are expressed as mean  $\pm$  SD calculated from five transductions and as a percentage of untreated control. \* $p < 0.05$  vs VA-lip-siRNA gp46. NS, not significant.

**Table 1** Tissue biodistribution of (<sup>3</sup>H)VA-lip-siRNAgp46 in rats

	Radioactivity (cpm/tissue)		p Value (normal vs DBTC rat)	Ratio (DBTC rat/ normal)
	Normal ( $\times 10^5$ )	DBTC rat ( $\times 10^5$ )		
Pancreas	0.481 $\pm$ 0.270	2.42 $\pm$ 0.840	0.001	5.03
Liver	77.2 $\pm$ 21.0	319 $\pm$ 105	0.002	4.14
Lung	3.83 $\pm$ 1.89	5.26 $\pm$ 3.66	NS	1.37
Spleen	1.75 $\pm$ 0.740	1.84 $\pm$ 1.11	NS	1.05
Intestine	4.23 $\pm$ 1.59	4.52 $\pm$ 1.00	NS	1.07

DBTC-treated rats (day 43 rat) and normal rats (n=3 per group) received a single intravenous injection of 200  $\mu$ Ci (<sup>3</sup>H)VA-lip-siRNAgp46 via the tail vein. Tissue biodistribution was analysed 24 h later. Data represent means $\pm$ SD (n=3). Similar results were obtained in two independent experiments. DBTC, dibutyltin dichloride; NS, not significant; VA-lip-siRNAgp46, vitamin A-coupled liposomes.

Azan-Mallory positive area was confirmed (figure 3B,C). Results were consistent with data showing substantial suppression of hydroxyproline levels (figure 3D).

#### Disappearance of aPSCs following treatment with VA-lip-siRNAgp46

To elucidate of disappearance of aPSCs after treatment, immunostaining for  $\alpha$ -SMA was performed. A substantially smaller area was stained for  $\alpha$ -SMA in rats treated with VA-lip-siRNAgp46 compared to the stained area in rats treated with Lip-siRNAgp46 or PBS (figure 3E,F).

#### Effect of VA-lip-siRNAgp46 treatment on pancreatic inflammation in DBTC-treated rats

Since it is known that PSCs play a role not only in fibrosis but also in the inflammatory response,<sup>21</sup> we explored the effect of VA-lip-siRNAgp46 treatment on inflammation of pancreas in DBTC-treated rats. The degree of inflammation, assessed by cell infiltration, fatty change, tubular complex formation and infiltration of macrophages and T cells were reduced after treatment with VA-lip-siRNAgp46 (see online supplementary figure S4 and table S1).

#### Resolution of hepatic fibrosis by VA-lip-siRNAgp46 in DBTC-treated rats

In the DBTC model, hepatic fibrosis occurs in addition to pancreatic fibrosis because DBTC causes obstruction of the common bile duct (see online supplementary figure S5A,B). The Azan-Mallory positive area was significantly reduced in specimens from VA-lip-siRNAgp46-treated rats as compared with that in control specimens (see online supplementary figure S5C).

#### Apoptosis of siRNAgp46-induced aPSCs

We examined the possibility that apoptotic death of rat aPSCs may be induced by siRNAgp46 treatment in vitro. The number of TUNEL positive cells was greater in siRNAgp46-transfected aPSCs than in control aPSCs (figure 4A). The ratio of apoptotic nuclei to total nuclei of siRNAgp46-transfected aPSCs was 38.3 $\pm$ 8.2% and was significantly increased relative to controls (figure 4B).

Apoptosis of aPSCs by siRNAHSP47 was also examined in human PSCs obtained by outgrowth from fibrotic human pancreas. TUNEL staining in apoptotic nuclei of siRNAHSP47-transfected aPSCs was higher than in control aPSCs (figure 4C). The ratio of apoptotic nuclei to total nuclei of siRNAHSP47-

transfected aPSCs was 18.3 $\pm$ 4.2% and was significantly increased relative to controls (figure 4D).

Furthermore, apoptosis of aPSCs by siRNAgp46 was examined in DBTC-treated rats. In DBTC pancreas, TUNEL positive cells in areas overlapping with aPSCs ( $\alpha$ -SMA positive) also increased after treatment with VA-lip-siRNAgp46 (figure 4E,F).

#### Resolution of pancreatic fibrosis by VA-lip-siRNAgp46 in cerulein-treated rats

In the cerulein induced pancreatitis model (see online supplementary figure S2 and figure 5A), the therapeutic effect of VA-lip-siRNAgp46 was also evident with respect to shrinkage of the fibrotic area (figure 5B) and suppression of hydroxyproline levels (figure 5C).

#### Changes in mRNA expression of fibrosis-related proteins before and after administration of VA-lip-siRNAgp46 in DBTC-treated rats

We measured mRNA levels of gp46, collagen I, MMP2, TIMP-1, and TGF $\beta$  in pancreas homogenates from normal rats and DBTC-treated rats (figure 6A). The mRNA ratios relative to GAPDH mRNA are shown in figure 6B. mRNAs of gp46, collagen I, MMP2 and TGF $\beta$  expressed in aPSCs were increased in fibrotic pancreas as compared to normal pancreas. TIMP-1 mRNA levels did not increase but instead showed a slight decrease in response to induction of fibrosis. However, after treatment with VA-lip-siRNAgp46, the expression of all mRNAs was significantly decreased, presumably due to the disappearance of aPSCs.

#### Elevated collagenase activity in the pancreas of DBTC-treated rats


To confirm that sufficient collagenase activity to resolve the predeposited collagen resides in fibrotic pancreas, we measured collagenase activity in the pancreas homogenates of DBTC-treated rats after three injections of VA-lip-siRNAgp46 (day 49), and found that it was significantly higher than that of control normal rat and became even higher after treatment with VA-lip-siRNA gp46 (figure 6C).

#### Changes in microvessel density in fibrotic pancreas tissue before and after treatment with VA-lip-siRNAgp46


In DBTC-treated pancreas before VA-lip-siRNAgp46 treatment, the mean microvessel counts were 39 $\pm$ 9.5. After 10 injections of VA-lip-siRNAgp46, the mean microvessel counts significantly reduced to 19 $\pm$ 9.2 (see online supplementary figure S6).

## DISCUSSION

PSCs have been shown to store vitamin A and synthesise collagen, and they can be activated by inflammatory cytokines, exerting functions similar to those of HSCs.<sup>1 2 7</sup> However, it has not been elucidated whether RBP is involved in vitamin A uptake. It is also not clear whether HSP47 is involved in collagen secretion by aPSCs. Therefore we first clarified that aPSCs indeed take up vitamin A in an RBP-mediated fashion by demonstrating that with VA-lip-siRNAgp46-FAM, but not with Lip-siRNAgp46-FAM, the fluorescence that appeared within 30 min of incubation was suppressed by anti-RBP antibody, and the amount of collagen on the culture plate of VA-lip-siRNAgp46-treated cells was significantly reduced (figure 1C,F). These findings verified the similarity of aPSCs to activated hepatic stellate cells in terms of RBP-mediated vitamin A uptake and gp46-assisted collagen secretion.



**Figure 3** Effect of intravenously injected vitamin A-coupled liposomes (VA-lip-siRNAgp46) on dibutyltin dichloride (DBTC)-induced pancreatic fibrosis. (A) Schedule of VA-lip-siRNAgp46 treatment in rats with DBTC-induced pancreatic fibrosis. Samples were obtained from DBTC-treated rats that received 10 injections of VA-lip-siRNAgp46, Lip-siRNA gp46 (siRNA doses of 0.75 mg/kg, 3 times every other day), or PBS alone (n=10 per group). (B) Representative photomicrographs of Azan-Mallory stained pancreas section. Pictures were taken at original magnification ( $\times 100$ ). Magnified images corresponding to the areas enclosed in boxes are presented as indicated. (C) Azan-Mallory positive staining area assessed by computerized image analysis. Data were obtained from six randomly selected fields in each of 10 rats from four groups and represent the mean  $\pm$  SD. (D) Hydroxyproline content in the pancreas. Mean  $\pm$  SD of 10 rats per group. (E) Representative immunohistochemical staining images of activated pancreatic stellate cells stained with anti- $\alpha$ -SMA antibody (brown). Pictures were taken at original magnification ( $\times 200$ ). Results from each group of 10 rats were essentially similar. (F)  $\alpha$ -SMA-positive staining area assessed by computerized image analysis. Data were obtained from six randomly selected fields in each of 10 rats from three groups and represent the mean  $\pm$  SD. \* $p < 0.05$ ; \*\* $p < 0.01$  vs VA-lip-siRNAgp46 treated-DBTC rat.



**Figure 4** Apoptosis of activated pancreatic stellate cells (aPSCs) induced by vitamin A-coupled liposomes (VA-lip-siRNA) treatment. (A) Representative photomicrographs of rat aPSCs undergoing apoptosis. Nuclei of aPSCs treated with VA-lip-siRNAgp46 or with control agents were stained with FITC-transferase-mediated deoxyuridine triphosphate nick-end labelling (TUNEL) (green) and DAPI (blue) 72 h after each treatment. Arrowheads indicate apoptotic cells. Pictures taken at original magnification ( $\times 200$ ) and magnified images corresponding to the areas enclosed in boxes are presented in the inset. (B) Quantification of apoptotic aPSCs shown in figure 4A. Apoptotic nuclei (green) and total nuclei (green + blue) were examined by fluorescent microscopy and counted in 10 randomly selected fields per slide for each indicated treatment; the ratios of apoptotic to normal nuclei were expressed as percentages. Results represent the mean  $\pm$  SD of three independent experiments.  $*p < 0.01$  vs VA-lip-siRNAgp46. (C) Representative photomicrographs of human aPSCs undergoing apoptosis. Nuclei of aPSCs treated with VA-lip-siRNAHSP47 or with control agents were stained with FITC-TUNEL (green) and DAPI (blue) 72 h after each treatment. Arrowheads indicate apoptotic cells. Pictures taken at original



Incidentally, regarding a possible role of other cellular components such as myofibroblasts in pancreatic fibrosis, it is rather difficult to distinguish it from that of aPSCs since when PSCs are activated, they become positive for  $\alpha$ -SMA, which is characteristic of myofibroblasts. In this context, it should be noted that quiescent PSCs store vitamin A, and when they become myofibroblasts, although no more vitamin A droplets in their cytoplasm are observed, they still can take up vitamin A as shown by *in vitro* transfection experiments using VA-lip-siRNA<sub>gp46</sub>. This was true for hepatic stellate cells too. This is carefully discussed in our previous paper.<sup>6</sup> It is speculated that deposited vitamin A in quiescent cells may be used up or partly excreted on myofibroblast transformation.

Prior to the examination of therapeutic effect, we assured specific delivery of VA-lip-siRNA<sub>gp46</sub> to aPSCs in fibrotic pancreas tissue, demonstrating the mergence of the green fluorescence of siRNA-FAM encapsulated in vitamin A liposomes with the red fluorescence of  $\alpha$ -SMA-positive cells (aPSCs), and higher accumulation of <sup>3</sup>H-VA-lip-siRNA in the pancreas of DBTC-treated rats than that in normal rats. The reason for 10-fold more radioactivity in the lung and intestine may be due to the higher content of macrophages in these organs. However, the 150-fold difference between liver and pancreas may not be simply ascribed to the difference in macrophage content, but more likely to the increased number of activated hepatic stellate cells since in this model, not only pancreas but also liver undergoes fibrosis because of obstruction of the common bile duct.

Collectively these results supported the notion that aPSCs preferentially take up siRNA<sub>gp46</sub> encapsulated in vitamin A liposomes *in vivo*. We also examined the kinetics of the effect *in vivo*, prior to assessment of therapeutic efficacy, and found that the suppression of gp46 expression lasted for 3 days after the injection of VA-lip-siRNA<sub>gp46</sub>. Thus, our therapy protocol involved injecting the drug three times per week.

The therapeutic effect of treatment, namely, histological improvement and normalised hydroxyproline levels in the pancreas, was observed following administration of drug 10 times. In our previous study using liver cirrhosis models, substantial resolution of fibrotic tissue was obtained after only five treatments.<sup>6</sup> In the present study, we also initially used only five treatments, but improvement of fibrosis was unsatisfactory. This may be due to the fact that the pancreas is relatively poor in blood supply compared with the liver, and thus the dosage of siRNA<sub>gp46</sub> at the fibrotic loci supplied by five injections may not have been sufficient to bring about effective resolution of pancreatic fibrosis.

Incidentally, it may be noteworthy to mention that our treatment brought about less vessel density since it promises wider implications such as depletion of tumour stroma in pancreas cancer. In this regard, it is nowadays a well accepted

concept that modality to normalise the vascularity by diminishing immature microvessels with anti-neoangiogenesis antibody, rather enhances the efficacy of chemotherapy when the antibody and anticancer drug are used in combination.<sup>22</sup>

Regarding the mechanism for the enhanced microvasculature in fibrotic tissue and decrement of vascularity after treatment, we assume, in good agreement with reports by and Hideshima and Okada,<sup>23, 24</sup> that aPSCs are producing angiogenesis factor(s) which will be eradicated from the tissue in accordance with apoptosis induction by treatment. This assumption is based on our recent finding that rat aPSCs are indeed expressing some angiogenesis factors, including fibroblast growth factor (unpublished observation).


The resolution of fibrosis appears to occur by a dual mechanism involving siRNA<sub>gp46</sub>, which inhibits *de novo* secretion of collagen, and tissue collagenase, which dissolves predeposited collagen matrix. This assumption is consistent with the fact that collagenase activity in the pancreas at day 42 and day 49 after DBTC treatment, when massive fibrosis occurred, was higher than that of normal pancreas. The reason for sustained high collagenase activity after VA-lip-siRNA<sub>gp46</sub> treatment, despite the suppressed MMP2 mRNA, may be that TIMP from aPSCs is also reduced due to apoptosis of aPSCs. In addition, a persistent extracellular matrix pool of collagenase after it is secreted from cells may account for the sustained collagenase activity.<sup>25, 26</sup> Activation of matrix metalloproteinases (MMPs) derived from inflammatory cells by reduced TIMP-1 expression may also account for the discrepancy.

Furthermore, apoptosis of aPSCs via transduction of siRNA<sub>gp46</sub> was confirmed in our *in vitro* experiment, indicating that a similar mechanism is involved in apoptosis of aHSCs and aPSCs. It is plausible that collagen secreted by aHSCs and aPSCs serves as a survival signal and that this signal is abrogated, leading to apoptosis, as a result of collagen degradation by collagenase (unpublished observation). In support of this possibility, our results demonstrated that high collagenase activity was maintained in the pancreas even after treatment (figure 6C). Incidentally, induction of apoptosis by inhibiting collagen secretion was confirmed to occur even in human PSCs treated with VA-lip-siRNA<sub>HSP47</sub>, indicating applicability of this approach in the clinical setting.

Interestingly, inflammation associated with fibrosis in this model was also decreased by this treatment, probably due to the fact that aPSCs, which were eradicated by our treatment, are involved not only in the development of fibrosis but also in the inflammatory reaction.<sup>21</sup>

Liver fibrosis in DBTC-treated rats was also successfully resolved with treatment in the present study. However, because the epithelial plug in the common duct was not removed by the treatment, increased serum amylase, alanine aminotransferase and bilirubin persisted (data not shown). In the DBTC

magnification ( $\times 200$ ) and magnified images corresponding to the areas enclosed in boxes are presented in the inset. (D) Quantification of apoptotic aPSCs shown in figure 4C. Apoptotic nuclei (green) and total nuclei (green + blue) were examined by fluorescent microscopy and counted in 10 randomly selected fields per slide for each indicated treatment; the ratios of apoptotic to normal nuclei were expressed as percentages. Results represent the mean  $\pm$  SD of three independent experiments. \* $p < 0.01$  vs VA-lip-siRNA<sub>HSP47</sub>. (E) Representative immunohistochemical staining of pancreas specimens from dibutyltin dichloride (DBTC) (5 mg/kg) treated rats (day 56) that received six injections of VA-lip-siRNA<sub>gp46</sub> (left panel) or VA-lip-siRNA<sub>random</sub> (right panel) (siRNA doses of 0.75 mg/kg, 3 times every other day) stained for TUNEL (brown, arrowhead) and  $\alpha$ -SMA (red). Magnified images of the corresponding areas in boxes indicated that the number of cells with TUNEL-positive overlapping of  $\alpha$ -SMA-positive PSCs was substantially increased in VA-lip-siRNA<sub>gp46</sub> specimens compared with VA-lip-siRNA<sub>random</sub> specimens. Pictures were taken at original magnification ( $\times 100$ ). (F) Quantification of TUNEL-positive PSCs in pancreas specimens from VA-lip-siRNA<sub>gp46</sub>-treated DBTC-rats and VA-lip-siRNA<sub>random</sub>. The number of TUNEL-positive cells in  $\alpha$ -SMA-positive areas was counted in 10 randomly selected fields per pancreas section of each rat. \* $p < 0.01$ . The results were essentially similar within each group of three rats.



**Figure 5** Effect of intravenously injected vitamin A-coupled liposomes (VA-lip-siRNAgp46) on cerulein-induced pancreatic fibrosis. (A) Schedule of VA-lip-siRNAgp46 treatment in rats with cerulein-induced pancreatic fibrosis. Samples were obtained from cerulein-treated rats that received 10 injections of VA-lip-siRNAgp46, Lip-siRNA gp46 (siRNA doses of 0.75 mg/kg, 3 times every other day) or PBS alone ( $n=10$  per group). (B) Representative photomicrographs of an Azan-Mallory-stained pancreas section. Pictures taken at original magnification ( $\times 100$ ). (C) Azan-Mallory-positive staining area assessed by computerized image analysis. Data were obtained from six randomly selected fields in each of 10 rats from three groups and represent the mean  $\pm$  SD. (D) Hydroxyproline content in the pancreas. Mean  $\pm$  SD of 10 rats per group. \* $p < 0.05$ ; \*\* $p < 0.01$  vs VA-lip-siRNAgp46 treated-dibutyltin dichloride rat.

model, unlike in advanced chronic pancreatitis in humans,  $\beta$ -cells remained intact and, therefore, no change in blood glucose levels was observed throughout the experimental period (data not shown).

To further evaluate the therapeutic effect of our modality in another model, we established a chronic pancreatitis model based on repeated administration of cerulein to rats.<sup>12, 18</sup> The results in this model were compatible with those of the DBTC model, confirming the general efficacy of our modality in dissimilar pancreatic fibrosis models.


Off-target effects and stimulation of toll-like receptors (TLRs) are both critical issues to be solved in the application of therapies using siRNA. Since we used the same siRNA as that of the previous study on liver cirrhosis in which we denied a

bystander effect by demonstrating comparable gene silencing efficacy and antifibrotic effect with two other independent siRNAs against the same target,<sup>6</sup> we believe it is unlikely to be the case.

Nonetheless, for future clinical applications, it will be necessary to conduct careful explorations of adverse effects, off-target effects and TLR stimulation.

In conclusion, the present data clearly indicate the therapeutic potential of our modality for pancreas fibrosis and suggest its potential for the treatment of organ fibrosis.

**Contributors** HI, YS and KM designed research, performed experiments and wrote the paper. AY, RF, HN, NB, TH, TS, KM, RT, MK, SA, YK, KH and JK performed experiments. YN designed research, wrote the paper and supervised the whole project. All authors discussed the results and commented on the manuscript.



**Figure 6** Effect of vitamin A-coupled liposomes (VA-lip-siRNAgp46) treatments on mRNA expression of fibrosis-related proteins and collagenase activity in pancreas homogenates. (A) Schedule of VA-lip-siRNAgp46 treatment in rats with dibutyltin dichloride (DBTC)-induced pancreatic fibrosis. Samples were obtained from DBTC-treated rats that received three injections of VA-lip-siRNAgp46, Lip-siRNA gp46 (siRNA doses of 0.75 mg/kg, 3 times every other day), or PBS alone ( $n=3$  per group). (B) The expression of gp46, procollagen I, MMP-2, TIMP-1, and TGF $\beta$  mRNA in normal rats, DBTC-treated rats treated with three injections of PBS, DBTC-treated rats treated with three injections of VA-lip-siRNArandom, and DBTC-treated rats treated with three injections of VA-lip-siRNAgp46 were quantitated by real-time PCR. Expression was normalised as the ratio to GAPDH mRNA, a housekeeping gene. \* $p<0.01$  vs DBTC rat (day42). \*\* $p<0.01$  vs PBS (day46). \*\*\* $p<0.05$  vs PBS (day46). (C) Total collagenase activity in pancreas homogenates from normal rats, DBTC-treated rats (day 42), DBTC-treated rats (day 49), and DBTC rats treated with three injections of VA-lip-siRNAgp46 (siRNA doses of 0.75 mg/kg, 3 times every other day, day 49) in the absence or presence of MMP inhibitor (1,10-phenanthroline). Results are the mean $\pm$ SD of five rats per group. \* $p<0.01$  vs without inhibitor. \*\* $p<0.05$  vs normal rat.

**Competing interests** None.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

- Bachem MG**, Schneider E, Gross H, *et al*. Identification, culture, and characterization of pancreatic stellate cells in rats and humans. *Gastroenterology* 1998;**115**:421–32.
- Schneider E**, Schmid-Kotsas A, Zhao J, *et al*. Identification of mediators stimulating proliferation and matrix synthesis of rat pancreatic stellate cells. *Am J Physiol Cell Physiol* 2001;**281**:C532–43.
- Shimizu K**, Shiratori K, Kobayashi M, *et al*. Troglitazone inhibits the progression of chronic pancreatitis and the profibrogenic activity of pancreatic stellate cells via a PPAR $\gamma$ -independent mechanism. *Pancreas* 2004;**29**:67–74.
- Yamada T**, Kuno A, Masuda K, *et al*. Candesartan, an angiotensin II receptor antagonist, suppresses pancreatic inflammation and fibrosis in rats. *J Pharmacol Exp Ther* 2003;**307**:17–23.
- Kuno A**, Yamada T, Masuda K, *et al*. Angiotensin-converting enzyme inhibitor attenuates pancreatic inflammation and fibrosis in male Wistar Bonn/Kobori rats. *Gastroenterology* 2003;**124**:1010–19.
- Sato Y**, Murase K, Kato J, *et al*. Resolution of liver cirrhosis using vitamin A-coupled liposomes to deliver siRNA against a collagen-specific chaperone. *Nat Biotechnol* 2008;**26**:431–42.
- Buchholz M**, Kestler HA, Holzmann K, *et al*. Transcriptome analysis of human hepatic and pancreatic stellate cells: organ-specific variations of a common transcriptional phenotype. *J Mol Med* 2005;**83**:795–805.
- Apte MV**, Haber PS, Applegate TL, *et al*. Periacinar stellate shaped cells in rat pancreas: identification, isolation, and culture. *Gut* 1998;**43**:128–33.
- Williams EJ**, Benyon RC, Trim N, *et al*. Relaxin inhibits effective collagen deposition by cultured hepatic stellate cells and decreases rat liver fibrosis in vivo. *Gut* 2001;**49**:577–83.
- Inoue M**, Ino Y, Gibo J, *et al*. The role of monocyte chemoattractant protein-1 in experimental chronic pancreatitis model induced by dibutyltin dichloride in rats. *Pancreas* 2002;**25**:e64–70.
- Merkord J**, Jonas L, Weber H, *et al*. Acute interstitial pancreatitis in rats induced by dibutyltin dichloride (DBTC): pathogenesis and natural course of lesions. *Pancreas* 1997;**15**:392–401.
- Elsasser HP**, Haake T, Grimmig M, *et al*. Repetitive cerulein-induced pancreatitis and pancreatic fibrosis in the rat. *Pancreas* 1992;**7**:385–90.
- Puig-Divi V**, Molero X, Salas A, *et al*. Induction of chronic pancreatic disease by trinitrobenzene sulfonic acid infusion into rat pancreatic ducts. *Pancreas* 1996;**13**:417–24.
- Merkord J**, Weber H, Jonas L, *et al*. Influence of ethanol on long-term effects of dibutyltin dichloride (DBTC) in pancreas and liver of rats. *Hum Exp Toxicol* 1998;**17**:144–50.
- Yamamoto M**, Otani M, Otsuki M. A new model of chronic pancreatitis in rats. *Am J Physiol Gastrointest Liver Physiol* 2006;**291**:G700–8.
- Tsuchitani M**, Saegusa T, Narama I, *et al*. A new diabetic strain of rat (WBN/Kob). *Lab Anim* 1985;**19**:200–7.
- Sparmann G**, Merkord J, Jaschke A, *et al*. Pancreatic fibrosis in experimental pancreatitis induced by dibutyltin dichloride. *Gastroenterology* 1997;**112**:1664–72.
- Ishibashi T**, Zhao H, Kawabe K, *et al*. Blocking of monocyte chemoattractant protein-1 (MCP-1) activity attenuates the severity of acute pancreatitis in rats. *J Gastroenterol* 2008;**43**:79–85.
- Iredale JP**, Benyon RC, Pickering J, *et al*. Mechanisms of spontaneous resolution of rat liver fibrosis. Hepatic stellate cell apoptosis and reduced hepatic expression of metalloproteinase inhibitors. *J Clin Invest* 1998;**102**:538–49.
- Weidenbach H**, Lerch MM, Turi S, *et al*. Failure of a prolyl 4-hydroxylase inhibitor to alter extracellular matrix deposition during experimental pancreatitis. *Digestion* 1997;**58**:50–7.
- Masamune A**, Watanabe T, Kikuta K, *et al*. Roles of pancreatic stellate cells in pancreatic inflammation and fibrosis. *Clin Gastroenterol Hepatol* 2009;**7**:S48–54.
- Jain RK**. Normalizing tumor vasculature with anti-angiogenic therapy: a new paradigm for combination therapy. *Nat Med* 2001;**7**:987–9.
- Hideshima K**, Seike J, Mimura S, *et al*. HSP47 and angiogenic factor expression and its implication for the healing of periosteal defects in the mouse cranium. *Oral Med Pathol* 2006;**11**:27–33.
- Okada H**, Inoue T, Kanno Y, *et al*. Selective depletion of fibroblasts preserves morphology and the functional integrity of peritoneum in transgenic mice with peritoneal fibrosing syndrome. *Kidney Int* 2003;**64**:1722–32.
- Zhou X**, Hovell CJ, Pawley S, *et al*. Expression of matrix metalloproteinase-2 and -14 persists during early resolution of experimental liver fibrosis and might contribute to fibrolysis. *Liver Int* 2004;**24**:492–501.
- Steffensen B**, Wallon UM, Overall CM. Extracellular matrix binding properties of recombinant fibronectin type II-like modules of human 72-kDa gelatinase/type IV collagenase. High affinity binding to native type I collagen but not native type IV collagen. *J Biol Chem* 1995;**270**:11555–66.

# Zoledronic Acid But Not Somatostatin Analogs Exerts Anti-Tumor Effects in a Model of Murine Prostatic Neuroendocrine Carcinoma of the Development of Castration-Resistant Prostate Cancer

Kohei Hashimoto,<sup>1</sup> Naoya Masumori,<sup>1\*</sup> Toshiaki Tanaka,<sup>1</sup> Toshihiro Maeda,<sup>1</sup> Ko Kobayashi,<sup>1</sup> Hiroshi Kitamura,<sup>1</sup> Koichi Hirata,<sup>2</sup> and Taiji Tsukamoto<sup>1</sup>

<sup>1</sup>Department of Urology, Sapporo Medical University School of Medicine, Japan

<sup>2</sup>First Department of Surgery, Sapporo Medical University School of Medicine, Japan

**BACKGROUND.** Since neuroendocrine (NE) cells play an important role in the development of castration-resistant prostate cancer (CRPC), target therapy to NE cells should be considered for treating CRPC. We investigated the effects zoledronic acid (ZOL) and two somatostatin analogs (octreotide: SMS, and pasireotide: SOM) on an NE allograft (NE-10) and its cell line (NE-CS), which were established from the prostate of the LPB-Tag 12T-10 transgenic mouse.

**METHODS.** We examined the *in vivo* effects of ZOL, SMS and SOM as single agents and their combinations on subcutaneously inoculated NE-10 allografts and the *in vitro* effects on NE-CS cells. Apoptosis and cell cycle activity were assessed by immunohistochemistry using TdT-mediated dUTP-biotin nick-end labeling (TUNEL) and a Ki-67 antibody, respectively.

**RESULTS.** *In vivo* growth of NE-10 tumors treated with ZOL, ZOL plus SMS, or ZOL plus SOM was significantly inhibited compared to the control as a consequence of induction of apoptosis and cell cycle arrest. ZOL induced time- and dose-dependent inhibition of *in vitro* proliferation of NE-CS cells, but the somatostatin analogs (SMS and SOM) did not. ZOL also inhibited migration of NE-CS cells. These effects were caused by inhibition of Erk1/2 phosphorylation via impairment of prenylation of Ras.

**CONCLUSIONS.** ZOL, but not SMS or SOM, induced apoptosis and inhibition of proliferation and migration through impaired prenylation of Ras in NE carcinoma models. Our findings support the possibility that ZOL could be used in the early phase for controlling NE cells, which may trigger progression to CRPC. *Prostate* © 2012 Wiley Periodicals, Inc.

**KEY WORDS:** zoledronic acid; somatostatin analog; neuroendocrine carcinoma; prostate cancer; anti-tumor effect

## INTRODUCTION

Prostate cancer is the most common non-cutaneous malignancy in men in developed countries [1]. Its incidence has been gradually increasing even in non-Caucasian men. As the growth of cancer cells is androgen-dependent, androgen deprivation therapy including surgical or chemical castration has been the mainstay of treatment for advanced prostate cancer. The response to this treatment lasts for a median of 36–48 months [2]. However, in most cases, the disease progresses despite the castration level of serum

testosterone, and results in castration-resistant prostate cancer (CRPC). The prognosis of CRPC is poor and a concrete treatment strategy has not yet been established.

---

\*Correspondence to: Naoya Masumori, Department of Urology, Sapporo Medical University School of Medicine S1, W16, Chuo-ku, Sapporo 060-8543, Japan. E-mail: masumori@sapmed.ac.jp  
Received 4 June 2012; Accepted 27 August 2012  
DOI 10.1002/pros.22590  
Published online in Wiley Online Library (wileyonlinelibrary.com).

The exact mechanisms behind progression to castration resistance remain poorly understood. However, recent studies suggest that neuroendocrine (NE) cells, as well as pathways involving or bypassing the androgen receptor (AR), may play an important role in the development of castration resistance [3]. NE cells are present in both the normal and neoplastic prostate. They regulate surrounding prostate cells by secreting growth-modulating neuropeptides such as chromogranin A, serotonin and parathyroid hormone-related protein (PTHrP) [4]. Increases in the NE phenotype and secretory products are thought to be closely associated with progression and castration resistance in prostate cancer [5,6]. Previously, we developed an NE allograft (NE-10) and its cell line (NE-CS) from the prostate of the LPB-Tag 12T-10 transgenic mouse [7–9]. We demonstrated that secretions from NE cells induced androgen-independent growth of human prostate cancer cell line LNCaP and promoted pulmonary metastasis [10]. Therefore, it is crucial to seek a new drug or drug combinations targeting these prostatic NE carcinoma models (NE-10 and NE-CS).

Zoledronic acid (ZOL) is a nitrogen-containing bisphosphonate that inhibits bone resorption of osteoclasts through the inhibition of farnesyl-pyrophosphate synthetase in the mevalonate pathway. This agent has been demonstrated to have beneficial effects in patients with bone metastases of prostate cancer, reducing bone pain and skeletal-related events [11]. ZOL was also shown to have direct anti-tumor activity in several cancer cell lines. It is suggested that ZOL inhibits proliferation and induces apoptosis by impairment of prenylation of Ras and other small GTP-binding proteins (G proteins) [12].

Somatostatin is a peptide hormone that regulates secretion of various exocrine and endocrine glands via specific somatostatin receptors (SSTR). Five different subtypes (SSTR1–5), which are coupled to G proteins, have been identified [13]. Since a majority of NE tumors predominantly express SSTR2, somatostatin analogs having high affinity for SSTR2a such as octreotide (SMS) are considered to be drugs for NE tumors [14,15]. Since several studies have reported that SSTR1 and SSTR5 are expressed in addition to SSTR2 in prostate cancer tissue [15,16], new somatostatin analogs such as pasireotide (SOM) that have high affinity for SSTR5 in addition to SSTR2 [17], may be useful as new drugs for prostatic NE carcinoma.

In the present study, we investigated whether the growth of NE carcinoma models (NE-10, NE-CS) could be influenced by ZOL and/or somatostatin analogs (SMS and SOM), having potential anti-tumor activities.

## MATERIALS AND METHODS

### Cell Lines and Cell Culture

NE-CS is a murine prostate neuroendocrine cancer cell line established in our institute [9]. It was derived from an NE-10 tumor [8]. Passage numbers between 12, and 16 were used in the study. The NE-CS cells were maintained in the culture medium described below in 5% CO<sub>2</sub> in a humidified incubator. The medium consisted of RPMI-1640 (Gibco BRL, Breda, The Netherlands) that was supplemented with MEM non-essential amino acid (10 ml/L, Gibco BRL), MEM sodium pyruvate, penicillin–streptomycin (10 ml/L, Gibco BRL), 10% fetal bovine serum (FBS, ICN Bio-medicals, Costa Mesa, CA), and 7.5% NaHCO<sub>3</sub>.

### Reverse Transcription Polymerase Chain Reaction (RT-PCR) Analysis

To investigate expression of SSTR in NE-10, RT-PCR analysis was performed. Tumor tissues in allografts of NE-10 were homogenized. Total RNA was extracted using an RNeasy kit (Qiagen, Valencia, CA) according to the manufacturer's instructions. A total of 2 µg of total RNA was reverse transcribed in a thermal cycles (Perkin–Elmer, Norwalk, CT) using SuperScript III (Invitrogen, Carlsbad, CA) and oligo (dT) 12–18 primers according to the manufacturer's instructions for 1 hr at 50°C in a 40 µl reaction mixture. Resulting cDNA (1 µl) was amplified with Taq polymerase and one set of oligonucleotide primers. Samples were denatured for 5 min at 94°C, and then amplified for 35 cycles at 94°C for 30 sec, 57°C for 30 sec, and 72°C for 1 min. Aliquots (9 µl) from each PCR sample were then analyzed by agarose-gel electrophoresis. Forward and reverse primer sequences were as follows: SSTR2a (5'-CAGCTGTACCATCAACTGGC, 5'-ATTGTCTCCTGCTTACTGTCTG), SSTR2b (5'-TGATCAATGTAGCTGTGTGG, 5'-CAAAGAACA-TTCTGGAAGC), SSTR5 (5'-TGCCTGATGGTCATGAGTGT, 5'-GGAAACTCTGGCGGAAGTTA), GAPDH (5'-TACAGCAACAGGGTGGTGGGA, 5'-ACCACAGTCCATGCCATCAC).

### Growth of NE-10 Allografts In Vivo

To examine the in vivo effects of ZOL, SMS and SOM as single agents and their combinatorial effects on prostatic NE carcinoma, we used NE-10 allografts. Six-week-old male BALB/c nude mice were castrated using the scrotal approach. After one week, 50 mg tissue fragments of the NE-10 allograft were inoculated subcutaneously (s.c.) into the flanks of mice. Two weeks after transplantation, NE-10 tumors grew to a volume of more than 100 mm<sup>3</sup>. The mice were then

randomized into six treatment groups (13 mice in each group): ZOL, SMS, SOM, ZOL plus SMS, ZOL plus SOM, and control. The groups were treated for 6 weeks with ZOL (1 µg/mouse, three times per week, s.c.), SMS (2 µg/mouse, once per day, s.c.), SOM (2 µg/mouse, twice per day, s.c.), ZOL plus SMS (1 µg/mouse, three times per week, s.c. plus 2 µg/mouse, once per day, s.c.), ZOL plus SOM (1 µg/mouse, three times per week, s.c. plus 2 µg/mouse, twice per day, s.c.), or saline (an equal volume of solvent/day, s.c.). These substances were dissolved in 100 µl saline. ZOL, SMS, and SOM were kindly provided by Novartis Pharma AG (Basel, Switzerland). These agents were soluble in saline. The body weights of mice were measured each week. The effects of treatments on tumor growth were determined by measuring tumor volume ( $0.523 \times \text{long diameter}^2 \times \text{short diameter}$ ). After 6 weeks of treatment, the mice were killed and the tumor and liver were removed. Liver weight was measured, and the numbers of metastatic nodules on the liver surface were macroscopically counted. The tissues were fixed in 10% formalin and embedded in paraffin. The 5 µm thick paraffin-embedded material was routinely processed for hematoxylin and eosin staining.

The Animal Experiment Committee of Sapporo Medical University approved the *in vivo* experiments. Animal care and housing followed the guidelines of the Animal Experiment Committee.

#### Cell Cycle and Apoptosis Analysis of Tissue Sections

Immunohistochemical staining was done with formalin-fixed paraffin-embedded tissue sections of NE-10 tumors. The 5 µm thick sections were deparaffinized in xylene and rehydrated in graded alcohol. Antigen retrieval was done by boiling sections for 20 min in a microwave oven in preheated 0.01 mol/L sodium citrate buffer (pH 6.0). Endogenous peroxidase activity was blocked by 3% hydrogen peroxide in ethanol for 10 min. After blocking with 1% non-fat dry milk in phosphate-buffered saline (PBS) (pH 7.4), the sections were reacted with a rabbit polyclonal anti-Ki-67 antibody (Abcam plc., Cambridge, UK) at 20 µg/ml or preimmune sera for 1 hr, followed by incubation with biotinylated goat anti-rabbit IgG (Nichirei, Tokyo, Japan) for 30 min. Subsequently, the sections were stained with streptavidin-biotin complex (Nichirei), followed by incubation with 3,3'-diaminobenzidine and counterstaining with hematoxylin. The same tissues were immunostained by TdT-mediated dUTP-biotin nick-end labeling (TUNEL) (In situ Apoptosis Detection Kit, Takara Bio, Inc., Otsu, Japan). The Ki-67 labeling index (KI) and apoptotic

index (AI) were determined as the ratios of immunohistochemically positive cells per 1,000 NE cells by using a fluorescence microscope (model BZ-9000; Keyence, Osaka, Japan).

#### Proliferation Assays

NE-CS cells ( $1 \times 10^4$ ) were suspended with 100 µl of culture medium in a 96-well plate for 24 hr, and then treated with the indicated concentrations (from 0.1 to 100 µmol/L) of ZOL, SMS and SOM for 24, 48, or 72 hr. For combination, the same concentrations of ZOL and SMS or SOM were used; for example, 1 µmol/L of ZOL to 1 µmol/L of SMS. In addition, they were treated with the indicated concentrations (from 0.1 to 100 µmol/L) of ZOL plus 1, 5, and 20 µmol/L of farnesyl-pyrophosphate ammonium salt (FOH) (Sigma-Aldrich, St. Louis, MO) for 48 hr. FOH is an isoprenoid to be involved in prenylation of several G proteins including Ras in the intracellular mevalonate pathway. Cell proliferation was assessed using a WST-8 (modified tetrazolium salt) cell proliferation kit (Cell Counting Kit-8, Dojin, Japan). Changes in absorbance at 450 nm were measured with a microplate reader. The growth inhibition was determined as the concentration inducing 50% inhibition (IC<sub>50</sub>). For analysis of the synergism between ZOL and somatostatin analogs (SMS and SOM), the combination indices (CI) were calculated by the isobologram equation method [18,19], and CI values of <1, 1, and >1 were considered to indicate synergistic, additive, and antagonistic effects, respectively [12].

#### Cell Cycle and Apoptosis Assays

NE-CS cells ( $3 \times 10^4$ ) were suspended with 100 µl of culture medium in a 96-well plate for 24 hr, and then treated with various concentrations of ZOL (from 10 to 100 µmol/L) for 48 hr. Then the cells were fixed with 4% paraformaldehyde, permeabilized with 0.1% Triton X-100, and labeled with the TUNEL technique (In situ Cell Death Detection Kit TMR red, Roche Diagnostics, Mannheim, Germany) and the primary anti-Ki-67 antibody at a 1/200 dilution (Abcam plc., Cambridge, UK). The Ki-67 antibody was detected with an Alexa Fluor 488 donkey anti-rabbit antibody, and nuclei were stained with 4,6-diamidino-2-phenylindole (DAPI) (Invitrogen, Carlsbad, CA). As in the *in vivo* study, the KI and the AI were measured by fluorescence immunohistochemistry using a fluorescence microscope (model BZ-9000; Keyence, Osaka, Japan).

#### Migration Assays

Cell migration analyses were performed as described previously (10). In a trans-well culture

chamber (Coster Science, Cambridge, MA), a polyvinylpyrrolidone-free polycarbonate filter with an 8.0  $\mu\text{m}$  pore size was precoated with 5  $\mu\text{g}$  of fibronectin (Biomedical Technologies, Stoughton, MA) on the lower surface. Two different experiments were performed. In experiment 1, NE-CS cells ( $1 \times 10^5$ ) were placed in the upper chamber with 100  $\mu\text{l}$  of culture medium with or without ZOL (10, 100  $\mu\text{mol/L}$ ). In the lower chamber, 600  $\mu\text{l}$  of culture medium was added. In experiment 2, NE-CS cells ( $1 \times 10^5$ ) were placed in the upper chamber with 100  $\mu\text{l}$  of culture medium adding 20  $\mu\text{mol/L}$  of FOH with or without ZOL (10, 100  $\mu\text{mol/L}$ ). In the lower chamber, 600  $\mu\text{l}$  of culture medium was added. The cells that migrated across the pores at 2, 4, 6, and 8 hr were counted under a microscope after hematoxylin and eosin staining. The experiments were carried out in triplicate. Data are shown as number of cells  $1 \text{ mm}^{-2}$  of membrane.

#### Pull Down and Western Blot Assays

NE-CS cells ( $1 \times 10^5$ ) were suspended with 2 ml of culture medium in a 6-well plate for 24 hr, and then treated with ZOL (10, 100  $\mu\text{mol/L}$ ) or ZOL (10, 100  $\mu\text{mol/L}$ ) + 20  $\mu\text{mol/L}$  FOH for 48 hr. Subsequent to the treatment, cells were washed three times with ice-cold PBS and solubilized in lysis buffer [RIPA buffer, 100 mmol/L PMSF, 500 mmol/L  $\text{Na}_3\text{VO}_4$ , 1 mol/L NaF, 2 mol/L Sigma 104 phosphatase substrate, Protease Inhibitor Mini Cocktail]. The total protein content of the cell lysates was determined by the BCA Protein Assay (Pierce, Rockford, IL).

Activated Ras was detected by pull-down assay. The GTP-bound form of Ras in the cell lysates was affinity-purified using the Raf1-Ras-binding domain (RBD)-GST complexed with glutathione beads following the manufacturer's instructions (Active Ras Pull Down and Detection Kit, Thermo Fisher Science, Waltham, MA). Complexes were analyzed by SDS-PAGE and immunoblotting with a Ras-specific antibody.

The cells lysates obtained were boiled in SDS sample buffer containing 0.5 mol/L 2-mercaptoethanol. Samples were separated by SDS-PAGE, transferred to polyvinylidene difluoride (PVDF) membranes and immunoblotted with rabbit monoclonal anti-Erk1/2 and anti-phospho-Erk1/2 antibodies (Cell Signaling Technology Inc., MA), and a mouse monoclonal anti- $\beta$ -actin antibody (Sigma-Aldrich). Separated proteins were visualized using horseradish peroxidase with enhancement by chemiluminescence (GE Healthcare Bio-Sciences Corp., NJ).

#### Statistical Analysis

We used the computer program StatView 5.0 for Windows (SAS Institute, Cary, NC). Student's *t*-test

was applied to compare results between two different groups. Repeated-measures ANOVA was used when comparing the in vivo tumor volume, and the in vitro cell migration in an individual group. One-way ANOVA was used when comparing the in vivo apoptosis, cell cycle progression, and liver metastases in an individual group. Statistical significance was assigned at  $P < 0.05$ .


## RESULTS

### Expression of SSTR2a and SSTR5 in NE-10 Allografts

Since effects of somatostatin analogs are mediated by expression of SSTR, we examined expression of the somatostatin receptor subtypes SSTR2 and SSTR5, to which SMS and SOM preferentially bind, respectively. Gene expression of SSTR2a and SSTR5 was observed in NE-10 allografts, but that of SSTR2b was not (Fig. 1).


### Effects of ZOL, SMS and SOM as Single Agents and in Combination on Subcutaneously Inoculated NE-10 Allografts

Growth of NE-10 tumors in mice treated with ZOL, ZOL plus SMS, and ZOL plus SOM was significantly slowed compared to the saline control ( $P = 0.003$ ,  $P < 0.001$ , and  $P = 0.001$ , respectively) (Fig. 2A). All treatments were well tolerated with maintenance of body weight (data not shown). We examined whether anti-tumor effects of each treatment were induced by apoptosis or cell cycle arrest by using TUNEL and Ki67 staining, respectively (Fig. 2B<sub>1</sub>). The AI was significantly increased in tumors of mice treated with ZOL, ZOL plus SMS, or ZOL plus SOM compared to the control (means: 9.2, 11.6, and 12.7, respectively, vs. 2.4) (Fig. 2B<sub>2</sub>). The KI was significantly decreased in tumors of mice treated with ZOL, ZOL plus SMS, or ZOL plus SOM compared to the control (means: 5.3, 8.3, and 4.2, respectively, vs. 15.9) (Fig. 2B<sub>3</sub>). The



**Fig. 1.** Expression of SSTR2a, SSTR2b, and SSTR5 in NE-10 allograft by RT-PCR. Gene expression of SSTR2a, and SSTR5 was observed in the NE-10 allograft, but that of SSTR2b was not.





**Fig. 2.** Effects of ZOL, SMS, and SOM as single agents and in combination on subcutaneous inoculated NE-10 allografts. Six-week-old male BALB/c nude mice were castrated. After one week, 50 mg tissue fragments from the NE-10 allograft model were subcutaneously inoculated into the backs of mice. For 2 weeks, NE-10 tumors were allowed to grow to approximately more than 100 mm<sup>3</sup> before randomization into six treatment groups: control, ZOL, SMS, SOM, ZOL plus SMS, and ZOL plus SOM (n = 13/group). NE-10 allografts in each group were treated for 6 weeks. **A:** Growth of NE-10 tumors in mice treated with ZOL, ZOL plus SMS, and ZOL plus SOM was significantly slowed compared to the saline control (P = 0.003, P < 0.001, and P = 0.001, respectively). Data are means; bars ± SE; \*, significantly different from control group (P < 0.05; repeated-measures ANOVA). **B:** Effects of ZOL, SMS and SOM as single agents and in combination on apoptosis and cell cycle progression. Immunohistochemical staining was done by using TUNEL and Ki67 staining (**B<sub>1</sub>**). Apoptotic effects were measured by the number of TUNEL-positive cells per 1,000 cells, apoptotic index (AI). The AI was significantly increased in tumors from mice treated with ZOL, ZOL plus SMS, or ZOL plus SOM compared to the control (means: 9.2, 11.6, and 12.7, respectively, vs. 2.4) (**B<sub>2</sub>**). Cell cycle progression was measured by the number of Ki67-positive cells per 1,000 cells (KI: Ki-67 labeling index). The KI was significantly decreased in tumors from mice treated with ZOL, ZOL plus SMS, or ZOL plus SOM compared to the control (means: 5.3, 8.3, and 4.2, respectively, vs. 15.9) (**B<sub>3</sub>**). Data are means; bars ± SD; \*, significantly different from control group (P < 0.05; one-way ANOVA). **C:** Effects of ZOL, SMS and SOM as single agents and in combination on liver metastases. The weights of livers having metastatic nodules in ZOL, ZOL plus SMS, or ZOL plus SOM were significantly lower than for the control (**C<sub>1</sub>**), but the numbers of metastatic nodules in these groups were not significantly different from the control (**C<sub>2</sub>**). Data are means; bars ± SD. \*, Significantly different from control group (P < 0.05; one-way ANOVA).

weights of livers having metastatic nodules in ZOL, ZOL plus SMS, or ZOL plus SOM were significantly lower than for the control (Fig. 2C<sub>1</sub>), but the numbers of metastatic nodules in these groups were not significantly different from the control (Fig. 2C<sub>2</sub>).

**Effects of ZOL, SMS and SOM as Single Agents and in Combination on Growth of NE-CS Cells In Vitro**


We investigated the inhibitory effects of ZOL, SMS, and SOM, alone and in combination on proliferation

of NE-CS cells. Cell viability was measured by the WST-8 assay when NE-CS cells were treated with various concentrations of ZOL, SMS and SOM (0.1–100  $\mu\text{mol/L}$ ) in the treatment groups for 24, 48 or 72 hr. For combinations, the same concentrations of ZOL and SMS or SOM were used. The IC<sub>50</sub> for ZOL at 72 hr was 15.7  $\mu\text{mol/L}$ , whereas those for ZOL plus SMS, and ZOL plus SOM were 14.1, and 13.5  $\mu\text{mol/L}$ , respectively (Fig. 3A). The combination of ZOL and somatostatin analogs did not demonstrate synergistic effects (CI: 0.57–1.00). ZOL induced time-

and dose-dependent proliferative inhibition of NE-CS cells (Fig. 3B). These effects of ZOL were reversed by 20  $\mu\text{mol/L}$  of FOH (Fig. 3C).

### ZOL Inhibits Cell Cycle Activity and Induces Apoptosis of NE-CS Cells

TUNEL-positive cells, indicated in red, increased with increased concentrations of ZOL. On the other hand, Ki-67-positive cells, colored green, decreased (Fig. 4A). We also analyzed the AI and KI with ZOL



**Fig. 3.** Effects of ZOL, SMS and SOM as single agents and in combination on growth of NE-CS cells. Cell viability was measured by WST-8 assay when NE-CS cells were treated with various concentrations of ZOL, SMS and SOM (0.1–100  $\mu\text{mol/L}$ ) for 24, 48 or 72 hr. For combination, the same concentrations of ZOL and SMS or SOM were used. Cell viability was also measured when NE-CS cells were treated for 48 hr with the indicated concentrations (from 0.1 to 100  $\mu\text{mol/L}$ ) of ZOL plus 1, 5, and 20  $\mu\text{mol/L}$  of farnesyl-pyrophosphate ammonium salt (FOH) ( $n = 5/\text{group}$ ). **A:** Cell viability of NE-CS cells at 72 hr in each treatment group. The IC<sub>50</sub> of ZOL at 72 hr for NE-CS cells was 15.7  $\mu\text{mol/L}$  for ZOL, whereas it was 14.1  $\mu\text{mol/L}$  for ZOL plus SMS, and 13.5  $\mu\text{mol/L}$  for ZOL plus SOM. The combination of ZOL and somatostatin analogs did not create synergistic effects. **B:** Cell viability of NE-CS cells in time- and dose-dependent manners. ZOL induced time- and dose-dependent proliferative inhibition of NE-CS cells. **C:** Cell viability of NE-CS cells at 48 hr in ZOL plus FOH. ZOL-induced inhibition was reversed by 20  $\mu\text{mol/L}$  of FOH.

concentrations of 0, 10, 50, and 100  $\mu\text{mol/L}$ . The AI was significantly increased in ZOL 50, and 100  $\mu\text{mol/L}$  compared to the control (means: 55.7 and 136.5, respectively, vs. 13.8) (Fig. 4B). The KI was significantly decreased in ZOL 10, 50, and 100  $\mu\text{mol/L}$  compared to the control (means: 37.6, 22.8, and 1.3, respectively, vs. 68.7) (Fig. 4C).


#### ZOL Inhibits Migration of NE-CS Cells

In addition to effects of ZOL on cell cycle activity and apoptosis, we examined whether ZOL inhibited migration of NE-CS cells, using a Boyden chamber assay. NE-CS cells, with or without ZOL concentrations of 10, and 100  $\mu\text{mol/L}$ , that migrated across the pores at 2, 4, 6 and 8 hr were counted. The numbers of cells migrating  $1\text{ mm}^{-2}$  of membrane were significantly decreased in ZOL 10, and 100  $\mu\text{mol/L}$  (Fig. 5A). When culture medium adding 20  $\mu\text{mol/L}$  of FOH


was incubated in upper chamber, the ZOL-induced inhibition was not appeared (Fig. 5B).

#### ZOL Utilizes the Ras/MAPK Pathway via the Mevalonate Pathway in NE-CS Cells

Since ZOL inhibits farnesyl-pyrophosphate synthase in the mevalonate pathway and impairs prenylation of Ras, we evaluated the effects of ZOL on Ras activity. We used FOH, which potentially induces farnesylation of Ras. As evaluated by pull-down assay, 10, and 100  $\mu\text{mol/L}$  inhibited Ras activation in NE-CS cells, and then the ZOL-induced inhibition was reversed by FOH (Fig. 6). We examined the effects of ZOL on Erk-1/2, which are the terminal proteins of the Ras/MAPK pathway. ZOL inhibited Erk1/2 phosphorylation as evaluated by Western blot assay.




**Fig. 4.** Effects of ZOL on apoptosis and cell activity of NE-CS cells. **A:** TUNEL and anti-Ki67 immunofluorescence were used for NE-CS cells treated with ZOL concentrations of 0, 10, 50, and 100  $\mu\text{mol/L}$  ( $n = 5/\text{group}$ ). DAPI was used to visualize cell nuclei. TUNEL-positive cells, colored red, increased with increased concentrations of ZOL. On the other hand, Ki-67-positive cells, colored green, decreased. **B:** The numbers of TUNEL-positive cells per 1,000 cells apoptotic index (AI) were significantly increased in ZOL 50, and 100  $\mu\text{mol/L}$  compared to the control (means: 55.7, and 136.5, respectively, vs. 13.8). Data are means; bars  $\pm$  SD; \*, significantly different from control group ( $P < 0.001$ ; Student's  $t$ -test). **C:** The numbers of Ki67-positive cells per 1,000 cells (KI: Ki-67 labeling index) were significantly decreased in ZOL 10, 50, and 100  $\mu\text{mol/L}$  compared to the control (means: 37.6, 22.8, and 1.3, respectively, vs. 68.7). Data are means; bars  $\pm$  SD; \*, significantly different from control group ( $P < 0.001$ ; Student's  $t$ -test).



**Fig. 5.** Effects of ZOL on migration of NE-CS cells. Migration assay was performed by using a Boyden chamber ( $n = 3/\text{group}$ ). **A:** In experiment 1, NE-CS cells ( $1 \times 10^5$ ) were placed in the upper chamber with 100  $\mu\text{l}$  of culture medium with or without ZOL (10, 100  $\mu\text{mol/L}$ ). In the lower chamber, 600  $\mu\text{l}$  of culture medium was added. The numbers of cells migrating per 1  $\text{mm}^{-2}$  of membrane were significantly decreased in ZOL 10, and 100  $\mu\text{mol/L}$ . Data are means; bars  $\pm$  SD; \*, significantly different from the control ( $P < 0.001$ ; repeated-measures ANOVA). **B:** In experiment 2, culture medium adding 20  $\mu\text{mol/L}$  of farnesyl-pyrophosphate ammonium salt (FOH) was incubated in the upper chamber. The numbers of cells migrating 1  $\text{mm}^{-2}$  of membrane were not significantly decreased in ZOL 10, and 100  $\mu\text{mol/L}$ . Data are means; bars  $\pm$  SD. \*, Significantly different from the control ( $P < 0.001$ ; repeated-measures ANOVA).

## DISCUSSION

Inappropriate NE regulation in the prostate might facilitate carcinogenesis, proliferation and other tissue changes such as loss of basal cells, angiogenesis, and piling up of prostatic luminal epithelium and invasion, which are characteristic of prostatic carcinoma



**Fig. 6.** Effects of ZOL on Ras/MAPK pathway of NE-CS cells. We used farnesyl-pyrophosphate ammonium salt (FOH), which potentially induces farnesylation of Ras. Ras activity was evaluated by pull-down assay, and Erk activity by Western blot assay. As evaluated by pull-down assay, 10, and 100  $\mu\text{mol/L}$  ZOL inhibited Ras activation in NE-CS cells, and then the ZOL-induced inhibition was reversed by FOH. ZOL inhibited Erk1/2 phosphorylation in NE-CS cells as evaluated by Western blot assay.

[20]. In addition, we previously demonstrated that secretions from NE cells stimulated prostatic cancer cells to achieve androgen-independent growth [21]. Androgen deprivation therapy induces an increased number of NE cells in prostate cancer and the frequency and density of NE cells are more pronounced in CRPC [22]. Thus, the control of NE cells might be important for establishing a treatment strategy for CRPC.

Somatostatin analogs have been used clinically used to treat NE tumors [23]. SMS and lanreotide, which have high affinity to SSTR2a, have been demonstrated to reduce excessive hormone production and accompanying symptoms from carcinoid tumors and pancreatic endocrine tumors such as glucagonoma, VIPoma and gastrinoma [14]. The anticancer effect may be the result of antiproliferative and apoptotic actions through direct and indirect mechanisms. The direct mechanism is mediated by SSTR on tumor cells, and suppression of secretion of several growth factors such as insulin-like growth factor-1 (IGF-1) may also indirectly inhibit the tumor growth [24–26]. In this study, in spite of the expression of SSTR2a and SSTR5 in our NE carcinoma models, we failed to find significant antiproliferative effects of SMS or SOM monotherapy in vitro or in vivo. In addition, the combination therapy with ZOL did not create a synergistic effect. Although, the exact reason is unclear, both