




ELSEVIER

Available online at www.sciencedirect.com

SciVerse ScienceDirect

journal homepage: www.elsevier.com/locate/jccase



Case Report

Acute pulmonary embolism induced by renal obstruction with benign prostatic hyperplasia: Case report

Masanari Umemura (MD, PhD)^{a,*}, David Ho (MD)^a, Naoki Nozawa (MD)^b, Erdene Balginnyam (MD, PhD)^a, Kousaku Iwatsubo (MD, PhD)^a, Toshihiko Saito (MD, PhD)^c, Tsutomu Endo (MD, PhD)^c, Yoshihiro Ishikawa (MD, PhD, FJCC)^d, Satoshi Umemura (MD, PhD, FJCC)^e, Kazuo Kimura (MD, PhD, FJCC)^f

^a Cardiovascular Research Institute, Department of Cell Biology and Molecular Medicine, New Jersey Medical School-University of Medicine and Dentistry of New Jersey, 185 South Orange Avenue, Newark, NJ 07103, USA

^b Nozwa naika Jyunkankika, Yokohama, Japan

^c Department of Cardiovascular Division, Saiseikai Yokohamashi Nanbu Hospital, Japan

^d Cardiovascular Research Institute, Yokohama City University School of Medicine, Graduate School of Medicine, Yokohama, Japan

^e Yokohama City University, Graduate School of Medicine, Department of Medical Science and Cardiorenal Medicine, Yokohama, Japan

^f Division of Cardiology, Yokohama City University Medical Center, Yokohama, Japan

Received 20 September 2011; received in revised form 13 October 2011; accepted 18 October 2011

KEY WORDS

Acute pulmonary embolism;
Urinary distension;
Prostatic hyperplasia;
Deep vein thrombosis

Summary

Background: We report a rare case of acute pulmonary embolism (PE) induced by urinary retention and bladder distention with benign prostatic hyperplasia (BPH).

Case report: A 76-year-old male with BPH presented to the hospital with anuria of 24 h duration and abdominal distention. Physical examination revealed tenderness and distention of the lower abdomen and a swollen right leg. Echocardiography after urethral catheterization showed a large free-floating thrombus traversing back and forth through the tricuspid orifice. Computed tomographic angiography demonstrated filling defects at the level of the right inter lobar pulmonary artery and the segmental branches of both pulmonary arteries, indicating acute PE. The patient was treated with heparin and warfarin for three weeks to ensure the resolution of the pulmonary embolus. After the resolution of all symptoms, the patient was discharged without further complication.

* Corresponding author. Tel.: +1 973 972 8925; fax: +1 973 972 7489.

E-mail addresses: umemura0807@gmail.com, m.umemura.0807@gmail.com (M. Umemura).



Conclusion: This case suggested that a distended bladder is a potential risk factor for the development of deep vein thrombosis and PE.

© 2011 Japanese College of Cardiology. Published by Elsevier Ltd. All rights reserved.

Introduction

Pulmonary embolism (PE) is a blockage of one or more arteries in lungs by air, fat, tumor tissue, or thrombus. Approximately 90% of PE arises from deep vein thrombosis (DVT) [1]. DVT risk factors include surgery, hospitalization, immobilization, smoking, obesity, age, medication, thrombophilia, or pregnancy [2]. Although bladder distention could also cause obstruction of veins in the pelvis, and thus subsequent DVT [3–5], it is rare for bladder distention to induce PE. In this regard, it remains unknown whether benign prostatic hyperplasia (BPH), which can also induce bladder distention [6], can lead to PE. Here, we report a PE case with a history of BPH accompanied by severe urinary retention and bladder distention.

Case report

A 76-year-old man was brought to our hospital with lower abdominal distention. He had noticed the symptoms 4 days prior to the visit. His past medical history was significant for chronic atrial fibrillation without chronic anticoagulation therapy. He had no prior history of heart failure or thrombosis. He never smoked and his family history was negative for clotting disorder or thrombosis.

On admission, his body mass index was 24 kg/m². Vital signs included a mildly elevated blood pressure at 134/79 mmHg, with irregular pulses at a rate of 70 beats/min, and peripheral oxygen saturation of 98% on room air. His body temperature at the time of presentation was 37°C. Physical examination was unremarkable except for a tender and rigid abdomen without guarding or rebound as well as a swollen right lower leg. Laboratory tests were consistent with dehydration, renal dysfunction, and inflammation as follows. Significant findings included a white blood cell count of 16,300 cells/ μ l. C-reactive protein was 9.50 mg/dl. Blood urea nitrogen was 88 mg/dl and serum creatinine was 2.34 mg/dl. Qualitative analysis of urine showed the presence of white blood cells. Prothrombin and partial thromboplastin times were normal, but D-dimer was increased (86 μ g/ml: normal range < 1 μ g/ml). The levels of plasma protein C and protein S were within normal limits, and anti-cardiolipin antibody was negative. The serum level of prostatic specific antigen was increased (14.1 μ g/ml: normal range < 4 μ g/ml). Chest X-ray showed no abnormality. Electrocardiogram was consistent with atrial fibrillation, and unchanged from prior recording taken a month earlier. Echocardiography was significant for mild concentric left ventricular hypertrophy with normal left ventricular function, and no thrombosis in his heart and no signs of right ventricular pressure overload were found. Abdominal ultrasound examination revealed an extended urinary bladder. Ultrasound examination for vein of lower extremity showed a thrombus in the right femoral vein and no thrombus in the left. Taken together, urinary distention secondary to

BPH, obstruction of pelvic vein within the pelvis by the distended urinary bladder, urinary tract infection, acute post-renal failure, and right DVT were suspected. Urinary catheterization was then performed to relieve the urinary obstruction.

However, when echocardiography was repeated 1 h after the urinary catheterization, we found a large floating thrombus within the right atrium, with frequent back and forth movement through the tricuspid orifice, which was not observed previously (Fig. 1A and B). Mild pressure overload of the right ventricle was also observed at this time. The patient was immediately brought to our intensive care unit. Follow-up echocardiography, however, showed that the observed thrombus in the right atrium had disappeared. Emergent computed tomographic (CT) angiography revealed clots at the level of right distal pulmonary artery, and the segmental branches of both pulmonary arteries (Fig. 2A–C). Pelvis CT showed that there was BPH. Chest and abdominal CT showed that there were no thrombi in either of the femoral veins or the inferior vena cava. Screening for malignancy with CT was negative.

Pulmonary perfusion scintigraphy showed diffuse filling defects in both lungs, which were consistent with findings of CT angiography (Fig. 3A and B). Lower extremity perfusion scintigraphy showed a delay of disappearance of ^{99m}Tc-MAA in his right leg compared to the left leg (Fig. 3C and D). Accordingly, we diagnosed that the patient had developed PE.

Treatment for PE with continuous intravenous infusion of heparin (14,000 units/day) and oral warfarin (3 mg/day) was started. We did not perform intravenous infusion of urokinase or tissue plasminogen activator because the patient showed no sign of hemodynamic instability or severe right ventricular pressure overload. On day 5 during the hospitalization, heparin was discontinued because the prothrombin time/international normalized ratio had reached 2.0. On day 14, CT showed reduced size of the thrombus (20% of the initial size). Regarding BPH, both suppressive therapy for BPH and intermittent self-catheterization were initiated. On day 19, the patient became symptom free and was discharged.

Discussion

There are few reports of urinary distention-induced DVT [3–5,7], although urinary retention itself can lead to increased venous pressure in the pelvis [7]. Accordingly, it is not well accepted that BPH can serve as a risk factor for PE through urinary retention and bladder distention. A review reported that 9 of 15 urinary distention cases, which led to DVT, had prostatic enlargement [8,9]. Meinardi et al. previously reported a case of BPH-induced DVT with protein S deficiency [5]. These authors concluded that BPH alone may be insufficient to induce PH unless other risk factors, such as thrombophilic defects, are accompanied. However,




Figure 1 Echocardiogram of axis view (A) and four chamber view (B). A giant floating thrombus (3 cm × 3 cm, white arrows) in the right atrium and ventricle frequently moved back and forth through the tricuspid orifice. RV, right ventricle.

as shown in the current and another case [10], BPH alone can induce DVT without any other risk factors. It is thus tentative to speculate that BPH should be recognized as a potential risk factor for DVT via urinary bladder distention. In this case, it should be noted that hypercoagulability, due to increased inflammation along with urinary tract infection,

might contribute to thrombus formation. More interestingly, bilateral lower extremity swelling is usual with such bladder distention; a previous review summarized that 14 of 15 patients had bilateral swelling [8] and that the occurrence of PE following DVT was uncertain. This is in contrast to the current case, which showed unilateral leg swelling




Figure 2 Chest computed tomography (CT) shows clots and disturbed blood flow in right distal pulmonary artery and left segmental branch (A–C). Clots are shown by arrowheads. Pelvic CT shows enlarged prostate (D, arrow).




Figure 3 Pulmonary perfusion scintigraphy showed filling defects in both lung [A (anterior view) and B (posterior view)]. Perfusion scintigraphy of lower extremities showed a delay of disappearance of ^{99m}Tc -MAA in his right leg compared to his left leg [C (early phase) and D (delay phase)].

followed by PE. When unilateral leg swelling is observed with urinary distention, DVT and potentially PE might be considered as well. Before urinary catheterization was performed, formation of thrombus due to distended bladder was not listed as a differential diagnosis because it is clinically rare. Placing temporary inferior vena cava might be effective for inhibition of PE, thus should be considered in the case of leg swelling concurrent with distended bladder.

Prognosis of PE is typically poor. The conventional treatments for PE include urokinase, heparin, tissue plasminogen activator, and warfarin [2]. We used warfarin in this case but not urokinase or tissue plasminogen activator because hemodynamic instability or severe right ventricular pressure overload were not found. Consequently, the patient was discharged without further complications. Factors that brought this fortunate outcome may include the PE onset within the hospital and thus the rapid initiation of heparin and warfarin therapy.

Conclusion

BPH could potentially cause DVT and thus PE via bladder distention.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.jccase.2011.10.004.

References

- [1] Enden T, Kløw NE. CT pulmonary angiography and suspected acute pulmonary embolism. *Acta Radiol* 2003;44:310–5.
- [2] Guidelines on diagnosis and management of acute pulmonary embolism. Task Force on Pulmonary Embolism, European Society of Cardiology. *Eur Heart J* 2000;21:1301–36.
- [3] Dyer JR, Molton M, Civil M, Davis TM. An unusual cause of deep venous thrombosis. *Med J Aust* 1993;158:648.
- [4] Andrew WK. An unusual cause of deep venous thrombosis of the lower limb. *S Afr Med J* 2000;90:42.
- [5] Meinardi JR, Kremer J, van der Meer J. Deep vein thrombosis associated with distension of the urinary bladder due to benign prostatic hypertrophy—a case report. *Neth J Med* 2004;62:137–8.
- [6] Meigs JB, Barry MJ, Giovannucci E, Rimm EB, Stampfer MJ, Kawachi I. Incidence rates and risk factors for acute urinary retention: the health professionals followup study. *J Urol* 1999;162:376–82.

- [7] Hopkins WF, Mattens WA, Pierce Jr JM. Increased venous pressure and edema in the lower extremities secondary to urinary retention. *Invest Urol* 1965;3:117–21.
- [8] Evans JM, Owens Jr TP, Zerbe DM, Rohren CH. Venous obstruction due to a distended urinary bladder. *Mayo Clin Proc* 1995;70:1077–9.
- [9] Ito M, Hatta K, Miyakawa K, Arai H. Pulmonary embolism from persistent dilatation of the bladder secondary to anticholinergic side effects. *Gen Hosp Psychiatry* 2009;31:187–9.
- [10] Karanth NS, Crausman RS, De Palo VA. Deep venous thrombosis as a consequence of benign prostatic hypertrophy. *Med Health R I* 2005;8:18–9.



Acute myocardial infarction with isolated conus branch occlusion

Masanari Umemura, MD, PhD,^{a,*} David Ho, MD,^a Naoki Nozawa, MD,^b
Erdene Balginnyam, MD, PhD,^a Kousaku Iwatsubo, MD, PhD,^a Thosihiko Saito, MD, PhD,^c
Tsutomu Endo, MD, PhD,^c Yoshihiro Ishikawa, MD, PhD,^d Satoshi Umemura, MD, PhD,^e
Kazuo Kimura, MD, PhD^f

^aCardiovascular Research Institute, Department of Cell Biology and Molecular Medicine,
New Jersey Medical School-University of Medicine and Dentistry of New Jersey, Newark, NJ, USA

^bNozwa naika Jyunkankika, Yokohama, Kanagawa, Japan

^cDepartment of Cardiovascular Division, Saiseikai Yokohamashi Nanbu Hospital, Yokohama, Kanagawa, Japan

^dCardiovascular Research Institute Yokohama City University School of Medicine, Graduate School of Medicine, Yokohama, Kanagawa, Japan

^eYokohama City University, Graduate School of Medicine Department of Medical Science and Cardiorenal Medicine, Yokohama, Kanagawa, Japan

^fDivision of Cardiology, Yokohama City University Medical Center, Yokohama, Kanagawa, Japan

Received 18 October 2011

Abstract

There are few reports of acute myocardial infarction (AMI) relating to the occlusion of the conus branch, most of which are iatrogenic in nature. So far as we are concerned, this is the first case of spontaneous AMI with isolated conus branch occlusion. Electrocardiogram (ECG) showed mild elevation of ST segment in leads V₁ through V₃. Cardiac markers of myocardial infarction were positive. Right coronary angiography revealed an isolated occlusion of the conus branch. Penetration of the guidewire in the occluded lesion was attempted, and recanalization was successfully achieved. The patient was discharged without any adverse events.

© 2012 Elsevier Inc. All rights reserved.

Keywords:

Acute myocardial infarction; Conus branch; Occlusion; ST elevation; Spontaneous; Isolated

Introduction

The conus branch of the right coronary artery (RCA) frequently provides blood supply to the outflow tract of right ventricle. Previous papers reported that characteristic electrocardiographic (ECG) changes (elevation of ST segment in the right precordial leads) appeared when the conus branch was occluded or narrowed. All these reported events occurred during coronary angiography/angioplasty or cardiac surgery in the RCA.^{1–5} Here, unlike these iatrogenic occlusions, we show the first case of spontaneous occlusion of the conus branch. The isolated conus branch occlusion led to acute myocardial infarction (AMI), and we treated the case with successful recanalization using penetration with a guidewire, instead of balloon/stent catheter.

Case report

A 78-year-old Japanese man awoke with retrosternal chest pain. The patient had been on home oxygen therapy for severe pulmonary emphysema and been on maintenance hemodialysis for the past 2 years. He had no history of coronary artery disease, diabetes, thrombosis, or atrial fibrillation and had never received anticoagulation therapy. Other risk factors included a history of tobacco use and smoking 20 cigarettes a day for 60 years. Family history was negative for clotting disorder, stroke, or coronary artery disease.

One hour after the onset of the chest pain, the patient visited an outside clinic. The electrocardiogram showed elevation of ST segment in the precordial leads (V₁ through V₃) (Fig. 1A). Sublingual nitroglycerin was given, and the patient was transferred to the emergency room of our hospital. At presentation, the patient had sustained chest pain that was not reduced by the nitroglycerin given at the outside clinic.

On admission, the blood pressure was 152/66 mm Hg and the heart rate was 82 beats per minute. The result of the physical examination was unremarkable. The electrocardiogram showed mild ST-segment elevation in the leads V₁ through V₃ (Fig. 1B). The maximum elevation of ST segment

* Corresponding author. Cardiovascular Research Institute, Department of Cell Biology and Molecular Medicine, New Jersey Medical School-University of Medicine and Dentistry of New Jersey, MSB G645, 185 South Orange Avenue, Newark, NJ 07103, USA.

E-mail address: m.umemura.0807@gmail.com




Fig. 1. (A) Electrocardiogram of the conus branch occlusion. (B) Electrocardiogram of the patient showed atrioventricular block 1 month before the onset of AMI. (C) Elevation of ST segment in leads V₁ through V₃ was seen in the outside clinic after sublingual nitroglycerin was given, and (D) resolution of ST-segment elevation after the recanalization using the guidewire.

was 2 mm in the lead V₂ (Fig. 1C) recorded 3 hours after the onset of the chest pain as compared with the ECG 1 month before the admission (Fig. 1A). White blood cell count was 3600 cells/ μ L. C-reactive protein level was 2.69 mg/dL. Serum creatinine level was 6.48 mg/dL. Cardiac markers were as follows: creatinine kinase level of 138 IU/L, creatinine kinase-MB fraction of 15 IU/L, and troponin T level of 0.37 ng/mL. Echocardiography showed relatively normal segmental wall motions of both right and left ventricles. At that point, acute coronary syndrome was strongly suspected, and coronary angiography was immediately performed. Complete occlusion of the conus branch of the RCA was found (Fig. 2A and B). Because coronary angioplasty was considered to be difficult in this lesion, penetration of guidewire was attempted. The guidewire was easily crossed through the occluded lesion (Fig. 2C), and Thrombolysis in Myocardial Infarction Trial (TIMI) grade III flow was restored as demonstrated in the following coronary angiography (Fig. 2D), suggesting successful recanalization. The resolution of chest pain was immediate. Intravenous infusion of heparin with oral warfarin (3 mg/day) and aspirin (100 mg/day) was initiated. The electrocardiogram taken at 24 hours after recanalization showed the resolution of ST elevations in leads V₁ through V₃ without the development of Q-wave or T-wave inversion (Fig. 1D). Creatinine kinase level peaked at 1066 IU/L with an MB fraction of 137 IU/L 1 day after the recanalization. On day 13, the patient was discharged with aspirin and warfarin. Although CHADS₂

score is 1, we added warfarin to the patient because we considered the possibility of his hypercoagulability due to severe emphysema and his maintenance of hemodialysis. It was also likely that thromboembolism caused myocardial infarction, not arteriosclerosis.

The characteristic ECG changes (elevation of ST segment in leads V₁ through V₃) observed in this case are consistent with those in the previous reports of conus branch occlusion.^{3,5} Matthews⁵ reported that conus branch was occluded by injection to RCA during angiography. Eichhofer et al⁴ reported that stent implantation in the right proximal coronary artery resulted in conus branch occlusion. The electrocardiogram showed similar pattern in the precordial leads (V₁ through V₃). These reports did not mention treatments, presumably because it is considered difficult to perform angioplasty or stent implantation for conus branch occlusion due to the small diameter of the artery. Successful balloon angioplasty in the conus branch was reported in a case of occlusion by suture material after mitral valve reconstruction.² On the contrary, occlusion of the conus branch is known to show similar ECG pattern as those seen with Brugada syndrome, that is, elevations of the ST segment in the right precordial leads (V₁ through V₃). There are 2 reports of Brugada-like ECG changes due to conus branch vasospasms with acetylcholine, which resulted in ventricular fibrillation or ventricular tachycardia.^{3,6} Because Brugada syndrome can lead to ventricular fibrillation,




Fig. 2. Coronary angiograms before and after the penetration of guidewire into the conus branch occlusion. The conus branch of the RCA showed total occlusion (arrows in A and B). The TIMI flow grade was 0 at this point. A guidewire was penetrated into the conus branch lesion (arrowhead in C), and grade III TIMI flow was observed (D). RAO indicates right anterior oblique; LAO, left anterior oblique.

and thus sudden death, treatment of conus branch occlusion is of clinical importance. In our case, penetration of a guidewire was attempted, and blood flow was restored without development of ventricular fibrillation or tachycardia. To our knowledge, this is the first report of successful recanalization with a guidewire in a conus branch occlusion. Because insertion of a guidewire into the conus branch may be easier than that of balloon/stent catheter, guidewire penetration should be considered as the first option for treatment of isolated conus branch occlusion. Furthermore, conus branch occlusion described in the previous reports were unexceptionally iatrogenic in nature; spontaneous occlusion like our case has not been reported. Nonetheless, all cases, including the current case, showed similar ECG changes, that is, ST elevation in the right precordial leads (V_1 through V_3) and moderate elevation in creatinine kinase (300–1000 IU/L). Meanwhile, the conus artery in about 50% of human hearts is not a branch of the RCA but arises from the right sinus ostium of Valsalva directly. Therefore, when ST elevations in the right precordial leads are observed, not only the conus branch of RCA but also the conus artery occlusion should be considered. Nonetheless, when angioplasty is considered to be difficult in either cases, penetration with a guidewire could be an alternative optional for

treatment. Because this artery is important in terms of critical ventricular arrhythmia development, such treatment should be performed immediately even though the putative irrigation area may not be very large.

References

1. Celik T, Yuksel US, Kursaklioglu H, et al. Precordial ST-segment elevation in acute occlusion of the proximal right coronary artery. *J Electrocardiol* 2006;39:301.
2. Hernández FH, Garrido-Lestache EB, Arribas P, et al. Recurrent ventricular fibrillation and ST segment elevation in the right precordial leads due to acute occlusion of the conus branch. *Rev Esp Cardiol* 2011 [in press].
3. Yamaki M, Sato N, Myojo T, et al. Possible contribution of ischemia of the conus branch to induction or augmentation of Brugada type electrocardiographic changes in patients with coronary artery disease. *Int Heart J* 2010;51:68.
4. Eichhofer J, Curzen N. Images in cardiovascular medicine. Unexpected profound transient anterior ST elevation after occlusion of the conus branch of the right coronary artery during angioplasty. *Circulation* 2005;111:e113.
5. Matthews RV, Oesterle SN. Precordial ST elevation with acute conus branch occlusion. *Cathet Cardiovasc Diagn* 1989;17:180.
6. Ogano M, Iwasaki Y, Morita N, et al. Proarrhythmic ECG deterioration caused by myocardial ischemia of the conus branch artery in patients with a Brugada ECG pattern. *Pacing Clin Electrophysiol* 2011;34:e26.



Mice Lacking Hypertension Candidate Gene ATP2B1 in Vascular Smooth Muscle Cells Show Significant Blood Pressure Elevation

Yusuke Kobayashi, Nobuhito Hirawa, Yasuharu Tabara, Hidenori Muraoka, Megumi Fujita, Nobuko Miyazaki, Akira Fujiwara, Yasuhiro Ichikawa, Yuichiro Yamamoto, Naoaki Ichihara, Sanae Saka, Hiromichi Wakui, Shin-ichiro Yoshida, Keisuke Yatsu, Yoshiyuki Toya, Gen Yasuda, Katsuhiko Kohara, Yoshikuni Kita, Kohtaro Takei, Yoshio Goshima, Yoshihiro Ishikawa, Hirotsugu Ueshima, Tetsuro Miki and Satoshi Umemura

Hypertension. 2012;59:854-860; originally published online February 6, 2012;

doi: 10.1161/HYPERTENSIONAHA.110.165068

Hypertension is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2012 American Heart Association, Inc. All rights reserved.

Print ISSN: 0194-911X. Online ISSN: 1524-4563

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://hyper.ahajournals.org/content/59/4/854>

Data Supplement (unedited) at:

<http://hyper.ahajournals.org/content/suppl/2012/02/03/HYPERTENSIONAHA.110.165068.DC1.html>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Hypertension* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the Permissions and Rights Question and Answer document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Hypertension* is online at:
<http://hyper.ahajournals.org//subscriptions/>

Mice Lacking Hypertension Candidate Gene ATP2B1 in Vascular Smooth Muscle Cells Show Significant Blood Pressure Elevation

Yusuke Kobayashi, Nobuhito Hirawa, Yasuharu Tabara, Hidenori Muraoka, Megumi Fujita, Nobuko Miyazaki, Akira Fujiwara, Yasuhiro Ichikawa, Yuichiro Yamamoto, Naoaki Ichihara, Sanae Saka, Hiromichi Wakui, Shin-ichiro Yoshida, Keisuke Yatsu, Yoshiyuki Toya, Gen Yasuda, Katsuhiko Kohara, Yoshikuni Kita, Kohtaro Takei, Yoshio Goshima, Yoshihiro Ishikawa, Hirotsugu Ueshima, Tetsuro Miki, Satoshi Umemura

Abstract—We reported previously that ATP2B1 was one of the genes for hypertension receptivity in a large-scale Japanese population, which has been replicated recently in Europeans and Koreans. ATP2B1 encodes the plasma membrane calcium ATPase isoform 1, which plays a critical role in intracellular calcium homeostasis. In addition, it is suggested that ATP2B1 plays a major role in vascular smooth muscle contraction. Because the ATP2B1 knockout (KO) mouse is embryo-lethal, we generated mice with vascular smooth muscle cell-specific KO of ATP2B1 using the Cre-loxP system to clarify the relationship between ATP2B1 and hypertension. The KO mice expressed significantly lower levels of ATP2B1 mRNA and protein in the aorta compared with control mice. KO mice showed significantly higher systolic blood pressure as measured by tail-cuff method and radiotelemetric method. Similar to ATP2B1, the expression of the Na⁺-Ca²⁺ exchanger isoform 1 mRNA was decreased in vascular smooth muscle cells of KO mice. However, ATP2B4 expression was increased in KO mice. The cultured vascular smooth muscle cells of KO mice showed increased intracellular calcium concentration not only in basal condition but also in phenylephrine-stimulated condition. Furthermore, phenylephrine-induced vasoconstriction was significantly increased in vascular rings of the femoral artery of KO mice. These results suggest that ATP2B1 plays important roles in the regulation of blood pressure through alteration of calcium handling and vasoconstriction in vascular smooth muscle cells. (*Hypertension*. 2012;59:854-860.) • **Online Data Supplement**

Key Words: hypertension ■ ATP2B1 ■ Cre-loxP system ■ blood pressure ■ Millennium Genome Project ■ Global Blood Pressure Genetics

Numerous studies have attempted to identify genetic markers for hypertension over the past 2 decades, but no cross-validated loci in different ethnic groups have thus far been identified except for the mendelian forms of hypertension.¹ In the Millennium Genome Project² we identified single nucleotide polymorphisms located upstream or within the ATP2B1 gene as strong susceptible polymorphisms for hypertension in Japanese. Some of these findings have been replicated in individuals of European descent in the Global Blood Pressure Genetics sample and have also been validated in other studies in individuals of European descent,³ Koreans,⁴⁻⁶ and Japanese.⁷ The single nucleotide polymorphisms of ATP2B1 identified in these studies showed a significant

association with hypertension in various large-scale study populations with different methods, genome-wide association study in the Cohorts for Heart and Aging Research in Genomic Epidemiology Consortium and the Korean study and candidate gene analysis in our previous study. However, the functional roles of ATP2B1 in blood pressure control have not yet been proven in vivo. The ATP2B1-null mutant mouse has been reported to be embryolethal⁸; thus, we need to make a conditional knockout (KO) mouse model of ATP2B1 using the Cre-loxP system to reveal the function of the gene. Because the ATP2B1 gene encodes one of the calcium pumps and plays an important role in contraction of bladder smooth muscle,⁹ we selected vascular smooth

Received October 25, 2010; first decision November 17, 2010; revision accepted January 15, 2012.

From the Departments of Medical Science and Cardiorenal Medicine (Y.K., M.F., N.M., A.F., N.I., S.S., H.W., S.Y., Y.T., S.U.) and Molecular Pharmacology and Neurobiology (H.M., K.T., Y.G.) and Cardiovascular Research Institute (Y.Ic., Y.Is.), Yokohama City University Graduate School of Medicine, Yokohama, Japan; Division of Nephrology and Hypertension (N.H., Y.Y., K.Y., G.Y.), Yokohama City University Medical Center, Yokohama, Japan; Department of Geriatric Medicine (Y.T., K.K., T.M.), Ehime University Graduate School of Medicine, Toon, Japan; Department of Health Science (Y.K., H.U.), Shiga University of Medical Science, Otsu, Japan.

The online-only Data Supplement is available with this article at <http://hyper.ahajournals.org/lookup/suppl/doi:10.1161/HYPERTENSIONAHA.110.165068/-DC1>.

Correspondence to Nobuhito Hirawa, Division of Nephrology and Hypertension, Yokohama City University Medical Center, 45-7 Urafune-cho, Minami-ku, Yokohama 232-0024, Japan. E-mail hirawa@yokohama-cu.ac.jp

© 2012 American Heart Association, Inc.

Hypertension is available at <http://hyper.ahajournals.org>

DOI: 10.1161/HYPERTENSIONAHA.110.165068

muscle cells (VSMCs) as target tissue of KO. Because we have already demonstrated that ATP2B1 mRNA expression in human umbilical artery smooth muscle cells was significantly lower in those having the risk allele for hypertension than in those with no risk allele,¹⁰ we hypothesized that VSMC ATP2B1 KO mice would exhibit high blood pressure. In the present study, we made the VSMC-specific ATP2B1 KO mice and evaluated their blood pressure and related mechanisms.

Materials and Methods

Animal Care

Animals were housed under a 12-hour light-dark cycle at a temperature of 25°C. Tap water was provided ad libitum. Experiments were conducted under the guidelines for animal experiments set by the animal experiment committee of Yokohama City University School of Medicine.

Creation of VSMC-Targeted ATP2B1 KO Mice

ATP2B1^{loxP/loxP} mice were generated using the Cre-loxP and flippase recombination enzyme-flippase recognition target (FLP-FRT) recombination system. ATP2B1 is encoded by 21 exons on chromosome 10, and mice systemically deficient in exon 10 are reported to be embryolethal.⁸ We, therefore, designed a vector to KO exon 10 of the ATP2B1 gene. The detailed technical strategy for conditional KO mouse generation is described in the Methods section of the online-only Data Supplement. To target inactivation of the ATP2B1 gene to VSMCs, ATP2B1^{loxP/loxP} mice were intercrossed with SM22-Cre transgenic mice (see details in the online-only Data Supplement) expressing Cre recombinase under control of the mouse transgelin (smooth muscle protein 22- α) promoter. The resulting ATP2B1^{loxP/-}/SM22-Cre animals were further mated with ATP2B1^{loxP/loxP} mice to generate ATP2B1^{loxP/loxP}/SM22-Cre (VSMC ATP2B1 KO) mice and ATP2B1^{loxP/loxP} mice without SM22-Cre (control mice). Animals used for experiments were backcrossed ≥ 6 times.

BP Measurement by Tail-Cuff Method and Radiotelemetric Method

Systolic blood pressure was measured by the tail-cuff method (BP-monitor MK-2000; Muromachi Kikai Co) at the age of 8 weeks and 22 weeks, as described previously.^{11,12} Furthermore, direct blood pressure measurement was performed by a radiotelemetric method in which a blood pressure transducer (PA-C10, Data Sciences International) was inserted into the left carotid artery. Ten days after transplantation, each mouse was housed individually in a standard cage on a receiver under a 12-hour light-dark cycle. Direct blood pressure was recorded every minute by radiotelemetry, as described previously.¹³

Real-Time Quantitative RT-PCR Analysis

Total RNA was extracted from the aorta or cultured VSMCs with ISOGEN (Nippon Gene), and cDNA was synthesized using the SuperScript III First Strand System (Invitrogen). Real-time quantitative RT-PCR was performed by incubating the reverse-transcription product with TaqMan PCR Master Mix and a designed TaqMan probe (Applied Biosystems).¹² RNA quantities were expressed relative to the 18S mRNA control.

Western Blot Analysis of ATP2B1

Western blot analysis was performed as described previously.¹¹ Further details of the Western blot analysis are described in the online-only Data Supplement.

Cell Culture of Mouse VSMCs and Measurement of Intracellular Calcium Concentration

VSMCs were aseptically isolated from thoracic aortic explants of an 8-week-old ATP2B1 KO mouse and its wild-type littermate, as described previously.¹⁴ Further details of the cell culture of mouse VSMCs are described in the online-only Data Supplement. Measurement of basal condition and phenylephrine-stimulated changes in intracellular calcium concentration were assessed by Fura-2 fluorescence ratio imaging using a microscopic digital imaging system (IX71, Olympus), as described previously.¹⁵ Briefly, ATP2B1 KO or control VSMCs grown on 25-mm coverslips were loaded with the calcium-specific dye Fura-2-acetoxymethyl ester (2.5 $\mu\text{mol/L}$, Invitrogen) and 0.01% Pluronic acid (Invitrogen) for 30 minutes at 37°C. After washing with the Hank balanced salt solution, cells were incubated for 20 minutes at 37°C in the Hank balanced salt solution to allow complete hydrolysis of Fura-2-acetoxymethyl ester to Fura-2. Emissions fluorescence was measured with a CCD camera (U-PMTVIX, Olympus) at a wavelength of 510 nm. Real-time shifts in Fura-2 ratio fluorescence (ratio of emissions: F340:F380), indicating changes in intracellular calcium concentration, were recorded before, during, and after stimulating VSMCs with 10⁻⁶ M phenylephrine (Sigma Aldrich), and we used calcium ionophore A23187 (Calbiochem) as positive control for the accuracy of the intracellular calcium concentrations. Summary data represent the average difference in basal condition and the peak increase in phenylephrine-induced intracellular calcium concentration.

Isometric Tension of Vascular Rings of Femoral Artery

We measured isometric tension of femoral artery vascular rings from KO mice and control mice, as described previously.¹⁶ Phenylephrine and potassium-enriched solution were added to stimulate vasoconstriction. Further details of the vasoconstriction assay are described in the online-only Data Supplement.

Statistical Analysis

For statistical analysis of differences between groups, Mann-Whitney U test or ANOVA followed by Bonferroni method was used. All of the quantitative data are expressed as mean \pm SE. Values of $P < 0.05$ were considered statistically significant.

Results

High Efficiency, VSMC-Selective Deletion of ATP2B1 Gene

Figure 1A shows the Southern blot analysis of tail DNA obtained from VSMCs. ATP2B1 KO mice demonstrated a deletion event occurring in VSMCs within the vascular bed of the tail. Quantitative RT-PCR analysis demonstrated that expression of ATP2B1 mRNA in isolated aorta of VSMC ATP2B1 KO mice was reduced by 80% to 90% compared with that in control mice (Figure 1B). Similarly, Western blot analysis showed that ATP2B1 protein in isolated aorta of VSMC ATP2B1 KO mice was reduced by 80% compared with that in control mice (Figure 1C and 1D).

VSMC ATP2B1 KO and Control Mice Were Both Born at Expected Mendelian Ratio

As shown in Figure 1E, both VSMC ATP2B1 KO mice and control mice, male and female, were born at the expected mendelian ratio and could not be distinguished from one another at birth.

No Difference in Growth Between VSMC ATP2B1 KO Mice and Control Mice

Body weight was measured at 8, 12, 22, and 36 weeks after birth. As seen in Figure 1F, there was no difference in




Figure 1. High efficiency, vascular smooth muscle cell (VSMC)-selective deletion of ATP2B1 gene. **A**, Tail DNA was prepared by standard methods, and the ATP2B1 gene was amplified by PCR using the forward primer 5'-CATCCTCTTAGTTATTAAGGAAGCAGT-3' (located in the intron before the first loxP site) and reverse primer 5'-GCCTTTTACAGCATGAACATAGCGA-3' (located in the intron after the second loxP site). The presence of wild-type ATP2B1 (W), floxed ATP2B1 (F), and recombined ATP2B1 (X) was determined using the forward primer and reverse primer, generating products of 1282 bp for W, 1442 bp for F, and 399 bp for X. **B**, ATP2B1 mRNA expression in aorta of 8-week-old mice ($n=6$ for each genotype) as quantified by quantitative RT-PCR using exon 10 and 11 amplification. Data are presented as mean and SE from 6 independent experiments ($P<0.0001$). **C** and **D**, ATP2B1 protein expression in aorta of VSMC ATP2B1 knockout (KO) mice and control mice estimated by immunoblot analysis ($n=5$ for each genotype). One representative of 5 independent experiments is shown. Data are presented as mean and SE. **E**, Birth rate of mating, which is expected to have the same ratio of births. The data were collected from 632 mice born by the mating of VSMC ATP2B1 KO mice and control mice. **F**, Growth curve of VSMC ATP2B1 KO mice and control mice. The data were all collected from male mice. Weights for each genotype represent mean and SE ($n=12-24$ for each genotype; C indicates control mice; K, VSMC ATP2B1 KO mice). The data collected show that there was no difference in body weight alterations.

alteration of body weight between VSMC ATP2B1 KO mice and control mice.

VSMC ATP2B1 KO Mice Showed Higher Blood Pressure Than Control Mice Under Resting Conditions

To ascertain whether deletion of ATP2B1 in VSMCs affects blood pressure, conscious VSMC ATP2B1 KO mice and control mice were subjected to blood pressure measurements by the tail-cuff method. All of the experiments were carried out in a blinded manner on male mice eating standard rodent chow (0.3% NaCl). Under resting conditions, VSMC ATP2B1 KO mice displayed higher systolic blood pressure than that of control mice at 8 and 22 weeks of age (Figure 2A). Heart rate did not differ significantly between the groups (data not shown).

VSMC ATP2B1 KO Mice Showed Higher Blood Pressure Assessed by 24-Hour Radiotelemetric System Than Control Mice

To confirm the effects of deletion of VSMC ATP2B1 on blood pressure and to analyze the circadian pattern of blood pressure, conscious VSMC ATP2B1 KO mice and control mice were subjected to blood pressure measurements by radiotelemetry. KO mice showed higher blood pressure than control mice at 14 weeks of age throughout the day (systolic

blood pressure, Figure 2B; diastolic blood pressure, Figure S9A; mean blood pressure, Figure S9B), whereas circadian variations in heart rate did not differ significantly between the groups (Figure 2C).

Expressions of Calcium-Regulatory Genes in Cultured VSMCs of ATP2B1 Mice

The same as for the *in vivo* results, expression of both ATP2B1 (0.07-fold $P<0.0001$; Figure 3A) and $\text{Na}^+-\text{Ca}^{2+}$ exchanger isoform 1 (NCX1) (0.3-fold $P<0.0001$; Figure 3B) mRNAs were decreased in cultured VSMCs of ATP2B1 KO mice aorta compared with those in the control mouse aorta. On the contrary, the expression of ATP2B4 mRNA was upregulated (1.9-fold $P<0.0001$; Figure 3C) in KO VSMCs.

Increased Intracellular Calcium Concentrations in VSMCs of ATP2B1 KO Mice

To investigate whether intracellular calcium concentration in VSMCs was altered through KO of the ATP2B1 gene, we used the Fura-2-acetoxymethyl ester fluorescence assay. As shown in Figure 3D, the intracellular calcium was higher in VSMCs of ATP2B1 KO mice (F340/F380 ratio of KO VSMCs: 0.631 ± 0.029 ; F340/F380 ratio of control VSMCs: 0.505 ± 0.022 ; $P<0.05$) at baseline condition. Furthermore, phenylephrine-induced peak increase in intracellular calcium concentration was also augmented in KO VSMCs than in




Figure 2. Blood pressure measured by tail-cuff method and radiotelemetric method. **A**, Systolic blood pressure was measured by tail-cuff method in 8-week-old knockout (KO) mice (117.7 ± 3.4 ; $n=15$) and control mice (105.3 ± 2.9 ; $n=13$), and 22-week-old vascular smooth muscle cell (VSMC) ATP2B1 KO mice (132.6 ± 3.3 ; $n=14$) and control mice (117.3 ± 5.4 ; $n=11$). Data are presented as mean and SE (C indicates control mice; K, VSMC ATP2B1 KO mice; 8w, 8-week-old; 22w, 22-week-old). **B**, Circadian patterns of systolic blood pressure in VSMC ATP2B1 KO mice ($n=9$) and control mice ($n=9$) on a 12-hour light (8:00 AM to 8:00 PM)/dark (8:00 PM to 8:00 AM) cycle are shown. Mice were studied on a normal-salt diet (0.3% NaCl). Values plotted are hourly means and SEs measured over 60 hours (C indicates control mice; K, VSMC ATP2B1 KO mice). **C**, Circadian patterns of heart rate in VSMC ATP2B1 KO mice ($n=9$) and control mice ($n=9$) on a 12-hour light (8:00 AM to 8:00 PM)/dark (8:00 PM to 8:00 AM) cycle are shown. Mice were studied on a normal-salt diet. Values plotted are hourly means and SEs measured over 60 hours.

control VSMCs (F340/F380 ratio of KO VSMCs: 1.187 ± 0.068 ; F340/F380 ratio of control VSMCs: 0.805 ± 0.034 ; $P < 0.001$).

Vasoconstriction Was Accelerated by Phenylephrine Loading in Femoral Artery of KO Mice

We examined the vasoconstrictor response of femoral artery rings to phenylephrine. As summarized in Figure 4, femoral artery rings of KO mice were hyperreactive to the maximum concentration of phenylephrine (10^{-5} M) compared with

those of control mice (KO: 84.1% KCl contraction; control: 54.4% KCl contraction; $P < 0.05$; $n=10$).

Discussion

Implication of ATP2B1 in Blood Pressure Control and Function of ATP2B1 in VSMCs

This study showed that blood pressure was significantly higher in mice lacking ATP2B1 in VSMCs than that in wild mice. These results confirm the importance of the ATP2B1 gene in regulation of blood pressure. The ATP2B1 gene is one of the genes that we reported in 2008 as a gene for hypertension receptivity in a large-scale Japanese population, which has been confirmed recently in individuals of European descent, Koreans, and other Japanese. We first paid attention to the gene and made a strategy for creating a conditional KO model of the gene to confirm the relation between ATP2B1 and hypertension. VSMC ATP2B1 KO mice showed no significant change in birth rate and growth, although their expressions of ATP2B1 in the aorta and primary cultured VSMCs were markedly reduced, and they showed significantly higher blood pressure. We confirmed that the elevation of blood pressure in ATP2B1 KO mice was certain, with no relation to age and light-dark cycle. Furthermore, alteration in calcium homeostasis in VSMCs and increased vasoconstriction of femoral artery were observed in ATP2B1 KO mice. Recently, we showed that single nucleotide polymorphisms in the ATP2B1 gene cause phenotypic changes in human tissue.¹⁰ ATP2B1 mRNA expression in human umbilical artery smooth muscle cells was significantly lower in those with a risk allele for hypertension than in those having no risk allele. The finding using human artery was consistent with those seen in mice lacking ATP2B1 in VSMCs. These findings support that KO of ATP2B1 in VSMCs caused blood pressure elevation.

The ATP2B1 gene encodes plasma membrane calcium ATPase isoform 1 (so-called PMCA1), which removes bivalent calcium ions from eukaryotic cells against very large concentration gradients and plays a critical role in intracellular calcium homeostasis. In mammals, calcium ATPase isoforms are encoded by ≥ 4 separate genes (ATP2B1 to B4),¹⁷ and organ-specific mRNA expression of the isoforms has been reported. Using bladder smooth muscle cells, contractility measurements have documented the important role of ATP2B1 in the extrusion of Ca^{2+} after carbachol stimulation or depolarization with potassium chloride.⁹ Although bladder smooth muscle expresses both ATP2B4 and ATP2B1, ATP2B1 inhibition caused 3-time increments in intracellular calcium concentration and contraction of bladder smooth muscle compared with ATP2B4 blockade. Thus, ATP2B1 rather than ATP2B4 may have an important role in calcium handling and regulation in contraction of smooth muscle cells. In vascular smooth muscle, ATP2B1 and ATP2B4 have been also shown to be expressed.¹⁸ However, there were few reports evaluating the role of ATP2B1 in VSMCs. Thus, we decided to knock out the ATP2B1 gene of VSMCs to clarify the function of the ATP2B1 gene in hypertension theoretically in humans. In fact, VSMC-specific KO of ATP2B1




Figure 3. Altered gene expression and calcium transient in primary cultured vascular smooth muscle cells (VSMCs) of knockout (KO) mice. **A**, ATP2B1 mRNA expression in primary cultured VSMCs ($n=8-10$ for each genotype) quantified by quantitative RT-PCR (qRT-PCR) using exon 10 and 11 amplification. Data are presented as mean and SE ($P < 0.0001$). **B**, NCX1 mRNA expression in primary cultured VSMCs ($n=8-10$ for each genotype) quantified by qRT-PCR. Data are presented as mean and SE ($P < 0.0001$). **C**, ATP2B4 mRNA expression in primary cultured VSMCs ($n=8-10$ for each genotype) quantified by qRT-PCR. Data are presented as mean and SE ($P < 0.0001$). **D**, Measurement of basal condition and phenylephrine-induced increase in intracellular calcium concentration of VSMCs were performed. Figure shows the time course of phenylephrine-stimulated change in intracellular calcium concentration in ATP2B1 KO VSMC and control VSMC mice ($n=26-30$ cells from 10 to 11 coverslips). ATP2B1 KO VSMC mice showed higher intracellular calcium concentration the entire time course before and after the phenylephrine stimulation. Data are displayed as ratio of F340/F380. Intracellular calcium concentration values over the entire cell were averaged to obtain the changes in the whole-cell calcium concentration. Data are presented as mean and SE (BL indicates baseline condition; PE, phenylephrine; K, ATP2B1 KO VSMC; C, control VSMC; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$).

showed an elevation in blood pressure associated with a rise in intracellular calcium and a decrease in NCX1 mRNA expression. Moreover, the vascular contractile response was increased in KO mice. These results suggest that ATP2B1 has important roles in calcium handling and contraction in VSMCs.

Important Relationship Between ATP2B1 and Other Calcium-Related Genes

NCX1, a calcium pump similar to ATP2B1, is known to play an important role in hypertension through its effect on VSMCs.^{19,20} Interestingly, a recent report revealed that calcium clearance proteins, such as plasma membrane Ca^{2+} -ATPase (PMCA), sarcoplasmic reticulum Ca^{2+} ATPase, and NCX1, showed coordinated expression.²¹ Furthermore, PMCA and NCX1 showed similar changes in expression in human arterial myocytes.²² These recent




Figure 4. Phenylephrine-induced vasoconstriction of femoral artery rings. Isolated femoral artery rings obtained from knockout (KO) mice and control mice were stimulated with the α_1 -adrenoceptor agonist phenylephrine. Data are presented as mean and SE of 9 to 10 independent experiments. Force is expressed as the percentage of maximal contraction obtained by potassium-enriched solution.

findings suggest that ATP2B1 and NCX1 have a strong relationship and are modulated by the same system. Moreover, NCX1 and PMCA showed colocalization in the basolateral membrane of mouse distal convoluted cells in several studies.^{23,24} Furthermore, several proteins associate with NCX1 and PMCA with an alteration in their activity.²⁵ On the contrary, ATP2B4 was upregulated in KO VSMCs. This upregulation of ATP2B4 seems to compensate for the decrease in expression of ATP2B1. These findings suggest that decreased expression of ATP2B1 and NCX1 is one of the possible mechanisms of increase in intracellular calcium concentration.

Possibility of Alteration in Intracellular Calcium Homeostasis

In a recent study, a novel PMCA1 selective inhibitor, caloxin1b3, raised cytosolic calcium concentration in endothelial cells.²⁶ This finding supports the results that KO of ATP2B1 in VSMCs causes significant alterations in calcium-related gene expression and in intracellular calcium concentration. In the present study, KO VSMCs showed higher intracellular calcium concentration compared with the control, and a higher response was observed in response to the stimulation of phenylephrine compared with that of control VSMCs. Because increased intracellular calcium concentration may lead to blood pressure elevation via vasoconstriction,²⁷ the increased calcium concentrations seen in ATP2B1 KO VSMCs may be one of the possible mechanisms of high blood pressure seen in AP2B1 KO mice.

Increased Vasoconstriction Is One of the Possible Mechanisms for Blood Pressure Elevation

In the present study, we confirmed an increased contractile response to phenylephrine in femoral artery rings of KO

mice. Because phenylephrine activates inositol 1,4,5-triphosphate-induced intracellular calcium release and also stimulates voltage-independent calcium-permeable channels,²⁸ the alteration in contractile response may be attributed to alteration in intracellular calcium homeostasis. As shown in the present study, increased intracellular calcium concentration would augment the contractile capacity, which might increase the blood pressure in ATP2B1 KO mice. These findings strongly support the hypothesis that ATP2B1 gene is associated with blood pressure control in vivo.

Conclusion

We revealed that ATP2B1 KO in VSMCs increases the blood pressure in vivo study. Lack of ATP2B1 in VSMCs also increased intracellular calcium concentration and augmented the vascular contractility in ex vivo study. Our results clearly demonstrated that ATP2B1 gene expression in VSMCs is important in blood pressure regulation. Because the ATP2B1 gene has been reported to be a hypertension-susceptible gene by our systemic multiple candidate gene analyses, the present data suggest not only the importance of the ATP2B1 gene as a hypertension-related gene but also the value of the systemic multiple candidate gene approach and genome-wide association study in finding disease-related genes.

Perspectives

We made mice with conditional KO of ATP2B1 in VSMCs. However, the role of ATP2B1 in cells other than VSMCs in blood pressure control is not known. Thus, we need to further investigate the role of the ATP2B1 gene using other types of Cre mice and should make new organ-specific KO mice to analyze the role of the ATP2B1 gene in other conditions.

Sources of Funding

This work was supported by grants for scientific research (KAKENHI and Priority Areas "Medical Genome Science [Millennium Genome Project]" and "Applied Genomics") from the Ministry of Education, Culture, Sports, Science, and Technology, Japan; a grant for a Science and Technology Incubation Program in Advanced Regions, Japan Science and Technology Agency; a grant for scientific research from Yokohama City University; and a research grant from Olympus Corporation.

Disclosures

N.H., Y.T., K.K., Y.K., H.U., T.M., and S.U. have been named as the inventors on a patent application by Ehime University, Shiga University of Medical Science, and Yokohama City University in work related to this study.

References

- Lifton RP. Molecular genetics of human blood pressure variation. *Science*. 1996;272:676–680.
- Kohara K, Tabara Y, Nakura J, Imai Y, Ohkubo T, Hata A, Soma M, Nakayama T, Umemura S, Hirawa N, Ueshima H, Kita Y, Ogiwara T, Katsuya T, Takahashi N, Tokunaga K, Miki T. Identification of hypertension-susceptibility genes and pathways by a systemic multiple candidate gene approach: the Millennium Genome Project for Hypertension. *Hypertens Res*. 2008;31:203–212.
- Levy D, Ehret GB, Rice K, Verwoert GC, Launer LJ, DeGhan A, Glazer NL, Morrison AC, Johnson AD, Aspelund T, Aulchenko Y, Lumley T, Kottgen A, Vasan RS, Rivadeneira F, Eiriksdottir G, Guo X, Arking DE, Mitchell GF, Mattace-Raso FU, Smith AV, Taylor K, Scharpf RB, Hwang SJ, Sijbrands EJ, Bis J, Harris TB, Ganesh SK, O'Donnell CJ, Hofman A, Rotter JI, Coresh J, Benjamin EJ, Uitterlinden AG, Heiss G, Fox CS, Witteman JC, Boerwinkle E, Wang TJ, Gudnason V, Larson MG, Chakravarti A, Psaty BM, van Duijn CM. Genome-wide association study of blood pressure and hypertension. *Nat Genet*. 2009;41:677–687.
- Cho YS, Go MJ, Kim YJ, Heo JY, Oh JH, Ban HJ, Yoon D, Lee MH, Kim DJ, Park M, Cha SH, Kim JW, Han BG, Min H, Ahn Y, Park MS, Han HR, Jang HY, Cho EY, Lee JE, Cho NH, Shin C, Park T, Park JW, Lee JK, Cardon L, Clarke G, McCarthy MI, Lee JY, Oh B, Kim HL. A large-scale genome-wide association study of Asian populations uncovers genetic factors influencing eight quantitative traits. *Nat Genet*. 2009;41:527–534.
- Hong KW, Go MJ, Jin HS, Lim JE, Lee JY, Han BG, Hwang SY, Lee SH, Park HK, Cho YS, Oh B. Genetic variations in ATP2B1, CSK, ARSG and CSMD1 loci are related to blood pressure and/or hypertension in two Korean cohorts. *J Hum Hypertens*. 2010;24:367–372.
- Hong KW, Jin HS, Lim JE, Kim S, Go MJ, Oh B. Recapitulation of two genomewide association studies on blood pressure and essential hypertension in the Korean population. *J Hum Genet*. 2010;55:336–341.
- Takeuchi F, Isono M, Katsuya T, Yamamoto K, Yokota M, Sugiyama T, Nabika T, Fujioka A, Ohnaka K, Asano H, Yamori Y, Yamaguchi S, Kobayashi S, Takayanagi R, Ogiwara T, Kato N. Blood pressure and hypertension are associated with 7 loci in the Japanese population. *Circulation*. 2010;121:2302–2309.
- Okunade GW, Miller ML, Pyne GJ, Sutliff RL, O'Connor KT, Neumann JC, Andringa A, Miller DA, Prasad V, Doetschman T, Paul RJ, Shull GE. Targeted ablation of plasma membrane Ca²⁺-ATPase (PMCA) 1 and 4 indicates a major housekeeping function for PMCA1 and a critical role in hyperactivated sperm motility and male fertility for PMCA4. *J Biol Chem*. 2004;279:33742–33750.
- Liu L, Ishida Y, Okunade G, Shull GE, Paul RJ. Role of plasma membrane Ca²⁺-ATPase in contraction-relaxation processes of the bladder: evidence from pmca gene-ablated mice. *Am J Physiol Cell Physiol*. 2006;290:C1239–1247.
- Tabara Y, Kohara K, Kita Y, Hirawa N, Katsuya T, Ohkubo T, Hiura Y, Tajima A, Morisaki T, Miyata T, Nakayama T, Takashima N, Nakura J, Kawamoto R, Takahashi N, Hata A, Soma M, Imai Y, Kokubo Y, Okamura T, Tomoike H, Iwai N, Ogiwara T, Inoue I, Tokunaga K, Johnson T, Caulfield M, Umemura S, Ueshima H, Miki T. Common variants in the ATP2B1 gene are associated with susceptibility to hypertension: the Japanese Millennium Genome Project. *Hypertension*. 2010;56:973–980.
- Tsurumi Y, Tamura K, Tanaka Y, Koide Y, Sakai M, Yabana M, Noda Y, Hashimoto T, Kihara M, Hirawa N, Toya Y, Kiuchi Y, Iwai M, Horiuchi M, Umemura S. Interacting molecule of AT1 receptor, ATRAP, is colocalized with AT1 receptor in the mouse renal tubules. *Kidney Int*. 2006;69:488–494.
- Sakai M, Tamura K, Tsurumi Y, Tanaka Y, Koide Y, Matsuda M, Ishigami T, Yabana M, Tokita Y, Hiroi Y, Komuro I, Umemura S. Expression of MAK-V/Hunk in renal distal tubules and its possible involvement in proliferative suppression. *Am J Physiol Renal Physiol*. 2007;292:F1526–F1536.
- Mills PA, Huetteman DA, Brockway BP, Zwiers LM, Gelsema AJ, Schwartz RS, Kramer K. A new method for measurement of blood pressure, heart rate, and activity in the mouse by radiotelemetry. *J Appl Physiol*. 2000;88:1537–1544.
- Kobayashi M, Inoue K, Warabi E, Minami T, Kodama T. A simple method of isolating mouse aortic endothelial cells. *J Atheroscler Thromb*. 2005;12:138–142.
- Zimmerman MC, Sharma RV, Davisson RL. Superoxide mediates angiotensin II-induced influx of extracellular calcium in neural cells. *Hypertension*. 2005;45:717–723.
- Akaike T, Jin MH, Yokoyama U, Izumi-Nakaseko H, Jiao Q, Iwasaki S, Iwamoto M, Nishimaki S, Sato M, Yokota S, Kamiya Y, Adachi-Akahane S, Ishikawa Y, Minamisawa S. T-type Ca²⁺ channels promote oxygenation-induced closure of the rat ductus arteriosus not only by vasoconstriction but also by neointima formation. *J Biol Chem*. 2009;284:24025–24034.
- Carafoli E. The Ca²⁺ pump of the plasma membrane. *J Biol Chem*. 1992;267:2115–2118.
- Pande J, Szewczyk MM, Kuszczak I, Grover S, Escher E, Grover AK. Functional effects of Caloxin 1c2, a novel engineered selective inhibitor of plasma membrane Ca²⁺-pump isoform 4, on coronary artery. *J Cell Mol Med*. 2008;12:1049–1060.

19. Iwamoto T, Kita S, Zhang J, Blaustein MP, Arai Y, Yoshida S, Wakimoto K, Komuro I, Katsuragi T. Salt-sensitive hypertension is triggered by Ca^{2+} entry via $\text{Na}^+/\text{Ca}^{2+}$ exchanger type-1 in vascular smooth muscle. *Nat Med*. 2004;10:1193–1199.
20. Blaustein MP. Sodium ions, calcium ions, blood pressure regulation, and hypertension: a reassessment and a hypothesis. *Am J Physiol*. 1977;232:C165–C173.
21. Pritchard TJ, Bowman PS, Jefferson A, Tosun M, Lynch RM, Paul RJ. Na^+/K^+ -atpase and Ca^{2+} clearance proteins in smooth muscle: a functional unit. *Am J Physiol Heart Circ Physiol*. 2010;299:H548–H556.
22. Baryshnikov SG, Pulina MV, Zulian A, Linde CI, Golovina VA. Orai1, a critical component of store-operated Ca^{2+} entry, is functionally associated with $\text{Na}^+/\text{Ca}^{2+}$ exchanger and plasma membrane Ca^{2+} pump in proliferating human arterial myocytes. *Am J Physiol Cell Physiol*. 2009;297:C1103–C1112.
23. Magyar CE, White KE, Rojas R, Apodaca G, Friedman PA. Plasma membrane Ca^{2+} -ATPase and NCX1 $\text{Na}^+/\text{Ca}^{2+}$ exchanger expression in distal convoluted tubule cells. *Am J Physiol Renal Physiol*. 2002;283:F29–F40.
24. Loffing J, Loffing-Cueni D, Valderrabano V, Klausli L, Hebert SC, Rossier BC, Hoenderop JG, Bindels RJ, Kaissling B. Distribution of transcellular calcium and sodium transport pathways along mouse distal nephron. *Am J Physiol Renal Physiol*. 2001;281:F1021–F1027.
25. Ruknudin AM, Lakatta EG. The regulation of the $\text{Na}^+/\text{Ca}^{2+}$ exchanger and plasmalemmal Ca^{2+} ATPase by other proteins. *Ann NY Acad Sci*. 2007;1099:86–102.
26. Szewczyk MM, Pande J, Akolkar G, Grover AK. Caloxin 1b3: a novel plasma membrane Ca^{2+} -pump isoform 1 selective inhibitor that increases cytosolic Ca^{2+} in endothelial cells. *Cell Calcium*. 2010;48:352–357.
27. Brenner R, Perez GJ, Bonev AD, Eckman DM, Kosek JC, Wiler SW, Patterson AJ, Nelson MT, Aldrich RW. Vasoregulation by the $\beta 1$ subunit of the calcium-activated potassium channel. *Nature*. 2000;407:870–876.
28. McDaniel SS, Platoshyn O, Wang J, Yu Y, Sweeney M, Krick S, Rubin LJ, Yuan JX. Capacitative Ca^{2+} entry in agonist-induced pulmonary vasoconstriction. *Am J Physiol Lung Cell Mol Physiol*. 2001;280:L870–L880.

Online Supplement

Title

Mice lacking hypertension candidate gene ATP2B1 in vascular smooth muscle cells show significant blood pressure elevation

Authors

Yusuke Kobayashi¹⁾, Nobuhito Hirawa²⁾, Yasuharu Tabara³⁾, Hidenori Muraoka⁴⁾, Megumi Fujita¹⁾, Nobuko Miyazaki¹⁾, Akira Fujiwara¹⁾, Yasuhiro Ichikawa⁵⁾, Yuichiro Yamamoto²⁾, Naoaki Ichihara¹⁾, Sanae Saka¹⁾, Hiromichi Wakui¹⁾, Shin-ichiro Yoshida¹⁾, Keisuke Yatsu²⁾, Yoshiyuki Toya¹⁾, Gen Yasuda²⁾, Katsuhiko Kohara³⁾, Yoshikuni Kita⁶⁾, Kohtaro Takei⁴⁾, Yoshio Goshima⁴⁾, Yoshihiro Ishikawa⁵⁾, Hirotsugu Ueshima⁶⁾, Tetsuro Miki³⁾, Satoshi Umemura¹⁾

Affiliations

1. Department of Medical Science and Cardiorenal Medicine, Yokohama City University Graduate School of Medicine, Yokohama, Japan
2. Division of Nephrology and Hypertension, Yokohama City University Medical Center, Yokohama, Japan
3. Department of Geriatric Medicine, Ehime University Graduate School of Medicine, Toon, Japan
4. Department of Molecular Pharmacology and Neurobiology, Yokohama City University Graduate School of Medicine, Yokohama, Japan
5. Cardiovascular Research Institute, Yokohama City University Graduate School of Medicine, Yokohama, Japan
6. Department of Health Science, Shiga University of Medical Science, Otsu, Japan

METHODS

Animal Care

Animals were housed under a 12-hour day/night cycle at a temperature of 25°C. Tap water was provided *ad libitum*. Experiments were conducted under the guidelines for animal experiments set by the Animal Experiment Committee of Yokohama City University School of Medicine. In order to clarify the importance of the ATP2B1 gene in blood pressure regulation, we attempt to knock out the function of the ATP2B1 gene

of vascular smooth muscle cells. To generate conditional ATP2B1 KO mice, we utilized the Cre-loxP and FLP-FRT recombination system.

Cre-Mice

SM22-Cre mice [Tg(Tagln-cre)¹Her/J, stock #004746] were obtained from The Jackson Laboratory (Bar Harbor, ME, USA). The SM22-Cre transgenic mice express Cre recombinase under control of the mouse transgelin (smooth muscle protein 22- α) promoter. Thus SM22-Cre mice knockout the gene that is sandwiched with loxP sites in vascular smooth muscle cell (VSMC) specifically. Mice engineered in this study were backcrossed onto the (C57BL/6J) genetic background for at least six generations.

ATP2B1 conditional knockout (KO) mouse

Conditional ATP2B1 KO mouse was generated by Cre/loxP and FLP-FRT recombination system. ATP2B1 is encoded by 21 exons on chromosome 10, and mice lacking exon 10 was reported to be an embryonic lethal. We therefore designed a new vector to knockout the exon 10 of ATP2B1 gene (Figure S1). This targeting vector contains a loxP-FRT-PGK-neo-FRT cassette with 5,572 bp of ATP2B1 homologous regions upstream of exon 10, and loxP site with 3,891 bp homologous regions downstream of exon 10. After linearization and electroporation into the C57BL/6 (RENKA) ES cell lines (Transgenic Inc., Kumamoto, Japan), 431 neo-resistant ES cell clones were obtained. Among the randomly selected 143 ES clones, homologous recombination was confirmed in six clones (#73583, #73584, #73588, #73612, #73693, #73697) by KpnI or PvuII digested Southern blot analysis using 5' probe, 3' probe or neo probe (Figure S2, S3, S4). To verify the recombination of loxP site downstream exon 10, PCR-amplified fragment length (using primer A and primer B) was analyzed by agarose-gel electrophoresis (Figure S5). Both wild-type allele (460 bp) and mutant allele (519 bp) was ascertained in all six clones. We next eliminated the PGK-neo region by transfection of Flp recombinase vector in to the #73588 clone (Figure S6). Flp/FRT recombination was verified by analyzing the length of PCR product amplified with primer C and primer D (Figure S7). Among 104 transfected clones, recombination was ascertained in the seven clones (#77743, #77775, #77936, #77937, #77956, #77961, #77964). KpnI or EcoRI digested Southern blot analysis using 5' probe, 3' probe, or DR probe further confirmed the Flp/FRT recombination in all seven clones (Figure S8). Three ATP2B1 floxed ES cell clones (#77743, #77775, #77936) were aggregated with 8-cell stage embryos (ICR strain). The embryo was transferred into uterus of recipient females, and chimeras were delivered by Caesarean section at embryonic 17 day. The

chimeras were mated with wild-type mouse, and germline transmission was ascertained in 14 (9 male, and 5 female) of 31 F1 mouse.

Creation of the Vascular Smooth Muscle Cell-targeted ATP2B1 KO Mice

ATP2B1^{loxP/loxP} mice were generated by the Cre-loxP and FLP-FRT recombination system. ATP2B1 is encoded by 21 exons on chromosome 10, and mice lacking exon 10 are reported to be embryo-lethal. We therefore designed a vector to knockout exon 10 of the ATP2B1 gene. To target inactivation of the ATP2B1 gene to VSMC, ATP2B1^{loxP/loxP} mice were intercrossed with SM22-Cre transgenic mice expressing Cre recombinase under control of the mouse transgelin (smooth muscle protein 22-alpha) promoter. The resulting ATP2B1^{loxP/-}/SM22-Cre animals were further mated with ATP2B1^{loxP/loxP} mice to generate ATP2B1^{loxP/loxP}/SM22-Cre (VSMC ATP2B1 KO) mice and ATP2B1^{loxP/loxP} mice without SM22-Cre (control mice). Animals used for experiments were backcrossed at least 6 times.

Blood Pressure Measured by Radiotelemetric Method

Direct blood pressure measurement was performed by a radiotelemetric method in which a blood pressure transducer (PA-C10, Data Sciences International, USA) was inserted into the left carotid artery. Ten days after transplantation, each mouse was housed individually in a standard cage on a receiver under a 12-hour light-dark cycle. Direct blood pressure was recorded every minute by radiotelemetry, as described previously¹⁾.

Cell Culture of Mouse VSMC

The VSMC were aseptically isolated from thoracic aortic explants of an 8-week-old ATP2B1 KO mouse and its wild-type littermate as described previously²⁾. VSMC were prepared by the explant method and cultured in Dulbecco's modified Eagle's Medium as described previously^{3,4)}. VSMC multiplying in a medium containing FBS rapidly lose their contractile features and become "synthetic" cells³⁾. Thus, the early passage cells (third to sixth passage) were used in the present study. Subconfluent VSMC were used in the following experiments. The serum of these cells were deprived for 24 hours and then stimulated as indicated. After the desired incubation period, cells were rinsed with ice-cold phosphate-buffered saline and then lysed and sonicated.

Western Blot Analysis of ATP2B1

Western blot analysis was performed as described previously^{5,6)}. Briefly, tissue extracts

were used for electrophoresis, and membranes (Millipore, USA) were incubated with rabbit polyclonal anti-ATP2B1 antibody (MBL, Japan) and subjected to enhanced chemiluminescence (GE Healthcare, US). The images were analyzed quantitatively using a FUJI LAS3000 Image Analyzer (FUJI Film, Japan) for determination of the ATP2B1 protein level. To measure the aortic expression ratio of ATP2B1/ β -actin, each ATP2B1 protein level was divided by the corresponding β -actin protein level obtained by re-probing, and thus derived from the same extract.

Isometric Tension of the Femoral Artery Vascular Rings

After the mice were anesthetized with pentobarbital, the femoral artery ring was isolated and placed in a tissue bath and kept at 37° C. Two tungsten wires (40 μ m in diameter) were threaded into the lumen, and the preparation was mounted in a two-channel myograph (Dual Wire myograph system 410A; Danish Myo Technology, Aarhus, Denmark). One tungsten wire was connected to a micro-manipulator, and the other was connected to a force transducer. All of the vascular rings were initially stabilized for at least 60 min with a modified Krebs-Henseleit solution (Sigma-Aldrich) whose temperature was maintained at 37 °C by a heated water jacket. Isometric tension was continuously monitored using a PowerLab/8 SP system (ADInstruments, Inc., Colorado Springs, CO). After the vascular ring was relaxed, the resting tension was adjusted to 5 mN. After normalization, phenylephrine (Sigma-Aldrich) was added to stimulate vasoconstriction. The concentration of phenylephrine was increased from 10^{-8} M to 10^{-5} M. After a plateau vasoconstriction had been attained, phenylephrine was washed out. At the end of all experiments, vasoconstriction of the femoral artery ring was induced by potassium-enriched solutions containing (in mM): KCl 120, CaCl₂ 2.5, MgSO₄ 1.17, NaHCO₃ 25, KH₂PO₄ 1.18, EDTA 0.027, Glucose 5.5.

RESULTS

VSMC ATP2B1 KO Mice showed Higher Blood Pressure Assessed by 24-hour Radiotelemetric System than Control Mice

To confirm the effects of deletion of vascular smooth muscle cell ATP2B1 on blood pressure and to analyze the circadian pattern of blood pressure, conscious VSMC ATP2B1 KO mice and control mice were subjected to blood pressure measurements by radiotelemetry. KO mice showed higher blood pressure than control mice at 14 weeks of age throughout the day (diastolic blood pressure; Figure S9A, mean blood pressure;