

- National Cancer Institute, 2003. <http://ctep.cancer.gov/forms/CTCAEv3.pdf>.
7. Altundag O, Gullu I, Altundag K *et al*. Induction chemotherapy with cisplatin and 5-fluorouracil followed by chemoradiotherapy or radiotherapy alone in the treatment of locoregionally advanced resectable cancers of the larynx and hypopharynx: results of single-center study of 45 patients. *Head Neck* 2005;**27**:15–21.
  8. Lambert L, Fortin B, Soulieres D *et al*. Organ preservation with concurrent chemoradiation for advanced laryngeal cancer: are we succeeding? *Int J Radiat Oncol Biol Phys* 2010;**76**:398–402.
  9. Fukada J, Shigematsu N, Takeda A *et al*. Weekly low-dose docetaxel-based chemoradiotherapy for locally advanced oropharyngeal or hypopharyngeal carcinoma: a retrospective, single-institution study. *Int J Radiat Oncol Biol Phys* 2010;**76**:417–24.
  10. Mendenhall WM, Parsons JT, Stringer SP *et al*. Radiotherapy alone or combined with neck dissection for T1–T2 carcinoma of the pyriform sinus: an alternative to conservation surgery. *Int J Radiat Oncol Biol Phys* 1993;**27**:1017–27.
  11. Nakamura K, Shioyama Y, Kawashima M *et al*. Multi-institutional analysis of early squamous cell carcinoma of the hypopharynx treated with radical radiotherapy. *Int J Radiat Oncol Biol Phys* 2006;**65**:1045–50.
  12. Vandenbrouck C, Eschwege F, De la Rochefordiere A *et al*. Squamous cell carcinoma of the pyriform sinus: retrospective study of 351 cases treated at the Institut Gustave-Roussy. *Head Neck Surg* 1987;**10**:4–13.
  13. Chen SW, Yang SN, Liang JA *et al*. Prognostic impact of tumor volume in patients with stage III–IVA hypopharyngeal cancer without bulky lymph nodes treated with definitive concurrent chemoradiotherapy. *Head Neck* 2009;**31**:709–16.
  14. Tai SK, Yang MH, Wang LW *et al*. Chemoradiotherapy laryngeal preservation for advanced hypopharyngeal cancer. *Jpn J Clin Oncol* 2008;**38**:521–7.
  15. Mekhail TM, Adelstein DJ, Rybicki LA *et al*. Enteral nutrition during the treatment of head and neck carcinoma: is a percutaneous endoscopic gastrostomy tube preferable to a nasogastric tube? *Cancer* 2001;**91**:1785–90.
  16. Fuwa N, Shibuya N, Hayashi N *et al*. Treatment results of alternating chemoradiotherapy for nasopharyngeal cancer using cisplatin and 5-fluorouracil – A phase II study. *Oral Oncology* 2007;**43**:948–55.
  17. Kohmura T, Hasegawa Y, Matsuura H *et al*. Clinical analysis of multiple primary malignancies of the hypopharynx and esophagus. *Am J Otolaryngol* 2001;**22**:107–10.
  18. Morimoto M, Nishiyama K, Nakamura S *et al*. Significance of endoscopic screening and endoscopic resection for esophageal cancer in patients with hypopharyngeal cancer. *Jpn J Clin Oncol* 2010;**40**:938–43.
  19. Muto M, Minashi K, Yano T *et al*. Early detection of superficial squamous cell carcinoma in the head and neck region and esophagus by narrow band imaging: a multi-center randomized controlled trial. *J Clin Oncol* 2010;**28**:1566–72.

