

14. この研究の研究資金^{けんきゆうしきん}について

研究費については、厚生労働科学研究費補助金^{こうせいろうどうか がくけんきゆうひ ほじょきん}がん臨床事業^{りんしやうじぎやう}「悪性神経膠腫^{あくせいしんけいこうしゆ}に対する

Temozolomide の治療効果を増強した標準治療確立に関する研究^{どくりつぎやうせいほうじん}」班、独立行政法人国立

がん研究センター^{けんきゆうかいはつひしていけんきゆう}がん研究開発費指定研究 20 指-4、^{きしやうあくせいしゆやう}「希少悪性腫瘍に対する標準治療確

立のための多施設共同研究^{たしせつきやうどうけんきゆう}」班(班研究代表者^{しぶいそういちろう} 渋井壮一郎)の研究費により負担することと

なっています。

15. 遺伝カウンセリングの体制

遺伝子解析研究に関して、不安に思うことがある場合や、相談したいことがある場合に備えて、遺伝カウンセリング部門、または窓口を設置しています。遺伝カウンセリング担当者があなたの相談に応じますので、担当医、同意のための説明担当者、あるいは^{いじかしやくいん}医事課職員にお申し出ください。

16. どんなことでも質問してください

ご質問があれば遠慮なくお尋ねください。これまでの説明をよくご理解いただいた上で、あなたがこの研究への参加に同意していただける場合には、「同意文書」にご署名の上、担当医までご提出くださいますようお願いいたします。

17. 担当医の連絡先、研究代表者、研究事務局

担当医:

施設研究責任者:

[参加施設の住所、電話番号を記入]

この臨床試験全体の責任者・連絡窓口は以下のとおりです。

研究事務局(臨床試験の実務責任者、連絡窓口)

なつめ あつし
夏目 敦至

名古屋大学大学院医学系研究科脳神経外科

〒466-8550 愛知県名古屋市昭和区鶴舞町 65

Tel: 052-744-2355 (ダイヤルイン) Fax: 052-744-2361

Email: anatsume@med.nagoya-u.ac.jp

研究代表者(臨床試験全体の責任者)

わかばやし としひこ
若林 俊彦

名古屋大学大学院医学系研究科脳神経外科

〒466-8550 愛知県名古屋市昭和区鶴舞町 65

Tel: 052-744-2355 (ダイヤルイン) Fax: 052-744-2361

Email: wakabat@med.nagoya-u.ac.jp

同意書

医療機関名 _____ 病院長 殿

研究名 : 「JCOG0911: 初発膠芽腫に対するインターフェロン- β + テモゾロミド併用化学放射線療法
のランダム化第 II 相試験」の附随研究: 化学療法、放射線療法を施行した膠芽腫例における
効果予測因子および予後因子に関する研究

説明内容:

- この研究が JCOG の臨床試験 (JCOG0911) の附随研究であること。
- 遺伝子とは
- 遺伝子と疾病
- この研究へのご協力について
- 研究協力の任意性と撤回の自由
- 研究の意義、目的と方法
- 予測される研究結果および提供者等に対して予測される危険や不利益
- 研究計画書の開示
- 試料の提供は無償であること
- 個人情報の保護について
- 研究の倫理審査
- 試料から得られた解析結果は、匿名化された上研究成果として公表
- 研究から生じる知的財産権の帰属
- この研究終了後の検体の取扱いの方針
- 研究資金について
- 遺伝カウンセリングについて
- 研究責任者の氏名と職名
- どんなことでも質問してください
- 担当医師の連絡先、研究代表者、試料解析研究事務局

上記の研究について、私が説明しました。

説明担当者署名:

説明年月日: _____ 年 _____ 月 _____ 日

上記の研究について、担当医師から説明を受けよく理解しましたので、研究への参加(腫瘍
組織、血液、および JCOG0911 試験の診療情報データの提供)に同意します。

患者本人署名:

同意年月日: _____ 年 _____ 月 _____ 日

研究成果の刊行に関する一覧

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
<u>Shibui S</u> , <u>Beppu T</u> , <u>Nishikawa R</u> , <u>Muragaki Y</u> , <u>Kuratsu J</u> , <u>Kumabe T</u> , <u>Kayama T</u> , <u>Nagane M</u> , <u>Wakabayashi T</u> , <u>Sumi M</u> , et al.	Randomized trial of chemoradiotherapy and adjuvant chemotherapy with nimustine (ACNU) versus nimustine plus procarbazine for newly diagnosed anaplastic astrocytoma and glioblastoma (JCOG0305)	Cancer Chemotherapy and Pharmacology	71	511-21	2013
<u>Okita Y</u> , <u>Kayama T</u> , <u>Shibui S</u> , et al.	Reactivation of cytomegalovirus following treatment of malignant glioma with temozolomide	Int Cancer Conf J	1	53-7	2012
<u>Ohno M</u> , <u>Shibui S</u> , et al.	Reactivation of hepatitis B virus after glioblastoma treatment with temozolomide	Neurol Med Chir	51	728-31	2011
<u>Sato A</u> , <u>Shibui S</u> , <u>Kayama T</u> , et al.	Pivotal role for ROS activation of p38 MAPK in the control of differentiation and tumor-initiating capacity of glioma-initiating cells	Stem Cell Res	12	119-31	2013

Sakurada K, <u>Kayama T</u> , et al.	Usefulness of multimodal examination and intraoperative magnetic resonance imaging system in glioma surgery	Neurol Med Chir	52	553-7	2012
Uzuka T, <u>Kumabe T</u> , <u>Beppu T</u> , <u>Kayama T</u> , et al.	Treatment outcomes in glioblastoma patients aged 76 years or older: a multicenter retrospective cohort study	J Neurooncol	116	299-306	2014
<u>Kumabe T</u> , <u>Kayama T</u> , et al.	Treatment results of glioblastoma during the last 30 years in a single Institute	Neurol Med Chir	53	786-96	2013
<u>Kumabe T</u> , et al.	Summary of 15 years experience of awake surgeries for neuroepithelial tumors in Tohoku university	Neurol Med Chir	53	455-66	2013
Hide T, <u>Kuratsu J</u> , et al.	New treatment strategies to eradicate cancer stem cells and niches in glioblastoma	Neurol Med Chir	53	764-72	2013

Takahashi Y, Kuratsu J, et al.	Prognostic value of isocitrate dehydrogenase 1, O ⁶ -methylguanine-DNA methyltransferase promoter methylation, and 1p19q co-deletion in Japanese malignant glioma patients	World J Surg Oncol	11	284	2013
Yamasaki F, Sugiyama K, et al.	Advantages of high b-value diffusion-weighted imaging to diagnose pseudo-responses in patients with recurrent glioma after bevacizumab treatment	Eur J Radiol	81	2805-10	2012
Saito T, Sugiyama K, et al.	Role of perfusion-weighted imaging at 3T in the histopathological differentiation between astrocytic and oligodendro-glial tumors	Eur J Radiol	81	1863-9	2012
Aoki T, Nishikawa R, Sugiyama K, Kumabe T, et al.	A multicenter phase I/II study of the BCNU implant (Gliadel® Wafer) for Japanese patients with malignant gliomas	Neurol Med Chir		in press	2013

<u>Beppu T, et al.</u>	Prediction of malignancy grading using computed tomography perfusion imaging in nonenhancing supratentorial gliomas	J Neurooncol	103	619-27	2011
<u>Beppu T, et al.</u>	Standardized uptake value in high uptake area on positron emission tomography with ¹⁸ F-FRP170 as a hypoxic cell tracer correlates with intratumoral oxygen pressure in glioblastoma	Mol Imag Biol		in press	
<u>Nagane M, Nishikawa R, Sugiyama K, Kuratsu J, Muragaki Y, et al.</u>	Phase II study of single-agent bevacizumab in Japanese patients with recurrent malignant glioma	Jpn J Clin Oncol	42	887-95	2012
<u>Nagane M, et al.</u>	Predictive value of mean apparent diffusion coefficient value for responsiveness of temozolomide-refractory malignant glioma to bevacizumab	Int J Clin Oncol	19	16-23	2014

Adachi J, <u>Nishikawa R</u> , et al.	O ⁶ - methylguanine-DNA methyltransferase promotor methylation in 45 primary central nervous system lymphomas: quantitative assessment of methylation and respoinsse to temozolomide treatment	J Neuro-Oncol	107	147-53	2012
Feng H, <u>Nishikawa R</u> , et al.	Phosphorylation of dedicator of cytokinesis 1 (Dock180) at tyroisine residue Y722 by Src family kinases mediates EGFR v III-driven Glioblastoma tumorigenesis	PNAS	109	3018-23	2012
Suzuki T, <u>Nishikawa R</u> , et al.	Cadherin 13 overexpression as an important factor related to the absence of tumor fluorescence in 5-aminolevulinic acid-guided resection of glioma	J Neurosurg	119	1331-9	2013
<u>Muragaki Y</u> , <u>Kayama T</u> , et al.	Phase II clinical study on intraoperative photodynamic therapy with talaporfin sodium and semiconductor laser in patients with malignant brain tumors	J Neurosurg	119	845-52	2013

Nitta M, Muragaki Y, et al.	Updated therapeutic strategy for adult low-grade glioma stratified by resection and tumor subtype	Neurol Med Chir	53	447-54	2013
Muragaki Y, et al.	Phase I/IIa trial of autologous formalin-fixed tumor vaccine concomitant with fractionated radiotherapy for newly diagnosed glioblastoma	J Neurosurg	115	248-55	2011
Wakabayashi T, Kayama T, Nishikawa R, Sugiyama K, et al.	A multicenter Phase I trial of combination therapy with interferon- β and temozolomide for high-grade gliomas (INTEGRA study): the final report	J Neurooncol	104	573-7	2011
Motomura K, Wakabayashi T, et al.	Benefits of interferon-beta and temozolomide combination therapy for newly diagnosed primary glioblastoma with the unmethylated MGMT	Cancer	117	1721-30	2011
Ohka F, Wakabayashi T, et al.	The global DNA methylation surrogate LINE-1 methylation is correlated with MGMT promoter methylation and is a better prognostic factor for glioma	PLoS One	6	e23332	2011

Mayahara H, Sumi M, et al.	Effect of chemotherapy on survival after whole brain radiation therapy for brain metastases: a single-center retrospective analysis	J Cancer Res Clin Oncol	138	1239-47	2012
Hashimoto K, Sumi M, Kayama T, Shibui S, et al.	Comparison of Clinical Outcomes of Surgery Followed by Local Brain Radiotherapy and Surgery Followed by Whole Brain Radiotherapy in Patients With Single Brain Metastasis: Single-Center Retrospective Analysis	Int J Radiat Oncol Biol Phys	81	475-80	2011
Yonemori K, et al.	Compliance with Good Clinical Practice in oncology registration trials in Japan	Ann Oncol	22	1451-6	2011

Randomized trial of chemoradiotherapy and adjuvant chemotherapy with nimustine (ACNU) versus nimustine plus procarbazine for newly diagnosed anaplastic astrocytoma and glioblastoma (JCOG0305)

Soichiro Shibui · Yoshitaka Narita · Junki Mizusawa · Takaaki Beppu · Kuniaki Ogasawara · Yutaka Sawamura · Hiroyuki Kobayashi · Ryo Nishikawa · Kazuhiko Mishima · Yoshihiro Muragaki · Takashi Maruyama · Junichi Kuratsu · Hideo Nakamura · Masato Kochi · Yoshio Minamida · Toshiaki Yamaki · Toshihiro Kumabe · Teiji Tominaga · Takamasa Kayama · Kaori Sakurada · Motoo Nagane · Keiichi Kobayashi · Hirohiko Nakamura · Tamio Ito · Takahito Yazaki · Hikaru Sasaki · Katsuyuki Tanaka · Hideaki Takahashi · Akio Asai · Tomoki Todo · Toshihiko Wakabayashi · Jun Takahashi · Shingo Takano · Takamitsu Fujimaki · Minako Sumi · Yasuji Miyakita · Yoichi Nakazato · Akihiro Sato · Haruhiko Fukuda · Kazuhiro Nomura

Received: 6 October 2012 / Accepted: 22 November 2012 / Published online: 11 December 2012
© Springer-Verlag Berlin Heidelberg 2012

Abstract

Purpose Glioblastoma (GBM) is one of the worst cancers in terms of prognosis. Standard therapy consists of resection with concomitant chemoradiotherapy. Resistance to nimustine hydrochloride (ACNU), an alkylating agent, has been linked to methylguanine DNA methyltransferase (MGMT). Daily administration of procarbazine (PCZ) has been reported to decrease MGMT activity. This study investigated the efficacy of ACNU + PCZ compared to ACNU alone for GBM and anaplastic astrocytoma (AA).

Methods Patients (20–69 years) who had newly diagnosed AA and GBM were randomly assigned to receive radiotherapy with ACNU alone or with ACNU + PCZ. The primary endpoint was overall survival (OS). This was designed as a phase II/III trial with a total sample size of 310 patients and was registered as UMIN-CTR C000000108.

Results After 111 patients from 19 centers in Japan were enrolled, this study was terminated early because temozolomide was newly approved in Japan. The median OS and median progression-free survival (PFS) with ACNU alone ($n = 55$) or ACNU + PCZ ($n = 56$) in the

S. Shibui (✉) · Y. Narita · Y. Miyakita · K. Nomura
Department of Neurosurgery and Neuro-Oncology,
National Cancer Center Hospital, 5-1-1,
Tsukiji, Chuo-ku, Tokyo 104-0045, Japan
e-mail: sshibui@ncc.go.jp

J. Mizusawa · A. Sato · H. Fukuda
Japan Clinical Oncology Group Data Center,
National Cancer Center, Tokyo, Japan

T. Beppu · K. Ogasawara
Department of Neurosurgery, Iwate Medical University,
Iwate, Japan

Y. Sawamura · H. Kobayashi
Department of Neurosurgery, Hokkaido University Graduate
School of Medicine, Hokkaido, Japan

R. Nishikawa · K. Mishima
Department of Neuro-Oncology/Neurosurgery,
International Medical Center, Saitama Medical University,
Saitama, Japan

Y. Muragaki · T. Maruyama
Department of Neurosurgery,
Tokyo Women's Medical University, Tokyo, Japan

J. Kuratsu · H. Nakamura · M. Kochi
Department of Neurosurgery,
Kumamoto University,
Kumamoto, Japan

Y. Minamida · T. Yamaki
Department of Neurosurgery,
Sapporo Medical University,
Sapporo, Japan

T. Kumabe · T. Tominaga
Department of Neurosurgery,
Tohoku University School of Medicine, Miyagi, Japan

T. Kayama · K. Sakurada
Department of Neurosurgery, Faculty of Medicine,
Yamagata University, Yamagata, Japan

intention-to-treat population were 27.4 and 22.4 months ($p = 0.75$), and 8.6 and 6.9 months, respectively. The median OS and median PFS of the GBM subgroup treated with ACNU alone ($n = 40$) or ACNU + PCZ ($n = 41$) were 19.0 and 19.5 months, and 6.2 and 6.3 months, respectively. Grade 3/4 hematologic adverse events occurred in more than 40 % of patients in both arms, and 27 % of patients discontinued treatment because of adverse events.

Conclusions The addition of PCZ to ACNU was not beneficial, in comparison with ACNU alone, for patients with newly diagnosed AA and GBM.

Keywords Glioblastoma · Anaplastic astrocytoma · Nimustine · ACNU · Procarbazine · MGMT

Abbreviations

GBM	Glioblastoma
AA	Anaplastic astrocytoma
ACNU	Nimustine hydrochloride
BCNU	Carmustine
TMZ	Temozolomide
MGMT	Methylguanine DNA methyltransferase
WHO	World Health Organization
PFS	Progression-free survival
OS	Overall survival
RT	Radiotherapy
HR	Hazard ratio
AE	Adverse event
ND	Not determined

CR	Complete response
PR	Partial response
SD	Stable disease
PD	Progressive disease
WBC	White blood cell
3D-CRT	Three-dimensional conformal radiotherapy
CT	Computed tomography
IMRT	Intensity-modulated radiation therapy
BEV	Beam's eye views
DVH	Dose–volume histograms
GTV	Gross tumor volume
CTV	Clinical target volume
PTV	Planning target volume
ICRU	International Commission on Radiation Units
FLAIR	Fluid-attenuated inversion recovery
OAR	Organ-at-risk

Introduction

Glioblastoma (GBM) is one of the worst cancers in terms of prognosis, with almost all patients experiencing progression without cure. According to the report of the Brain Tumor Registry of Japan, the %5-year survival of World Health Organization (WHO) grade IV GBM is 6.9 % and that of WHO grade III anaplastic astrocytoma (AA) is 33.9 % [1].

Standard therapy against GBM consists of the maximal resection that is safely possible, with concomitant chemo-radiotherapy. Currently, temozolomide (TMZ) is the

M. Nagane · K. Kobayashi
Department of Neurosurgery,
Kyorin University Faculty of Medicine, Tokyo, Japan

H. Nakamura · T. Ito
Department of Neurosurgery,
Nakamura Memorial Hospital, Hokkaido, Japan

T. Yazaki · H. Sasaki
Department of Neurosurgery,
Keio University School of Medicine, Tokyo, Japan

K. Tanaka
Department of Neurosurgery, St. Marianna University
School of Medicine, Kanagawa, Japan

H. Takahashi
Department of Neurosurgery, Brain Research Institute,
Niigata University, Niigata, Japan

A. Asai
Department of Neurosurgery,
Saitama Medical Center, Saitama, Japan

T. Todo
Department of Neurosurgery, University of Tokyo, Tokyo, Japan

T. Wakabayashi
Department of Neurosurgery, Nagoya University Graduate
School of Medicine, Nagoya, Japan

J. Takahashi
Department of Neurosurgery,
Kyoto University Graduate School of Medicine,
Kyoto, Japan

S. Takano
Department of Neurosurgery, Tsukuba University,
Tsukuba, Japan

T. Fujimaki
Department of Neurosurgery,
Teikyo University School
of Medicine, Tokyo, Japan

M. Sumi
Department of Radiation-Oncology,
National Cancer Center Hospital,
Tokyo, Japan

Y. Nakazato
Department of Pathology,
Gunma University, Gunma, Japan

standard agent used in the treatment of GBM. However, before the TMZ era, nitrosourea had been widely used for GBM and AA. The Glioma Meta-analysis Trialists Group described that chemotherapy including nitrosourea showed significant prolongation of survival, with a hazard ratio of 0.85 ($p < 0.0001$) [2].

Nimustine hydrochloride (ACNU) was developed in Japan, and for more than 20 years since 1980, it has been the standard chemotherapeutic agent against gliomas [3]. Wolff et al. [4] analyzed 364 studies, including a total of 24,193 patients with high-grade glioma, and reported that the survival gain in the 15 ACNU-treated cohorts was 8.9 months, compared to those who received different drugs or no chemotherapy. Takakura et al. [5] reported that the overall survival (OS) of AA and GBM treated by radiotherapy (RT) and concomitant ACNU were 36 and 12 months, respectively. Furthermore, the response rate of a more than 50 % reduction in tumor size was 46.2 % in both AA and GBM. Alkylating agents, including ACNU and procarbazine (PCZ), confer cytotoxic effects on glioma cells by alkylation at the O^6 -position of guanine in DNA. This results in the formation of DNA cross-links [6]. Methylguanine DNA methyltransferase (MGMT) removes methylation damage induced by nitrosourea from the O^6 -position of DNA guanines before cell injury, and this enzyme was detectable in 76 % of glioma tissues [7]. MGMT in glioma cells is a primary defense against nitrosourea, but the cellular methyltransferase activity of MGMT is exhausted after MGMT takes effect. Daily administration of PCZ for 10 days was reported to cause the accumulation of O^6 -methylguanine; it also decreased MGMT activity in rat liver [8] and lymphocytes in lymphoma patients [9]. Inhibition of MGMT by O^6 -benzylguanine increased the cytotoxicity of TMZ and carmustine (BCNU) to tumor cells [10]. From these results, it can be predicted that daily administration of PCZ, by depleting MGMT activity, will increase the efficacy of ACNU against AA and GBM.

To prove this hypothesis and establish a more potent standard therapy for AA and GBM, the Brain Tumor Study Group of the Japan Clinical Oncology Group (JCOG) conducted this clinical trial. The study was terminated at the end of the phase II part. The current report describes the final outcome of the study.

Subjects and methods

Patient eligibility criteria

Patients aged 20 to less than 70 years of age who had newly diagnosed and histologically proven supratentorial GBM or AA were eligible for this study. Patients were

enrolled between 3 and 14 days after their operation. To be eligible, a patient's preoperative MRI had to show that more than 50 % of the tumor was located in supratentorial areas, except the optic nerve, olfactory nerve, or pituitary gland. Eligible patients had Eastern Cooperative Oncology Group (ECOG) performance status (PS) of 0–2 or 3 (only in cases with neurologic symptoms caused by a tumor) and adequate hematologic, pulmonary, renal, and hepatic function, defined as follows: white blood cell (WBC) count $\geq 3.0 \times 10^3/\text{mL}$, hemoglobin level $\geq 8.0 \text{ g/dL}$, platelets count $\geq 1.0 \times 10^6/\text{mL}$, aspartate transaminase (AST) level $\leq 100 \text{ IU/L}$, alanine transaminase (ALT) level $\leq 100 \text{ IU/L}$, serum creatinine level $\leq 1.0 \text{ mg/dL}$. Additionally, written informed consent was obtained from all the participating patients. We excluded patients with multiple or disseminated tumors or large tumors in which the planned target volume for irradiation exceeded 1/3 of the whole-brain volume. Additionally, we also classified as ineligible any patient who was pregnant, had meningitis, pneumonia, diabetes mellitus with insulin injection, myocardial infarction, or unstable angina pectoris within the last 3 months, mental disorders, a history of pulmonary fibrosis or interstitial pneumonia, or other forms of active cancer occurring within 5 years of treatment. The study protocol was approved by JCOG Protocol Review Committee and institutional review board at each center.

Treatment

After the confirmation of the eligibility criteria, registration was made by telephone or fax to the JCOG Data Center. Patients were randomized within 14 days of surgery to either ACNU with RT (the control arm, A) or to ACNU + PCZ with RT (the experimental arm, B) (Fig. 1a) by a minimization method with adjustment factors consisting of histology (GBM vs. AA), age (younger than 60 vs. 60 years or older), residual tumor (presence vs. absence), and institution. Residual tumor was assessed using a gadolinium-enhanced MRI obtained within 72 h of the surgery.

Radiotherapy with concomitant chemotherapy was started within 3 weeks after the surgery. Patient positioning and immobilization with an individual head mask and computed tomography (CT)-based planning were required. Treatment was delivered using linear accelerators with nominal energies $\geq 4 \text{ MV}$. Intensity-modulated radiation therapy (IMRT) technique was not permitted. All fields were to be treated every day. Three-dimensional conformal radiotherapy (3D-CRT) planning including the use of beam's eye views (BEV) and dose–volume histograms (DVH) were recommended for volumetric dose evaluation. Quality assurance reviews were done at the Radiotherapy Support Centre in Tokyo, Japan, with feedback sent to each




Fig. 1 a Study design of JCOG 0305: RT + ACNU versus RT + ACNU + PCZ; 40 patients with GBM and 15 patients with AA were assigned to *arm A*, and 41 patients with GBM and 15 patients with AA were assigned to *arm B*. b Treatment schedule of RT + ACNU (*Arm A*) and RT + ACNU + PCZ (*Arm B*)

institution by the radiotherapy study coordinator (Minako Sumi). The minimum and maximum dose to the PTV should be comprised between 95 and 107 % of the International Commission on Radiation Units (ICRU) reference point dose. The gross tumor volume (GTV) was defined as the primary tumor with or without enhancement on CT or magnetic resonance imaging (MRI). The clinical target volume 1 (CTV1) included GTV, the resection cavity and surrounding edema (high-intensity area on T2-weighted or fluid-attenuated inversion recovery (FLAIR) image) plus a 1.5-cm margin. The CTV2 included GTV and the resection cavity plus a 1.5-cm margin. Planning target volume (PTV) was defined as CTV plus a margin of 0.5 cm or more. The doses for PTV1 and PTV2 were 50 and 10 Gy, respectively. The protocol required contouring organ-at-risk (OAR), including optic chiasm, brain stem, and retina. Cumulative doses to the optic chiasm and brainstem were limited to a maximum dose of 50 and 45 Gy for the retina.

In the control arm A, 80 mg/m² of ACNU was administered intravenously on days 1 and 36 during RT (Fig. 1b). In the experimental arm B, 80 mg/m² of oral PCZ was administered daily from days 1 to 10 and days 36 to 45, and given together with intravenous ACNU (80 mg/m²) on

days 8 and 43. Adjuvant therapy consisting of 80 mg/m² of ACNU alone in arm A or ACNU plus PCZ (PCZ: 80 mg/m² orally on days 1–10, ACNU: 80 mg/m² intravenously on day 8) in arm B started 56 days from the final administration of ACNU and was given every 8 weeks, for up to 12 cycles. Doses of ACNU and PCZ were calculated using actual body surface area, reduced for toxicity, and were not escalated.

Evaluations and follow-up

Baseline and follow-up examinations included vital signs, subjective symptoms, neurologic examination, MRI scan, and blood and serum laboratory examinations. For each patient, these examinations were performed weekly, with the exception of MRI scans, which were performed between the end of the initial chemoradiotherapy and the beginning of adjuvant therapy. All examinations were performed before each cycle of adjuvant chemotherapy, at a frequency of nearly every 2 months. After completion of the treatment protocol, patients were assessed every 3 months until progression. Toxicity was graded using the National Cancer Institute Common Toxicity Criteria (version 2). Findings of radiation necrosis were also assessed on MRI. Each patient was required to undergo a follow-up examination for at least 2 years from the date of randomization.

Tumor progression on MRI was defined according to Response Evaluation Criteria in Solid Tumors (RECIST), version 1.0 [11]. Progression of disease was defined as a 20 % increase in tumor size, as shown by contrast-enhanced imaging, or the development of new lesions, neurologic deterioration, or death by any cause. Further treatment at recurrence or progression was discretionary, but recorded.

A central pathology review by 3 independent pathologists (Yoichi Nakazato, a member of the Working Group for WHO 2007 classification; Nobuaki Funata; and Toru Iwaki) was performed and determinations given. A central review of radiological response was also performed.

Statistical analysis

When we planned this study, TMZ had been widely approved and was used worldwide. However, TMZ was not available in Japan. ACNU remained the standard therapy in Japan, but there was no sufficient data regarding this treatment. We planned a phase II/III clinical trial, with the phase II part designed to confirm the feasibility of ACNU and ACNU + PCZ.

The primary and secondary endpoints for the phase II part were %6-month survival and adverse events (AEs) in ACNU + PCZ arm. The primary endpoint of the phase III part was OS, while the secondary endpoints were PFS, response rate, complete response rate, and AEs.

Overall survival was calculated from the date of random assignment to the date of death from any cause and censored at the last follow-up for event-free patients. PFS was calculated from the date of randomization to the date of progression or death from any cause and censored at the last verifiable progression-free date for event-free patients. OS and PFS were estimated by the Kaplan–Meier method. OS was analyzed by the stratified log-rank test for eligible patients with adjustment factors, excluding institution. Unstratified log-rank tests were used for the analysis of PFS and subgroup analyses of OS and PFS. Fisher’s exact test was used for categorical data. All *p* values are two-sided, except for primary analysis of OS.

We assumed %2-year survivals in AA and GBM for arm A were 50 and 20 %, and the ratio of those patients with AA or GBM enrolled in this study was expected to be 2:3. The phase III study was designed to enroll 155 patients per arm with 5 years of accrual and 2 years of follow-up, including those for the phase II part and about 10 % of ineligible patients, to achieve at least 75 % power to detect a hazard ratio (HR) of 0.74, with a one-sided alpha of 0.05 [12].

Three interim analyses were planned. The first was planned during phase II to test whether %6-month survival in arm A was superior to the predefined threshold (80 %), with a one-sided alpha of 0.1 and beta of 0.2, when 56 patients were included in ACNU + PCZ arm. The second and third interim analyses of OS were planned during phase III. For analyses of phase III part, multiplicity was adjusted by the Lan and DeMets alpha-spending function with the O’Brien and Fleming stopping boundary to control the type I error for primary endpoint.

In March 2007, protocol was amended to stop patient accrual after 111 patients had enrolled and to carry out the final analysis without planned interim analyses for both of phase II and phase III part. This was done because toxicity of both arms was unexpectedly high in phase II and because TMZ became available in Japan.

All statistical analyses were performed using SAS software, release 9.1 (SAS Institute, Cary, NC).

This trial is registered with UMIN-CTR (www.umin.ac.jp/ctr/), number C000000108.

Results

Patient characteristics

A total of 111 patients from 19 centers were randomly assigned to arm A (*n* = 55) or arm B (*n* = 56) from March 2004 to September 2006. Primary analyses were performed in September 2007, and the updated analyses were completed in July 2009. All patients were eligible for this study. Baseline characteristics were well balanced between the arms

(Table 1). The median ages for arms A and B were 56 and 54 years, respectively. Total numbers for AA and GBM patients were 15 (27.3 %) and 40 (72.7 %) in arm A and 15 (26.8 %) and 41 (73.2 %) in arm B. PS 0 to 1 in arms A and B were 45 (81.8 %) and 41 (73.2 %), respectively. Eighteen (32.7 %) patients in arm A and 21 (37.5 %) patients in arm B underwent gross total removal, and no residual tumor was confirmed on MRI scans. The median duration of follow-up was 20.2 (range 0–48.0 months) for all eligible patients.

Treatment

Patient compliance with the treatment regimen is depicted in Table 2. Among the 111 total patients, 1 patient in arm A died from pulmonary embolism before the beginning of initial chemoradiotherapy. Fifty-three (96.3 %) patients in arm A completed initial chemoradiotherapy and received ACNU twice. In arm B, 48 out of 56 (85.7 %) patients received 2 cycles of PCZ + ACNU, and 8 patients (14.3 %) received 1 cycle of PCZ + ACNU in initial chemoradiotherapy. Eighteen (32.7 %) patients in arm A and 20 (35.7 %) patients in arm B failed to start adjuvant chemotherapy. Furthermore, 14 (25.5 %) patients in arm A and 23 (41.1 %) patients in arm B discontinued protocol therapy by the fourth cycle of adjuvant chemotherapy. The numbers of patients who received 4 cycles of chemotherapy or more were 23 (41.8 %) and 13 (23.2 %) in arm A and B, respectively. Only 5 (9.1 %) patients in arm A and 2 (3.6 %) patients in arm B completed the full protocol therapy. Nineteen (34.5 %) patients in arm A and 22 (39.3 %) patients in arm B discontinued the protocol for reasons other than completion of protocol or disease progression [arm A: 31 (56.4 %), arm B: 32 (57.1 %)]. Reasons for discontinuation were as follows: AEs [arm A: 6 (10.9 %), arm B: 13 (23.2 %)]; patient refusal related to AE

Table 1 Baseline characteristics in the ITT population

	Arm A (<i>n</i> = 55) (RT + ACNU)	Arm B (<i>n</i> = 56) (RT + PCZ + ACNU)
Age	56 (24–69)	54 (24–69)
Sex		
Male	32 (58.2 %)	33 (58.9 %)
Female	23 (41.8 %)	23 (41.1 %)
PS		
0, 1	45 (81.8 %)	41 (73.2 %)
2, 3	10 (18.2 %)	15 (26.8 %)
Histology		
Grade 3 (AA)	15 (27.3 %)	15 (26.8 %)
Grade 4 (GBM)	40 (72.7 %)	41 (73.2 %)
Surgery		
Gross total removal	18 (32.7 %)	21 (37.5 %)
Partial removal	30 (54.5 %)	26 (46.4 %)
Biopsy only	7 (12.7 %)	9 (16.1 %)

Table 2 Compliance

	Arm A (n = 55) (RT + ACNU)	Arm B (n = 56) (RT + PCZ + ACNU)
RT (completion)	54 (98.2 %)	56 (100 %)
Initial chemotherapy		
1 cycle	1 (1.8 %)	8 (14.3 %)
2 cycles	53 (96.4 %)	48 (85.7 %)
Adjuvant chemotherapy		
None	18 (32.7 %)	20 (35.7 %)
1–3 cycles	14 (25.5 %)	23 (41.1 %)
4–6 cycles	9 (16.4 %)	7 (12.5 %)
7–11 cycles	9 (16.4 %)	4 (7.1 %)
12 cycles (completion)	5 (9.1 %)	2 (3.6 %)

[arm A: 3 (5.5 %), arm B: 8 (14.3 %)]; and patient refusal not related to AE [arm A: 7 (12.7 %), arm B: 1 (1.8 %)].

After discontinuation of the protocol, 28 (50.9 %) patients in arm A and 29 (51.8 %) patients in arm B received TMZ as further treatment.

Central review of histology

The central pathology review diagnosis of all cases was performed according to the WHO 2007 classification (Table 3). Among 81 GBM in the intention-to-treat (ITT) population, 69 (85.2 %), 4 (4.9 %), 3 (3.7 %), 2 (2.5 %), and 3 (3.7 %) were diagnosed as GBM, anaplastic oligoastrocytoma (AOA), anaplastic oligodendroglioma (AO), AA, and others, respectively. Only 10 cases (33.3 %) were diagnosed as AA among the 30 AA in the ITT population, while 8 (26.7 %), 5 (16.7 %), 3 (10.0 %), 2 (6.7 %), and 2 (6.7 %) were confirmed as GBM, AOA, diffuse astrocytoma, pilocytic astrocytoma, and others, respectively. Among all 111 cases, phenotype change of astrocytic to oligodendroglial tumor occurred in 14 cases (12.6 %). Finally, 77 and 12 patients were diagnosed with GBM and AA, respectively, by central pathology review.

Table 3 Local diagnosis and central pathology review

Grade	Histology	Local diagnosis	Central pathology review		
			Total	Arm A	Arm B
IV	<i>Glioblastoma</i>	83	77	37	40
III	<i>Anaplastic astrocytoma</i>	30	12	6	6
III	Anaplastic oligoastrocytoma		9	3	6
III	Anaplastic oligodendroglioma		3	2	1
III	Anaplastic ependymoma		1	1	0
II	Diffuse astrocytoma		4	1	3
II	Oligoastrocytoma		1	1	0
II	Oligodendroglioma		1	1	0
I	Pilocytic astrocytoma		2	2	0
	Sarcoma		1	1	0
	Total	111	111	55	56

Response rate

The overall radiographic response rate for 66 measurable diseases after surgery, based on RECIST guideline, was assessed by Diagnostic Radiology Committee. The response rate was 21.2 % (7/33) in arm A and 6.1 % (2/33) in arm B. In GBM patients from the ITT population, response rates were 25.0 % in arm A [3 CR (complete response), 2 PR (partial response), 1 SD (stable disease), 14 PD (progressive disease)] and 9.1 % in arm B [1 CR, 1 PR, 1 SD, 19 PD] ($p = 0.23$). In AA patients from the ITT population, response rates were 15.4 % in arm A (2 PR, 1 SD, 10 PD) and 0 % in arm B (11 PD) ($p = 0.48$).

Progression-free survival

In the entire ITT population, PFS was 8.6 months [95 % confidence interval (CI); 5.1–20.5] in arm A ($n = 55$, 44 events), compared with 6.9 months (95 % CI 5.1–9.0, $p = 0.36$) in arm B ($n = 56$, 47 events) (Fig. 2a). According to grades in the ITT population, PFS of GBM in arm A ($n = 40$, 33 events) and B ($n = 41$, 35 events) was 6.2 (95 % CI 4.2–13.2) and 6.3 months (95 % CI 3.0–8.9), respectively ($p = 0.35$) (Fig. 2b). PFS of AA in arm A ($n = 15$, 11 PD) and B ($n = 15$, 12 events) was 8.6 (95 % CI 5.1–35.4) and 9.0 months (95 % CI 5.8–21.8), respectively ($p = 0.83$) (Fig. 2c). No difference was observed between the arms in any subgroup defined by histology, presence of remaining tumor, or age under or over 60.

In the subgroup defined by central pathology review, the PFS of GBM in arms A ($n = 37$, 33 events) and B ($n = 40$, 36 events) was 5.1 (95 % CI 4.2–10.3) and 5.7 months (95 % CI 2.7–8.4), respectively ($p = 0.49$) (Fig. 2d). The PFS of AA and AOA in arm A ($n = 9$, 4 events) and B ($n = 12$, 9 events) was ND (not determined) and 7.9 months (95 % CI 5.2–22.5), respectively ($p = 0.21$).




Fig. 2 Progression-free survival in the intention-to-treat (ITT) population (a), the GBM subgroup (b), the AA subgroup (c), and the GBM subgroup with central pathology review (d). Overall survival in

the ITT population (e), the GBM subgroup (f), the AA subgroup (g), and the GBM subgroup with central pathology review (h). RT + ACNU alone (solid line), RT + ACNU + PCZ (dotted line)

Overall survival and cause of death

From the entire ITT population, 35 patients died in each group. In arm A (n = 55), OS was 27.4 months (95 % CI 16.2–35.4), compared with 22.4 months (95 % CI 16.4–

29.2) in arm B (n = 56) (Fig. 2e). The %2-year survival in arms A and B was 51.9 % and 46.2 %, respectively. There was no difference between the 2 arms (p = 0.75 and pre-planned, one-sided p = 0.62, by stratified log-rank test).

The OS of GBM subgroup in arms A ($n = 40$, 28 death) and B ($n = 41$, 27 death) was 19.0 (95 % CI 15.2–33.3) and 19.5 months (95 % CI 15.8–29.2), respectively ($p = 0.90$) (Fig. 2f). The %2-year survival in arms A and B was 48 and 41 %, respectively. The OS of AA subgroup in arms A ($n = 15$, 7 death) and in arm B ($n = 15$, 8 death) was 35.4 [95 % CI 15.7–not estimated (NE)] and 27.4 months (95 % CI 17.8–NE), respectively ($p = 0.88$) (Fig. 2g). There were no differences between the arms of any subgroup.

In the subgroups defined by central pathology review, the OS of GBM in arm A ($n = 37$, 28 death) and in arm B ($n = 40$, 29 death) was 16.6 (95 % CI 13.3–29.5) and 18.7 months (95 % CI 15.4–23.4), respectively ($p = 0.92$) (Fig. 2h). The %2-year survival in arms A and B was 38 and 34 %, respectively. The OS of AA and AOA in arm A ($n = 9$, 3 death) and B ($n = 12$, 4 death) was 33.3 months (95 % CI 15.7–33.3) and NE, respectively ($p = 0.83$).

Among the 70 total deaths, 31/35 (88.6 %) patients in arm A and 32/35 (91.4 %) in arm B experienced neuronal death of an original tumor. One patient (2.9 %) in arm A and 2 (5.7 %) patients in arm B contracted treatment-related pneumonia and died from that illness. Other causes of death were pulmonary embolism (1), pneumonia (2), and unknown (1).

Toxicity

Toxicity was assessed in 110 patients receiving initial therapy and in 73 patients receiving adjuvant chemotherapy. The most frequent grade 3/4 toxicities, experienced by more than 10 % of patients, were hematologic, neurologic, gastrointestinal, and hepatic AEs (Table 4). Patients in both arms frequently experienced leukopenia and neutropenia; more than half of the patients in arm B experienced these AEs during adjuvant therapy as well as during initial therapy. More than 40 % of patients in arm A also experienced these hematologic events even during adjuvant therapy. Grade 4 neutropenia was observed in 5.6 and 39.3 % of patients in arms A and B during initial chemoradiotherapy and 11.1 and 15.6 % during adjuvant therapy. Grade 3/4 nausea and anorexia were seen in 10.7 and 16.1 % of patients in initial therapy in arm B, but were rare in the adjuvant-therapy subgroups in both arms. One patient in arm B had cerebral infarction. Extrapyrimalidal signs, including tremors or involuntary movements, occurred in 2 patients in each arm.

Grade 3/4 pneumonitis occurred in 1 patient in arm A and 2 in arm B during the entire treatment period. Opportunistic infections—including 2 cases of *Pneumocystis jirovecii* pneumonia (PCP), 1 case of oral candidiasis, and 2 case of herpes zoster—occurred in arm B. One patient (1.8 %) in arm A and 2 (3.6 %) patients in arm B

died from treatment-related pneumonia, and 1 of these patients in arm B had PCP. One patient in arm B died from sepsis and acute respiratory distress syndrome after initial therapy. One patient in arm A died from pulmonary embolism before starting chemoradiotherapy, and 1 patient in arm A and 2 patients in arm B died from pneumonia following tumor progression.

Radiation necrosis was observed in 2 out of 54 (3.7 %) patients in arm A and 1 out of 56 (1.8 %) patients in arm B. During surgery, 1 patient in arm A was found to have radiation necrosis. Pseudo-progression within 3 months after chemoradiotherapy was not suspected in any patient.

Discussion

This study aimed to evaluate the efficacy and safety of treatment with ACNU + PCZ compared to ACNU alone as concomitant chemoradiotherapy against AA and GBM. We found no obvious differences in OS or PFS for AA and GBM between the treatment groups, but patients treated with ACNU + PCZ experienced more adverse effects than those treated with ACNU alone. TMZ is an effective regimen for malignant gliomas with less toxicity than our ACNU regimens, but it was not approved in Japan when this study began. At the end of the phase II part of this study, TMZ became available even in Japan, so this study was terminated at that point.

Methylguanine DNA methyltransferase is a major DNA repair protein and is implicated in resistance of glioma cells to alkylating agents [13]. Transcriptional silencing by MGMT promoter methylation results in inhibition of MGMT expression [14], and thus MGMT promoter methylation is strongly associated with survival in glioma patients treated with either nitrosourea or TMZ [15–17]. The status of the promoter of MGMT in primary tumors was frequently observed to change from methylated to unmethylated in recurrent tumors following ACNU or TMZ treatment [18, 19], which constitutes one of the mechanisms behind malignant gliomas' resistance to nitrosourea and TMZ. The rationale for treatment with ACNU + PCZ is that daily application of PCZ depletes MGMT activity, increasing sensitivity of AA and GBM to ACNU. Dose-dense TMZ therapy based on the theory of depletion of MGMT [20, 21], or BCNU or TMZ with direct inhibition of MGMT by *O*⁶-benzylguanine [22, 23] has been shown in previous studies to be effective for GBMs. However, there was no difference in OS found between standard and dose-dense TMZ for newly diagnosed GBMs [20].

While we were conducting this study, the European Organization for Research and Treatment of Cancer (EORTC) Brain Tumor and Radiotherapy Groups and the

Table 4 Toxicity

Grade 3/4 adverse events	Initial therapy with RT (<i>n</i> = 110) (%)		Adjuvant therapy (<i>n</i> = 73) (%)	
	Arm A	Arm B	Arm A	Arm B
Hematologic				
Leukopenia	38.9	73.2	40.5	69.4
Neutropenia	38.9	76.8	44.4	56.3
Thrombocytopenia	5.6	50.0	40.5	50.0
Anemia	0	8.9	10.8	8.3
Neurologic				
Seizure	9.3	7.1	5.4	8.8
Speech impairment	11.1	10.7	5.4	2.9
Neuropathy-motor	11.1	12.5	0	0
Extrapyramidal sign	0	0	5.4	2.7
Pulmonary (pneumonitis)	0	3.6	2.7	0
Gastrointestinal				
Nausea	0	10.7	0	0
Anorexia	1.9	16.1	0	2.9
Hepatic				
AST	3.7	12.5	2.9	2.9
ALT	3.7	16.1	2.9	8.8
Total bilirubin	1.9	5.4	0	0
Renal (creatinine)	0	0	0	0
Metabolic				
Hyponatremia	1.9	8.9	5.9	2.9
Hypokalemia	1.9	7.1	2.9	2.9
Fever	0	3.6	0	0
Dermatologic: erythema	3.7	5.4	0	2.9

National Cancer Institute of Canada (NCIC) Clinical Trials Group (EORTC/NCIC TMZ study) reported, in 2005, that RT + TMZ significantly prolonged the survival of GBM patients compared to RT alone [24]. The median PFS, OS, and 2-year survival for RT + TMZ were 6.9, 14.6 months, and 26.5 %, respectively [24]. Although our results compared favorably with the EORTC/NCIC TMZ study, the PFS of RT + ACNU alone for GBMs in our ITT population and in GBM subgroups in central pathology review were 6.2 and 5.1 months, shorter than those from the EORTC/NCIC TMZ study. Since more than half of the patients in our study underwent TMZ treatment following disease progression, it is possible that TMZ rescued these patients with progression after ACNU regimens and prolonged the survival of these patients.

The incidence of grade 3/4 hematologic AEs—such as leukopenia, neutropenia, and thrombocytopenia—were reported to be 5, 4, and 11 %, respectively, in adjuvant TMZ therapy in the EORTC/NCIC TMZ trial [24]. Compared to TMZ, even ACNU alone caused severe hematologic AEs in 40 % of the patients in our study, and most of those patients in both arms discontinued the treatment

protocol due to AEs or patient refusal related to AEs. It is noteworthy that approximately 30 % of patients in both arms failed to start adjuvant chemotherapy. The low completion rate of our protocol might explain the lack of differences in PFS and OS between the arms. After 2 patients in arm B experienced PCP, prophylactic use of cotrimoxazole (trimethoprim–sulfamethoxazole) against PCP was recommended in this study and was found to be useful.

Radiation necrosis has been reported in 2.5–21 % of patients undergoing chemoradiotherapy against malignant gliomas [25]. This complication was observed in 2.7 % of the patients in our study, but was tolerable. “Pseudo-progression” is the phenomenon of transient early disease progression after treatment with chemoradiotherapy consisting of TMZ for GBM progressive and enhancing lesions, as shown on MRI images taken immediately after treatment [25]. No patients in our study were suspected of pseudo-progression within 6 months after beginning chemoradiotherapy.

In general, the difference in histological diagnosis for local versus central pathology review is a major problem in the conduct of clinical trials on gliomas [26]. In our study, the concordance of GBM and AA between local and central diagnosis was low, but nearly identical to previous reports. In the EORTC/NCIC TMZ trial, central pathology review was performed in 85 % of cases, which confirmed the diagnosis of GBM in 93 % of the reviewed cases; 3 % had AA or AOA. In the phase III study of RT versus RT + BCNU + dibromodulcitol (EORTC 26882), of the 193 cases of AA diagnosed by the local pathologist, 176 were reviewed by the central pathologist. At review, 61 patients (35 %) were diagnosed with AA, 13 (8 %) with AOA, 4 (2 %) with AO, 44 (25 %) with GBM, 41 (23 %) with low-grade gliomas, and 13 (7 %) with another diagnosis [27].

The WHO classification system reflects the prognoses depending on grade I–IV tumors, or astrocytic or oligodendroglial tumors. However, it is based on morphological descriptions and contains subjective elements; thus, inter-observer variation occurs. The boundaries between grades II, III, and IV in gliomas are unclear, and there is a trend toward a more frequent diagnosis of oligodendroglial tumors [28]. Central pathological review before inclusion of a patient into clinical study is ideal, but it is very difficult to complete for aggressive grade III/IV tumors. Even if central review before enrollment is difficult in a multi-institutional setting, it is indispensable to perform post hoc central review at least in order to appropriately interpret the results of clinical studies of gliomas. A consensus meeting might also be useful before commencing clinical studies in order to gain concordance between local and central diagnoses. More objective classification of tumors based on

genotype, such the IDH1/2 mutation or 1p/19q codeletion, should be included in at least the stratification factor and subgroup analysis.

Conclusions

No significant differences in OS or PFS were found between ACNU alone and ACNU + PCZ in either AA or GBM. We found that ACNU + PCZ treatment was more toxic in our treatment schedule. Therefore, we conclude that the addition of PCZ to ACNU was not beneficial for newly diagnosed, high-grade astrocytomas as compared to ACNU alone. Considering the greater number of AEs associated with ACNU regimens, RT + TMZ should serve as a standard therapeutic regimen in the treatment of newly diagnosed AA and GBM.

Acknowledgments We thank all the members of the JCOG Brain Tumor Study Group and the staff of the JCOG Data Center. We appreciate Dr. Nobuaki Funata and Dr. Toru Iwaki for pathological review, Dr. Satoshi Ishikura for quality assurance of radiation therapy, and Dr. Hiroshi Katayama, Dr. Kenichi Nakamura, and Mr. Hidenobu Yamada for review of the manuscript. This work was supported in part by the National Cancer Center Research and Development Fund (23A-16 and 23A-20), the Health and Labour Sciences Research Grants (H14-032, H15-025, H16-005, H17-005), and the Grant-in Aid for Cancer Research (20S-4, 20S-6) from the Ministry of Health, Labour and Welfare.

Conflict of interest The authors declare that they have no conflict of interest.

References

1. The Committee of Brain Tumor Registry of Japan (2003) Report of brain tumor registry of Japan (1984–2000) 12th edition. *Neurol Med Chir (Tokyo)* 49(Suppl):1–101
2. Glioma Meta-analysis Trialists (GMT) Group (2002) Chemotherapy in adult high-grade glioma: a systematic review and meta-analysis of individual patient data from 12 randomised trials. *Lancet* 359(9311):1011–1018. doi:10.1016/S0140-6736(02)08091-1
3. Matsutani M, Nakamura O, Nakamura M, Nagashima T, Asai A, Fujimaki T, Tanaka H, Ueki K, Tanaka Y (1994) Radiation therapy combined with radiosensitizing agents for cerebral glioblastoma in adults. *J Neurooncol* 19(3):227–237
4. Wolff JE, Berrak S, Koontz Webb SE, Zhang M (2008) Nitrosourea efficacy in high-grade glioma: a survival gain analysis summarizing 504 cohorts with 24193 patients. *J Neurooncol* 88(1):57–63. doi:10.1007/s11060-008-9533-5
5. Takakura K, Abe H, Tanaka R, Kitamura K, Miwa T, Takeuchi K, Yamamoto S, Kageyama N, Handa H, Mogami H et al (1986) Effects of ACNU and radiotherapy on malignant glioma. *J Neurosurg* 64(1):53–57. doi:10.3171/jns.1986.64.1.0053
6. Pegg AE (1990) Mammalian O⁶-alkylguanine-DNA alkyltransferase: regulation and importance in response to alkylating carcinogenic and therapeutic agents. *Cancer Res* 50(19):6119–6129
7. Silber JR, Bobola MS, Ghatan S, Blank A, Kolstoe DD, Berger MS (1998) O⁶-methylguanine-DNA methyltransferase activity in adult gliomas: relation to patient and tumor characteristics. *Cancer Res* 58(5):1068–1073
8. Valavanis C, Souliotis VL, Kyrtopoulos SA (1994) Differential effects of procarbazine and methylnitrosourea on the accumulation of O⁶-methylguanine and the depletion and recovery of O⁶-alkylguanine-DNA alkyltransferase in rat tissues. *Carcinogenesis* 15(8):1681–1688
9. Souliotis VL, Kaila S, Boussiotis VA, Pangalis GA, Kyrtopoulos SA (1990) Accumulation of O⁶-methylguanine in human blood leukocyte DNA during exposure to procarbazine and its relationships with dose and repair. *Cancer Res* 50(9):2759–2764
10. Wedge SR, Porteus JK, May BL, Newlands ES (1996) Potentiation of temozolomide and BCNU cytotoxicity by O(6)-benzylguanine: a comparative study in vitro. *Br J Cancer* 73(4):482–490
11. Therasse P, Arbuck SG, Eisenhauer EA, Wanders J, Kaplan RS, Rubinstein L, Verweij J, Van Glabbeke M, van Oosterom AT, Christian MC, Gwyther SG (2000) New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. *J Natl Cancer Inst* 92(3):205–216
12. Bernstein D, Lagakos SW (1978) Sample size and power determination for stratified clinical trials. *J Stat Comput Simul* 8:65–73
13. Bobola MS, Tseng SH, Blank A, Berger MS, Silber JR (1996) Role of O⁶-methylguanine-DNA methyltransferase in resistance of human brain tumor cell lines to the clinically relevant methylating agents temozolomide and streptozotocin. *Clin Cancer Res* 2(4):735–741
14. Esteller M, Garcia-Foncillas J, Andion E, Goodman SN, Hidalgo OF, Vanaclocha V, Baylin SB, Herman JG (2000) Inactivation of the DNA-repair gene MGMT and the clinical response of gliomas to alkylating agents. *N Engl J Med* 343(19):1350–1354. doi:10.1056/NEJM200011093431901
15. Hegi ME, Diserens AC, Gorlia T, Hamou MF, de Tribolet N, Weller M, Kros JM, Hainfellner JA, Mason W, Mariani L, Bromberg JE, Hau P, Mirimanoff RO, Cairncross JG, Janzer RC, Stupp R (2005) MGMT gene silencing and benefit from temozolomide in glioblastoma. *N Engl J Med* 352(10):997–1003. doi:10.1056/NEJMoa043331
16. Sonoda Y, Yokosawa M, Saito R, Kanamori M, Yamashita Y, Kumabe T, Watanabe M, Tominaga T (2010) O(6)-Methylguanine DNA methyltransferase determined by promoter hypermethylation and immunohistochemical expression is correlated with progression-free survival in patients with glioblastoma. *Int J Clin Oncol* 15(4):352–358. doi:10.1007/s10147-010-0065-6
17. van den Bent MJ, Gravendeel LA, Gorlia T, Kros JM, Lapre L, Wesseling P, Teepen JL, Idbaih A, Sanson M, Smitt PA, French PJ (2011) A hypermethylated phenotype is a better predictor of survival than MGMT methylation in anaplastic oligodendroglial brain tumors: a report from EORTC study 26951. *Clin Cancer Res* 17(22):7148–7155. doi:10.1158/1078-0432.CCR-11-1274
18. Brandes AA, Franceschi E, Tosoni A, Bartolini S, Bacci A, Agati R, Ghimenton C, Turazzi S, Talacchi A, Skrap M, Marucci G, Volpin L, Morandi L, Pizzolitto S, Gardiman M, Andreoli A, Calbucci F, Ermani M (2010) O(6)-methylguanine DNA-methyltransferase methylation status can change between first surgery for newly diagnosed glioblastoma and second surgery for recurrence: clinical implications. *Neuro Oncol* 12(3):283–288. doi:10.1093/neuonc/nop050
19. Okita Y, Narita Y, Miyakita Y, Ohno M, Fukushima S, Kayama T, Shibui S (2012) Pathological findings and prognostic factors in recurrent glioblastomas. *Brain Tumor Pathol.* doi:10.1007/s10014-012-0084-2
20. Aldape KD, Wang M, Sulman EP, Hegi M, Colman H, Jones G, Chakravarti A, Mehta MP, Andrews DW, Long L, Diefes K,