

文 献

1. Civantos FJ, Stoeckli SJ, Takes RP, Woolgar JA, de Bree R, Paleri V, et al. What is the role of sentinel lymph node biopsy in the management of oral cancer in 2010? *Eur Arch Otorhinolaryngol* 2010; 267: 839–844.
2. Stoeckli SJ, Alkureishi LW, Ross GL. Sentinel node biopsy for early oral and oropharyngeal squamous cell carcinoma. *Eur Arch Otorhinolaryngol* 2009; 266: 787–793.
3. Stoeckli SJ, Pfaltz M, Ross GL, Steinert HC, MacDonald DG, Wittekind C, et al. The second international conference on sentinel node biopsy in mucosal head and neck cancer. *Ann Surg Oncol* 2005; 12: 919–924.
4. Paleri V, Rees G, Arullendran P, Shoaib T, Krishman S. Sentinel lymph node biopsy in squamous cell cancer of the oral cavity and oral pharynx: a diagnostic meta-analysis. *Head Neck* 2005; 27: 739–747.
5. Stark DD, Weissleder R, Elizondo G, Hahn PF, Saini S, Todd LE, et al. Superparamagnetic iron oxide: clinical applications as a contrast agent for MR imaging of the liver. *Radiology* 1988; 168: 297–301.
6. Weissleder R, Hahn PF, Stark DD, Elizondo G, Saini S, Todd LE, et al. Superparamagnetic iron oxide: enhanced detection of splenic tumors with MR imaging. *Radiology* 1988; 169: 399–403.
7. Torchia MG, Nason R, Danzinger R, Lewis JM, Thlivertis JA. Interstitial MR lymphography for the detection of sentinel lymph nodes. *J Surg Oncol* 2001; 78: 151–156.
8. Nakagawa T, Minamiya Y, Katayose Y, Saito H, Taguchi K, Imano H, et al. A novel method for sentinel lymph node mapping using magnetite in patients with non-small cell lung cancer. *J Thorac Cardiovasc Surg* 2003; 126: 563–567.
9. Sheng F, Inoue Y, Kiryu S, Watanabe M, Ohtomo K. Interstitial MR lymphography in mice with gadopentetate dimeglumine and gadoxetate disodium. *J Magn Reson Imaging* 2011; 33: 490–497.
10. Raynal I, Prigent P, Peyramaure S, Najid A, Rebuzzi C, Corot C. Macrophage endocytosis of superparamagnetic iron oxide nanoparticles. Mechanisms and comparison of ferumoxides and ferumoxtran-10. *Invest Radiol* 2004; 39: 56–63.
11. Sun R, Dittrich J, Le-Huu M, Mueller MM, Bedke MM, Kartenbeck J, et al. Physical and biological characterization of superparamagnetic iron oxide- and ultrasmall superparamagnetic iron oxide-labeled cells. A comparison. *Invest Radiol* 2005; 40: 504–513.
12. Hayakawa T, Iwaki T. *A color atlas of sectional anatomy of the rat*. Tokyo: Addthree, 2008. pp. 18–30. (in Japanese)
13. Vogl TJ, Mack MG, Juergens M, Stark M, Pegios W, Bergman C, et al. MR diagnosis of head and neck tumors: comparison of contrast enhancement with triple-dose gadodiamide and standard-dose gadopentetate dimeglumine in the same patients. *AJR Am J Roentgenol* 1994; 163: 425–432.
14. Carroll KW, Feller JF, Tirman PF. Useful internal standards for distinguishing infiltrative marrow pathology from hematopoietic marrow at MRI. *J Magn Reson Imaging* 1997; 7: 394–398.
15. Terada A, Hasegawa Y, Goto M, Sato E, Hyodo I, Ogawa T, et al. Sentinel lymph node radiolocalization in clinically negative neck oral cancer. *Head Neck* 2006; 28: 114–120.
16. Bach-Gansmo T, Fahlvik AK, Ericsson A, Hemmingsson A. Superparamagnetic iron oxide for liver imaging. Comparison among three different preparations. *Invest Radiol* 1994; 29: 339–344.
17. Kosuda S, Kusano S, Kohno N, Ohno Y, Tanabe T, Kitahara S, et al. Feasibility and cost-effectiveness of sentinel lymph node radiolocalization in stage N0 head and neck cancer. *Arch Otolaryngol Head Neck Surg* 2003; 129: 1105–1109.
18. Van Beers BE, Grandin C, Pauwels S, Mottet I, Goudemant JF, Delos M, et al. Gd-EOB-DTPA enhancement pattern of hepatocellular carcinomas in rats: comparison with Tc-99m-IDA uptake. *J Magn Reson Imaging* 1994; 4: 351–354.
19. Raynal I, Prigent P, Peyramaure S, Najid A, Rebuzzi C, Corot C. Macrophage endocytosis of superparamagnetic iron oxide nanoparticles. *Invest Radiol* 2004; 39: 56–63.

