

研究成果の刊行に関する一覧表

書籍

| 著者氏名 | 論文タイトル名 | 書籍全体の編集者名 | 書籍名 | 出版社名 | 出版地 | 出版年 | ページ |
|-------|--|--------------|-------------------------|--------|-----|------|---------|
| 松下由実 | 健康日本21(第2次) | 田中平三 | これからのかく衆衛生学 社会・環境と健康 | 南江堂 | 東京 | 2013 | 98-103 |
| 松下由実 | 内臓脂肪面積とメタボリックリスクファクターの集積： The Hitachi Health Study から | 岡 芳知 谷澤幸生 | 糖尿病学2011 | 診断と治療社 | 東京 | 2011 | 134-139 |
| 山本修一郎 | 内臓脂肪蓄積、インスリン抵抗性と大腸腫瘍の関連： The Hitachi Health Studyより | 岡 芳知 谷澤幸生 | 糖尿病学2011 | 診断と治療社 | 東京 | 2011 | 140-146 |

雑誌

| 発表者氏名 | 論文タイトル名 | 発表誌名 | 巻号 | ページ | 出版年 |
|---|--|-----------------------------------|----|-----|----------|
| Iwata M, Matsushita Y, Fukuda K, Wakura T, Okabe K, Koshimizu Y, Fukushima Y, Kobashi C, Yamazaki Y, Honoki H, Suzuki H, Tobe K. | Secretory units of islets in transplantation index (SUIT) is a useful predictor of insulin requirement in Japanese type 2 diabetic patients. | J Diabetes Investigation. | | | in press |
| Matsushita Y, Nakagawa T, Shinohara M, Yamamoto S, Takahashi Y, Mizoue T, Yokoyama T, Noda M. | How can waist circumference predict the body composition? | Diabetology & Metabolic Syndrome. | 6 | 11 | 2014 |

| | | | | | |
|--|---|-----------------------------------|----|-----------|------|
| Matsushita Y, Nakagawa T, Yamamoto S, Kato T, Ouchi T, Kikuchi N, Takahashi Y, Yokoyama T, Mizoue T, Noda M. | Adiponectin and visceral fat associate with cardiovascular risk factors. | Obesity. | 22 | 287-291 | 2014 |
| Okumura A, Unoki-Kubota H, Matsushita Y, Shiga T, Moriyoshi Y, Yamagoe S, Kaburagi Y. | Increased serum leukocyte cell-derived chemotaxin 2 (LECT2) levels in obesity and fatty liver. | Biosci Trends. | 7 | 276-283 | 2013 |
| Okazawa T, Iwata M, Matsushita Y, Kamura Y, Kato H, Okazawa S, Kigawa M, Tobe K.. | Aging attenuates the association of central obesity with the accumulation of metabolic risk factors when assessed using the waist circumference measured at the umbilical level (the Japanese standard method). | Nutr Diabetes. | 3 | e96 | 2013 |
| Matsushita Y, Nakagawa T, Yamamoto S, Takahashi Y, Yokoyama T, Mizoue T, Noda M. | Effect of longitudinal changes in visceral fat area on incidence of metabolic risk factors: the Hitachi Health Study. | Obesity. | 21 | 2126-2129 | 2013 |
| Yi S, Nakagawa T, Yamamoto S, Mizoue T, Takahashi Y, Noda M, Matsushita Y. | Short sleep duration in association with CT-scanned abdominal fat areas: The Hitachi Health Study. | International Journal of Obesity. | 37 | 129-134 | 2013 |
| Akter S, Nanri A, Yi S, Pham NM, Kurotani K, Kimura Y, Matsushita Y, Mizoue T. | Dietary patterns and C-peptide concentrations in a Japanese working population. | Nutrition. | 28 | e29-35 | 2012 |

| | | | | | |
|--|--|-----------------------|-----|-----------|------|
| Yi S, Nanri A, Matsushita Y, Kasai H, Kawai K, Mizoue T. | Depressive symptoms and oxidative DNA damage in Japanese municipal employees. | Psychiatry Res. | 200 | 318-322 | 2012 |
| Nanri A, Tomita K, Matsushita Y, Ichikawa F, Yamamoto M, Nagafuchi Y, Kakumoto Y, Mizoue T. | Effect of six months lifestyle intervention in Japanese men with metabolic syndrome: randomized controlled trial. | J Occup Health. | 54 | 215-222 | 2012 |
| Matsushita Y, Nakagawa T, Yamamoto S, Takahashi Y, Yokoyama T, Mizoue T, Noda M. | Effect of longitudinal Changes in visceral fat area and other anthropometric indices to the changes in metabolic risk factors in Japanese men. The Hitachi Health Study. | Diabetes Care. | 35 | 1139-1143 | 2012 |
| Poudel-Tandukar K, Sato M, Ejima Y, Nanri A, Matsushita Y, Imaizumi K, Mizoue T. | Relationship of serum fatty acid composition and desaturase activity to C-reactive protein in Japanese men and women. | Atherosclerosis. | 220 | 520-524 | 2012 |
| Matsushita Y, Nakagawa T, Yamamoto S, Takahashi Y, Yokoyama T, Mizoue T and Noda M. | Visceral fat area cutoff for the detection of multiple risk factors of metabolic syndrome in Japanese: The Hitachi Health Study. | Obesity. | 20 | 1744-1749 | 2012 |
| Yi S, Nanri A, Poudel-Tandukar K, Nonaka D, Matsushita Y, Hori A, Mizoue T. | Association between serum ferritin concentrations and depressive symptoms in Japanese municipal employees. | Psychiatry Res. | 189 | 368-372 | 2011 |
| Kimura Y, Nanri A, Matsushita Y, Sasaki S, Mizoue T. | Eating behavior in relation to prevalence of overweight among Japanese men. | Asia Pac J Clin Nutr. | 20 | 29-34 | 2011 |

| | | | | | |
|---|---|--|-------------|---------|------|
| Kimura Y, Matsushita Y, Nanri A, Mizoue T. | Metabolic syndrome and depressive symptoms among Japanese men and women. | Environmental Health and Preventive Medicine. | 16 | 363-368 | 2011 |
| Matsushita Y, Nakagawa T, Yamamoto S, Takahashi Y, Noda M, Mizoue T. | Associations of smoking cessation with visceral fat area and prevalence of metabolic syndrome in men: the Hitachi Health Study. | Obesity. | 19 | 647-651 | 2011 |
| 松下由実 | 内臓脂肪蓄積と癌の関連 The Hitachi Health Studyよ り | 内分泌・糖尿病・代 謝内科. | 35 | 189-194 | 2012 |
| 松下由実 | 糖尿病の予防・管理・治療 糖 尿病非薬物療法 食事療法 糖尿病治療における食事療法 の疫学研究 | 日本臨牀. | 70 (増刊3) | 750-753 | 2012 |