

Table 2 The associations between systolic and diastolic blood pressure at early pregnancy and educational level (and 95% CIs) (n=923)*

Educational level	Unadjusted Model	Basic Model†	Basic Model† +BMI	Basic Model† +Smoking	Basic Model† +Salt intake	Basic Model† +Alcohol intake	Basic Model† +Income	Basic Model† +BW gain	Full Model‡
SBP, mmHg									
High	Reference	Reference	Reference	Reference	Reference	Reference	Reference	Reference	Reference
Mid	-0.04 (-1.84 to 1.76)	-0.28 (-2.08 to 1.52)	-0.46 (-2.23, 1.30)	-0.15 (-1.96, 1.67)	-0.41 (-2.25, 1.42)	-0.49 (-2.32, 1.34)	-0.81 (-2.69, 1.06)	-0.32 (-2.15, 1.51)	-1.06 (-2.96, 0.84)
Low	2.39 (0.59 to 4.19)	2.42 (0.61 to 4.23)	1.19 (-0.62, 3.00)	2.85 (0.94, 4.76)	2.37 (0.52, 4.23)	2.23 (0.38, 4.08)	1.46 (-0.47, 3.38)	2.21 (0.36, 4.06)	0.13 (-1.94, 2.20)
DBP, mmHg									
High	Reference	Reference	Reference	Reference	Reference	Reference	Reference	Reference	Reference
Mid	0.16 (-1.09 to 1.41)	-0.05 (-1.29 to 1.20)	-0.16 (-1.38, 1.06)	0.13 (-1.12, 1.38)	-0.23 (-1.49, 1.03)	-0.32 (-1.58, 0.93)	-0.088 (-1.39, 1.21)	-0.10 (-1.37, 1.16)	-0.29 (-1.60, 1.03)
Low	0.74 (-0.52 to 1.99)	0.79 (-0.46 to 2.04)	0.041 (-1.21, 1.30)	1.31 (-0.0043, 2.63)	0.62 (-0.66, 1.90)	0.56 (-0.71, 1.83)	0.42 (-0.92, 1.75)	0.75 (-0.53, 2.02)	-0.26 (-1.69, 1.17)

* Estimates denote difference in BP (mmHg) between low/mid and high educational level.

† Basic model: adjusted for gestational age at BP measurement, maternal age, parity, past medical history of hypertension, pregnancy-induced hypertension, diabetes mellitus, renal disease, family history of hypertension.

‡ Full model: Basic model+ pre-pregnancy BMI, smoking, salt intake, alcohol intake, income, BW gain until mid-pregnancy.

Abbreviations: SBP, systolic blood pressure; DBP, diastolic blood pressure, BMI, pre-pregnancy body mass index; BW gain, body weight gain until mid-pregnancy.

The associations between systolic and diastolic BP at mid-pregnancy and educational level are shown in Table 3. In the basic model, the BP in the low educational group was significantly higher than that in the high educational group (difference = 1.84 mmHg, 95% CI: 0.057 to 3.61). Similar to the results shown in Table 2, after adding pre-pregnancy BMI to the basic model, BP differences became smaller and were not anymore statistically significant. Interestingly, adding smoking to the basic model increased the BP difference which resulted in statistical significance (difference = 2.53 mmHg, 95% CI: 0.66 to 4.40). With regard to the diastolic BP, educational levels were not statistically associated. Similar to early pregnancy, adding BMI to the basic model attenuated the differences in diastolic BP.

Table 3 The associations between systolic and diastolic blood pressure at mid-pregnancy and educational level (and 95% CIs) (n=923)*

Educational level	Unadjusted Model	Basic Model†	Basic Model† +BMI	Basic Model† +Smoking	Basic Model† +Salt intake	Basic Model† +Alcohol intake	Basic Model† +Income	Basic Model† +BW gain	Basic Model† Full Model‡
SBP, mmHg									
High	Reference	Reference	Reference	Reference	Reference	Reference	Reference	Reference	Reference
Mid	0.29 (-1.50, 2.07)	-0.0063 (-1.77, 1.76)	-0.20 (-1.92, 1.52)	0.17 (-1.60, 1.95)	-0.14 (-1.94, 1.66)	-0.18 (-1.98, 1.62)	-0.31 (-2.16, 1.54)	0.32 (-1.46, 2.10)	0.085 (-1.75, 1.92)
Low	1.52 (-0.27, 3.30)	1.84 (0.057, 3.61)	0.58 (-1.20, 2.35)	2.53 (0.66 to 4.40)	1.87 (0.045, 3.69)	1.76 (-0.057, 3.59)	1.17 (-0.73, 3.06)	1.77 (-0.035, 3.57)	0.53 (-1.48, 2.53)
DBP, mmHg									
High	Reference	Reference	Reference	Reference	Reference	Reference	Reference	Reference	Reference
Mid	0.81 (-0.42, 2.03)	0.58 (-0.62, 1.78)	0.47 (-0.71, 1.66)	0.73 (-0.48, 1.94)	0.40 (-0.82, 1.62)	0.37 (-0.86, 1.59)	0.14 (-1.10, 1.37)	0.69 (-0.53, 1.90)	0.17 (-1.08, 1.41)
Low	0.83 (-0.39, 2.06)	0.99 (-0.22 to 2.20)	0.30 (-0.92, 1.52)	1.44 (0.17 to 2.72)	1.10 (-0.13 to 2.34)	1.02 (-0.22, 2.25)	0.42 (-0.85, 1.69)	1.00 (-0.23, 2.22)	0.06 (-1.31, 1.42)

* Estimates denote difference in BP (mmHg) between low, mid and high educational level.

† Basic model: adjusted for gestational age at BP measurement, maternal age, parity, past medical history of hypertension, pregnancy-induced hypertension, diabetes mellitus, renal disease, family history of hypertension

‡ Full model: Basic model+ pre-pregnancy BMI, smoking, salt intake, alcohol intake, income, BW gain until mid-pregnancy.

Abbreviations: SBP, systolic blood pressure; DBP, diastolic blood pressure, BMI, pre-pregnancy body mass index; BW gain, body weight gain until mid-pregnancy.

The odds ratio (OR) of PIH by educational level is shown in Table 4. In total, PIH was diagnosed in 23 cases (2.5%). In the unadjusted model, the point estimate of the OR of PIH of the low educational group was 1.24 (95% CI: 0.44 to 3.44) in comparison with high educational group. In the full model adjusted for all confounders and mediators, the point estimate of the OR of PIH of the low educational group was 1.19 (95% CI: 0.31 to 4.60) in comparison with the high educational group, suggesting that the low educational group was 1.19 times more likely to develop PIH than the high educational group, even though the OR was not statistically significant.

Table 4 Odds ratio (and 95% confidence intervals) of pregnancy-induced hypertension stratified by educational level (n=923)

Educational Level	N (%)	Unadjusted	Full Model‡
PIH	23 (2.5)		
High (n=467)	10 (2.1)	Reference	Reference
Mid (n=228)	7 (3.1)	1.45 (0.54, 3.85)	0.44 (0.076, 2.60)
Low (n=228)	6 (2.6)	1.24 (0.44, 3.44)	1.19 (0.31, 4.60)

‡Full model: adjusted for maternal age, parity, past medical history of hypertension, pregnancy-induced hypertension.

Diabetes mellitus, renal disease, Family history of hypertension, pre-pregnancy body mass index, smoking, Salt intake.

Alcohol intake, income, body weight gain until mid-pregnancy.

Discussion

In our study, we found that among Japanese pregnant women the low educational group had significantly higher systolic BP than the high educational groups in early pregnancy and that this group maintained higher systolic and diastolic BPs toward mid-pregnancy. Furthermore, we found that the pre-pregnancy BMI mediated the association between educational levels and BP. On the other hand, smoking, alcohol intake, salt intake, and body weight gain until mid-pregnancy did not mediate this association.

The results of the current study are consistent with those of a previous study on SES and BP during pregnancy [13]. Silva et al. reported that the low educational group had significantly higher BPs throughout pregnancy. They categorized the educational level into 4 levels: high (university or higher), mid-high (higher vocational training), mid-low (less than 3 years of general secondary school, intermediate vocational training completed, or first year of higher vocational training), and low (no education, primary school, lower vocational training, intermediate general school, or ≤ 3 years of general secondary school). The systolic and diastolic BP difference between the low and high educational groups was 2.67 mmHg (95% CI: 0.66 to 4.40) and 0.53 mmHg (95% CI: -0.58 to 1.64), respectively, at early pregnancy (less than 18 weeks of gestation). Even though the categorization of educational level in our study is different from their study, our results are similar; in our study, the systolic and diastolic BP difference between the low and high educational groups was 2.39 mmHg (95% CI: 0.59 to 4.19) and 0.74 mmHg (95% CI: -0.52 to 1.99), respectively, at early pregnancy.

A decrease in systolic and diastolic BP from early to mid-pregnancy, known as a protective factor for PIH [13,23], was observed among every educational group in our study. In Silva's study, no decrease in diastolic BP from early to mid-pregnancy was reported among the low educational group. This inconsistency of BP decrease by educational groups might be due to

different characteristics of the study population. Compared with Silva's study, our sample population was less likely to be overweight (pre-pregnancy BMI; this study vs. Silva's study, 20.2 ± 2.3 vs. 23.2 ± 3.9) and to consume alcohol during pregnancy (this study vs. Silva's study, 15.6% vs. 47.1%). Furthermore, differences in the health care system for pregnant women might contribute to these differences, i.e., prenatal care in Japan might be associated with BP control, and quick access to a hospital with universal health insurance might be considered to have a beneficial effect on BP control.

Regarding mediators, the same previous study reported that the pre-pregnancy BMI mediates the association between educational level and BP [13], whereas smoking, alcohol intake, and body weight gain until mid-pregnancy do not mediate this association. Our results confirmed the results of the previous study and added to the literature that salt intake do not mediate the association between educational levels and BP during pregnancy.

From the results of our study, smoking showed an inverse association with BP, i.e., smoking pregnant women showed lower systolic and diastolic BP. Yet, smoking mediates, in part, the association between SES and cardiovascular events in adults [24]. Habitual smokers generally have a lower BP than non-smokers [25], which is related to lower body weight [26]. In pregnant women, it is reported that smoking is associated with lower diastolic BP until mid-pregnancy compared with non-smoking women [27]. Our results are also in line with those results. Smoking was associated with lower systolic and diastolic BP until mid-pregnancy, and this association did not change among the different educational groups.

Although not statistically significant, the point estimate of the OR for PIH in the low educational group was 1.19 compared with the high educational group, suggesting that the educational level might be associated with PIH. The non-significant result could be due to small sample size, i.e., the number of PIH cases was small in our study. Previous reports on the association between low educational level and PIH are inconsistent. On the basis of results of a population-based, prospective cohort study, Silva et al. reported that low educational level was significantly associated with the occurrence of preeclampsia [28] and gestational hypertension [29]. Haelterman et al. conducted a case-control study of 99 severe preeclampsia cases and reported that individuals with a low educational level (primary school or below) had a statistically higher OR (OR = 2.3, 95% CI: 1.2 to 4.4) than those who attended a primary school or higher [30]. On the other hand, by using birth records from North Carolina, Savitz et al. reported that maternal education did not differ with regard to PIH occurrence, although the process of diagnosing PIH might be potentially fallible with possible underascertainment [31]. In our study, the diagnosis of PIH is considered accurate because it was made by a single obstetrician (S.J.) on the basis of the same criteria. Further research is necessary to confirm the association between maternal educational level and PIH using a larger sample size.

There are several limitations in our study. First, this study was conducted retrospectively at a single center with a relatively small sample size. Because of the small number of participants with PIH, we were not able to separate PIH into true preeclampsia (hypertension with proteinuria). To analyze the subtype, a study with a larger sample size should be performed. Second, the percentage of participants who graduated from high school or less was also relatively small and the proportion of the high educational group was high in comparison with the general population among those who delivered babies in Japan [22] suggesting sampling bias. However, we demonstrated an association between educational level and BP even among higher SES pregnant mothers. Third, unmeasured possible mediators and

confounders, e.g., maternal birth weight [32], exercise during pregnancy [33], and neighborhood effects [34], might exist. Fourth, since some of the participants were recruited beyond the first trimester, and pre-pregnancy body weight and height were self-reported by questionnaire, pre-pregnancy BMI might be underestimated. Thus, a further study that replicates these study findings using a population-based, multicenter, large prospective cohort study is essential.

On the basis of our study results, a policy on body weight control may be effective targeting people who do not attend junior college or university in order to prevent higher BP during pregnancy. In the United State, where child and adolescent obesity is a growing problem, health education on body weight control is implemented among high school students aiming for future BP control [35]. A similar health policy might be effective in Japan.

Conclusion

In conclusion, among Japanese pregnant women, the low educational group had significantly higher systolic BP than the mid and high educational groups in early pregnancy and mid-pregnancy. The pre-pregnancy BMI mediated the association between educational levels and BP. Thus, education on body weight control in high schools might be useful to control BP during pregnancy, which would prevent the onset of PIH.

Abbreviations

BP, Blood pressure; BMI, Body mass index; CI, Confidence interval; DM, diabetes mellitus; OR, Odds ratio; PIH, Pregnancy induced hypertension; SD, Standard deviation; SES, Socioeconomic status

Competing interests

The authors declared no competing interest.

Authors' contributions

SJ analyzed data and wrote first draft of manuscript. TF conceived design, interpreted the results, and finalized the manuscript. AH and HS contributed interpretation of data. NA and YO contributed acquisition of data. All authors read and approved the final manuscript.

Acknowledgements

We thank the principal members of the T-CHILD study, especially Dr. H. Saito, N. Sakamoto, and S. Aizawa. We also appreciate the help of Ms. S. Saito and Ms. C. Matsuzawa in the management of the cohort data. This study was supported by a Grant of the National Center for Child Health and Development (20A-1).

References

1. Churchill D, Perry IJ, Beevers DG: **Ambulatory blood pressure in pregnancy and fetal growth.** *Lancet* 1997, **349**(9044):7–10.
2. Waugh J, Perry IJ, Halligan AW, De Swiet M, Lambert PC, Penny JA, Taylor DJ, Jones DR, Shennan A: **Birth weight and 24-hour ambulatory blood pressure in nonproteinuric hypertensive pregnancy.** *Am J Obstet Gynecol* 2000, **183**(3):633–637.
3. Poon LC, Kametas NA, Pandeva I, Valencia C, Nicolaides KH: **Mean arterial pressure at 11(+0) to 13(+6) weeks in the prediction of preeclampsia.** *Hypertension* 2008, **51**(4):1027–1033.
4. Cnossen JS, Vollebregt KC, de Vrieze N, ter Riet G, Mol BW, Franx A, Khan KS, van der Post JA: **Accuracy of mean arterial pressure and blood pressure measurements in predicting pre-eclampsia: systematic review and meta-analysis.** *BMJ* 2008, **336**(7653):1117–1120.
5. Schroeder BM: **ACOG practice bulletin on diagnosing and managing preeclampsia and eclampsia.** American college of obstetricians and gynecologists. *Am Fam Physician* 2002, **66**(2):330–331.
6. Arnadottir GA, Geirsson RT, Arngrimsson R, Jonsdottir LS, Olafsson O: **Cardiovascular death in women who had hypertension in pregnancy: a case-control study.** *BJOG* 2005, **112**(3):286–292.
7. Vikse BE, Irgens LM, Leivestad T, Skjaerven R, Iversen BM: **Preeclampsia and the risk of end-stage renal disease.** *N Engl J Med* 2008, **359**(8):800–809.
8. Colhoun HM, Hemingway H, Poulter NR: **Socio-economic status and blood pressure: an overview analysis.** *J Hum Hypertens* 1998, **12**(2):91–110.
9. Diez Roux AV, Chambless L, Merkin SS, Arnett D, Eigenbrodt M, Nieto FJ, Szklo M, Sorlie P: **Socioeconomic disadvantage and change in blood pressure associated with aging.** *Circulation* 2002, **106**(6):703–710.
10. Loucks EB, Abrahamowicz M, Xiao Y, Lynch JW: **Associations of education with 30 year life course blood pressure trajectories: framingham offspring study.** *BMC Public Health* 2011, **11**:139.
11. Banyard VL, Williams LM, Siegel JA: **The impact of complex trauma and depression on parenting: an exploration of mediating risk and protective factors.** *Child Maltreat* 2003, **8**(4):334–349.
12. Wilson BJ, Watson MS, Prescott GJ, Sunderland S, Campbell DM, Hannaford P, Smith WC: **Hypertensive diseases of pregnancy and risk of hypertension and stroke in later life: results from cohort study.** *BMJ* 2003, **326**(7394):845.

13. Silva LM, Steegers EA, Burdorf A, Jaddoe VW, Arends LR, Hofman A, Mackenbach JP, Raat H: **No midpregnancy fall in diastolic blood pressure in women with a low educational level: the generation R study.** *Hypertension* 2008, **52**(4):645–651.
14. Stamler R, Shipley M, Elliott P, Dyer A, Sans S, Stamler J: **Higher blood pressure in adults with less education. Some explanations from INTERSALT.** *Hypertension* 1992, **19**(3):237–241.
15. Siegrist J, Bernhardt R, Feng ZC, Schettler G: **Socioeconomic differences in cardiovascular risk factors in china.** *Int J Epidemiol* 1990, **19**(4):905–910.
16. Steptoe A, Shamaei-Tousi A, Gylfe A, Henderson B, Bergstrom S, Marmot M: **Socioeconomic status, pathogen burden and cardiovascular disease risk.** *Heart* 2007, **93**(12):1567–1570.
17. Molarius A, Seidell JC, Sans S, Tuomilehto J, Kuulasmaa K: **Educational level, relative body weight, and changes in their association over 10 years: an international perspective from the WHO MONICA project.** *Am J Public Health* 2000, **90**(8):1260–1268.
18. Tada N, Maruyama C, Koba S, Tanaka H, Birou S, Teramoto T, Sasaki J: **Japanese dietary lifestyle and cardiovascular disease.** *J Atheroscler Thromb* 2011.
19. Kagamimori S, Gaina A, Nasermoaddeli A: **Socioeconomic status and health in the japanese population.** *Soc Sci Med* 2009, **68**(12):2152–2160.
20. JSSHP: *Guideline 2009 for care and treatment of hypertension in pregnancy.* 1st edition. Tokyo MEDICAL VIEW; 2009.
21. Kobayashi S, Murakami K, Sasaki S, Okubo H, Hirota N, Notsu A, Fukui M, Date C: **Comparison of relative validity of food group intakes estimated by comprehensive and brief-type self-administered diet history questionnaires against 16 d dietary records in japanese adults.** *Public Health Nutr* 2011:1–12.
22. Ito J, Fujiwara T, Barr RG: **Is paternal infant care associated with breastfeeding? a population-based study in japan.** *J Hum Lact.* in press.
23. Jwa SC, Arata N, Sakamoto N, Watanabe N, Aoki H, Kurauchi-Mito A, Dongmei Q, Ohya Y, Ichihara A, Kitagawa M: **Prediction of pregnancy-induced hypertension by a shift of blood pressure class according to the JSH 2009 guidelines.** *Hypertens Res* 2011, **34**(11):1203–1208.
24. Albert MA, Glynn RJ, Buring J, Ridker PM: **Impact of traditional and novel risk factors on the relationship between socioeconomic status and incident cardiovascular events.** *Circulation* 2006, **114**(24):2619–2626.
25. Mikkelsen KL, Wiinberg N, Hoegholm A, Christensen HR, Bang LE, Nielsen PE, Svendsen TL, Kampmann JP, Madsen NH, Bentzon MW: **Smoking related to 24-h ambulatory blood pressure and heart rate: a study in 352 normotensive danish subjects.** *Am J Hypertens* 1997, **10**(5 Pt 1):483–491.

26. Perkins KA, Epstein LH, Marks BL, Stiller RL, Jacob RG: **The effect of nicotine on energy expenditure during light physical activity.** *N Engl J Med* 1989, **320**(14):898–903.
27. Bakker R, Steegers EA, Mackenbach JP, Hofman A, Jaddoe VW: **Maternal smoking and blood pressure in different trimesters of pregnancy: the generation R study.** *J Hypertens* 2010, **28**(11):2210–2218.
28. Silva LM, Coolman M, Steegers EA, Jaddoe VW, Moll HA, Hofman A, Mackenbach JP, Raat H: **Low socioeconomic status is a risk factor for preeclampsia: the generation R study.** *J Hypertens* 2008, **26**(6):1200–1208.
29. Silva L, Coolman M, Steegers E, Jaddoe V, Moll H, Hofman A, Mackenbach J, Raat H: **Maternal educational level and risk of gestational hypertension: the generation R study.** *J Hum Hypertens* 2008, **22**(7):483–492.
30. Haelterman E, Qvist R, Barlow P, Alexander S: **Social deprivation and poor access to care as risk factors for severe pre-eclampsia.** *Eur J Obstet Gynecol Reprod Biol* 2003, **111**(1):25–32.
31. Savitz DA, Zhang J: **Pregnancy-induced hypertension in north carolina, 1988 and 1989.** *Am J Public Health* 1992, **82**(5):675–679.
32. Innes KE, Byers TE, Marshall JA, Baron A, Orleans M, Hamman RF: **Association of a woman's own birth weight with her subsequent risk for pregnancy-induced hypertension.** *Am J Epidemiol* 2003, **158**(9):861–870.
33. Saftlas AF, Logsden-Sackett N, Wang W, Woolson R, Bracken MB: **Work, leisure-time physical activity, and risk of preeclampsia and gestational hypertension.** *Am J Epidemiol* 2004, **160**(8):758–765.
34. Vinikoor-Imler LC, Messer LC, Evenson KR, Laraia BA: **Neighborhood conditions are associated with maternal health behaviors and pregnancy outcomes.** *Soc Sci Med* 2011.
35. CDC: **School health guidelines to promote healthy eating and physical activity.** *MMWR Recomm Rep* 2011, **60**(RR-5):1–76.

Molecular and Clinical Studies in 138 Japanese Patients with Silver-Russell Syndrome

Tomoko Fuke^{1,2}, Seiji Mizuno³, Toshiro Nagai⁴, Tomonobu Hasegawa², Reiko Horikawa⁵, Yoko Miyoshi⁶, Koji Muroya⁷, Tatsuro Kondoh⁸, Chikahiko Numakura⁹, Seiji Sato¹⁰, Kazuhiko Nakabayashi¹¹, Chiharu Tayama¹¹, Kenichiro Hata¹¹, Shinichiro Sano^{1,12}, Keiko Matsubara¹, Masayo Kagami¹, Kazuki Yamazawa¹, Tsutomu Ogata^{1,12*}

1 Department of Molecular Endocrinology, National Research Institute for Child Health and Development, Tokyo, Japan, **2** Department of Pediatrics, Keio University School of Medicine, Tokyo, Japan, **3** Department of Pediatrics, Central Hospital, Aichi Human Service Center, Aichi, Japan, **4** Department of Pediatrics, Dokkyo Medical University Koshigaya Hospital, Saitama, Japan, **5** Division of Endocrinology and Metabolism, National Center for Child Health and Development, Tokyo, Japan, **6** Department of Pediatrics, Osaka University Graduate School of Medicine, Suita, Japan, **7** Department of Endocrinology and Metabolism, Kanagawa Children's Medical Center, Kanagawa, Japan, **8** Division of Developmental Disability, Misakaenosono Mutsumi Developmental, Medical, and Welfare Center, Isahaya, Japan, **9** Department of Pediatrics, Yamagata University School of Medicine, Yamagata, Japan, **10** Department of Pediatrics, Saitama Municipal Hospital, Saitama, Japan, **11** Department of Maternal-Fetal Biology, National Research Institute for Child Health and Development, Tokyo, Japan, **12** Department of Pediatrics, Hamamatsu University School of Medicine, Hamamatsu, Japan

Abstract

Background: Recent studies have revealed relative frequency and characteristic phenotype of two major causative factors for Silver-Russell syndrome (SRS), i.e. epimutation of the *H19*-differentially methylated region (DMR) and uniparental maternal disomy 7 (upd(7)mat), as well as multilocus methylation abnormalities and positive correlation between methylation index and body and placental sizes in *H19*-DMR epimutation. Furthermore, rare genomic alterations have been found in a few of patients with idiopathic SRS. Here, we performed molecular and clinical findings in 138 Japanese SRS patients, and examined these matters.

Methodology/Principal Findings: We identified *H19*-DMR epimutation in cases 1–43 (group 1), upd(7)mat in cases 44–52 (group 2), and neither *H19*-DMR epimutation nor upd(7)mat in cases 53–138 (group 3). Multilocus analysis revealed hyper- or hypomethylated DMRs in 2.4% of examined DMRs in group 1; in particular, an extremely hypomethylated *ARHI*-DMR was identified in case 13. Oligonucleotide array comparative genomic hybridization identified a ~3.86 Mb deletion at chromosome 17q24 in case 73. Epigenotype-phenotype analysis revealed that group 1 had more reduced birth length and weight, more preserved birth occipitofrontal circumference (OFC), more frequent body asymmetry and brachydactyly, and less frequent speech delay than group 2. The degree of placental hypoplasia was similar between the two groups. In group 1, the methylation index for the *H19*-DMR was positively correlated with birth length and weight, present height and weight, and placental weight, but with neither birth nor present OFC.

Conclusions/Significance: The results are grossly consistent with the previously reported data, although the frequency of epimutations is lower in the Japanese SRS patients than in the Western European SRS patients. Furthermore, the results provide useful information regarding placental hypoplasia in SRS, clinical phenotypes of the hypomethylated *ARHI*-DMR, and underlying causative factors for idiopathic SRS.

Citation: Fuke T, Mizuno S, Nagai T, Hasegawa T, Horikawa R, et al. (2013) Molecular and Clinical Studies in 138 Japanese Patients with Silver-Russell Syndrome. PLoS ONE 8(3): e60105. doi:10.1371/journal.pone.0060105

Editor: Monica Miozzo, Università degli Studi di Milano, Italy

Received: September 7, 2012; **Accepted:** February 21, 2013; **Published:** March 22, 2013

Copyright: © 2013 Fuke et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was funded by Grants-in-Aid for Scientific Research (A) (22249010) and Research (B) (21028026) from the Japan Society for the Promotion of Science (<http://www.jsps.go.jp/english/index.html>), by Grant for Research on Rare and Intractable Diseases (H24-042) from the Ministry of Health, Labor and Welfare (<http://www.mhlw.go.jp/english/>), and by Grant for National Center for Child Health and Development (23A-1) (<http://www.ncchd.go.jp/English/Englishtop.htm>). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: tomogata@hama-med.ac.jp

Introduction

Silver-Russell syndrome (SRS) is a rare congenital developmental disorder characterized by pre- and postnatal growth failure, relative macrocephaly, triangular face, hemihypotrophy, and fifth-finger clinodactyly [1]. Recent studies have shown that epimutation (hypomethylation) of the paternally derived differentially methylated region (DMR) in the upstream of *H19* (*H19*-DMR) on

chromosome 11p15.5 and maternal uniparental disomy for chromosome 7 (upd(7)mat) account for ~45% and 5–10% of SRS patients, respectively [1,2]. In this regard, phenotypic assessment has suggested that birth length and weight are more reduced and characteristic body features are more frequent in patients with *H19*-DMR epimutation than in those with upd(7)mat, whereas developmental delay tends to be more

frequent in patients with upd(7)mat than in those with *H19*-DMR epimutation [3,4]. Furthermore, consistent with the notion that imprinted genes play an essential role in placental growth and development [5], placental hypoplasia has been found in both *H19*-DMR epimutation and upd(7)mat [4,6], although comparison of placental weight has not been performed between *H19*-DMR hypomethylation and upd(7)mat. In addition, multilocus hypo- or hypermethylation and positive correlation between methylation index (MI, the ratio of methylated clones) and body and placental sizes have been reported in patients with *H19*-DMR epimutation [4,7–9], and several types of rare genomic alterations have been identified in a few of SRS patients [1,10–12].

Here, we report on molecular and clinical findings in 138 Japanese SRS patients, and discuss on the results obtained in this study.

Patients and Methods

Ethics statement

This study was approved by the Institutional Review Board Committee at the National Center for Child Health and Development. The parents of the affected children and the adult patients who can express an intention by themselves have given written informed consent to participate in this study and to publish their molecular and clinical data.

Patients

This study consisted of 138 Japanese patients (66 males and 72 females) with SRS phenotype aged 0–30 years (median 4.1 years), including 64 previously reported patients (20 patients with variable degrees of *H19*-DMR epimutation, three patients with upd(7)mat, one patient with 46,XY/46,XY,upd(7)mat mosaicism in whom upd(7)mat cells accounted for 91–92% of leukocytes and salivary cells and for 11% of placental tissue, and 40 patients of unknown cause) [4,6,13]. The 138 patients had a normal karyotype in all the ≥ 50 lymphocytes examined, and satisfied the selection criteria proposed by Netchine et al. [14], i.e., birth length and/or birth weight ≤ -2 standard deviation score (SDS) for gestational age as a mandatory criteria plus at least three of the following five features: (i) postnatal short stature (≤ -2 SDS) at 2 year of age or at the nearest measure available, (ii) relative macrocephaly at birth, i.e., SDS for birth length or birth weight minus SDS for birth occipitofrontal circumference (OFC) ≤ -1.5 , (iii) prominent forehead during early childhood, (iv) body asymmetry, and (v) feeding difficulties during early childhood. Birth and present length/height, weight, and OFC were assessed by the gestational/postnatal age- and sex-matched Japanese reference data from the Ministry of Health, Labor, and Welfare and the Ministry of Education, Science, Sports and Culture. Placental weight was assessed by the gestational age-matched Japanese reference data [15]. Clinical features were evaluated by clinicians at different hospitals who participated in this study, using the same clinical datasheet. The SRS patients were classified into three groups by the molecular studies, i.e., those with *H19*-DMR hypomethylation (epimutation) (group 1), those with upd(7)mat (group 2), and the remaining patients (group 3).

Primers and samples

Primers utilized in this study are shown in Table S1. Leukocyte genomic DNA samples were examined in this study.

Methylation analysis

We performed pyrosequencing analysis for the *H19*-DMR encompassing the 6th CTCF (CCCTC-binding factor) binding site

that functions as the primary regulator for the monoallelic *IGF2* and *H19* expressions [16–18], using bisulfite treated leukocyte genomic DNA samples of all the 138 patients. The procedure was as described in the manufacturer's instructions (Qiagen, Valencia, CA, USA). In brief, a 279 bp region was PCR-amplified with a primer set (PyF and PyR) for both methylated and unmethylated clones, and a sequence primer (SP) was hybridized to a single-stranded PCR products. Subsequently, the MIs were obtained for four CpG dinucleotides (CG5–CG7 and CG9), using PyroMark Q24 (Qiagen) (the MI for CG8 was not obtained, because the “C” residue of CG8 constitutes a C/T SNP) (Figure 1A). The PyF/PyR and SP were designed by PyroMark Assay Design Software Ver2.0. While the PyF sequence contains a SNP (*rs11564736*) with a mean minor allele frequency of 5% in multiple populations, the minor allele frequency is 0% in the Japanese as well as in the Asian populations (http://browser.1000genomes.org/Homo_sapiens/Variation/Population?db=core;r=11:2020801–2021801;v=rs11564736;vdb=variation;vf=7864021). Thus, we utilized this PyF.

We also carried out combined bisulfite restriction analysis (COBRA) for the *H19*-DMR. The methods were as described previously [4]. In short, a 435 bp region was PCR-amplified with a primer set (CoF and CoR) that hybridize to both methylated and unmethylated clones, and MIs were obtained for two CpG dinucleotides (CG5 and CG16) after digestion of the PCR products with methylated allele-specific restriction enzymes (*Hpy188I* and *AflIII*) (Figure 1A).

Thus, we could examine CG5 by both pyrosequencing and COBRA. While we also attempted to analyze CG16 by both methods, it was impossible to design an SP for the analysis of CG16 (although we could design an SP between CG11 and CG12, clear methylation data were not obtained for CG16, probably because of the distance between the SP and CG16).

In addition, we performed COBRA for the KvDMR1 in all the 138 patients (Figure S1A) because of the possibility that epimutation of the KvDMR1 could lead to SRS phenotype via some mechanism(s) such as overexpression of a negative growth regulator *CDKN1C* [19], and for multiple DMRs on various chromosomes in patients in whom relatively large amount of DNA samples were available, as reported previously [4,20,21]. To define the reference ranges of MIs (minimum ~ maximum), 50 control subjects were similarly studied with permission.

To screen upd(7)mat, PCR amplification was performed for the *MEST*-DMR on chromosome 7q32.2 in all the 138 patients, using methylated and unmethylated allele-specific PCR primer sets, as reported previously [6] (Figure 2A). In addition, bisulfite sequencing and direct sequencing for the primer binding sites for the *ARHI*-DMR analysis were performed in a patient (case 13) with an extremely low MI for the *ARHI*-DMR.

Microsatellite analysis

Microsatellite analysis was performed for four loci within a ~4.5 Mb telomeric 11p region (*D11S2071*, *D11S922*, *D11S1318*, and *D11S988*) in patients with hypomethylated *H19*-DMR, to examine the possibility of upd(11p)mat involving the *H19*-DMR. Microsatellite analysis was also carried out for nine loci widely dispersed on chromosome 7 (Table S2) in patients with abnormal methylation patterns of the *MEST*-DMR, to examine the possibility of upd(7)mat and to infer the underlying causes for upd(7)mat, i.e., trisomy rescue, gamete complementation, monosomy rescue, and post-fertilization mitotic error [22]. The methods have been reported previously [4,6].




Figure 1. Methylation analysis of the *H19*-DMR, using bisulfite-treated genomic DNA. A. Schematic representation of a segment encompassing 21 CpG dinucleotides (CG1→CG21) within the *H19*-DMR. The cytosine residues at the CpG dinucleotides are usually methylated after paternal transmission (filled circles) and unmethylated after maternal transmission (open circles). The CTCF binding site 6 (CTCF6) is indicated with a blue rectangle; the cytosine residue at CG8 constitutes a C/T SNP (indicated with a gray rectangle). For pyrosequencing analysis, a 279 bp segment was PCR-amplified with PyF & PyR primers, and a sequence primer (SP) was hybridized to a single-stranded PCR products. Subsequently, the MIs were obtained for four CpG dinucleotides (CG5–CG7 and CG9) (indicated with a yellow rectangle). For COBRA, a 435 bp region was PCR-amplified with CoF & CoR primers, and the PCR product was digested with methylated allele-specific restriction enzymes to examine the methylation pattern of CG5 and CG16 (the PCR products is digested with *Hpy188I* when the cytosine residue at CG5 is methylated and with *AflIII* when the cytosine residue at CG16 is methylated) (indicated with orange rectangles). *IGF2* is a paternally expressed gene, and *H19* is a maternally expressed gene. The stippled ellipse indicates the enhancer for *IGF2* and *H19*. B. Pyrosequencing data. Left part: Representative results indicating the MIs for CG5–CG7 and CG9. CG5–CG7 and CG9 are hypomethylated in case 1, and similarly methylated between case 53 and a control subject. Right part: Histograms showing the distribution of the MIs (the horizontal axis: the methylation index; and the vertical axis: the patient number). Forty-three SRS patients with low MIs are shown in red. C. COBRA data. Left part: Representative findings of PCR products loaded onto a DNA 1000 LabChip (Agilent, Santa Clara, CA, USA) after digestion with *Hpy188I* or *AflIII*. U: unmethylated clone specific bands; M: methylated clone specific bands; and BWS: Beckwith-Wiedemann syndrome patient with upd(11p15)pat. Right part: Histograms showing the distribution of the MIs.
doi:10.1371/journal.pone.0060105.g001

Oligoarray comparative genomic hybridization (CGH)

We performed oligoarray CGH in the 138 SRS patients, using a genome-wide 4×180K Agilent platform catalog array and a custom-build high density oligoarray for the 11p15.5, 7p12.2, 12q14, and 17q24 regions where rare copy number variants have been identified in several SRS patients [1,10–12] as well as for the 7q32–qter region involved in the segmental upd(7)mat in four SRS patients [23–25]. The custom-build high density oligoarray contained 3,214 probes for 7p12.2, 434 probes for 7q32, 23,162

probes for 12q14, and 39,518 probes for 17q24, together with ~10,000 reference probes for other chromosomal region (Agilent Technologies, Palo Alto, CA, USA). The procedure was as described in the manufacturer's instructions.

Statistical analysis

After examining normality by χ^2 test, the variables following the normal distribution were expressed as the mean±SD, and those not following the normal distribution were expressed with the




Figure 2. Methylated and unmethylated allele-specific PCR analysis for the *MEST*-DMR. A. Schematic representation of the *MEST*-DMR. The cytosine residues at the CpG dinucleotides are usually unmethylated after paternal transmission (open circles) and methylated after maternal transmission (filled circles). The PCR primers have been designed to hybridize either methylated or unmethylated clones. B. The results of methylation analysis with methylated and unmethylated allele-specific primers.

doi:10.1371/journal.pone.0060105.g002

median and range. Statistical significance of the mean was analyzed by Student's *t*-test or Welch's *t*-test after comparing the variances by *F* test, that of the median by Mann-Whitney's *U*-test, that of the frequency by Fisher's exact probability test, and that of the correlation by Pearson's correlation coefficient after confirming the normality of the variables. $P < 0.05$ was considered significant.

Results

Identification of *H19*-DMR hypomethylation

Representative findings are shown in Figure 1B and 1C, and the MIs are summarized in Table 1. Overall, the MIs obtained by the pyrosequencing analysis tended to be lower and distributed more narrowly than those obtained by the COBRA. Despite such difference, the MIs obtained by the pyrosequencing analysis for CG5–CG7 and CG9 and by the COBRA for CG5 and CG16 were invariably below the normal range in the same 43 patients (cases 1–43) (group 1). By contrast, the MIs were almost invariably within the normal range in the remaining 95 patients, while the MIs obtained by the pyrosequencing analysis slightly (1–2%) exceeded the normal range in the same three patients (cases 136–138).

In the 43 cases of group 1, microsatellite analysis for four loci at the telomeric 11p region excluded maternal upd in 14 cases in whom parental DNA samples were available; in the remaining 29 cases, microsatellite analysis identified two alleles for at least one locus, excluding maternal uniparental isodisomy for this region. Furthermore, oligoarray CGH for the chromosome 11p15.5 region showed no copy number alteration such as duplication of maternally derived *H19*-DMR and deletion of paternally derived

Table 1. The methylation indices (%) for the *H19*-DMR.

	Cases 1–43	Cases 44–138	Control subjects
Pyrosequencing analysis			
CG5	4–24	35–50	33–48
CG6	5–26	36–53	34–51
CG7	4–24	35–49	30–47
CG9	5–23	34–48	30–46
COBRA			
CG5 (<i>Hpy</i> 188I)	3.3–35.1	37.8–60.8	36.2–58.5
CG16 (<i>Afl</i> III)	4.1–35.0	43.0–59.4	38.7–60.0

The position of examined CpG dinucleotides (CG5–7, CG9, and CG16) is shown in Figure 1A.

COBRA: combined bisulfite restriction analysis.

doi:10.1371/journal.pone.0060105.t001

H19-DMR. For the KvDMR1, the MIs of the 138 patients remained within the reference range (Fig. S1B and C).

Identification of upd(7)mat

Methylation analysis for the *MEST*-DMR revealed that unmethylated bands were absent from eight patients and remained faint in a single patient (cases 44–52) (group 2) (Figure 2B). Subsequent microsatellite analysis confirmed upd(7)mat in the eight patients and mosaic upd(7)mat in the remaining one patient, and indicated trisomy rescue or gamete complementation type upd(7)mat in cases 44–48, monosomy rescue or post-fertilization mitotic error type upd(7)mat in cases 49–51, and post-fertilization mitotic error type mosaic upd(7)mat in case 52 (Table S2).

Multiple DMR analysis

We examined 17 autosomal DMRs other than the *H19*-DMR in 14 patients in group 1, four patients in group 2, and 20 patients in group 3, and the *XIST*-DMR in eight female patients in group 1, one female patient in group 2, and five female patients in group 3 (Table S3). The MIs outside the reference ranges were identified in five of 14 examined cases (35.7%) and six of a total of 246 examined DMRs (2.4%) in group 1. In particular, a single case with the mean MI value of 23 obtained by the pyrosequencing analysis for CG5–CG7 and CG9 had an extremely low MI for the *ARHI*-DMR (case 13 of group 1). This extreme hypomethylation was confirmed by bisulfite sequencing, and direct sequencing showed normal sequences of the primer-binding sites, thereby excluding the possibility that such an extremely low MI could be due to insufficient primer hybridization because of the presence of a nucleotide variation within the primer-binding sites (Figure 3). Furthermore, no copy number variation involving the *ARHI*-DMR was identified by CGH analysis using a genome-wide catalog array. Consistent with upd(7)mat, three DMRs on chromosome 7 were extremely hypermethylated in four examined cases of group 2. Only a single DMR was mildly hypermethylated in a total of 345 examined DMRs in group 3. The abnormal MIs, except for those for the *H19*-DMR in group 1 and for the three DMRs on chromosome 7 in group 2, were confirmed by three times experiments.

Oligonucleotide array CGH

A ~3.86 Mb deletion at chromosome 17q24 was identified in a single patient (case 73 of group 3) (Figure 4).




Figure 3. Analysis of the ARHI-DMR in case 13. For bisulfite sequencing, each line indicates a single clone, and each circle denotes a CpG dinucleotide; the cytosine residues at the CpG dinucleotides are usually unmethylated after paternal transmission (open circles) and methylated after maternal transmission (filled circles). Electrochromatograms delineate the sequences of the primer binding sites utilized for the methylation analysis. doi:10.1371/journal.pone.0060105.g003

Epigenotype-phenotype analysis

Clinical findings of SRS patients in groups 1–3 are summarized in Table 2. All the patients met the mandatory criteria, and most patients in each group had severely reduced birth length and weight (both ≤ -2 SDS). For the five clinical features utilized as scoring system criteria, while 23.2% of patients in group 1 and 22.2% of patients in group 2 exhibited all the five features, there was no patient in group 3 who was positive for all the five features. By contrast, while 39.5% of patients in group 1 and 33.3% of patients in group 2 manifested just three of the five features, 77.6% of patients in group 3 were positive for just three features. In particular, the frequencies of relative macrocephaly at birth and body asymmetry were low in group 3, while those of the remaining three scoring system criteria including prominent forehead during early childhood were similar among groups 1–3.

Phenotypic comparison between groups 1 and 2 revealed that birth length and weight were more reduced and birth OFC was

more preserved in group 1 than in group 2, despite comparable gestational age. In the postnatal life, present height and weight became similar between the two groups, whereas present OFC became significantly smaller in group 1 than in group 2. Body asymmetry and brachydactyly were more frequent and speech delay was less frequent in group 1 than in group 2. Placental weight was similar between the two groups, and became more similar after excluding case 52 with mosaic upd(7)mat (see legends for Table 2). Parental age at childbirth was also similar between the two groups. In group 2, placental weight was grossly similar among examined cases, as was parental age at childbirth (see legends for Table 2).

Case 13 with an extremely low MI for the ARHI-DMR and case 73 with a cryptic deletion at chromosome 17q24 had no specific phenotype other than SRS-like phenotype (Table S4). However, of the five clinical features utilized as scoring system criteria, all the five features were exhibited by case 13 and just three features were




Figure 4. Oligonucleotide array CGH in case 73, showing a ~3.86 Mb deletion at chromosome 17q24. The black, the red, and the green dots denote signals indicative of the normal, the increased ($>+0.5$), and the decreased (<-1.0) copy numbers, respectively. The horizontal bar with arrowheads indicates a ~2.3 Mb deletion identified in a patient with Carney complex and SRS-like phenotype [44], and the black square represent a ~65 kb segment harboring the breakpoint of a *de novo* translocation 46,XY,t(1;17)(q24;q23-q24) identified in a patient with SRS phenotype [45,46]. doi:10.1371/journal.pone.0060105.g004

Table 2. Phenotypic comparison in three groups of patients with Silver-Russell syndrome.

	<i>H19-DMR</i> hypomethylation	Upd(7)mat	Unknown	<i>P</i> -value		
	(Group 1)	(Group 2)	(Group 3)	G1 vs. G2	G1 vs. G3	G2 vs. G3
Patient number	43 (31.2%)	9 (6.5%)	85 (62.0%)			
Mandatory criteria	43/43 (100%)	9/9 (100%)	85/85 (100%)	1.000	1.000	1.000
Scoring system criteria (5/5)	10/43 (23.2%)	2/9 (22.2%)	0/85 (0.00%)	0.965	1.52 × 10⁻⁴	2.58 × 10⁻²
Scoring system criteria (4/5)	16/43 (37.2%)	4/9 (44.4%)	19/85 (22.4%)	0.792	1.45 × 10⁻²	0.145
Scoring system criteria (3/5)	17/43 (39.5%)	3/9 (33.3%)	66/85 (77.6%)	0.821	7.17 × 10⁻⁴	0.161
Gestational age (weeks:days)	38:0 (34:3~40:0) (n = 36)	38:0 (34:4~40:0) (n = 9)	37:6 (27:1~41:4) (n = 65)	0.877	0.120	0.450
BL (SDS)	-4.13 ± 2.01 (n = 31)	-3.18 ± 1.16 (n = 9)	-2.93 ± 1.43 (n = 60)	2.67 × 10⁻²	6.69 × 10⁻⁵	0.619
BW (SDS)	-3.50 ± 0.85 (n = 42)	-2.90 ± 0.64 (n = 9)	-2.71 ± 1.14 (n = 64)	3.28 × 10⁻²	5.87 × 10⁻⁴	0.640
BL ≤ -2 SDS and/or BW ≤ -2 SDS*	43/43 (100%)	9/9 (100%)	85/85 (100%)	1.000	1.000	1.000
BL ≤ -2 SDS and BW ≤ -2 SDS	39/43 (90.7%)	7/9 (77.8%)	76/85 (89.4%)	0.474	0.821	0.304
BOFC (SDS)	-0.54 ± 1.22 (n = 29)	-1.44 ± 0.47 (n = 9)	-1.92 ± 1.09 (n = 48)	3.74 × 10⁻²	1.52 × 10⁻⁶	0.202
BL (SDS) - BOFC (SDS)	-3.70 ± 2.02 (n = 27)	-1.73 ± 1.20 (n = 9)	-0.943 ± 1.48 (n = 43)	1.02 × 10⁻²	3.40 × 10⁻⁹	0.111
BW (SDS) - BOFC (SDS)	-3.21 ± 1.20 (n = 27)	-1.53 ± 0.57 (n = 9)	-1.04 ± 1.55 (n = 48)	0.326	7.38 × 10⁻⁹	0.331
Relative macrocephaly at birth† BL or BW (SDS) - BOFC (SDS) ≤ -1.5	29/29 (100%)	7/9 (77.8%)	16/45 (35.6%)	0.341	3.67 × 10⁻⁸	2.05 × 10⁻²
Present age (years:months)	4.1 (0:6~30:6) (n = 31)	4.8 (2:4~25:2) (n = 9)	4.3 (0:1~18:6) (n = 60)	0.437	0.813	0.335
PH (SDS)	-3.58 ± 1.65 (n = 35)	-3.77 ± 1.13 (n = 9)	-3.17 ± 1.50 (n = 61)	0.757	0.218	0.253
PH ≤ -2 SDS (≥ 2 years)†	29/35 (82.5%)	8/9 (88.9%)	52/61 (85.2%)	0.760	0.758	0.772
PW (SDS)	-3.15 ± 1.16 (n = 32)	-2.77 ± 0.76 (n = 9)	-2.77 ± 1.34 (n = 59)	0.362	0.144	0.968
POFC (SDS)	-1.16 ± 1.18 (n = 21)	-0.01 ± 0.91 (n = 9)	-1.81 ± 1.57 (n = 35)	2.01 × 10⁻³	0.107	3.08 × 10⁻³
PH (SDS) - POFC (SDS)	-2.47 ± 1.63 (n = 16)	-3.62 ± 1.38 (n = 8)	-1.55 ± 1.82 (n = 35)	0.103	4.39 × 10⁻²	1.64 × 10⁻²
PW (SDS) - POFC (SDS)	-2.84 ± 1.31 (n = 21)	-2.69 ± 1.36 (n = 9)	-1.08 ± 1.71 (n = 35)	0.782	2.54 × 10⁻²	1.90 × 10⁻⁴
Relative macrocephaly at present PH or PW (SDS) - POFC (SDS) ≤ -1.5	20/21 (95.2%)	8/8 (100%)	29/43 (67.4%)	0.223	4.77 × 10⁻³	0.156
Triangular face during early childhood	42/43 (97.7%)	8/9 (88.9%)	65/65 (100%)	0.442	0.0773	5.98 × 10⁻³
Prominent forehead during early childhood†	31/37 (83.8%)	7/9 (100%)	41/53 (77.4%)	0.200	0.456	0.978
Ear anomalies	14/35 (40.0%)	3/9 (33.3%)	15/55 (27.3%)	0.717	0.290	0.823
Irregular teeth	12/26 (46.2%)	4/9 (44.4%)	12/45 (26.7%)	0.930	0.0968	0.291
Body asymmetry†	30/37 (81.1%)	3/9 (33.3%)	19/59 (32.2%)	4.77 × 10⁻³	3.51 × 10⁻⁶	0.947
Clinodactyly	29/37 (78.4%)	5/9 (55.6%)	50/58 (86.2%)	0.167	0.323	2.68 × 10⁻²
Brachydactyly	30/38 (78.9%)	2/9 (22.2%)	34/56 (60.7%)	1.16 × 10 ⁻³	0.0642	3.24 × 10⁻²
Syndactyly	3/36 (8.3%)	0/9 (0.00%)	3/52 (5.77%)	0.375	0.641	0.464
Simian crease	4/26 (15.4%)	2/7 (28.6%)	6/49 (12.2%)	0.429	0.705	0.252
Muscular hypotonia	17/32 (53.1%)	5/9 (55.6%)	12/50 (24.0%)	0.898	7.49 × 10⁻³	0.0564
Developmental delay	18/37 (48.6%)	6/9 (66.7%)	25/54 (46.3%)	0.337	0.826	0.262
Speech delay	8/31 (25.8%)	6/9 (66.7%)	18/43 (41.9%)	2.55 × 10⁻²	0.156	0.179
Feeding difficulty†	16/34 (47.1%)	6/9 (66.7%)	25/51 (49.0%)	0.301	0.860	0.333
Placental weight (SDS)	-2.10 ± 0.74 (n = 14)	-1.72 ± 0.74 (n = 6) ^a	-1.02 ± 0.86 (n = 18)	0.312	4.12 × 10⁻³	8.24 × 10⁻³
Paternal age at childbirth (years:months)	32:0 (19:0~52:0) (n = 24)	35:0 (27:0~48:0) (n = 9)	32:0 (25:0~46:0) (n = 45)	0.223	1.00	0.105
Maternal age at childbirth (years:months)	32:0 (19:0~43:0) (n = 25)	33:0 (25:0~42:0) (n = 9) ^b	30:0 (22:0~43:0) (n = 46)	0.275	0.765	0.117

BL: birth length; BW: birth weight; BOFC: birth occipitofrontal circumference; PH: present height; PW: present weight; POFC: present occipitofrontal circumference, and SDS: standard deviation score.

For body features, the denominators indicate the number of patients examined for the presence or absence of each feature, and the numerators represent the number of patients assessed to be positive for that feature.

*Mandatory criteria and †five clinical features utilized as selection criteria for Silver-Russell syndrome proposed by Netchine et al. [14].

Significant *P*-values (<0.05) are boldfaced.

^aPlacental weight SDS is -1.68, -2.55, -2.24, -1.12, -2.14 and -0.60 in case 46, 47, 49, 50, 51 and 52, respectively; the placental weight SDS is -1.95 ± 0.57 in five cases except for case 52 with mosaic upd(7)mat.

^bMaternal childbearing age is 32, 32, 33, 33, 42, 32, 34, 33, 25 and 36 years in case 44-52, respectively.

doi:10.1371/journal.pone.0060105.t002

manifested by case 73. In addition, cases 136–138 with slightly elevated MIs for CG5–CG7 and CG9, and cases with multilocus methylation abnormalities, had no particular phenotype other than SRS-compatible clinical features.

Correlation analysis

In group 1, the mean value of the MIs for CG5–CG7 and CG9 obtained by pyrosequencing analysis was positively correlated with the birth length and weight, the present height and weight, and the placental weight, but with neither the birth nor the present OFC (Table 3). Such correlations with the growth parameters were grossly similar but somewhat different for the MIs obtained by COBRA (Table S5). Furthermore, the placental weight was positively correlated with the birth weight and length, but not with the birth OFC. Such positive correlations were not found in groups 2 and 3.

Discussion

The present study identified hypomethylation of the *H19*-DMR and *upd(7)mat* in 31.2% and 6.5% of 138 Japanese SRS patients, respectively. In this regard, the normal KvDMR1 methylation patterns indicate that the aberrant methylation in 43 cases of group 1 is confined to the *H19*-DMR. Furthermore, oligoarray CGH excludes copy number variants involving the *H19*-DMR, and microsatellite analysis argues against segmental maternal isodisomy that could be produced by post-fertilization mitotic error [26]. These findings imply that the *H19*-DMR hypomethylation is due to epimutation (hypomethylation of the normally methylated *H19*-DMR of paternal origin).

The frequency of epimutations detected in this study is lower than that reported in Western European SRS patients [1,2,14], although the frequency of *upd(7)mat* is grossly similar between the two populations [2,11,14,27,28]. In this context, it is noteworthy that, of the five scoring system criteria, the frequencies of relative macrocephaly at birth and body asymmetry were low in group 3, while those of the remaining three scoring system criteria were similar among groups 1–3. Since relative macrocephaly and body asymmetry are characteristic of *H19*-DMR epimutation, the lack of these two features in a substantial fraction of cases in group 3 would primarily explain the low frequency of *H19*-DMR

epimutations in this study. In group 3, furthermore, the low prevalence of relative macrocephaly at birth appears to be discordant with the high prevalence of prominent forehead during early childhood. Since relative macrocephaly was evaluated by an objective method (SDS for birth length or birth weight minus SDS for birth OFC ≤ -1.5) and prominent forehead was assessed by a subjective impression of different clinicians, it is recommended to utilize relative macrocephaly as a more important and reliable feature in the scoring system than prominent forehead. In addition, the difference in the ethnic group might also be relevant to the low frequency of *H19*-DMR epimutations in this study.

Epigenotype-phenotype correlations in this study are grossly similar to those previously reported in Western European SRS patients [1–3]. Cases 1–43 in group 1 with *H19*-DMR epimutation had more reduced birth weight and length, more preserved birth OFC and more reduced present OFC, more frequent body features, and less frequent speech delay than case 44–52 in group 2 with *upd(7)mat*, although the difference in the prevalence of somatic features appears to be less remarkable in this study than in the previous studies [3,4]. This provides further support for the presence of relatively characteristic clinical features in *H19*-DMR epimutation and *upd(7)mat* [1–3]. In this context, previous studies have indicated biallelic *IGF2* expression in the human fetal choroid plexus, cerebellum, and brain, and monoallelic *IGF2* expression in the adult brain, while the precise brain tissue(s) with such a unique expression pattern remains to be clarified [29,30,31]. This may explain why the birth OFC is well preserved and the present OFC is reduced in group 1. However, since the difference in present OFC between groups 1 and 2 is not necessarily significant in the previous studies [32], the postnatal OFC growth awaits further investigations.

Placental weight was similarly reduced in groups 1 and 2. Thus, placental weight is unlikely to represent an indicator for the discrimination between the two groups, although the present data provide further support for imprinted genes being involved in placental growth, with growth-promoting effects of *PEGs* and growth-suppressing effects of *MEGs* [5,6]. It should be pointed out, however, that the placental hypoplasia could be due to some other genetic or environmental factor(s). In particular, while placental weight was apparently similar among cases of group 2, possible confined placental mosaicism [33,34] with trisomy for chromosome 7 may have exerted some effects on placental growth in cases with trisomy rescue type *upd(7)mat*.

Correlation analysis would imply that the *IGF2* expression level, as reflected by the MI of the *H19*-DMR, plays a critical role in the determination of pre- and postnatal body (stature and weight) and placental growth in patients with *H19*-DMR epimutation. Since the placental weight was positively correlated with the birth length and weight, the reduced *IGF2* expression level appears to have a similar effect on the body and the placental growth. Furthermore, the lack of correlations between the MI and birth and present OFC and between placental weight and birth OFC would be compatible with the above mentioned *IGF2* expression pattern in the central nervous system [29]. Although the MI would also reflect the *H19* expression level, this would not have a major growth effect. It has been implicated that *H19* functions as a tumor suppressor [35,36].

Multilocus analysis revealed co-existing hyper- and hypomethylated DMRs predominantly in cases of group 1, with frequencies of 35.7% of examined patients and 2.4% of examined DMRs. The results are grossly consistent with the previous data indicating that co-existing abnormal methylation patterns of DMRs are almost exclusively identified in patients with *H19*-DMR epimutation with frequencies of 9.5–30.0% of analyzed patients and 1.8–5.2% of a

Table 3. Correlation analyses in patients with *H19*-DMR hypomethylations.

Parameter 1	Parameter 2	r	P-value
Methylation index (%)* vs.	Birth length (SDS)	0.647	6.70 × 10 ⁻³
	Birth weight (SDS)	0.590	7.80 × 10 ⁻³
	Birth OFC (SDS)	0.190	0.498
	Present height (SDS)	0.612	5.33 × 10 ⁻³
	Present weight (SDS)	0.605	7.81 × 10 ⁻³
	Present OFC (SDS)	-0.166	0.647
Placental weight (SDS) vs.	Birth weight (SDS)	0.717	8.64 × 10 ⁻³
	Birth length (SDS)	0.636	2.63 × 10 ⁻²
	Birth OFC (SDS)	0.400	0.198

SDS: standard deviation score; and OFC: occipitofrontal circumference.

*The mean value of MIs for CG5, CG6, CG7, and CG9 obtained by pyrosequencing analysis.

Significant P-values (<0.05) are boldfaced.

doi:10.1371/journal.pone.0060105.t003

total of analyzed DMRs [7–9]. Notably, the co-existing methylation abnormalities were predominantly observed as mild hypermethylations of maternally methylated DMRs and were restricted to a single DMR or two DMRs in patients with multilocus abnormalities. Such findings are obviously inexplicable not only by assuming a ζ FP57 mutation that is known to cause severely abnormal methylation patterns of multiple DMRs or a ζ AC1 mutation that may affect methylation patterns of multiple DMRs [37–39], but also by assuming defective maintenance of methylation in the postzygotic period [7]. Thus, some factor(s) susceptible to the co-occurrence of hypomethylation of the *H19*-DMR and hypermethylation of other DMR(s) might be operating during a gametogenic or postzygotic period in cases with *H19*-DMR epimutation.

The patients with multilocus methylation abnormalities had no specific clinical features other than SRS-compatible phenotype. Previous studies have also indicated grossly similar SRS-like phenotype between patients with monolocus and multilocus hypomethylations [7], although patients with multilocus hypomethylation occasionally have apparently severe clinical phenotype [7]. These findings would argue for the notion that the *H19*-DMR epimutation has an (epi)dominant clinical effect. Indeed, *H19*-DMR hypomethylation has led to SRS-like phenotype in a patient with parthenogenetic chimerism/mosaicism [21], whereas *H19*-DMR hypermethylation has resulted in Beckwith-Wiedemann syndrome-like phenotype in patients with androgenetic mosaicism [40].

An extremely hypomethylated *ARHI*-DMR was found in case 13. In this regard, it is known that *ARHI* with a potentially cell growth suppressor function is normally expressed from paternally inherited chromosome with unmethylated *ARHI*-DMR [41]. Indeed, hypermethylation of the *ARHI*-DMR, which is predicted to result in reduced expression of *ARHI*, has been identified as a tumorigenic factor for several cancers with an enhanced cell growth function [42,43]. Thus, it is possible that hypomethylation of the *ARHI*-DMR has led to overexpression of *ARHI*, contributing to the development of typical SRS phenotype in the presence of a low but relatively preserved MI of the *H19*-DMR in case 13.

Oligonucleotide array CGH identified a ~3.86 Mb deletion at chromosome 17q24 in case 73 of group 3. This provides further support for the presence of rare copy number variants in several SRS patients and the relevance of non-imprinted gene(s) to the development of SRS [10]. Interestingly, the microdeletion overlap with that identified in a patient with Carney complex and SRS-like features [44], and the overlapping region encompasses a ~65 kb segment defining the breakpoint of a *de novo* reciprocal translocation involving 17q23–q24 in a patient with SRS-like phenotype (Figure 4) [45,46]. Furthermore, the translocation breakage has affected *KPNA2* involved in the nuclear transport of proteins [46–48]. Thus, *KPNA2* has been regarded as a candidate gene for SRS, although mutation analysis of *KPNA2* has failed to detect a disease-causing mutation in SRS patients [49].

Lastly, it would be worth discussing on the comparison between pyrosequencing analysis and COBRA. Since the same 43 patients were found to have low MIs by both analyses, this implies that both methods can be utilized as a diagnostic tool. While the distribution of the MIs was somewhat different between the two methods, this would primarily be due to the difference in the employed methods such as the hybridization efficiency of utilized primers. Importantly, pyrosequencing analysis was capable of studying plural CpG dinucleotides at the C/TCF6 binding site, whereas COBRA examined only single CpG dinucleotides outside the C/TCF6 binding site. Thus, the MIs obtained by pyrosequencing analysis would be more accurate than those obtained by

COBRA in terms of *IGF2* expression levels, and this would underlie the reasonable correlations of MIs yielded by pyrosequencing analysis with body and placental growth parameters.

In summary, the present study provides useful information for the definition of molecular and clinical findings in SRS. However, several matters still remain to be elucidated, including underlying mechanisms in SRS patients with no *H19*-DMR epimutation or upd(7)mat and the DMR(s) and imprinted gene(s) responsible for the development of SRS in patients with upd(7)mat. Furthermore, while advanced maternal age at childbirth has been shown to be a predisposing factor for the development of upd(15)mat because of increased non-disjunction at meiosis 1 [50], such studies remain fragmentary for upd(7)mat, primarily because of the relative paucity of upd(7)mat. Further studies will permit a better characterization of SRS.

Supporting Information

Figure S1 Methylation analysis of the KvDMR1 using COBRA. A. Schematic representation of the KvDMR1. A 326 bp region harboring 24 CpG dinucleotides was studied. The cytosine residues at the CpG dinucleotides are usually methylated after paternal transmission (filled circles) and unmethylated after maternal transmission (open circles); after bisulfite treatment, this region is digested with *Hpy188I* when the cytosine at the 5th CpG dinucleotide (indicated with a green rectangle) is methylated and with *EcoI* when the cytosines at the 22nd CpG dinucleotide (indicated with a pink rectangle) is methylated. *KCNQ1* is a paternally expressed gene, and *KCNQ1* and *CDKN1C* are maternally expressed genes. B. Representative COBRA results. U: unmethylated clone specific bands; M: methylated clone specific bands; and BWS: Beckwith-Wiedemann syndrome patient with upd(11p15)pat. C. Histograms showing the distribution of the MIs (the horizontal axis: the methylation index; and the vertical axis: the patient number). (TIF)

Table S1 Primers utilized in the methylation analysis and microsatellite analysis. (XLS)

Table S2 The results of microsatellite analysis. (XLSX)

Table S3 Methylation indices for multiple differentially methylated regions (DMRs) obtained by COBRA in 38 patients with Silver-Russell syndrome. (XLSX)

Table S4 Clinical findings in two unique patients. (DOC)

Table S5 Correlation analyses in patients with *H19*-DMR hypomethylations. (DOC)

Acknowledgments

We would like to thank the following doctors for providing us with blood samples and clinical data: Toshio Yamazaki, Yukihiro Hasegawa, Daisuke Ariyasu, Michiko Hayashidani, Hiroshi Yoshihashi, Tomoki Kosho, Rika Kosaki, Yasuhiro Naiki, Hideki Fujita, Aya Yamashita, Katsuhiro Maeyama, Sayu Otori, Takashi Koike, Makoto Ono, Sachiko Kitanaka, Jun Mori, Shinichiro Miyagawa, Masamichi Ogawa, Toshiaki Tanaka, Naomi Hizuka, Yoko Fujimoto, Zenro Kizaki, Fumio Katsushima, Takashi Iwamoto, Wakako Yamamoto, Hotaka Kamasaki, Shun Soneda, Takahiro Tajima, Yoriko Watanabe, Kanako Ishii, Noriko Kinoshita, Tohyju Tanaka, Eriko Nishi, Tohru Yorifuji, Yoshio Makita, Yoshinobu Honda, Junko Tsubaki, Yoko Shimabukuro, Yoko Hiraki, Kazushige

Dobashi, Kazumichi Onigata, Hisaaki Kabata, Shigeki Ishii, Kimiko Taniguchi, and Masaoki Sugita. We are also grateful to Drs. Kyoko Tanabe and Kentaro Matsuoka for their technical assistance.

References

- Eggermann T (2010) Russell-Silver syndrome. *Am J Med Genet C Semin Med Genet* 154C: 355–364.
- Binder G, Begemann M, Eggermann T, Kannenberg K (2011) Silver-Russell syndrome. *Best Pract Res Clin Endocrinol Metab* 25: 153–160.
- Wakeling EL, Amcro SA, Alders M, Blick J, Forsythe E, et al. (2010) Epigenotype-phenotype correlations in Silver-Russell syndrome. *J Med Genet* 47: 760–768.
- Yamazawa K, Kagami M, Nagai T, Kondoh T, Onigata K, et al. (2008) Molecular and clinical findings and their correlations in Silver-Russell syndrome: implications for a positive role of IGF2 in growth determination and differential imprinting regulation of the IGF2-H19 domain in bodies and placentas. *J Mol Med (Berl)* 86: 1171–1181.
- Fowden AL, Sibley C, Reik W, Constancia M (2006) Imprinted genes, placental development and fetal growth. *Horm Res* 65 Suppl 3: 50–58.
- Yamazawa K, Kagami M, Ogawa M, Horikawa R, Ogata T (2008) Placental hypoplasia in maternal uniparental disomy for chromosome 7. *Am J Med Genet A* 146A: 514–516.
- Azzi S, Rossignol S, Steunou V, Sas T, Thibaud N, et al. (2009) Multilocus methylation analysis in a large cohort of 11p15-related foetal growth disorders (Russell Silver and Beckwith Wiedemann syndromes) reveals simultaneous loss of methylation at paternal and maternal imprinted loci. *Hum Mol Genet* 18: 4724–4733.
- Turner CL, Mackay DM, Callaway JL, Docherty LE, Poole RL, et al. (2010) Methylation analysis of 79 patients with growth restriction reveals novel patterns of methylation change at imprinted loci. *Eur J Hum Genet* 18: 648–655.
- Hiura H, Okae H, Miyauchi N, Sato F, Sato A, et al. (2012) Characterization of DNA methylation errors in patients with imprinting disorders conceived by assisted reproduction technologies. *Hum Reprod* 27: 2541–2548.
- Abu-Amero S, Monk D, Frost J, Preece M, Stanier P, et al. (2008) The genetic aetiology of Silver-Russell syndrome. *J Med Genet* 45: 193–199.
- Hitchins MP, Stanier P, Preece MA, Moore GE (2001) Silver-Russell syndrome: a dissection of the genetic aetiology and candidate chromosomal regions. *J Med Genet* 38: 810–819.
- Spengler S, Schönherr N, Binder G, Wollmann HA, Fricke-Otto S, et al. (2010) Submicroscopic chromosomal imbalances in idiopathic Silver-Russell syndrome (SRS): the SRS phenotype overlaps with the 12q14 microdeletion syndrome. *J Med Genet* 47: 356–360.
- Fuke-Sato T, Yamazawa K, Nakabayashi K, Matsubara K, Matsuoka K, et al. (2012) Mosaic upd(7)mat in a patient with Silver-Russell syndrome. *Am J Med Genet A* 158A: 465–468.
- Netchine I, Rossignol S, Dufourg MN, Azzi S, Rousseau A, et al. (2007) 11p15 imprinting center region 1 loss of methylation is a common and specific cause of typical Russell-Silver syndrome: clinical scoring system and epigenetic-phenotypic correlations. *J Clin Endocrinol Metab* 92: 3148–3154.
- Kagami M, Yamazawa K, Matsubara K, Matsuo N, Ogata T (2008) Placentomegaly in paternal uniparental disomy for human chromosome 14. *Placenta* 29: 760–761.
- Bell AC, Felsenfeld G (2000) Methylation of a CTCF-dependent boundary controls imprinted expression of the *Igf2* gene. *Nature* 405: 482–485.
- Hark AT, Schoenherr CJ, Katz DJ, Ingram RS, LeVorse JM, et al. (2000) CTCF mediates methylation-sensitive enhancer-blocking activity at the H19/Igf2 locus. *Nature* 405: 486–489.
- Takai D, Gonzales FA, Tsai YC, Thayer MJ, Jones PA (2001) Large scale mapping of methylcytosines in CTCF-binding sites in the human H19 promoter and aberrant hypomethylation in human bladder cancer. *Hum Mol Genet* 10: 2619–2626.
- Fisher AM, Thomas NS, Cockwell A, Stecko O, Kerr B, et al. (2002) Duplications of chromosome 11p15 of maternal origin result in a phenotype that includes growth retardation. *Hum Genet* 111: 290–296.
- Brena RM, Auer H, Kornacker K, Hackanson B, Raval A, et al. (2006) Accurate quantification of DNA methylation using combined bisulfite restriction analysis coupled with the Agilent 2100 Bioanalyzer platform. *Nucleic Acids Res* 34: e17.
- Yamazawa K, Nakabayashi K, Kagami M, Sato T, Saitoh S, et al. (2010) Parthenogenetic chimaerism/mosaicism with a Silver-Russell syndrome-like phenotype. *J Med Genet* 47: 782–785.
- Shaffer LG, Agan N, Goldberg JD, Ledbetter DH, Longshore JW, et al. (2001) American College of Medical Genetics statement on diagnostic testing for uniparental disomy. *Genet Med* 3: 206–211.
- Hannula K, Lipsanen-Nyman M, Kontiokari T, Kere J (2001) A narrow segment of maternal uniparental disomy of chromosome 7q31-qter in Silver-Russell syndrome delimits a candidate gene region. *Am J Hum Genet* 68: 247–253.
- Eggermann T, Schönherr N, Jager S, Spaich C, Ranke MB, et al. (2008) Segmental maternal UPD(7q) in Silver-Russell syndrome. *Clin Genet* 74: 486–489.
- Begemann M, Spengler S, Kordass U, Schroder C, Eggermann T (2012) Segmental maternal uniparental disomy 7q associated with DLK1/GTL2 (14q32) hypomethylation. *Am J Med Genet A* 158A: 423–428.
- Yamazawa K, Ogata T, Ferguson-Smith AC (2010) Uniparental Disomy and Human Disease: An Overview. *Am J Med Genet C* 154C: 329–334.
- Gicquel C, Rossignol S, Cabrol S, Houang M, Steunou V, et al. (2005) Epimutation of the telomeric imprinting center region on chromosome 11p15 in Silver-Russell syndrome. *Nat Genet* 37: 1003–1007.
- Blick J, Terhal P, van den Bogaard MJ, Maas S, Hamel B, et al. (2006) Hypomethylation of the H19 gene causes not only Silver-Russell syndrome (SRS) but also isolated asymmetry or an SRS-like phenotype. *Am J Hum Genet* 78: 604–614.
- Ulaner GA, Yang Y, Hu JF, Li T, Vu TH, et al. (2003) CTCF binding at the insulin-like growth factor-II (IGF2)/H19 imprinting control region is insufficient to regulate IGF2/H19 expression in human tissues. *Endocrinology* 144: 4420–4426.
- Pham NV, Nguyen MT, Hu JF, Vu TH, Hoffman AR (1998) Dissociation of IGF2 and H19 imprinting in human brain. *Brain Res* 810: 1–8.
- Albrecht S, Waha A, Koch A, Kraus JA, Goodyer CG, et al. (1996) Variable imprinting of H19 and IGF2 in fetal cerebellum and medulloblastoma. *J Neuropathol Exp Neurol* 55: 1270–1276.
- Kotzot D (2008) Maternal uniparental disomy 7 and Silver-Russell syndrome - clinical update and comparison with other subgroups. *Eur J Med Genet* 51: 444–451.
- Kotzot D, Balmer D, Baumer A, Chrzanoska K, Hamel BC, et al. (2000) Maternal uniparental disomy 7-review and further delineation of the phenotype. *Eur J Pediatr* 159: 247–256.
- Robinson WP (2000) Mechanisms leading to uniparental disomy and their clinical consequences. *Bioessays* 22: 452–459.
- Hao Y, Crenshaw T, Moulton T, Newcomb E, Tycko B (1993) Tumour-suppressor activity of H19 RNA. *Nature* 365: 764–767.
- Juan V, Crain C, Wilson C (2000) Evidence for evolutionarily conserved secondary structure in the H19 tumor suppressor RNA. *Nucleic Acids Res* 28: 1221–1227.
- Arima T, Kamikihara T, Hayashida T, Kato K, Inoue T, et al. (2005) ZAC, LIT1 (KCNQ1OT1) and p57KIP2 (CDKN1C) are in an imprinted gene network that may play a role in Beckwith-Wiedemann syndrome. *Nucleic Acids Res* 33: 2650–2660.
- Varrault A, Gueydan C, Delalbre A, Bellmann A, Houssami S, et al. (2006) Zac1 regulates an imprinted gene network critically involved in the control of embryonic growth. *Dev Cell* 11: 711–722.
- Quenneville S, Verde G, Corsinotti A, Kapopoulou A, Jakobsson J, et al. (2011) In embryonic stem cells, ZFP57/KAP1 recognize a methylated hexanucleotide to affect chromatin and DNA methylation of imprinting control regions. *Mol Cell* 44: 361–372.
- Yamazawa K, Nakabayashi K, Matsuoka K, Matsubara K, Hata K, et al. (2011) Androgenetic/biparental mosaicism in a girl with Beckwith-Wiedemann syndrome-like and upd(14)pat-like phenotypes. *J Hum Genet* 56: 91–93.
- Huang J, Lin Y, Li L, Qing D, Teng XM, et al. (2009) ARHI, as a novel suppressor of cell growth and downregulated in human hepatocellular carcinoma, could contribute to hepatocarcinogenesis. *Mol Carcinog* 48: 130–140.
- Feng W, Marquez RT, Lu Z, Liu J, Lu KH, et al. (2008) Imprinted tumor suppressor genes ARHI and PEG3 are the most frequently down-regulated in human ovarian cancers by loss of heterozygosity and promoter methylation. *Cancer* 112: 1489–1502.
- Tang HL, Hu YQ, Qin XP, Jazag A, Yang H, et al. (2012) Aplasia ras homolog member I is downregulated in gastric cancer and silencing its expression promotes cell growth in vitro. *J Gastroenterol Hepatol* 27: 1395–1404.
- Blyth M, Huang S, Maloney V, Crolla JA, Karen Temple I (2008) A 2.3 Mb deletion of 17q24.2-q24.3 associated with 'Carney Complex plus'. *Eur J Med Genet* 51: 672–678.
- Midro AT, Debek K, Sawicka A, Marcinkiewicz D, Rogowska M (1993) Second observation of Silver-Russell syndrome in a carrier of a reciprocal translocation with one breakpoint at site 17q25. *Clin Genet* 44: 53–55.
- Dörr S, Midro AT, Farber C, Giannakudis J, Hansmann I (2001) Construction of a detailed physical and transcript map of the candidate region for Russell-Silver syndrome on chromosome 17q23-q24. *Genomics* 71: 174–181.
- Cuomo CA, Kirch SA, Gyuris J, Brent R, Oettinger MA (1994) Rch1, a protein that specifically interacts with the RAG-1 recombination-activating protein. *Proc Natl Acad Sci U S A* 91: 6156–6160.

Author Contributions

Conceived and designed the experiments: TF KY TO. Performed the experiments: TF KN CT S. Sano K. Matsubara MK KY. Analyzed the data: TF KN KH KY. Contributed reagents/materials/analysis tools: SM TN TH RH YM K. Muroya TK CN S. Sato TO. Wrote the paper: TO.

48. Weis K, Mattaj IW, Lamond AI (1995) Identification of hSRP1 alpha as a functional receptor for nuclear localization sequences. *Science* 268: 1049–1053.
49. Dörr S, Schlicker M, Hansmann I (2001) Genomic structure of karyopherin alpha2 (KPNA2) within a low-copy repeat on chromosome 17q23-q24 and mutation analysis in patients with Russell-Silver syndrome. *Hum Genet* 109: 479–486.
50. Matsubara K, Murakami N, Nagai T, Ogata T (2011) Maternal age effect on the development of Prader-Willi syndrome resulting from upd(15)mat through meiosis I errors. *J Hum Genet* 56: 566–571.

特集

DOHaD アップデート—胎生期から乳児期までの環境と成人期の健康・疾患

6. 糖尿病と DOHaD

やす ひ ちろ ぶ
安日一郎

国立病院機構長崎医療センター産婦人科

要旨

「糖尿病と DOHaD」というテーマは、胎内の低栄養環境(すなわち低出生体重児)が児の将来の糖尿病発症に関連するという従来の Barker 仮説にとどまらず、胎内の過剰栄養環境(すなわち巨大児)と児の将来の糖尿病発症との関連というユニークな視点が加わる。糖尿病合併妊娠(母体の糖尿病)-糖尿病性胎児病-巨大児-児の将来の糖尿病発症という、疾患の世代間連鎖モデルという新しい概念として興味深い。

KeyWords DOHaD, 糖尿病, 巨大児

はじめに

DOHaD (developmental origins of health and disease) は、Barker が提唱した胎児起源説 (fetal origin hypothesis)¹⁾ を発展させた概念である。Barker 仮説は、低出生体重で生まれた児が将来の成人期の心血管イベントのリスク因子であることを明らかにしたことに端を発する²⁾。その後、成人期の肥満、高血圧、高脂血症、そして耐糖能異常のいわゆるメタボリック症候群あるいはインスリン抵抗性症候群と称される病態の発症と低出生体重との関連が明らかとなった³⁾。病態的に心血管イベントの上流に位置するメタボリック症候群は、その別称のとおりインスリン抵抗性が基本病態である⁴⁾。成人期のインスリン抵抗性の病態の発症起源が、胎児期

のみならず新生児期の栄養状態にも起因するという概念へ発展したものが DOHaD である⁵⁾。

本稿では、糖尿病と DOHaD の関連について最新のエビデンスをもとに概説する。糖尿病(2型糖尿病)は、インスリン抵抗性の代表的疾患であり DOHaD の基本疾患の一つである。母体の糖尿病に起因する子宮内環境(高血糖状態)が胎児の将来の糖尿病の発症起源となるという、遺伝要因とは異なる概念としての同一疾病の世代間連鎖という意味で、他のメタボリック症候群とは一線を画して興味深い。さらに、母体の妊娠中の血糖管理によって児の将来の糖尿病発症を予防できるという観点からもユニークである。世界的な肥満と糖尿病のパンデミックとも称される今日、糖尿病とその背景となるインスリン抵抗性の代表的な表現型である肥満に対する予防戦略としても極めて重要なテーマである。