

In 2008, mutations in the *EYS* gene were reported in patients with autosomal recessive retinitis pigmentosa (arRP) [1, 2]. The *EYS* mutations, which have been shown to be predominantly truncating mutations, have been described in patients with different ethnic origins and account for 5–16 % of arRP [3–7]. Thus, disruption of the *EYS* function has been identified as a frequent cause of arRP worldwide. With the exception for arRP, to date there have been no *EYS* mutations reported for any other phenotype.

Here, we report clinical and genetic features of a patient with autosomal recessive cone–rod dystrophy (arCRD) associated with compound heterozygous *EYS* mutations.


### Case report

A 31-year-old male patient (JU#0659) was referred to our hospital with a complaint of loss of visual acuity. He first reported a decreased visual acuity at the age of 29 years. Family history indicated that his parents had no previous reports of any ocular symptoms (Fig. 1). At his initial examination, decimal best-corrected visual acuity (BCVA) was 0.9 [with  $-6.00$  diopter (dpt), cylinder (cyl)  $-1.25$  dpt axis (Ax)  $130^\circ$ ] in his right eye and 0.6 (with  $-4.50$  dpt, cyl  $-0.75$  dpt Ax  $180^\circ$ ) in his left eye. Anterior segment examination showed no remarkable findings. Intraocular pressures were 16 mmHg in the right and 15 mmHg in the left eye. Funduscopy showed retinal degenerations within the vascular arcade in both eyes (Fig. 2a). Neither retinal degeneration nor attenuation of retinal vessels


was observed in the periphery. Fluorescein angiography (VISUCAM NM/FA; Carl Zeiss Meditec AG, Dublin, CA, USA) showed a hyperfluorescence pattern due to a window defect within the vascular arcades of both eyes (Fig. 2b). At the age of 32 years, visual field testing using Goldmann kinetic perimetry (GP; Haag-Streit, Bern, Switzerland) showed bilateral central scotomas of the I-3e and I-4e isopters with relative sparing of the center, but preserved peripheral visual fields of the V-4e and I-4e isopters in both eyes (Fig. 3). Full-field electroretinography (ERG) was performed according to the protocols of the International Society for Clinical Electrophysiology of Vision. The procedure and conditions for ERG recording have been reported previously [8]. The ERG showed the rod, standard combined, cone, and 30-Hz flicker responses were bilaterally reduced to about one-third of those in a control, but the peak implicit time of each response was not delayed (Fig. 4).

At the age of 36 years, his BCVA decreased to 0.2 in both eyes. Funduscopy images using the Optos 200Tx imaging system (Optos PLC, Dunfermline, United Kingdom) showed retinal degenerations within the vascular arcades in both eyes, but relatively preserved mid-peripheral to peripheral retinal findings with no apparent attenuation of the retinal vessels (Fig. 5a). Fundus autofluorescence imaging (FAI) (Optos PLC) showed decreased autofluorescence within the vascular arcades but increased autofluorescence of the foveal area, and increased autofluorescence outside the vascular arcades in both eyes (Fig. 5b). Optical coherence tomography (OCT) (Cirrus HD-OCT; Carl Zeiss Meditec AG) showed retinal thinning with a visible foveal external limiting membrane line (Fig. 5c), and entire macular thinning in both eyes (Fig. 5d).

To identify disease-causing gene mutations, we performed whole-exome sequencing analysis as per a previously described method [9]. The obtained sequence data in the patient were compared with reference human genome sequences. Initially, we focused on only variants that could change the amino acid sequence. Subsequently, we filtered the remaining variants based on the criteria that a frequency of mutation was less than 1 % in the 1000-genome database (<http://www.1000genomes.org>). Finally, we screened variants residing within 207 retinal disease-associated genes published in the November 15, 2013




**Fig. 1** Pedigree of a Japanese family. Unaffected family members (males, open squares; females, open circles) and an affected proband (male, solid square) are shown



**Fig. 2** Fundus photographs and fluorescein angiography images (FA) of the patient at the age of 31 years. **a** Fundus photographs show retinal degenerations within the vascular


arcades in both eyes. **b** FA shows a hyperfluorescence pattern due to a window defect within the vascular arcades of both eyes



**Fig. 3** Visual field testing using Goldmann kinetic perimetry (GP) at the age of 32 years. GP shows bilateral central scotomas of the I-3e and I-4e isopters with relative sparing of the center,

but with preserved peripheral visual fields of the V-4e and I-4e isopters in both eyes

**Fig. 4** Full-field electroretinography (ERG) at the age of 32 years. ERG shows that the rod, standard combined, cone, and 30-Hz flicker responses are bilaterally reduced to about one-third of those in a control, but the peak implicit time of each response is not delayed



RetNet database (<https://sph.uth.edu/retnet/>). Based on the obtained data, known *EYS* mutations were identified in a compound heterozygous state as disease-causing mutations. In the other 206 genes, there were no mutations found in compound heterozygous or homozygous states. The identified *EYS* gene mutations were c.C8805A and c.4957dupA, which result in the truncating mutations p.Y2935X and p.S1653KfsX2, respectively. The findings were confirmed by Sanger sequencing. The patient's unaffected parents were heterozygous for each mutation. The compound heterozygous mutations (p.Y2935X and p.S1653KfsX2) have been previously reported as a cause of arRP [7]. The accession number of the *EYS* mRNA reference sequence we used was NM\_001142800.1 from the National Center for Biotechnology Information.

## Discussion


In this report, whole-exome sequencing analysis identified *EYS* mutations in a Japanese patient with arCRD. *EYS* mutations have previously only been reported in patients with arRP [1, 2]. This is the first report that describes arCRD associated with compound heterozygous mutations in the *EYS* gene.

Previous studies have revealed that the *EYS* protein is expressed specifically in the human retina and is

localized in the outer segment of the photoreceptor layers of the porcine retina [1, 2]. Although the function and structure of human *EYS* protein remain unclear, it has been suggested to be essential for photoreceptor morphogenesis [2]. In fact, *EYS* mutations give rise to RP phenotypes with thinning of the outer retinal layers [5, 7], which results from the degeneration of both the rod and cone photoreceptors. Clinical features of patients with *EYS* mutations include a typical form of RP that is characterized by a progressive constricted visual field, bone pigmentations and attenuation of the retinal vessels [3–5]. The pattern of the ERG in RP patients shows non-recordable or markedly decreased responses [3–5]. On the other hand, our patient exhibited retinal degenerations that were predominantly seen within the vascular arcades (Fig. 5a, b), central scotomas and preserved peripheral visual fields (Fig. 3), and decreased responses in the both rod and cone ERG (Fig. 4) in both eyes. Generally, CRD exhibits several features such as decreased central vision, a predominant degeneration of the macular region as compared with the mid-peripheral region, and decreased amplitudes in the cone ERG that are equal to or worse than the decreased rod ERG amplitudes [10, 11]. These characteristics of CRD were clearly consistent with the phenotype of our patient. Taken together, these findings indicated that the patient diagnosis was arCRD and not arRP.

**Fig. 5** Fundus photographs, fundus autofluorescence images (FAI), and optic coherence tomography images (OCT) at the age of 36 years.

**a** Fundus photographs show retinal degenerations within the vascular arcades in both eyes, but relatively preserved mid-peripheral to peripheral retinal findings with no apparent attenuation of the retinal vessels. **b** FAI shows decreased autofluorescence within the vascular arcades but increased autofluorescence of the foveal area, and increased autofluorescence outside the vascular arcades in both eyes. **c** OCT (HD 5-line raster scan) shows retinal thinning with a visible foveal external limiting membrane line in both eyes. **d** OCT (Macular cube scan) shows entire macular thinning in both eyes



Whole-exome sequencing analysis disclosed that our patient had the *EYS* mutations, which demonstrates that the *EYS* mutations can be responsible for both the arCRD and the arRP phenotypes. Interestingly, mutations in the *ABCA4* [12–14], *CERKL* [15–18], and *C8orf37* [19, 20] genes have also been reported to be disease-causing mutations of both the arCRD and arRP phenotypes. With regard to the *ABCA4* gene mutations, the degree of functional damage caused by the various *ABCA4* mutation types can underlie the different degeneration patterns, for example, Stargardt disease (a type of macular dystrophy), arCRD or arRP [12–14]. The majority of patients with *CERKL* mutations exhibit arCRD [17, 18] and less frequently arRP [15, 16]. This is consistent with the fact that the *CERKL* protein is predominantly expressed in the cone photoreceptors [21]. In addition, different *C8orf37* mutations can cause either the arCRD or arRP phenotypes, which is consistent with the fact that the *C8orf37* protein is expressed in both the rod and cone photoreceptors [19, 20]. However, this does not explain the pattern of the photoreceptor degeneration. On the other hand, the compound heterozygous *EYS* mutations (p.Y2935X and p.S1653KfsX2) that were found in our patient have also been reported in an arRP patient [7]. Although it is not understood why the same compound heterozygous mutations would underlie either the arCRD or arRP phenotypes, this finding suggests there is the presence of different modifier alleles between the arCRD or arRP patients with the compound heterozygous *EYS* mutations. Even so, our whole-exome sequencing analysis did not demonstrate any compound heterozygous or homozygous mutations in other 206 retinal disease-associated genes published in the RetNet database.

In conclusion, we demonstrated that *EYS* mutations are the cause of not only arRP but also arCRD. Further investigations will need to be undertaken in order to clarify the prevalence of *EYS* mutations among arCRD patients, and to determine the genotype–phenotype correlations between the arCRD and *EYS* mutations.

**Acknowledgments** This study was supported by grants to T.I. from the Ministry of Health, Labor and Welfare of Japan (13803661), to M.A. and T.H. from the Ministry of Education, Culture, Sports, Science and Technology of Japan (Grant-in-Aid for Scientific Research C, 25462744 and 25462738), and to T.H. from the Vehicle Racing Commemorative Foundation.

**Conflict of interest** The authors declare there are no conflicts of interest for this study.

## References

1. Abd El-Aziz MM, Barragan I, O'Driscoll CA, Goodstadt L, Prigmore E, Borrego S, Mena M, Pieras JI, El-Ashry MF, Safieh LA, Shah A, Cheetham ME, Carter NP, Chakarova C, Ponting CP, Bhattacharya SS, Antinolo G (2008) *EYS*, encoding an ortholog of *Drosophila* spacemaker, is mutated in autosomal recessive retinitis pigmentosa. *Nat Genet* 40:1285–1287
2. Collin RW, Littink KW, Klevering BJ, van den Born LI, Koenekoop RK, Zonneveld MN, Blokland EA, Strom TM, Hoyng CB, den Hollander AI, Cremers FP (2008) Identification of a 2 Mb human ortholog of *Drosophila* eyes shut/spacemaker that is mutated in patients with retinitis pigmentosa. *Am J Hum Genet* 83:594–603
3. Abd El-Aziz MM, O'Driscoll CA, Kaye RS, Barragan I, El-Ashry MF, Borrego S, Antinolo G, Pang CP, Webster AR, Bhattacharya SS (2010) Identification of novel mutations in the ortholog of *Drosophila* eyes shut gene (*EYS*) causing autosomal recessive retinitis pigmentosa. *Invest Ophthalmol Vis Sci* 51:4266–4272
4. Audo I, Sahel JA, Mohand-Said S, Lancelot ME, Antonio A, Moskova-Doumanova V, Nandrot EF, Doumanov J, Barragan I, Antinolo G, Bhattacharya SS, Zeitz C (2010) *EYS* is a major gene for rod-cone dystrophies in France. *Hum Mutat* 31:E1406–E1435
5. Bandah-Rozenfeld D, Littink KW, Ben-Yosef T, Strom TM, Chowers I, Collin RW, den Hollander AI, van den Born LI, Zonneveld MN, Merin S, Banin E, Cremers FP, Sharon D (2010) Novel null mutations in the *EYS* gene are a frequent cause of autosomal recessive retinitis pigmentosa in the Israeli population. *Invest Ophthalmol Vis Sci* 51:4387–4394
6. Hosono K, Ishigami C, Takahashi M, Park DH, Hirami Y, Nakanishi H, Ueno S, Yokoi T, Hikoya A, Fujita T, Zhao Y, Nishina S, Shin JP, Kim IT, Yamamoto S, Azuma N, Terasaki H, Sato M, Kondo M, Minoshima S, Hotta Y (2012) Two novel mutations in the *EYS* gene are possible major causes of autosomal recessive retinitis pigmentosa in the Japanese population. *PLoS ONE* 7:e31036
7. Iwanami M, Oshikawa M, Nishida T, Nakadomari S, Kato S (2012) High prevalence of mutations in the *EYS* gene in Japanese patients with autosomal recessive retinitis pigmentosa. *Invest Ophthalmol Vis Sci* 53:1033–1040
8. Hayashi T, Gekka T, Kozaki K, Ohkuma Y, Tanaka I, Yamada H, Tsuneoka H (2012) Autosomal dominant occult macular dystrophy with an *RP1L1* mutation (R45W). *Optom Vis Sci* 89:684–691
9. Katagiri S, Yoshitake K, Akahori M, Hayashi T, Furuno M, Nishino J, Ikeo K, Tsuneoka H, Iwata T (2013) Whole-exome sequencing identifies a novel *ALMS1* mutation (p.Q2051X) in two Japanese brothers with Alström syndrome. *Mol Vis* 19:2393–2406
10. Yagasaki K, Jacobson SG (1989) Cone-rod dystrophy. Phenotypic diversity by retinal function testing. *Arch Ophthalmol* 107:701–708

11. Szlyk JP, Fishman GA, Alexander KR, Peachey NS, Derlacki DJ (1993) Clinical subtypes of cone-rod dystrophy. *Arch Ophthalmol* 111:781–788
12. Allikmets R, Singh N, Sun H, Shroyer NF, Hutchinson A, Chidambaram A, Gerrard B, Baird L, Stauffer D, Peiffer A, Rattner A, Smallwood P, Li Y, Anderson KL, Lewis RA, Nathans J, Leppert M, Dean M, Lupski JR (1997) A photoreceptor cell-specific ATP-binding transporter gene (*ABCR*) is mutated in recessive Stargardt macular dystrophy. *Nat Genet* 15:236–246
13. Klevering BJ, Yzer S, Rohrschneider K, Zonneveld M, Allikmets R, van den Born LI, Maugeri A, Hoyng CB, Cremers FP (2004) Microarray-based mutation analysis of the *ABCA4* (*ABCR*) gene in autosomal recessive cone-rod dystrophy and retinitis pigmentosa. *Eur J Hum Genet* 12:1024–1032
14. Klevering BJ, Deutman AF, Maugeri A, Cremers FP, Hoyng CB (2005) The spectrum of retinal phenotypes caused by mutations in the *ABCA4* gene. *Graefes Arch Clin Exp Ophthalmol* 243:90–100
15. Bayes M, Goldaracena B, Martinez-Mir A, Iragui-Madoz MI, Solans T, Chivelet P, Bussaglia E, Ramos-Arroyo MA, Baiget M, Vilageliu L, Balcells S, Gonzalez-Duarte R, Grinberg D (1998) A new autosomal recessive retinitis pigmentosa locus maps on chromosome 2q31–q33. *J Med Genet* 35:141–145
16. Tuson M, Marfany G, Gonzalez-Duarte R (2004) Mutation of *CERKL*, a novel human ceramide kinase gene, causes autosomal recessive retinitis pigmentosa (RP26). *Am J Hum Genet* 74:128–138
17. Auslender N, Sharon D, Abbasi AH, Garzozzi HJ, Banin E, Ben-Yosef T (2007) A common founder mutation of *CERKL* underlies autosomal recessive retinal degeneration with early macular involvement among Yemenite Jews. *Invest Ophthalmol Vis Sci* 48:5431–5438
18. Ali M, Ramprasad VL, Soumitra N, Mohamed MD, Jafri H, Rashid Y, Danciger M, McKibbin M, Kumaramanickavel G, Inglehearn CF (2008) A missense mutation in the nuclear localization signal sequence of *CERKL* (p.R106S) causes autosomal recessive retinal degeneration. *Mol Vis* 14:1960–1964
19. Estrada-Cuzcano A, Neveling K, Kohl S, Banin E, Rotenstreich Y, Sharon D, Falik-Zaccari TC, Hipp S, Roepman R, Wissinger B, Letteboer SJ, Mans DA, Blokland EA, Kwint MP, Gijzen SJ, van Huet RA, Collin RW, Scheffer H, Veltman JA, Zrenner E, den Hollander AI, Klevering BJ, Cremers FP (2012) Mutations in *C8orf37*, encoding a ciliary protein, are associated with autosomal-recessive retinal dystrophies with early macular involvement. *Am J Hum Genet* 90:102–109
20. van Huet RA, Estrada-Cuzcano A, Banin E, Rotenstreich Y, Hipp S, Kohl S, Hoyng CB, den Hollander AI, Collin RW, Klevering BJ (2013) Clinical characteristics of rod and cone photoreceptor dystrophies in patients with mutations in the *C8orf37* gene. *Invest Ophthalmol Vis Sci* 54:4683–4690
21. Vekslin S, Ben-Yosef T (2010) Spatiotemporal expression pattern of ceramide kinase-like in the mouse retina. *Mol Vis* 16:2539–2549

# Retinal angiomatous proliferation associated with risk alleles of *ARMS2/HTRA1* gene polymorphisms in Japanese patients

This article was published in the following Dove Press journal:

Clinical Ophthalmology

27 December 2013

[Number of times this article has been viewed](#)

Yasuhiro Ohkuma<sup>1</sup>  
Takaaki Hayashi<sup>1</sup>  
Tsutomu Sakai<sup>1</sup>  
Akira Watanabe<sup>1</sup>  
Hisashi Yamada<sup>2</sup>  
Masakazu Akahori<sup>3</sup>  
Takeshi Itabashi<sup>3</sup>  
Takeshi Iwata<sup>3</sup>  
Toru Noda<sup>4</sup>  
Hiroshi Tsuneoka<sup>1</sup>

<sup>1</sup>Department of Ophthalmology,

<sup>2</sup>Department of Molecular Genetics, Institute of DNA Medicine, The Jikei University School of Medicine,

<sup>3</sup>Division of Molecular and Cellular Biology, National Institute of Sensory Organs, <sup>4</sup>Division of Ophthalmology, National Hospital Organization Tokyo Medical Center, Tokyo, Japan

**Background:** The purpose of this study was to investigate the association between *ARMS2/HTRA1*, *CFH*, and *C3* gene polymorphisms and retinal angiomatous proliferation (RAP), an infrequent and severe form of exudative age-related macular degeneration, which is characterized by intraretinal neovascularization.

**Methods:** Diagnosis of RAP was based on fundus photographs, images of fluorescein and indocyanine green angiographies, and optical coherence tomography findings. Six single nucleotide polymorphisms (SNPs), A69S (rs10490924) in *ARMS2*, rs11200638 in *HTRA1*, I62V (rs800292) in *CFH*, Y402H (rs1061170) in *CFH*, R80G (rs2230199) in *C3*, and rs2241394 in *C3*, were genotyped in eight Japanese patients with RAP.

**Results:** The two SNPs in the *ARMS2/HTRA1* were in complete linkage disequilibrium. The frequency of the risk T allele in *ARMS2* (the risk A allele in *HTRA1*) was 93.8% in the RAP patients. The frequency of homozygosity for the risk genotype TT of *ARMS2* (the risk genotype AA of *HTRA1*) was 87.5%. The frequency of the non-risk allele (A) of I62V was 100%. The frequencies of risk alleles of Y402H, R80G, and rs2241394 were 12.5%, 0%, and 18.8%, respectively.

**Conclusion:** Our results suggest that the risk alleles of the *ARMS2/HTRA1* SNPs may be associated with development of RAP and play a major role in the pathogenesis of intraretinal angiogenesis.

**Keywords:** age-related macular degeneration, retinal angiomatous proliferation, single nucleotide polymorphisms, *ARMS2/HTRA1* genes, components of the complement system

## Introduction

Age-related macular degeneration (AMD) is the most common cause of legal blindness in the elderly, affecting more than 50 million people worldwide.<sup>1</sup> In Japan, the prevalence of AMD has risen from 0.87% in 1988 to 1.4% in 2007.<sup>2,3</sup> Maruko et al have classified exudative AMD patients into three subtypes, namely typical wet-type AMD, polypoidal choroidal vasculopathy (PCV), and retinal angiomatous proliferation (RAP).<sup>4</sup>

AMD is a multifactorial disease with genetic, behavioral, and environmental factors.<sup>5</sup> Recently, genetic association studies have revealed that single nucleotide polymorphisms (SNPs) in *CFH* (1q32), *ARMS2/HTRA1* (10q26), and *C3* (19p13) have been identified as major contributors to the pathogenesis of AMD.<sup>6-17</sup> Among various SNPs of those genes, the Y402H (rs1061170) and I62V (rs800292) variants in the *CFH* gene and the A69S (rs10490924) variant in the *ARMS2* gene have been investigated in detail.<sup>6, 15, 18-27</sup> The differences in genotypes associated with AMD

Correspondence: Takaaki Hayashi  
Department of Ophthalmology,  
The Jikei University School of  
Medicine, 3-25-8 Nishi-shimbashi,  
Minato-ku, Tokyo 105-8461, Japan  
Tel +81 3 3433 1111 ext 3581  
Fax +81 3 3433 1936  
Email taka@jikei.ac.jp

have been investigated among various ethnic groups and by subtypes of exudative AMD,<sup>16,18–27</sup> showing that the I62V and A69S variants are associated with AMD in both Caucasian and Asian subjects.<sup>18–27</sup> The Y402H and R80G (in the *C3* gene) variants have been associated with AMD in Caucasians<sup>6–15</sup> but not in Asians.<sup>18,19,21–25,28</sup> The C allele of the Y402H variant and the G allele of the R80G variant are infrequent in Asians.

The term RAP was first coined by Yannuzzi et al in 2001.<sup>29</sup> They suggested the retinal origin of this neovascularization, which proceeds posteriorly and finally forms a retinal-choroidal anastomosis. RAP is sometimes called type 3 neovascularization to distinguish it from type 1 neovascularization (choroidal neovascularization under the retinal pigment epithelium) and type 2 neovascularization (choroidal neovascularization that penetrates the retinal pigment epithelium).<sup>30,31</sup> RAP accounts for 4.5% of all exudative AMD in Japanese patients<sup>4</sup> and 15% of exudative AMD in Caucasian patients.<sup>32</sup> RAP is characterized by bilateral, multiple soft drusen, intraretinal hemorrhages, and intraretinal edema. The natural history of RAP is characterized by a rapid loss of vision.<sup>33</sup> RAP resists various treatments and recurs persistently.<sup>34–40</sup>

The phenotypic diversity of AMD is thought to be related to differences in genetic backgrounds.<sup>20,24–27</sup> Various reports have examined genetic backgrounds in PCV. Lee et al reported that the I62V and A69S variants, but not the Y402H variant, were related to PCV in Chinese patients.<sup>23</sup> Hayashi et al reported that all three of these SNPs (I62V, Y402H, and A69S) were related to PCV in Japanese patients.<sup>26</sup> Goto et al reported that rs2241394 in the *C3* gene was associated with PCV.<sup>25</sup>

Wegscheider et al reported that the Y402H polymorphism was associated with RAP in Caucasians.<sup>20</sup> However, the genetic association with RAP has not been evaluated sufficiently because of the rarity of RAP in Japan. There are only a few reports about associations between A69S and RAP in the Japanese population.<sup>26,27</sup> The purpose of the current study was to investigate the involvement of genetic factors in not only the *ARMS2/HTRA1* but also the *CFH* and *C3* genes in Japanese patients with RAP.

## Materials and methods

The study was approved by the institutional review board of The Jikei Medical University, and all procedures were conducted in accordance with the principles of the Declaration of Helsinki. Eight unrelated Japanese patients with RAP were recruited from the Department of Ophthalmology at

The Jikei Medical University and the National Hospital Organization Tokyo Medical Center. Informed consent was obtained from all subjects.

All patients with RAP underwent a full ophthalmic examination, including slit-lamp biomicroscopy, funduscopy, optical coherence tomography, and fluorescein and indocyanine green fundus angiographies. The diagnosis of RAP was based on the criteria of Yannuzzi et al<sup>29</sup> and was classified as a defined anastomosis connecting the retinal circulation to a vascular complex within the retina, usually with surrounding intraretinal blood and intraretinal or cystoid macular edema.

Genomic DNA was extracted from the peripheral blood of each individual. A total of six SNPs consisting of A69S (rs10490924) in *ARMS2*, rs11200638 in *HTRA1*, Y402H (rs1061170) in *CFH*, I62V (rs800292) in *CFH*, R80G (rs2230199) in *C3*, and rs2241394 in *C3* were genotyped. Polymerase chain reaction amplification was performed using LA Taq polymerase (Takara Bio Inc, Ohtsu, Japan) and primers for *ARMS2* (forward primer: 5'-GCCTATACCCAGGACCGATG-3', reverse primer: 5'-CATGTTCTCAGCATCTCCAAGTC-3'), *HTRA1* (forward primer: 5'-TCTCTGCGAATACGGACACG-3', reverse primer: 5'-ACT GTG TCCATT CAG CTC CTA A-3'), *CFH* Y402H (forward primer: 5'-CAGAAATAGGGCCAAGAAAAGAGT-3', reverse primer: 5'-ATGTAAGTGTGGTCTGCGC-3'), *CFH* I62V (forward primer: 5'-GATTGCAATGAACTTCTCCAAG-3', reverse primer: 5'-GGATTAAGAGCAACCATTCTCC-3'), *C3* R80G (forward primer: 5'-CCTCGCACCTCCTTACA-3', reverse primer: 5'-TCTGGCTGGCACCTCAAT-3'), and *C3* rs2241394 (forward primer: 5'-GGCTGGGTGACTGTACCTCTTC-3', reverse primer: 5'-CATGTTCTCAGCATCTCCAAGTC-3') to amplify these regions. Polymerase chain reaction products were used as the templates for direct DNA sequencing kits (Applied Biosystems, Foster City, CA, USA) on an automated sequencer (3730xl DNA analyzer; Applied Biosystems).

## Results

### Genetic analysis

Five men and three women were analyzed in the study. The mean patient age was 82.6±4.6 years (range 76–91 years). Both eyes were affected in four patients (50.0%). All SNPs were successfully genotyped in all patients (Table 1). The two SNPs in the *ARMS2/HTRA1* were in complete linkage disequilibrium. The frequency of the risk T allele in



**Table 1** Polymorphisms in *ARMS2/HTRA1/CFH/C3* genes: genotypes in Japanese patients with retinal angiomatous proliferation

Patient number	Age	Sex	Affected eye	ARMS2	HTRA1	CFH	C3		
				rs10490924 (A69S)	rs1120638	rs800292 (I62V)	rs1061170 (Y402H)	rs2230199 (R80G)	rs2241394
1	83	F	Bilateral	TT	AA	AA	TT	CC	CG
2	79	M	Unilateral	TT	AA	AA	CT	CC	CC
3	84	M	Unilateral	TT	AA	AA	TT	CC	CC
4	87	F	Unilateral	TT	AA	AA	CT	CC	CG
5	82	M	Bilateral	TT	AA	AA	TT	CC	CC
6	78	M	Unilateral	TT	AA	AA	TT	CC	CC
7	76	F	Bilateral	TT	AA	AA	TT	CC	CG
8	91	M	Bilateral	TG	AG	AA	TT	CC	CC

**Note:** Risk alleles are shown in bold.

**Abbreviations:** M, male; F, female.

the *ARMS2* gene (the risk A allele in the *HTRA1* gene) was 93.8%. The frequency of homozygosity for the risk genotype (TT) of the *ARMS2* gene was 87.5%. The frequency of the non-risk allele (A) of I62V was 100%. The frequencies of risk alleles of Y402H in the *CFH* gene, rs2230199 (R80G) and rs2241394 in the *C3* gene were 12.5%, 0%, and 18.8%, respectively.


### Representative case (patient 1)

An 83-year-old woman with homozygosity for the risk genotype (TT) of the *ARMS2* gene presented with bilateral RAP (Figure 1). She had undergone cataract surgery in both eyes prior to diagnosis of RAP. Her decimal best-corrected visual acuity was 0.3 in the right eye and 0.07 in the left eye. Both eyes were treated by standard-fluence photodynamic therapy with verteporfin (Visudyne<sup>®</sup>, Novartis Pharma AG, Basel, Switzerland) in combination with 1.25 mg (0.05 mL) of intravitreal bevacizumab (Avastin<sup>®</sup>, Genentech, San Francisco, CA, USA), and her vision improved to 0.5 in the right eye

and 0.15 in the left eye, with a rapid resolution of intraretinal edema. There was no recurrence of intraretinal edema or hemorrhages, and her vision remained stable for 2 years following the combination therapy.

### Discussion

In this study we genotyped six SNPs in RAP patients that were highly representative of the common genetic variations of exudative AMD. Our results raise the possibility of an association between *ARMS2* (A69S)/*HTRA1* (rs1120638) variants and RAP, but a weak association for the other SNPs. Hayashi et al recently demonstrated that the A69S, Y402H, and I62V variants are associated with RAP and that the A69S variant has the strongest association for RAP among the three exudative AMD subtypes.<sup>26</sup> Tanaka et al also reported that A69S might serve as strong genetic markers of RAP.<sup>27</sup> Our findings are consistent with their findings<sup>26,27</sup> regarding the A69S variant, but we had negative results for the I62V (*CFH*) variant (Table 2).



**Figure 1** Color fundus photographs (A and D), indocyanine green fundus angiographies (B and E), and optical coherence tomography images (C and F) from the right eye (A–C) and the left eye (D–F) of an 83-year-old woman (patient 1).

**Notes:** We diagnosed her right eye with stage II retinal angiomatous proliferation and her left eye with stage III retinal angiomatous proliferation. (A and D) Fundus image shows intraretinal hemorrhages with a large number of soft drusen and pigment epithelial detachment. (B) Indocyanine green fundus angiographies shows some hotspots. One of them connects retinal vessels (arrow), corresponding to the intraretinal neovascularization. (C) A vertical optical coherence tomography image shows a pigment epithelial detachment, cystoid macular edema, and retinal angiomatous proliferation lesion (arrow). (E) Indocyanine green fundus angiographies shows choroidal neovascularization (arrow) that connects retinal vessels, corresponding to retinal-choroidal anastomosis. (F) A vertical optical coherence tomography image shows a pigment epithelial detachment, cystoid macular edema, and a retinal pigment epithelium line that has ruptured (arrow).

**Table 2** Genotype frequency of A69S (*ARMS2*), I62V (*CFH*), and Y402H (*CFH*) polymorphisms in Japanese controls and Japanese patients with retinal angiomatous proliferation

Genotype	rs10490924 (A69S)			rs800292 (I62V)			rs1061170 (Y402H)		
	TT	TG	GG	GG	GA	AA	CC	CT	TT
Controls (n=1,351) <sup>26</sup>	196 (14.6%)	638 (47.8%)	502 (37.6%)	456 (34.1%)	649 (48.5%)	233 (17.4%)	8 (0.6%)	160 (11.9%)	1,174 (87.5%)
Patients (n=36) <sup>26</sup>	31 (86.1%)	3 (8.3%)	2 (5.6%)	20 (55.6%)	11 (30.6%)	5 (13.8%)	0 (0%)	5 (14.3%)	30 (85.7%)
Patients (n=51) <sup>27</sup>	39 (76.5%)	10 (19.6%)	2 (3.9%)	29 (56.9%)	20 (39.2%)	2 (3.9%)	ND	ND	ND
Patients (n=8) (in present study)	7 (87.5%)	1 (12.5%)	0 (0%)	0 (0%)	0 (0%)	8 (100%)	0 (0%)	2 (25%)	6 (75%)

**Note:** Risk alleles are shown in bold.

**Abbreviation:** ND, not described.

Components of the complement system have been identified in drusen, indicating a potential role of the complement system in the pathogenesis of AMD.<sup>41,42</sup> C3 and CFH are key components of the alternative complement pathway. C3 is the most abundant complement component and is synthesized predominantly in the liver. Cleavage of C3 into C3a and C3b is the central step in complement activation and can be initiated by the classic antibody-mediated pathway, the lectin pathway, or the alternative complement pathway. CFH is a critical negative regulator of the alternative pathway of the complement system. It binds to C3b, promotes the decay of C3 convertase, and serves as a cofactor for the factor I-mediated proteolytic inactivation of C3b, resulting in inhibition of the complement cascade.

Because there are bilateral multiple soft drusen in the presence in RAP, we suspected that RAP would be more strongly associated with genetic abnormalities in the complement system than other AMD subtypes. However, we hardly detected the risk alleles of the *CFH* and *C3* genes. One reason for these results is the infrequency of the C allele in Y402H and the G allele in R80G in Asians. Reticular pseudodrusen (RPD) are defined as “drusen that form ill-defined networks of broad interlacing ribbons” in the Wisconsin grading system for maculopathy.<sup>43</sup> RPD have been recognized as a distinctive morphologic feature observed in exudative AMD.<sup>44</sup> Recent studies have demonstrated the association between RPD and reduced macular sensitivity.<sup>45,46</sup> Importantly, it is reported that the prevalence of RPD was high in patients with RAP and the risk genotype (TT) in A69S,<sup>47</sup> and RPD usually occurs bilaterally,<sup>48</sup> suggesting the impact of genetic background for RPD.

It was an unexpected result that we did not detect the risk G allele in I62V (Table 2). The I62V variant has been associated with exudative AMD in both Caucasian and Asian patients.<sup>18,19,21–25,27</sup> In Japanese population samples, it has been demonstrated that the risk genotype (GG) in I62V is

significantly associated with RAP (Table 2).<sup>26,27</sup> However, the risk genotype in I62V was not detected in our RAP patients (Table 2). Our findings suggest that the presence of the risk genotype (I62V) may not be necessarily associated with development of RAP. A larger sample size will be required to determine whether the risk genotype in I62V is eventually associated with RAP.

RAP is characterized by intraretinal neovascularization above the retinal pigment epithelium. Two different origins of this neovascularization have been proposed. Yannuzzi et al suggested that the neovascularization in RAP originates from the neural retina.<sup>29</sup> On the other hand, Freund et al proposed type 3 neovascularization that originates not only from deep retinal capillaries but also from the choroid.<sup>31</sup> As for the location of gene expression, the *CFH* gene is expressed primarily in the retinal pigment epithelium, drusen, and choroidal capillaries;<sup>6</sup> the *C3* gene is expressed in the neural retina, choroid, and retinal pigment epithelium;<sup>41</sup> and the *ARMS2* gene is expressed in the ellipsoid region of the photoreceptor cells.<sup>14</sup> Since it seems that the location of characteristic neovascularization corresponds to the location of susceptible gene expression in RAP, our results support the hypothesis by Yannuzzi et al<sup>29</sup> that the origin of neovascularization in RAP is in the neural retina.

In conclusion, our results suggest that the risk alleles/genotypes of the *ARMS2/HTRA1* SNPs may be strongly associated with development of RAP and that they play a major role in the pathogenesis of intraretinal angiogenesis.

## Acknowledgments

This work was supported by grants from The Jikei University Research Fund (TH) and the Vehicle Racing Commemorative Foundation (TH and HT).

## Disclosure

The authors report no conflicts of interest in this work.