

- [37] Prasad SS, Kojic L, Wen YH, Chen Z, Xiong W, Jia W, et al. Retinal gene expression after central retinal artery ligation: effects of ischemia and reperfusion. *Invest Ophthalmol Vis Sci* 2010;51:6207–19.
- [38] Sato AK, Viswanathan M, Kent RB, Wood CR. Therapeutic peptides: technological advances driving peptides into development. *Curr Opin Biotechnol* 2006;17:638–42.
- [39] Schaar KL, Breneman MM, Savitz SI. Functional assessments in the rodent stroke model. *Exp Transl Stroke Med* 2010;2:13.
- [40] Sims NR, Muyderman H. Mitochondria, oxidative metabolism and cell death in stroke. *Biochim Biophys Acta* 2010;1802:80–91.
- [41] Siren AL, Fratelli M, Brines M, Goemans C, Casagrande S, Lewczuk P, et al. Erythropoietin prevents neuronal apoptosis after cerebral ischemia and metabolic stress. *Proc Natl Acad Sci USA* 2001;98:4044–9.
- [42] Skopeliti M, Kratzer U, Altenberend F, Panayotou G, Kalbacher H, Stevanovic S, et al. Proteomic exploitation on prothymosin alpha-induced mononuclear cell activation. *Proteomics* 2007;7:1814–24.
- [43] Skopeliti M, Iconomidou VA, Derhovanessian E, Pawelec G, Voelter W, Kalbacher H, et al. Prothymosin alpha immunoreactive carboxyl-terminal peptide TKKQKTDEDD stimulates lymphocyte reactions, induces dendritic cell maturation and adopts a beta-sheet conformation in a sequence-specific manner. *Mol Immunol* 2009;46:784–92.
- [44] Ueda H, Fujita R. Cell death mode switch from necrosis to apoptosis in brain. *Biol Pharm Bull* 2004;27:950–5.
- [45] Ueda H, Fujita R, Yoshida A, Matsunaga H, Ueda M. Identification of prothymosin-alpha1, the necrosis-apoptosis switch molecule in cortical neuronal cultures. *J Cell Biol* 2007;176:853–62.
- [46] Ueda H. Prothymosin alpha plays a key role in cell death mode-switch, a new concept for neuroprotective mechanisms in stroke. *Naunyn Schmiedeberg Arch Pharmacol* 2008;377:315–23.
- [47] Ueda H. Prothymosin alpha and cell death mode switch, a novel target for the prevention of cerebral ischemia-induced damage. *Pharmacol Ther* 2009;123:323–33.
- [48] Ueda H, Matsunaga H, Uchida H, Ueda M. Prothymosin alpha as robustness molecule against ischemic stress to brain and retina. *Ann NY Acad Sci* 2010;1194:20–6.
- [49] Ueda H, Matsunaga H, Halder SK. Prothymosin α plays multifunctional cell robustness roles in genomic, epigenetic, and nongenomic mechanisms. *Ann NY Acad Sci* 2012;1269:34–43.
- [50] Ueda H, Matsunaga H, Halder SK. Prothymosin α – a novel endogenous neuroprotective polypeptide against ischemic damages. In: Nyberg FJ, editor. *Neuropeptides in neuroprotection and neuroregeneration*. CRC Press; 2012. p. 128–43.
- [51] Witmer MT, Pavan PR, Fouraker BD, Levy-Clarke GA. Acute retinal necrosis associated optic neuropathy. *Acta Ophthalmol* 2011;89:599–607.
- [52] White BC, Sullivan JM, DeGracia DJ, O'Neil BJ, Neumar RW, Grossman LI, et al. Brain ischemia and reperfusion: molecular mechanisms of neuronal injury. *J Neurol Sci* 2000;179:1–33.
- [53] Yin KJ, Deng Z, Huang H, Hamblin M, Xie C, Zhang J, et al. miR-497 regulates neuronal death in mouse brain after transient focal cerebral ischemia. *Neurobiol Dis* 2010;38:17–26.

ORIGINAL
ARTICLENovel neuroprotective action of prothymosin
alpha-derived peptide against retinal and brain
ischemic damagesSebok Kumar Halder, Hayato Matsunaga, Haruka Yamaguchi and
Hiroshi Ueda*Department of Molecular Pharmacology and Neuroscience, Nagasaki University Graduate School of
Biomedical Sciences, Nagasaki, Japan***Abstract**

Prothymosin alpha (ProT α), a nuclear protein, is implicated in the inhibition of ischemia-induced necrosis as well as apoptosis in the brain and retina. Although ProT α has multiple biological functions through distinct regions in its sequence, it has remained which region is involved in this neuroprotection. This study reported that the active core peptide sequence P₃₀ (amino acids 49–78) of ProT α exerts its full survival effect in cultured cortical neurons against ischemic stress. Our *in vivo* study revealed that intravitreal administration of P₃₀ at 24 h after retinal ischemia significantly blocks the ischemia-induced functional damages of retina at day 7. In addition, P₃₀ completely rescued the retinal ischemia-induced ganglion cell damages at day 7 after the ischemic stress, along with partial

blockade of the loss of bipolar, amacrine, and photoreceptor cells. On the other hand, intracerebroventricular (3 nmol) or systemic (1 mg/kg; *i.v.*) injection of P₃₀ at 1 h after cerebral ischemia (1 h tMCAO) significantly blocked the ischemia-induced brain damages and disruption of blood vessels. Systemic P₃₀ delivery (1 mg/kg; *i.v.*) also significantly ameliorated the ischemic brain caused by photochemically induced thrombosis. Taken together, this study confers a precise demonstration about the novel protective activity of ProT α -derived small peptide P₃₀ against the ischemic damages *in vitro* and *in vivo*.

Keywords: blood vessel, brain ischemia, neuroprotective peptide, prothymosin alpha, retinal ischemia.
J. Neurochem. (2013) **125**, 713–723.

Ischemic damages in the central nervous system including brain and retina are associated with the rapid and severe loss of functional and cellular responses through the mechanisms of necrosis as well as apoptosis by several types of cytotoxic mediators (White *et al.* 2000; Paolucci *et al.* 2003; Ueda and Fujita 2004; Feigin 2005; Flynn *et al.* 2008; Fornage 2009; Dvorianchikova *et al.* 2010; Neroev *et al.* 2010; Sims and Muyderman 2010; Yin *et al.* 2010; Iadecola and Anrather 2011; Witmer *et al.* 2011). At the same time, several neuroprotective molecules such as brain-derived neurotrophic factor (BDNF), fibroblast growth factor (FGF), and erythropoietin (EPO) are produced upon ischemia to play limited attenuation of ischemic damages through the anti-apoptosis mechanisms, without exerting protective activity against necrosis (Siren *et al.* 2001; Korada *et al.* 2002; Maiese *et al.* 2004; Blanco *et al.* 2008; Fujita *et al.* 2009; Madinier *et al.* 2009; Ueda *et al.* 2010; Bejot *et al.* 2011).

Prothymosin alpha (ProT α) has been identified in the conditioned medium of serum-free primary culture of cortical

neurons, as an anti-necrosis factor (Ueda *et al.* 2007). In addition, ProT α potently inhibits the ischemia-induced damages in brain and retina (Fujita and Ueda 2007; Fujita *et al.* 2009; Ueda *et al.* 2010). It is interesting that ProT α

Received October 25, 2012; revised manuscript received December 3, 2012; accepted December 4, 2012.

Address correspondence and reprint requests to Dr. Hiroshi Ueda, Department of Molecular Pharmacology and Neuroscience, Nagasaki University Graduate School of Biomedical Sciences, 1-14 Bunkyo-machi, Nagasaki 852-8521, Japan.

E-mail: ueda@nagasaki-u.ac.jp

Abbreviations used: a.a., amino acid; BDNF, brain-derived neurotrophic factor; Chx10, ceh-10 homeodomain-containing homolog; ERG, electroretinogram; GCL, ganglion cell layer; H&E, hematoxylin and eosin; *i.c.v.*, intracerebroventricularly; *i.v.*, intravenously; *i.vt.*, intravitreally; INL, inner nuclear layer; IPL, inner plexiform layer; NeuN, neuronal nuclei; ONL, outer nuclear layer; P₃₀, peptide sequence comprised of 30 amino acids; PIT, photochemically induced thrombosis; ProT α , prothymosin alpha; tMCAO, transient middle cerebral artery occlusion; TTC, 2,3,5-triphenyltetrazolium chloride.

has distinct actions, which are all related to the cell survival (Jiang *et al.* 2003; Ueda 2008; Mosoian *et al.* 2010; Ueda *et al.* 2012). Some studies revealed that different peptide sequences in ProT α are implicated with these survival actions. The peptide sequence in the central domain of ProT α (amino acids; a.a. 32–52) is related to the interaction with Kelch-like ECH-associated protein 1 (Keap1), which play roles in the induction of oxidative stress-protecting genes expression by liberating Nrf2 from the Nrf2-Keap1 inhibitory complex (Karapetian *et al.* 2005). The N-terminal sequence in ProT α (a.a. 2–29), corresponding to thymosin alpha 1, which has an ability to induce anti-cancer effects (Garaci *et al.* 2007; Danielli *et al.* 2012). In addition, thymosin alpha 1 has been approved in 35 countries for the treatment of hepatitis B and C, and as an immune stimulant and adjuvant (Goldstein and Goldstein 2009; Pierluigi *et al.* 2010). Previous reports suggested that C-terminal region (a.a. 89–109, 99–109 and 100–109) of human ProT α exerts immunoenhancing effects including pro-inflammatory activity through the stimulation of monocytes via toll-like receptor (TLR) signaling, induces dendritic cell maturation and adopts β -sheet conformation (Skopeliti *et al.* 2009). Most recently, there is a report about the survival activity of the middle part (a.a. 41–83) of human ProT α against mutant huntington-caused cytotoxicity in the cultured cells (Dong *et al.* 2012). However, it remains to be elucidated which region is responsible for the neuroprotection against ischemia-induced neuronal damages. In this study, we have attempted to see the neuroprotective activity of ProT α -derived small peptide against ischemic damages *in vitro* and *in vivo*.

Materials and methods

Animals

Male C57/BLJ mice weighing 20–25 g were purchased from Tagawa Experimental Animals (Nagasaki, Japan) and used for all the experiments. Mice were kept in a room maintained at constant temperature ($21 \pm 2^\circ\text{C}$) and relative humidity ($55 \pm 5\%$) with an automatic 12 h light/dark cycle with free access to standard laboratory diet and tap water. Animal care and all experimental procedures were formally approved by Nagasaki University Animal Care and Use Committee (Animal Experiments Approval Number: 1104190914).

Expression constructs and purification procedures for GST-fusion rat ProT α deletion mutants

The rat ProT α gene was amplified from cDNA derived from rat embryonic brain. The gene constructions for expression of recombinant GST-ProT α deletion mutants (Full-length, Δ 1–29, Δ 1–48, Δ 1–68, Δ 1–86, Δ 30–112, Δ 58–112, Δ 79–112, and Δ 102–112) were previously described (Ueda *et al.* 2007; Matsunaga and Ueda 2010). Here, we newly made a GST-ProT α -49–78. The amplified genes blunted at their 5'-ends and cloned in-frame into the BamHI (blunted)-EcoRI sites of pGEX-5X-1 (GE Healthcare Bio-Science Corp, Piscataway, NJ, USA). The PCR primers used were as

follows: Δ 1–48-F, 5'-AGGGATCCAATGGCTGACAATGAGGT AGATG-3' and Δ 79–112-R, 5'-TTGAATTCCTAATCTCCATC TTCTTCCTC-3'. F-primer contains a BamHI site, while all R-primer contains a stop codon and an EcoRI site. The recombinant proteins were purified using Glutathione-Sepharose™ (GE Healthcare Bio-Science Corp).

Identification of functional active core domain in ProT α

To determine the active core domain in ProT α , we measured the survival activity in primary cultured rat cortical neurons. The preparation and culture of cortical neurons were previously described (Ueda *et al.* 2007). The culture of neurons was started at low density (1×10^5 cells/cm²) under the serum-free conditions in the presence or absence of GST and GST-ProT α deletion mutants (100 nM). After 12 h from the start of culture, survival activity was evaluated by WST-8 reduction activity (Cell Counting Kit-8; DOJINDO, Kumamoto, Japan). Finally, we successfully obtained the functional active core domain comprised of 30 amino acids in ProT α (a.a. 49–78) and referred as P₃₀ according to number of amino acids.

Peptide administration

Intravitreal injection was performed using a 33-gauge needle connected to a microsyringe and the needle was inserted approximately 1 mm behind the corneal limbus, guided under a stereoscopic microscope to avoid lens and retinal injury. Peptide P₃₀ was dissolved in 0.05% dimethyl sulfoxide (DMSO), which was diluted with 0.1 M potassium-free phosphate buffered saline (K⁺-free PBS). P₃₀ was injected intravitreally (i.vt.) in the eye with doses of 1, 3 and 10 pmol/ μL at 24 h after retinal ischemia ($n = 5$, $n = 6$ and $n = 7$, respectively). Vehicle was treated with equal volume of 0.05% DMSO in a similar manner. On the other hand, P₃₀ was injected intracerebroventricularly (0.03 and 3 nmol/5 μL , i.c.v.; $n = 6$ and $n = 7$, respectively) in the brain at 1 h after cerebral ischemia (1 h tMCAO). P₃₀ (0.3 and 1 mg/kg) was administered intravenously (i.v.) at 1 h after cerebral ischemia. In addition, P₃₀ was delivered (1 mg/kg, i.v.) at 3 and 6 h after the cerebral ischemic stress.

Retinal ischemia

Retinal ischemia was performed following the method as described previously (Fujita *et al.* 2009). Briefly, mice were anesthetized with an intraperitoneal (i.p.) injection of sodium pentobarbital (50 mg/kg) and pupils were fully dilated with 1% atropine sulfate drops (Nitten, Nagoya, Japan). The anterior chamber of the eye was cannulated with a 33-gauge needle attached to an infusion container of sterile intraocular irrigating solution (BSS PLUS dilution buffer; Alcon, Fort Worth, TX, USA). Retinal ischemia was induced by elevating the IOP to generate a hydrostatic pressure of 130 mm Hg for 45 min by lifting the container. Following 45 min after retinal ischemic stress, the needle was withdrawn and 0.3% ofloxacin (Santen Pharmaceutical Co. Ltd., Osaka, Japan) was applied topically into the eye to avoid infection.

Electroretinogram

Electroretinogram (ERG) study was performed following the protocol as previously described (Fujita *et al.* 2009). Briefly, mice were dark-adapted for 3–4 h, then anesthetized with intraperitoneal injection of pentobarbital sodium (50 mg/kg) and pupils were

dilated with 1% atropine. A contact electrode (KE-S; Kyoto contact lenses, Kyoto, Japan) was placed topically on the corneal apex and reference electrode was placed near the ipsilateral eye. The ground was a subdermal platinum needle electrode near the abdominal area. ERGs were produced by 20 J flash intensities. The flash stimulus source (SLS-3100; Nihon Kohden, Tokyo, Japan) illuminated the eye by diffuse reflection off the interior surface of the ganzfeld. Maximum flash luminance was measured with detector (MEB-9104; Nihon Kohden). After the intensity series, an incandescent background light sufficient to desensitize the rod system was turned on, and ERGs produced by the standard stimulus were recorded every 2 min for 20 min. The background was then turned off, and ERGs were produced by the standard stimulus every 2 min for the first 30 min of dark adaptation. The a- and b-wave amplitudes were measured online (Neuropack m, QP-903B; Nihon Kohden). ERG was performed at day 7 after retinal ischemia.

Middle cerebral artery occlusion model

The transient middle cerebral artery occlusion (tMCAO) model was induced following the method as described previously (Halder *et al.* 2012). Briefly, mice were anesthetized by 2% isoflurane (Mylan, Tokyo, Japan), and body temperature was monitored and maintained at 37°C during surgery. After a midline neck incision, the middle cerebral artery was occluded transiently using 8-0 in size monofilament nylon surgical suture (Natsume Co. Ltd., Tokyo, Japan) coated with silicon (Xantopren, Bayer dental, Osaka, Japan) that was inserted through the left common carotid artery and advanced into the left internal carotid artery. Following 1 h tMCAO, the animals were briefly re-anesthetized with isoflurane and the monofilament was withdrawn for reperfusion studies. Cerebral blood flow was monitored by laser Doppler flowmeter (ALF21; Advance Co., Tokyo, Japan) using a probe (diameter 0.5 mm) of a laser Doppler flowmeter (ALF2100; Advance Co.) inserted into the left striatum (anterior: -0.5 mm, lateral: 1.8 mm from bregma; depth: 4.2 mm from the skull surface) through a guide cannula.

Photochemically induced middle cerebral artery thrombosis

Photochemically induced thrombosis (PIT) was produced following the protocol as described previously (Nagai *et al.* 2007). Briefly, anesthesia was induced with 3% isoflurane, and the rectal temperature was maintained at 37°C. The temporal muscle was dissected, the skull was exposed, and a 1.5-mm opening was made over the middle cerebral artery (MCA). Photo-illumination of green light (wavelength: 540 nm) was achieved with a xenon lamp (model L-4887, Hamamatsu Photonics, Hamamatsu, Japan) with heat-absorbing and green filters, via an optic fiber with a focus of 1 mm, placed on the opening in the skull. Rose Bengal (Wako, Osaka, Japan) was injected (3 mg/kg, i.v.) in mice, and photo-illumination (5000 lx) was applied for 10 min, after which the temporal muscle and skin were replaced. The MCA occlusion time (from the start of light exposure until the flow in the MCA is stopped) was monitored by observation in real time under the microscope.

Retinal and brain tissue preparation

For retinal tissue preparation, mice were deeply anaesthetized with sodium pentobarbital (50 mg/kg, i.p.). Eye was quickly isolated, washed with saline and 4% paraformaldehyde (PFA). Eye was then nicked through pupil, post-fixed in 4% PFA for 24 h and finally

transferred to 25% sucrose solution (in 0.1 M K⁺-free PBS) overnight for cryoprotection. Following frozen in cryoembedding compound, retinal sections were prepared at 10 μ m thickness. For brain tissue preparation, mice were deeply anesthetized with sodium pentobarbital (50 mg/kg, i.p.) and perfused transcardially with 0.1 M PBS, followed by 4% PFA. Brain was then quickly removed, post-fixed in 4% PFA and transferred immediately to 25% sucrose solution overnight. Brain was frozen in cryoembedding compound and coronal sections were cut at 30 μ m thickness for immunohistochemical analysis.

Hematoxylin and eosin staining

For hematoxylin and eosin (H&E) staining, frozen retinal sections were washed with 0.1 M K⁺-free PBS, immersed in Mayer's hematoxylin solution (Wako) for 5 min at 25°C and then washed with tap water for 20 min. Following brief treatment with 95% ethanol, sections were immersed in eosin-alcohol solution (Wako) for 4 min at 25°C. Sections were dehydrated through a series of ethanol solutions, xylene, and over-slipped with Permount (Fisher Scientific, Waltham, MA, USA). Sections were then analyzed using a BZ-8000 microscope with BZ Image Measurement Software (KEYENCE, Osaka, Japan).

Immunohistochemical analysis

To perform fluorescence immunohistochemistry, retinal sections were washed with 0.1 M K⁺-free PBS and incubated with 50% methanol followed by 100% methanol for 10 min. Following treatment with blocking buffers [bovine serum albumin (BSA) as well as 10% goat serum with 0.1% Triton X-100 in phosphate buffered saline (PBST)], retinal sections were incubated overnight at 4°C with following primary antibodies: anti-NeuN (1 : 100; mouse monoclonal IgG₁, clone A60; Chemicon, Temecula, CA, USA); anti-syntaxin-1 (1 : 500; mouse monoclonal; Sigma-Aldrich, St. Louis, MO, USA); and anti-Chx10 (1 : 300; sheep polyclonal; Exalpha Biologicals Inc., MA, USA). Sections were then incubated with Alexa Fluor 488-conjugated anti-mouse IgG and Alexa Fluor 488-conjugated anti-sheep IgG secondary antibodies (1 : 300; Molecular Probes, Eugene, OR, USA). The nuclei were visualized with Hoechst 33342 (1 : 10 000; Molecular Probes). Samples were then washed thoroughly with PBS and cover-slipped with Perma Fluor (Thermo Shandon, Pittsburgh, PA, USA). Images were collected using a BZ-8000 microscope with BZ Image Measurement Software.

For blood vessels staining, biotinylated *Lycopersicon esculentum* (tomato) lectin (Vector Laboratories, Burlingame, CA, USA) is diluted with PBS. Biotinylated tomato lectin was injected (1 mg/mL, 100 μ L, i.v.) at 24 h after cerebral ischemia (1 h tMCAO). Mice were perfused 5 min after tomato lectin injection. Following tissue preparation as described in the method section, coronal brain section was blocked with 2% BSA in 0.1% PBST for 2 h, and then incubated with Alexa Fluor 488 streptavidin conjugates for 2 h at 25°C. Sections were washed with PBS and cover-slipped with Perma Fluor. Images were collected using an LSM 710 confocal microscope with ZEN Software (Carl Zeiss, Oberkochen, Germany).

TTC staining

For 2,3,5-triphenyltetrazolium chloride (TTC) staining, brain was quickly removed at 24 h after cerebral ischemia (1 h tMCAO)

followed by P₃₀ administration ($n = 7$), sectioned coronally with a 1-mm thickness and washed with K⁺-free PBS. Brain slices were incubated in 2% TTC (Sigma-Aldrich) in 0.9% NaCl in dark place for 15–20 min at 25°C and transferred in 4% PFA overnight. Images of brain slices were then collected by scanner, and infarct volume was calculated by Image J software (NIH, Bethesda, MD, USA).

Behavioral assessments

Following P₃₀ administration with doses of 0.03 and 3 nmol/5 μ L (i.c.v., $n = 6$ and $n = 7$, respectively), 0.3 and 1 mg/kg (i.v., $n = 5$ and $n = 7$, respectively) at 1 h as well as 1 mg/kg (i.v.) at 3 and 6 h ($n = 5$ and $n = 7$, respectively) after cerebral ischemia (1 h tMCAO), behavioral studies were assessed through 14 days. Clinical score was evaluated from day 1 after ischemia in the following way: 0, no observable deficits; 1, failure to extend the forepaw fully; 2, circling; 3, falling to one side; 4, no spontaneous movement; 5, death. In this study, 0.5 point was added to each score when the motor dysfunction was severe for scores between 1 and 4. Survival rate was evaluated from day 1 after tMCAO and calculated by the percentage of vehicle or P₃₀ post-treated mice that were alive through 14 days after ischemia.

Statistical analysis

All results are shown as means \pm SEM. Two independent groups were compared using the Student's *t*-test. Multiple groups were compared using Dunnett's multiple comparison test after a one-factor ANOVA or a repeated measure ANOVA. Survival rate was compared using Logrank test after Kaplan–Meyer method. $p < 0.05$ was considered significant.




Fig. 1 The central core peptide sequence of prothymosin alpha (ProT α) is an essential domain for survival activity of ProT α against serum-free ischemic stress. (a) The amino acid sequence of rat ProT α . Red colored sequence indicates functional core domain (a.a. 49–78 referred as P₃₀) in ProT α . (b) Schematic drawings of GST-fusion of full length ProT α and its deletion mutants. (c) Identification of essential domain of ProT α for its survival activity. The primary cultured cortical neurons were incubated with GST and GST-fusion ProT α mutants under the serum-free condition. The survival activity was measured at 12 h after the start of culture. Data represent the means \pm SEM. (* $p < 0.01$, vs. Veh).

ipsilateral eye at 24 h after retinal ischemia. The hematoxylin and eosin (H&E) staining data showed that the number of cells in different retinal layers as well as the retinal thickness is significantly decreased in the vehicle-treated mice at day 7 after the ischemic stress, whereas 10 pmol P₃₀ maximally and significantly inhibited this cellular loss in retina and decrease in retinal thickness at day 7 (Fig. 2a, b).

In electroretinogram (ERG), the amplitude called a-wave represents the functional activity of photoreceptor cells, whereas b-wave indicates the functions of mixture of cells including bipolar, Muller, amacrine, and ganglion cells (Asi and Perlman 1992; Fujita *et al.* 2009). Following after retinal ischemia and reperfusion, the ERGs analysis showed that a- and b-wave amplitudes are significantly decreased in the vehicle-treated mice at day 7 after retinal ischemia, compared with the control (Fig. 2c, d). Following P₃₀ treatment, dose-dependent increase in a- and b-wave amplitudes were observed at day 7 after the retinal ischemic stress, and

10 pmol P₃₀ exerted its maximum protective effect against the ischemic damages (Fig. 2c, d). On the other hand, no significant protective effect of thymosin alpha 1 (a.a. 2–29) corresponding to N-terminal sequence of ProT α and the C-terminal peptide (a.a. 102–112) against retinal ischemic damages were observed at day 7 after ischemia (data are not shown).

P₃₀-induced cell type-specific survival against retinal ischemic damages

To examine the cell type-specific protective activity of ProT α -derived peptide in ischemic retina, P₃₀ was injected (10 pmol/ μ L, i.vt.) in the ipsilateral eye at 24 h after retinal ischemia. The immunohistochemical analysis showed that NeuN-positive neurons (Buckingham *et al.* 2008) in the ganglion cell layer (GCL) are significantly diminished at day 7 after retinal ischemic stress, compared to the control (Fig. 3a). Following P₃₀ treatment at 24 h after retinal




Fig. 2 Prothymosin alpha (ProT α)-derived peptide protects the retinal ischemia-induced functional damages. P₃₀ is injected intravitreally (i.vt.) at the doses of 1, 3, and 10 pmol/ μ L in the ipsilateral eye at 24 h after retinal ischemia. Vehicle is treated with 0.05% dimethyl sulfoxide (DMSO) in a similar manner. (a–d) Protective activity of P₃₀ is a dose-dependent manner. Following P₃₀ injection at 24 h after retinal ischemia, (a) hematoxylin and eosin (H&E) staining of retinal section

is performed at day 7 (right panel). (b–d) Measurement of retinal thickness (b) as well as the a-wave (c) and b-wave (d) amplitudes of ERG analysis are done at day 7 after retinal ischemia in P₃₀ post-treated mice. Data are mean \pm SEM. (* p < 0.05, vs. Control, # p < 0.05, vs. Veh) from experiments using five to seven mice for each group.




Fig. 3 Cell type-specific protection by P_{30} against retinal ischemic damages. P_{30} is injected (10 pmol/ μ L; i.v.t.) in the ipsilateral eye at 24 h after retinal ischemic stress, and the immunohistochemical analysis of retinal sections is performed at day 7. (a–d) The staining of ganglion cells (a) in the ganglion cell layer (GCL) (NeuN: green), bipolar cells (b) in the inner nuclear layer (INL) (Chx10: green), amacrine cells (c)

(syntaxin-1: green), photoreceptor cells (d) in the outer nuclear layer (ONL) (Hoechst: blue) are done at day 7 after retinal ischemia. The higher magnification views of lower panels in (a–d) indicate the expression of retinal cell types noted by dotted rectangles (respective upper panels). Scale bars: 10 μ m. Experiments were performed using five to eight mice for each group.

ischemia, the complete recovery of NeuN-positive neuronal cells was observed in the GCL of ischemic retina at day 7 after the ischemic stress (Fig. 3a). On the other hand, treatment of P_{30} partially, but significantly blocked the loss of Chx10-positive bipolar cells (Rhee *et al.* 2007) in the inner nuclear layer (INL) (Fig. 3b), syntaxin-1-positive amacrine cells (Sherry *et al.* 2006), of which the cell bodies and processes are located in the INL and inner plexiform layer (IPL), respectively (Fig. 3c), and photo-receptor cells in the outer nuclear layer (ONL) (Fig. 3d), compared with the respective controls and vehicles.


Inhibition of cerebral ischemia-induced brain damages by P_{30}

To evaluate the protective activity of P_{30} against ischemic brain damages, mice were post-treated with P_{30} in time- and dose-dependent manner following different routes of administration, and subsequent 2,3,5-triphenyl tetrazolium chloride (TTC) staining at 24 h and behavioral assessments through

14 days were performed after cerebral ischemia (1 h tMCAO). The TTC staining data showed that the infarct volume is significantly decreased at 24 h in the ischemic brain by intracerebroventricular (i.c.v.) injection of 3 nmol P_{30} at 1 h after tMCAO (Fig. 4a, b), but not by 0.03 nmol (data are not shown). We also observed that the clinical score is significantly decreased at day 1 after 1 h tMCAO in mice injected with 3 nmol P_{30} (i.c.v.) at 1 h after the ischemic stress (Fig. 4c). In addition, significant decrease in clinical score and increase in survival rate were observed through 14 days after i.c.v. delivery (1 h after ischemia) of 3 nmol P_{30} , an indication of long-lasting protective effect of P_{30} against ischemic brain damages (Fig. 4d, e).

On the other hand, P_{30} was injected intravenously (i.v.) with doses of 0.3 and 1 mg/kg at 1 h after cerebral ischemia (1 h tMCAO). Our TTC staining data revealed that the infarct volume is significantly decreased at 24 h in the ischemic brain treated with 1 mg/kg of P_{30} treatment at 1 h after the ischemic stress (Fig. 4f, g). Following

Fig. 4 P₃₀ inhibits cerebral ischemia-induced brain damages. (a–e) Intracerebroventricular (i.c.v.) delivery with P₃₀ protects ischemic brain damages. P₃₀ is injected (3 nmol, i.c.v.) in the brain at 1 h after the cerebral ischemia [1 h transient middle cerebral artery occlusion (tMCAO)], (a–c) TTC staining (a), measurement of infarct volume (b), and clinical scores (c) are performed at day 1 after tMCAO. Data represent the means \pm SEM. (* p < 0.05, vs. Veh). (d, e) Assessment of the clinical score (d) and survival rate (e) are done through 14 days after the tMCAO mice post-treated with P₃₀. The group of P₃₀ treatment was significant compared to group of Veh treatment (* p < 0.01, vs. Veh). Survival rate of P₃₀ treatment tended to be significant compared to Veh treatment. Experiments were performed using five to eight mice for each group. (f–k) Blockade of cerebral ischemia-induced brain damages by systemic administration of P₃₀. P₃₀ is delivered intravenously (i.v.) with doses of 0.3 and 1 mg/kg at 1 h after cerebral ischemia (1 h tMCAO). (f–h) TTC staining (f), measurement of infarct volume (g) and clinical scores (h) are performed at day 1 after tMCAO. (i, j) The clinical score (i) and survival rate (j) are measured for 7 and 14 days, respectively. The group of P₃₀ treatment (1 mg/kg) was significant compared to group of Veh treatment in clinical score (* p < 0.01, vs. Veh). Survival rate of P₃₀ treatment tended to be significant compared to Veh treatment. (k) Time-course systemic injection of P₃₀ (1 mg/kg, i.v.) at 1, 3, and 6 h after cerebral ischemia (1 h tMCAO). Data represent the means \pm SEM. (* p < 0.05, vs. Veh). Experiments are performed using six to eight mice for each group.



post-treatment (i.v.) with 1 mg/kg of P₃₀ at the same time point, the clinical score was significantly declined through 7 days and survival rate was maximally increased through 14 days after tMCAO, compared with the vehicle and ischemic mice treated with 0.3 mg/kg of P₃₀ (Fig. 4h–j). The behavioral study also confirmed that systemic (i.v.) P₃₀ delivery with the dose of 1 mg/kg at 1 h after ischemia induces its maximum protective effect at day 7 against the ischemic brain damages, compared to P₃₀ treatment at 3 or 6 h after cerebral ischemia (Fig. 4k).

P₃₀ inhibits the cerebral ischemia-induced blood vessel damages

In the ischemic stroke and cerebrovascular disease, vascular defect is occurred along with neuronal damages (Paul *et al.* 2001; Fujita and Ueda 2007). To investigate whether P₃₀ protects the ischemia-induced blood vessel damages, P₃₀ was injected (1 mg/kg; i.v.) at 1 h after cerebral ischemia (1 h tMCAO). Following blood vessel immunostaining using biotinylated tomato lectin and Alexa Fluor 488 streptavidin at 24 h after ischemia, the findings revealed that the number

blood vessels are markedly decreased in somatosensory cortex in the brain of vehicle-treated mice, compared with the control (Fig. 5a, b). In addition, the decrease in lengths of the blood vessels was observed at 24 h after tMCAO (Fig. 5a, c). This ischemia-induced loss of tomato lectin-stained blood vessels in terms of number and lengths was completely recovered in the somatosensory cortex at 24 h after the ischemic stress in mice post-treated with P₃₀, but the lengths were relatively larger than the vessels in the control brain, an indication of the protective role of P₃₀ against ischemia-induced blood vessel damages (Fig. 5a–c). Similar results of the recovery of cerebral ischemia-induced blood vessels damages by P₃₀ were observed in the striatum and hippocampus at 24 h after 1 h tMCAO (data are not shown).

P₃₀ ameliorates the ischemic brain caused by photochemically induced thrombosis

It is well known that ischemic model because of middle cerebral artery (MCA) occlusion with photochemically induced thrombosis (PIT) is analogous to clinical condition




Fig. 5 P₃₀ inhibits the cerebral ischemia-induced blood vessels damages. (a–c) P₃₀ is injected (1 mg/kg, i.v.) at 1 h after cerebral ischemia [1 h transient middle cerebral artery occlusion (tMCAO)]. Following administration with biotinylated tomato lectin (1 mg/mL, 100 μ L, i.v.) at 24 h after cerebral ischemia (1 h tMCAO), and subsequent perfusion 5 min after biotinylated tomato lectin injection, immunostaining of blood vessels by Alexa Fluor 488 streptavidin (a) as well as measurement of number (b) and length (c) of blood vessels is performed at day 1 after the ischemic stress. Data represent the means \pm SEM. (* p < 0.05, vs. Control, # p < 0.05, vs. Veh) from experiments using five to seven mice for each group.

(Tanaka *et al.* 2007). In this ischemic mouse model, there was a significant behavioral damage evaluated by clinical score (Fig. 6a). This damage was significantly attenuated by systemic post-treatment with P₃₀ (1 mg/kg, i.v.) at 1 h after PIT (Fig. 6a). Following behavioral study after PIT stress, neurological assessments using TTC staining were performed at 24 h. The TTC staining data revealed that there was a marked increase in cerebral infarction observed at 24 h after PIT in vehicle-treated mice (Fig. 6b), but this cerebral brain damage in terms of infarct volume and hemisphere expansion was significantly inhibited by systemic treatment of P₃₀ (Fig. 6c, d).

Discussion

This study demonstrates three major findings. First, active core peptide domain P₃₀ (a.a. 49–78) derived from ProT α retains the original survival activity in cultured neuronal cells against ischemic (serum-free) stress. Second, characterizations of P₃₀ actions reveal that it potently inhibits the ischemia-induced damages in retina and brain. Third, P₃₀




Fig. 6 P₃₀ improves the ischemic brain damages caused by photochemically induced thrombosis. P₃₀ is administered (1 mg/kg, i.v.) at 1 h after photochemically induced thrombosis (PIT) in mice. (a) Clinical scores at day 1 after ischemia. (b) Representative picture of TTC staining at 24 h after PIT. (c, d) Measurement of infarct volume (c) and hemispheric expansion (d) at 24 h after PIT. Data represent the means \pm SEM. (* p < 0.05, ** p < 0.01, vs. Veh) from experiments using five to seven mice for each group.

induces protective action against ischemia-induced disruption of cerebral blood vessels.

Several *in vitro* studies reported about the different sequence-specific functions of ProT α , which is also involved in the mechanisms of cell survival (Jiang *et al.* 2003; Karapetian *et al.* 2005; Skopeliti *et al.* 2007; Ueda *et al.* 2007; Ueda 2009; Mosoian *et al.* 2010; Danielli *et al.* 2012; Dong *et al.* 2012). On the basis of previous information, we firstly designed *in vitro* experiments to find out the sequence-specific neuroprotective actions of ProT α using various deletion mutants of GST-ProT α in neuronal cells culture under ischemic stress. The peptides lacking sequence (a.a. 1–29), which belongs to thymosin alpha 1 (a.a. 2–29), sequence (a.a. 1–48), which mostly covers the binding region for Keap1, or C-terminal sequences (a.a. 79–112 and 102–112) completely retained the original survival activity as like ProT α . However, the significant decrease in survival effect was observed by the deficiency of parts of the central core peptide sequence comprised of 30 amino acids in ProT α (a.a. 49–78). Interestingly, this central active core peptide of ProT α referred as P₃₀ (a.a. 49–78) itself exerts full survival action in neuronal cells against ischemia. Retinal ischemia causes the functional and cellular damages in different layers of retina through several destructive cascade of mechanisms, as consequence of visual impairment and blindness (Osborne *et al.* 2004). Our recent *in vivo* studies suggested that ProT α potentially inhibits this ischemia-induced functional and cellular damages of retina (Fujita *et al.* 2009; Ueda *et al.* 2010). To evaluate the *in vivo* protective effect of P₃₀ against ischemic damages, ischemic retina was post-treated with P₃₀. The findings using H&E staining and ERG study revealed that P₃₀ significantly blocks the retinal ischemia-induced decrease in cells number of different layers and retinal thickness. In addition, immunohistochemical analysis clarified that P₃₀ completely rescues the retinal ischemia-induced ganglion cell damages, along with the partial but significant blockade of the loss of bipolar, amacrine, and photoreceptor cells. Stroke following cerebral ischemia (tMCAO) or photothrombotic brain ischemia causes the neuronal damages, along with adequate disruption of cerebral blood vessels (Beck and Plate 2009; Hofmeijer and van Putten 2012; Krysl *et al.* 2012). We previously explained the protective role of ProT α against cerebral ischemia-induced brain damages (Fujita and Ueda 2007; Ueda 2009; Ueda *et al.* 2010). The present findings of TTC staining and neurological assessment suggested that intracerebroventricular (3 nmol, i.c.v.) or systemic (1 mg/kg, i.v.) treatment with P₃₀ at 1 h after cerebral ischemia (1 h tMCAO) significantly blocks ischemia-induced brain damages. Following immunostaining with tomato lectin in P₃₀-treated (1 mg/kg, i.v.) ischemic mice, the complete recovery of ischemia-induced (tMCAO) cerebral blood vessels damages was observed through day 1, a consideration of P₃₀ as a new angiogenic factor. In addition, systemic administration with P₃₀ (1 mg/

kg, i.v.) significantly ameliorated the ischemic brain caused by photochemically induced thrombosis (PIT), a representative clinical model of cerebral ischemia.

The present investigations were performed following several routes of the administration of P₃₀. According to the fact that retinal ischemia possesses high reproducibility and quantitation to understand the pathophysiological changes and signaling pathways under ischemic condition (Prasad *et al.* 2010), we used this ischemic injury as a simple model for screening of survival activity by i.vt. administration of P₃₀. We already reported that i.v. administration with full-length ProT α induces protective effect against retinal ischemia (Fujita *et al.* 2009). In brain ischemia, we firstly decided to perform i.c.v. administration of P₃₀ to evaluate the improvement of ischemic injury, and successfully confirmed against ischemic brain damages. Our recent studies revealed that myc-tagged ProT α (1 mg/kg) is penetrated to the damaged area of brain at least 3 h after brain ischemia by intraperitoneal (i.p.) administration, and that systemic administration (i.p. and i.v.) of ProT α ameliorates brain ischemia-induced functional and cellular damages (Fujita and Ueda 2007). It is well known that brain ischemic stress disrupts the blood–brain barrier (BBB) (Paul *et al.* 2001; Fujita and Ueda 2007). Thus, we presume that like ProT α , systemic administered P₃₀ would penetrate to the damaged brain through the disrupted BBB. Although relationship between route of administration and penetrated amounts of P₃₀ to the brain are not clear, isotope and/or fluorescence labeling might be useful method for the calculation of penetration. In the systemic administration, ProT α and P₃₀ exercise the maximum improvement effect against brain ischemia in 100 μ g/kg (equivalent 8.08 nmoles/kg) and 1 mg/kg (equivalent 0.30 μ moles/kg), respectively. This difference of efficacy between ProT α and P₃₀ might be because of the stability of P₃₀ *in vivo*, though GST-ProT α and GST-P₃₀ (a.a. 49–78) showed similar survival activity in this *in vitro* study. However, the modification of amino acid and/or mutation in sequence of P₃₀ may provide a better solution to improve the stability and survival activity of P₃₀. This should be the next issue to address.

Cortical neurons in serum-free primary culture rapidly die by necrosis, which is completely inhibited by ProT α (Fujita and Ueda 2003; Ueda *et al.* 2007). As ProT α also protects the retinal ischemia-induced necrosis and apoptosis through the up-regulation of BDNF and EPO, and this retinal protection is completely abolished by antisense oligodeoxynucleotide or antibody treatment against ProT α (Fujita *et al.* 2009; Ueda *et al.* 2010), it should be an interesting next subject to investigate whether the same mechanisms are involved in the P₃₀-induced functional and cellular protection against ischemic damages. Despite of being neuroprotective activity of several proteins, peptides have been detected as a new class of attractable therapeutic molecule owing to their diversity, synthesis, and higher