

to the Rex response element (RxRE) of viral transcripts and promotes selective nuclear export of viral genomic RNA [32,36]. P30II is also an RNA-binding protein that binds mainly to double-spliced *tax/rex* mRNA and reduces translation from this mRNA by retaining it to the nucleolus [37–39]. In a series of experiments testing the influence of these viral factors on NMD activity, only p27Rex and p21Rex displayed a robust time- and dose-dependent inhibition of NMD activity (Fig. 4). The most important and well-studied function of Rex is to promote nuclear export and enhance the stability of unspliced and singly spliced HTLV-1 mRNAs that encode viral structural proteins, by interacting with Rex Responsive Element (RxRE) situated at the 3' end of all HTLV-1 mRNAs [18,19,40–42], and by engagement with the cellular nuclear export factor CRM1 [18,43,44]. The molecular mechanism of Rex in protection of unspliced viral RNA has not been fully clarified, however, partially explained by active export of mRNA from the nucleus, thereby escaping from the site of splicing. Nevertheless, the fate of viral RNA in the cytoplasm has not been methodically explored. From our data, we propose that Rex assists viral RNA transit within the cytoplasm and ensures efficient translation by subduing NMD and the host mRNA surveillance systems.

We demonstrated that the NMD activity of TL-Om1 cells was significantly suppressed (Fig. 3B), although these cells do not express any of viral proteins at detectable levels. The molecular mechanism of NMD inhibition in TL-Om1 cells is apparently different from that of Rex expressing cells, however, it can be conceived that disturbances of cellular homeostasis caused by HTLV-1 during the early stages of infection, including inhibition of NMD by Rex, might have been conserved after malignant transformation of the infected cells. To gain further insights into the mechanisms, comparative analyses of cell lines that were HTLV-1 immortalized in vitro, such as MT-2, and those derived from ATL and without Rex expression, such as TL-Om1, may provide valuable information as model cells in the early stages of infection and immortalization, and those in the later transformed phase, respectively.

During the completion of our study, Mocquet et al. [45] demonstrated that HTLV-1 Tax interacts with UPP1 and exerts an inhibitory effect on NMD activity. In the present study, Tax also consistently exhibited an inhibitory effect on NMD activity, but both p21Rex and p27Rex consistently outperformed Tax in our assays (Figs. 4 and 6A). Therefore, the studies described here specifically focused on the inhibition of NMD activity exerted by both Rex alleles. In our view, it is not

surprising that retroviruses have evolved multiple and overlapping mechanisms to overcome the restrictive properties of NMD. This is especially relevant at early times of viral infection or alternatively, when the provirus is emerging from latency. Under such conditions, there may be insufficient Rex concentration to saturate the NMD pathway. Therefore, another mechanism that augments the NMD inhibitory activity of Rex; such as that presented by Tax, may be required to cooperatively incapacitate NMD. Significantly, recent studies have demonstrated that *tax/rex* mRNAs are the most abundant viral transcripts at early times of infection or following proviral reactivation, and that viral gene expression follows a two-phase pattern that is determined principally by Rex [46].

Our data are consistent with the view that increased luciferase activity of NMD-sensitive reporters by Rex was due to the suppression of overall cellular NMD activity but not due to RxRE-mediated nuclear export of reporter mRNAs by Rex. This is because (1) none of the NMD reporter mRNAs used in this study contain RxRE; (2) the amounts of unspliced and spliced forms of β -globin mRNA in nuclear and cytosolic fractions of the cells were not influenced by Rex expression (Fig. S3A); and (3) the insertion of RxRE after the β -globin fragment of the NMD reporter plasmids had no impact on the NMD inhibitory effect of Rex (Fig. S3B). The data that p21Rex, lacking NLS and RxRE-binding domain at the N-terminal region, is also capable of suppressing NMD strongly support the notion that NMD inhibition is genetically separable from the RxRE-binding function of Rex (Fig. 4). Indeed, we demonstrated that Rex stabilizes a series of NMD target mRNA with NMD-inducing features such as uORFs and 3' UTR introns (Fig. 5), providing strong evidence that Rex blocks overall cellular NMD activity. On the other hand, it is widely accepted that NMD is involved in cellular homeostasis not only by eliminating PTC-containing aberrant mRNAs but also by controlling the levels of natural transcripts. Approximately 1–10% of eukaryotic transcripts are known to be targeted by NMD because of their potentially NMD-sensitive structures, i.e., PTCs encoded by alternative exons, introns in 3' UTRs, or uORFs [2,47]. Thus, NMD dysfunction can lead to the destruction of cell homeostasis and may even trigger tumorigenesis [48]. Clearly, overall NMD inhibition by Rex is beneficial to HTLV-1 as a direct mechanism to enhance the stability of HTLV-1 genomic RNA. In contrast, it may perturb cellular gene expression and discrimination of aberrant mRNAs, which are critically regulated by NMD under normal conditions. Indeed, NMD targets many mRNAs, of which products participate in cell cycle regulation, T cell

with that in control cells (half-life = 2.3 h). In contrast, Tax and p30II did not significantly stabilize HTLV-1 unspliced RNA (half-lives = 2.8 and 3.4 h, respectively). The half-life was calculated based on the regression curve (dashed line) ($n = 4$, mean \pm SE). The right side panel shows mRNAs levels of *tax*, *p27rex*, *p21rex*, and *p30ii* at 0 h measured by semi-quantitative RT-PCR in HTLV-1 infected cells expressing the indicated viral proteins, ectopically. (C) Expression of HTLV-1 structural proteins, Gag p53 (precursor), p24, and p19, in HTLV-1 infected 293FT cells co-cultured with MT-2 after NMD suppression by *As-upf2* mRNA overexpression to knockdown UPP2 protein expression (top panel). The expression levels of all Gag proteins, p53, p24, and p19, were significantly elevated in the 293FT cells after NMD suppression. (D) The influence of p27Rex and p21Rex on the expression of HTLV-1 Gag structural proteins, p53, p24, and p19, in HTLV-1 infected 293FT cells co-cultured with MT-2. The results showed that the expression levels of HTLV-1 Gag proteins, especially those of Gag p53 and p19, were significantly increased by p27Rex and p21Rex overexpression.

development, and inflammation [2]. The potential pleiotropic impact of Rex mediated through the NMD pathway represents an important and novel aspect of host–retrovirus interaction in viral leukemogenesis.

In conclusion, this study provides evidence that the host-encoded NMD pathway restricts viral RNA expression, thereby reducing HTLV-1. Furthermore, we demonstrated that a novel function for HTLV-1 Rex in inhibition of the host NMD machinery. Rex-mediated inhibition of NMD activity is pleiotropic and may affect both viral and cellular target RNA. Together with the data presented here, there is accumulating evidence of a dynamic interplay between the cellular NMD pathway and pathways for viral gene expression and replication. Clarifying underlying molecular mechanism in inhibition of NMD by Rex is essential to understand this new aspect of host–pathogen interaction. Finally we suggest that the NMD pathway may exert anti-viral effects on other RNA viruses and may participate generally in the innate immune response to viral pathogens. It is therefore likely that other viruses have evolved mechanisms to silence NMD activity during viral infection.

Acknowledgment

This work was supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan, to TW (No. 19659241) and to KN (Nos. 22700863, 24501304).

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.micinf.2013.03.006>.

References

- [1] R.T. Hillman, R.E. Green, S.E. Brenner, An unappreciated role for RNA surveillance, *Genome Biol.* 5 (2004) R8.
- [2] J.T. Mendell, N.A. Sharifi, J.L. Meyers, F. Martinez-Murillo, H.C. Dietz, Nonsense surveillance regulates expression of diverse classes of mammalian transcripts and mutates genomic noise, *Nat. Genet.* 36 (2004) 1073–1078.
- [3] F.J. Iborra, A.E. Escargueil, K.Y. Kwek, A. Akoulitchev, P.R. Cook, Molecular cross-talk between the transcription, translation, and nonsense-mediated decay machineries, *J. Cell Sci.* 117 (2004) 899–906.
- [4] L.E. Maquat, Nonsense-mediated mRNA decay: splicing, translation and mRNP dynamics, *Nat. Rev.* 5 (2004) 89–99.
- [5] L. Balvay, M.L. Lastra, B. Sargueil, J. Darlix, T. Ohlmann, Translational control of retroviruses, *Nat. Rev.* 5 (2007) 128–140.
- [6] N.J. McGlincy, C.W. Smith, Alternative splicing resulting in nonsense-mediated mRNA decay: what is the meaning of nonsense? *Trends Biochem. Sci.* 33 (2008) 385–393.
- [7] E.P. Plant, P. Wang, J.L. Jacobs, J.D. Dinman, A programmed-1 ribosomal frameshift signal can function as a *cis*-acting mRNA destabilizing element, *Nucleic Acids Res.* 32 (2004) 784–790.
- [8] A.L. Saltzman, Y.K. Kim, Q. Pan, M.M. Fagnani, L.E. Maquat, B.J. Blencowe, Regulation of multiple core spliceosomal proteins by alternative splicing-coupled nonsense-mediated mRNA decay, *Mol. Cell Biol.* 28 (2008) 4320–4330.
- [9] C. Theis, J. Reeder, R. Giegerich, KnotInFrame: prediction of -1 ribosomal frameshift events, *Nucleic Acids Res.* 36 (2008) 6013–6020.
- [10] A.M. Dickson, J. Wilusz, Strategies for viral RNA stability: live long and prosper, *Trends Genet.* 27 (2011) 286–293.
- [11] S. Boelz, G. Neu-Yilik, N.H. Gehring, M.W. Hentze, A.E. Kulozik, A chemiluminescence-based reporter system to monitor nonsense-mediated mRNA decay, *Biochem. Biophys. Res. Commun.* 349 (2006) 186–191.
- [12] T. Ohsugi, T. Kumasaka, T. Urano, Construction of a full-length human T cell leukemia virus type I genome from MT-2 cells containing multiple defective proviruses using overlapping polymerase chain reaction, *Anal. Biochem.* 329 (2004) 281–288.
- [13] N.H. Gehring, G. Neu-Yilik, T.W. Schell, M. Hentze, A.E. Kulozik, Y14 and hUpf3b form an NMD-activating complex, *Mol. Cell* 11 (2003) 939–949.
- [14] Y.K. Kim, L. Furic, L. DesGroseillers, L.E. Maquat, Mammalian stau-phen1 recruits Upf1 to specific mRNA 3'UTRs so as to elicit mRNA decay, *Cell* 120 (2005) 195–208.
- [15] B. Lee, Y. Tanaka, H. Tozawa, Monoclonal antibody defining Tax1 protein of human T-cell leukemia virus type-I, *Tohoku J. Exp. Med.* 157 (1989) 1–11.
- [16] Y. Tanaka, B. Lee, T. Inoi, H. Tozawa, N. Yamamoto, Y. Hinuma, Antigens related to three core proteins of HTLV-I (p24, p19 and p15) and their intracellular localizations, as defined by monoclonal antibodies, *Int. J. Cancer* 37 (1986) 35–42.
- [17] J. Zhao, B.K. Sun, J.A. Erwin, J. Song, J.T. Lee, Polycomb proteins targeted by a short repeat RNA to the mouse X chromosome, *Science* 322 (2008) 750–756.
- [18] M. Gröne, C. Koch, R. Grassmann, The HTLV-1 Rex protein induces nuclear accumulation of unspliced viral RNA by avoiding intron excision and degradation, *Virology* 218 (1996) 316–325.
- [19] M. Hidaka, J. Inoue, M. Yoshida, M. Seiki, Post-transcriptional regulator (rex) of HTLV-1 initiates expression of viral structural proteins but suppresses expression of regulatory proteins, *EMBO J.* 7 (1988) 519–523.
- [20] J. Inoue, M. Yoshida, M. Seiki, Transcriptional (p40x) and post-transcriptional (p27x-III) regulators are required for the expression and replication of human T-cell leukemia virus type I genes, *Proc. Natl. Acad. Sci. U. S. A.* 84 (1987) 3653–3657.
- [21] A. Eulalio, I. Behm-Ansmant, E. Izaurralde, P-Bodies: at the crossroads of post-transcriptional pathways, *Nat. Rev.* 8 (2007) 9–22.
- [22] T.M. Franks, G. Singh, J. Lykke-Andersen, UPF1 ATPase dependent mRNP disassembly is required for completion of nonsense-mediated mRNA decay, *Cell* 143 (2010) 938–950.
- [23] O. Isken, L.E. Maquat, The multiple lives of NMD factors: balancing roles in gene and genome regulation, *Nat. Rev.* 9 (2008) 699–712.
- [24] R. Parker, U. Sheth, P bodies and the control of mRNA translation and degradation, *Mol. Cell* 25 (2007) 635–646.
- [25] J. Rehwinkel, I. Behm-Ansmant, D. Gatfield, E. Izaurralde, A crucial role for GW182 and the DCP1:DCP2 decapping complex in miRNA-mediated gene silencing, *RNA* 11 (2005) 1640–1647.
- [26] U. Sheth, R. Parker, Targeting of aberrant mRNAs to cytoplasmic processing bodies, *Cell* 125 (2006) 1095–1109.
- [27] L. Unterholzner, E. Izaurralde, SMG7 acts as a molecular link between mRNA surveillance and mRNA decay, *Mol. Cell* 16 (2004) 587–596.
- [28] J.E. Weil, K.L. Beemon, A 3' UTR sequence stabilizes termination codons in the unspliced RNA of Rous sarcoma virus, *RNA* 12 (2007) 102–110.
- [29] E.G. Lee, D. Kuppers, M. Horn, J. Roy, C. May, M.L. Linial, A premature termination codon mutation at the C terminus of foamy virus *gag* downregulates the levels of spliced *pol* mRNA, *J. Virol.* 82 (2008) 1656–1664.
- [30] L. Ajamian, L. Abrahamyan, M. Milev, P.V. Ivanov, A.E. Kulozik, N.H. Gehring, A.J. Mouland, Unexpected roles for UPF1 in HIV-1 RNA metabolism and translation, *RNA* 14 (2008) 914–927.
- [31] J.R. Hogg, S.P. Goff, UPF1 senses 3' UTR length to potentiate mRNA decay, *Cell* 143 (2010) 379–389.
- [32] O.U. Susova, V.E. Gurtsevich, The role of region pX in the life cycle of HTLV-I and in carcinogenesis, *Mol. Biol.* 37 (2003) 334–344.
- [33] G. Franchini, R. Fukumoto, J.R. Fullen, T-cell control by human T-cell leukemia/lymphoma virus type-1, *Int. J. Hematol.* 78 (2003) 280–296.
- [34] M. Boxus, J.-C. Twizere, S. Legros, J.-F. Dewulf, R. Kettmann, L. Willems, The HTLV-1 Tax interactome, *Retrovirology* 5 (2008) 76.

- [35] F. Kashanchi, J.N. Brady, Transcriptional and post-transcriptional gene regulation of HTLV-1, *Oncogene* 24 (2005) 5938–5951.
- [36] I. Younis, P.L. Green, The human T-cell leukemia virus Rex protein, *Front. Biosci.* 10 (2005) 431–445.
- [37] C. Nicot, M. Dunder, J.M. Johnson, J.R. Fullen, N. Alonzo, R. Fukumoto, G.L. Princler, D. Derse, T. Misteli, G. Franchini, HTLV-1-encoded p30II is a post-transcriptional negative regulator of viral replication, *Nat. Med.* 10 (2004) 197–201.
- [38] U. Sinha-Datta, A. Datta, S. Ghorbel, M. Duc Dodon, C. Nicot, Human T-cell lymphotropic virus type I Rex and p30 interactions govern the switch between virus latency and replication, *J. Biol. Chem.* 282 (2007) 14608–14615.
- [39] H.H. Baydoun, M. Bellon, C. Nicot, HTLV-1 yin and yang: Rex and p30 master regulators of viral mRNA trafficking, *AIDS Rev.* 10 (2008) 195–204.
- [40] Y.F. Ahmed, G.M. Gilmartin, S.M. Hanly, J.R. Nevins, W.C. Greene, Structure–function analyses of the HTLV-I Rex and HIV-1 Rev RNA response elements: insights into the mechanism of Rex and Rev action, *Gene Develop.* 4 (1990) 1014–1022.
- [41] Y.F. Ahmed, G.M. Gilmartin, S.M. Hanly, J.R. Nevins, W.C. Greene, The HTLV-I Rex response element mediates a novel form of mRNA polyadenylation, *Cell* 64 (1991) 727–737.
- [42] Y. Adachi, T. Nosaka, M. Hatanaka, Protein kinase inhibitor H-7 blocks accumulation of unspliced mRNA of human T-cell leukemia virus type I (HTLV-I), *Biochem. Biophys. Res. Commun.* 169 (1990) 469–475.
- [43] Y. Hakata, T. Umemoto, S. Matsushita, H. Shida, Involvement of human CRM1 (exportin 1) in the export and multimerization of the Rex protein of human T-cell leukemia virus type 1, *J. Virol.* 72 (1998) 6602–6607.
- [44] Y. Hakata, M. Yamada, Rat CRM1 is responsible for the poor activity of human t-cell leukemia virus type 1 Rex protein in rat cells, *J. Virol.* 75 (2001) 11515–11525.
- [45] V. Mocquet, J. Neusiedler, F. Rende, D. Cluet, J.-P. Robin, J.-M. Terme, M. Duc Dodon, J. Wittmann, C. Morris, H.L. Hir, V. Ciminale, P. Jalinot, The human t-lymphotropic virus type 1 Tax protein inhibits nonsense-mediated mRNA decay by interacting with INT6/EIF3E and UPF1, *J. Virol.* 86 (2012) 7530–7543.
- [46] F. Rende, I. Cavallari, A. Corradin, M. Silic-Benussi, F. Toulza, G.M. Toffolo, Y. Tanaka, S. Jacobson, G.P. Taylor, D.M. D’Agostino, C.R.M. Bangham, V. Ciminale, Kinetics and intracellular compartmentalization of HTLV-1 gene expression: nuclear retention of HBZ mRNAs, *Blood* 117 (2011) 4855–4859.
- [47] P. Nicholson, H. Yepiskoposyan, S. Metze, R.Z. Orozco, N. Kleinschmidt, O. Mühlemann, Nonsense-mediated mRNA decay in human cells: mechanistic insights, functions beyond quality control and the double-life of NMD factors, *Cell. Mol. Life Sci.* 67 (2010) 677–700.
- [48] L.B. Gardner, Nonsense-mediated RNA decay regulation by cellular stress: implications for tumorigenesis, *Mol. Cancer Res.* 8 (2010) 295–308.

Polycomb-Mediated Loss of miR-31 Activates NIK-Dependent NF- κ B Pathway in Adult T Cell Leukemia and Other Cancers

Makoto Yamagishi,^{1,3} Kazumi Nakano,¹ Aiko Miyake,¹ Tadanori Yamochi,¹ Yayoi Kagami,¹ Akihisa Tsutsumi,¹ Yuka Matsuda,¹ Aiko Sato-Otsubo,⁴ Satsuki Muto,^{1,4} Atae Utsunomiya,⁵ Kazunari Yamaguchi,⁶ Kaoru Uchimaru,² Seishi Ogawa,⁴ and Toshiki Watanabe^{1,*}

¹Graduate School of Frontier Sciences

²Institute of Medical Science

The University of Tokyo, Tokyo, 108-8639, Japan

³Japan Foundation for AIDS Prevention, Tokyo, 101-0061, Japan

⁴Cancer Genomics Project, Graduate School of Medicine, The University of Tokyo, Tokyo, 113-8655, Japan

⁵Department of Haematology, Imamura Hospital, Bun-in, Kagoshima, 890-0064, Japan

⁶Department of Safety Research on Blood and Biologics, NIID, Tokyo, 208-0611, Japan

*Correspondence: tnabe@ims.u-tokyo.ac.jp

DOI 10.1016/j.ccr.2011.12.015

SUMMARY

Constitutive NF- κ B activation has causative roles in adult T cell leukemia (ATL) caused by HTLV-1 and other cancers. Here, we report a pathway involving Polycomb-mediated miRNA silencing and NF- κ B activation. We determine the miRNA signatures and reveal miR-31 loss in primary ATL cells. MiR-31 negatively regulates the noncanonical NF- κ B pathway by targeting NF- κ B inducing kinase (NIK). Loss of miR-31 therefore triggers oncogenic signaling. In ATL cells, miR-31 level is epigenetically regulated, and aberrant upregulation of Polycomb proteins contribute to miR-31 downregulation in an epigenetic fashion, leading to activation of NF- κ B and apoptosis resistance. Furthermore, this emerging circuit operates in other cancers and receptor-initiated NF- κ B cascade. Our findings provide a perspective involving the epigenetic program, inflammatory responses, and oncogenic signaling.

INTRODUCTION

Adult T cell leukemia (ATL) is an aggressive T cell neoplasm with very poor prognosis (Yamaguchi and Watanabe, 2002). Human T cell leukemia virus type I (HTLV-I) is recognized as an etiological factor in T cell malignancy. Although mounting molecular evidence has contributed to our ability to cure several cancers and other diseases, the genetic background of ATL leukemogenesis is not yet fully understood. Thus, it is an urgent request to clarify the molecular mechanism of ATL development.

Constitutive activation of nuclear factor- κ B (NF- κ B) is observed in the ATL cell lines and primary isolated tumor cells from ATL patients, although the viral oncoprotein Tax, a powerful activator of NF- κ B, is not expressed in these malignant cells

(Hironaka et al., 2004; Watanabe et al., 2005). NF- κ B activation aberrantly contributes to cell propagation and anti-apoptotic responses in ATL and other cancers (Prasad et al., 2010). In our previous study, inhibition of NF- κ B activity with a specific inhibitor, DHMEQ, drastically impaired the levels of ATL cell growth and resistance to apoptosis (Watanabe et al., 2005), suggesting that the molecular background of aberrant NF- κ B activation may give us potential therapeutic targets. A recent report provided a new readout that NF- κ B-inducing kinase (NIK) has a causal role in tumor progression and the aggressive phenotypes of various cancers, including ATL (Saitoh et al., 2008). NIK plays a pivotal role in the noncanonical (alternative) NF- κ B pathway as a crucial kinase in receptor-initiating signaling, including signaling from CD40, LTBR, and BAFFR.

Significance

Here, we propose a molecular perspective of the onset of oncogenic signaling. NIK overexpression is a major driving force for constitutive NF- κ B activation in various types of cancers. Using ATL cells as a model of NF- κ B-addiction, we identified miR-31 as a suppressor of NIK that is completely silenced in ATL cells. Furthermore, an oncogenic function of a subset of Polycomb is implicated in NF- κ B signaling via miRNA regulation. This study introduces a fundamental link between the Polycomb-mediated epigenetic regulation and the NF- κ B signaling, allowing us to attribute the constitutive activation of NF- κ B to epigenetic reprogramming.

Several studies have recently implicated another functional significance of NIK protein in epithelial cell proliferation, inflammatory response, and oncogenic signaling (for review, see Thu and Richmond, 2010). Although the expression level of NIK is strictly maintained by proteasomal degradation in normal cells (Liao et al., 2004), increased level of NIK transcript are observed in some cancers, causing inappropriate accumulation of NIK protein without stimuli (Annunziata et al., 2007; Saitoh et al., 2008). Overexpression of NIK leads to aberrant phenotypes in several cell types; however, little is known about the abnormal accumulation of NIK in malignant cells.

Recent advances have led to deeper understanding of a new aspect of posttranscriptional gene regulation, i.e., regulation by a class of noncoding RNAs. MicroRNAs (miRNAs) are functional RNAs with 18–25 nt in length that contribute to a class of cellular functions by negatively controlling targeted gene expression via base-pairing to 3' untranslated region (3' UTR). A single miRNA regulates the expression of multiple genes, and the functions of miRNAs therefore need to be orchestrated for cellular homeostasis (Ventura and Jacks, 2009). In the context of cancer pathology, many studies have provided evidences that miRNAs can act as either oncogenes or tumor suppressors. Although the relationship between miRNA deregulation and oncogenes has been clarified in several cancer cells, there has been no integrated analysis of gene expression in ATL. Since miRNAs have important functions in living cells, miRNA expression needs to be tightly regulated. Our knowledge about the regulatory mechanisms of miRNA expression is very inadequate because research effort has focused mainly on the role of miRNAs, which remains one of the most intriguing questions. miRNA regulation involves multiple steps. miRNA maturation has been identified as an important step, and its deregulation leads to progression and development of cancer (Davis et al., 2008; Trabucchi et al., 2009). Genetic deletion in cancer cells has also been reported to account for specific miRNA defect (Varambally et al., 2008). In addition, miRNA expression seems to be epigenetically programmed. DNA methylation and histone modification are strong candidates for miRNA regulation and their abnormalities, therefore, have causal roles in cancer initiation, development, and progression. In particular, Polycomb group proteins have central functions in cellular development and regeneration by controlling histone methylation, especially at histone H3 Lys27 (H3K27), which induces chromatin compaction (Simon and Kingston, 2009). Recent studies have revealed that the amount of Polycomb family is closely associated with cancer phenotypes and malignancy in breast cancer, prostate cancer, and other neoplasms (Sparmann and van Lohuizen, 2006). However, the substantial status of Polycomb family and their epigenetic impact in ATL cells have not been documented. Furthermore, the general roles of Polycomb proteins in miRNA regulation are mostly unknown. As described above, since miRNAs are multifunctional molecules in gene regulation, it is of pivotal importance to clarify the miRNAs functions and their regulatory circuit in order to formulate therapeutic strategies.

In the present study, we first performed global miRNA and mRNA profilings of the ATL cells derived from patients to precisely define the significance of miRNA expressions and functions.

RESULTS

miRNA Expression Signature in Primary ATL Cells

To characterize the miRNA expression signature in the primary ATL cells, we first performed a miRNA expression microarray analysis. For results with physiological significance, we used total RNA prepared from clinical ATL samples ($n = 40$, Table S1 available online) and control CD4+ T cells from healthy donors ($n = 22$) aged 50–70 years. A strict threshold ($p < 1 \times 10^{-5}$) and two-dimensional hierarchical clustering analysis revealed 61 miRNAs that showed significantly altered levels of expression in ATL cells compared with those of control CD4+ T cells (Figure 1A). It is noteworthy that 59 miRNAs out of 61 (96.7%) showed decreased expression in the primary ATL cells. Among them, we identified miR-31 as one of the most profoundly repressed miRNAs in all ATL individuals (fold change, 0.00403; Figure 1B). miR-31 was recently reported as a tumor suppressor and/or metastasis-associated miRNA in metastatic breast cancer. However, the biological functions of miR-31 in lymphocytes have not been studied. We therefore focused on the biological significance and regulatory mechanisms of miR-31 expression in T cells as well as in solid cancers.

miR-31 Negatively Regulates NF- κ B Signaling via NIK Expression

To study the functional significance of miR-31 loss, we attempted to identify the target genes of miR-31 using four computational algorithms. We also performed gene expression microarray analysis of the primary ATL cells ($n = 52$, Table S1) and normal CD4+ T cells ($n = 21$) in order to detect aberrations in gene expression. Selected putative target genes are known to be involved in cell-cycle regulation and T cell development (Table S2). To experimentally identify the target genes, we performed reporter-based screens as described below. Luciferase-3' UTR reporter assays demonstrated a remarkable negative effect against upstream gene expression by the *MAP3K14* 3' UTR sequence (Figure S1B), which is consistent with an initial cloning report (Malinin et al., 1997). *MAP3K14*, also known as NIK, has a central role in noncanonical NF- κ B signaling by phosphorylation of IKK α . A previous report (Saitoh et al., 2008) and the present results (Table S2) show that NIK is overexpressed in ATL cells, leading to constitutive NF- κ B activation. As shown in Figure 2A, treatment with a miR-31 inhibitor increased *NIK* 3' UTR reporter activity, suggesting the involvement of endogenous miR-31 in NIK downregulation. A computational search predicted one site each of miR-31 and miR-31 antisense (miR-31*) binding sites in the *NIK* 3' UTR (Figure 2B). To identify the regulatory sequence in 3' UTR of *NIK*, we established additional two reporters with mutated sequence in each potential seed region (Figure 2C; Figure S1C). Mutant 1, which contains mutated sequence in the miR-31 seed region, partially canceled the negative effect of endogenous miR-31 (Figure S1D) and prevented the effect of Anti-miR-31 treatment (Figure 2D). On the contrary, our results suggest that miR-31* does not participate in NIK regulation. miR-31-mediated reporter regulation was also observed in T cell lines (Figure S1E). To confirm the results, we repeated the experiment to examine whether miR-31 could inhibit NIK expression through seed sequence. We made expression plasmid vectors carrying NIK, NIK-3'




Figure 1. Global Profiling of Cellular miRNA on Primary ATL Cells

(A) Two-dimensional hierarchical clustering analysis and Pearson correlation as similarity measure on the miRNAs expressed at significantly different levels between the ATL (n = 40) and the control (n = 22) groups. Sixty-one miRNAs were identified ($p < 1 \times 10^{-5}$) and by filtering on more than 5-fold changes. A vertical branch shows the expression pattern of the selected miRNAs in each individual.

(B) Fold changes in the 61 miRNAs between ATL and Normal ($p < 10^{-5}$, fold change >5-fold). Selected miRNAs are arranged according to p values. See also Table S1.

UTRWT, or NIK-3' UTRMu1 and tested their expressions in 293T cells. Results demonstrated that expression of NIK-3' UTRWT was inhibited by simultaneous introduction of miR-31 (Figure 2E). miR-31 inhibition inversely rescued the NIK level, revealing that the cellular miR-31 level negatively affected that of the NIK protein through its 3' UTR sequence. These lines of evidence collectively demonstrated that miR-31 recognizes and regulates NIK mRNA through specific binding to its 3' UTR.

Transient introduction of the miR-31 precursor in TL-Om1 cells, which were established from an ATL patient, resulted in downregulation of NIK at the mRNA and protein levels, associated with downregulation of the phospho-IKKα/β level and NF-κB activity (Figures S1F and S1G). In contrast, miR-31 inhibition resulted in accumulation of NIK mRNA and protein in HeLa cells (Figure 2F). Manipulation of the miR-31 level clearly indicated that the miR-31 level negatively correlates with cellular




Figure 2. miR-31 is a Negative Regulator of NF- κ B Pathway by Inhibiting NIK Expression

(A) Reporter-based miR-31's target gene screening. A series of 3' UTR-luciferase reporters was transfected in HeLa cells together with or without miR-31 specific inhibitory RNA (Anti-miR-31) or control RNA (Anti-control). Relative values of Dual-luciferase assay are presented. "Non-target" represents reporter without any 3' UTR. "di-miR-31" reporter contains two perfect match sequences. The data are presented as mean \pm SD of three independent experiments.

(B) Schematic of miR-31 target sites in the NIK 3' UTR.

(C) Mutation-induced reporters. Red box stands for mutated target region (see Figure S1C).

(D) miR-31 negatively regulates NIK 3' UTR analyzed by reporter assay (n = 4, mean \pm SD). Luciferase activities of reporter series were tested in a presence or absence of miR-31 inhibitor.

(E) FLAG-tagged NIK protein is negatively regulated through its 3' UTR and miR-31 binding. Plasmid vectors and miR-31 precursor or miR-31 inhibitor are cotransfected in 293T cells. Western blots showed levels of NIK and endogenous p52. Asterisk indicates nonspecific bands.

(F) NIK mRNA (left) and protein (right) levels in HeLa cells measured by quantitative RT-PCR (n = 3, mean \pm SD) and western blotting, respectively. Treatment of miR-31 inhibitor resulted in NIK accumulation. Result of densitometry is shown in the bottom panel. Asterisk indicates nonspecific bands.

(G) Cellular NF- κ B activity in HeLa cells (n = 5, mean \pm SD) in a presence or absence of miR-31 precursor or inhibitor.

NF- κ B activity (Figure 2G). Furthermore, enforced miR-31 expression in B cells attenuated both BAFF and CD40L-mediated NIK accumulation and the subsequent NF- κ B signaling (Figures 2H–2K). Consistent with previous reports (Ramakrishnan et al., 2004; Zarnegar et al., 2008b), we also found decreased levels of I κ B α phosphorylation. On the other hand, TNF- α -triggered canonical NF- κ B activation was not affected by miR-31 in Jurkat cells (Figures S1H–S1K). These results collectively show that miR-31 inhibits the basal and receptor-initiated activities of noncanonical NF- κ B pathway. With genetic evidence and an experimental approach, we further demonstrated that the function of miR-31 is well conserved among several classes of species (Figures S1L–S1O). Taking together all these results, miR-31, which is almost completely absent in primary ATL cells, appears to play a critical role in negative regulation of the NF- κ B pathway by manipulating the expression of NIK.

miR-31 Suppresses ATL Cell Growth and Promotes Apoptosis by Inhibiting NF- κ B

Although it was documented that abnormal NIK accumulation in ATL cells acts as a constitutive activator of the NF- κ B pathway, the mechanism underlying overproduction of NIK remains to be elucidated. The results described in the previous section indicated that the amount of miR-31 is linked to the level of NIK, and we therefore speculated that downregulation of miR-31 expression is at least partially responsible for the constitutive activation of NF- κ B in ATL cells. Quantitative RT-PCR revealed that *NIK* mRNA levels were negatively correlated with miR-31 levels in primary ATL cell samples (Figure 3A). To investigate the functional roles of NIK and miR-31, we established TL-Om1 cells stably expressing the miR-31 or NIK specific shRNA (shNIK) by retroviral vectors. RT-PCR and western blots showed that expression of miR-31 or shNIK reduced NIK at mRNA and protein levels as well as the levels of phospho-IKK α / β , p52, and I κ B α (Figures 3B and 3C). Decreased levels of nuclear RelA and RelB are considered to represent repressed activities of the canonical and noncanonical NF- κ B pathways, respectively (Figure 3D). EMSA and NF- κ B reporter assays also revealed the repressive function of miR-31 and shNIK on the NF- κ B activity (Figures 3E and 3F; Figures S2A, S2B, S5B, and S5C). Re-expression of NIK led to NF- κ B activation that was inhibited by miR-31, suggesting a reciprocal relationship between the level of miR-31 and that of NIK.

We and others previously showed that constitutive NF- κ B activation is a strong driver of ATL proliferation and prosurvival properties. Here, we examined the effects of miR-31 loss on ATL cell growth. We found that TL-Om1 cells expressing miR-31 or shNIK showed a significant attenuation of cell proliferation compared with control cells. In addition, serum starvation experiments showed greater sensitivity to induced cell death in NIK-repressed cells (Figure 3G). miR-31 expression showed the same phenotypic results in other ATL-derived cell lines

(Figures S2C, S2D, and S5E). Jurkat cells do not have significant basal activity of NF- κ B, and showed no significant difference in cell growth with or without induced expression of miR-31 (Figure S2E).

Next, we hypothesized that miR-31-mediated NF- κ B modulation may affect cellular apoptosis, because numerous studies have demonstrated that NF- κ B activation is a strong antiapoptotic factor in ATL and other cancer cells. We found that repression of NIK by miR-31 or shNIK resulted in downregulation of a subset of genes involved in resistance to apoptosis such as BCL-XL, XIAP, and FLIP (Figure 3H), suggesting that miR-31 has a role in proapoptosis through inhibition of NF- κ B activity. To assess the biological function of miR-31 in apoptosis signals, we utilized a lentivirus gene transfer system for cell lines and freshly isolated tumor cells. The lentivirus vector is competent to infect nondividing cells and the infected cells can be monitored by the fluorescence of Venus. We found that lentivirus-mediated miR-31 expression promoted basal and Fas-directed apoptosis in TL-Om1 cells (Figure 3I). Venus-negative population showed no significant changes, demonstrating the specificity of miR-31 activity. To confirm the relationship among miR-31, NIK, and NF- κ B signaling, we also prepared another retroviral vector encoding NIK without its 3' UTR sequence. As results, re-expression of NIK reversed the miR-31-mediated apoptosis. In addition, miR-31 expression led to caspase 3 activation (Figure 3J). Collectively, these findings indicate that miR-31 mediates apoptosis through repression of NIK in ATL cell lines.

Tumor cells from ATL patients primarily represent the malignant characteristics. In fact, miR-31 loss is found from patient samples (Figures 1 and 3A). To demonstrate the responsibility of miR-31 for tumor cell survival, we tested whether lentivirus-mediated miR-31 expression has a killing effect against tumor cells. After establishment of lentivirus infection, the apoptotic cells were determined by flow cytometry. The results revealed that expression of miR-31 facilitated tumor cell death. Since NIK repression by shRNA lentivirus also showed a strong killing effect, NIK and NF- κ B activity are suggested as crucial players for survival in ATL tumor cells (Figure 3K). Strong toxicities were not observed in normal resting lymphocytes that express low levels of NIK. Taken together, these lines of experimental evidence, including data from cell lines and primary ATL cells, definitively support two notions that (1) miR-31 acts as a tumor suppressor in T cells, and (2) NIK-regulated NF- κ B has pivotal importance in cancer cell survival.

Loss of miR-31 Occurs in T Cells with Genetic and Epigenetic Abnormalities

The results described above together with previous publications indicate that regulation of miR-31 expression has profound impacts on multiple functions in human tumors as well as in normal cells. However, little is known about the regulatory mechanism of miR-31 expression. The human gene that encodes miR-31, *hsa-miR-31*, is located at 9p21.3, which is

(H–K) miR-31 attenuates signal-dependent NF- κ B activation in B cells. (H and I) BJAB cells expressing miR-31 or control RNA were treated with BAFF (0.2 μ g/ml) or CD40L (0.5 μ g/ml) for indicated time periods. The protein levels of NIK, phospho-IKK α / β , p100/p52 (arrowheads indicate active p52), and phospho-I κ B α were shown. Actin was detected as control. (J) NF- κ B activity ($n = 5$, mean \pm SD) evaluated by NF- κ B-luciferase reporter assay at 24 hr after cytokine treatments. (K) NF- κ B-dependent gene expressions were inhibited by miR-31 ($n = 3$, mean \pm SD). See also Table S2 and Figure S1.




Figure 3. Loss of miR-31 is responsible for constitutive NF- κ B activation, abnormal cell growth, and resistance to apoptotic cell death in ATL cells

(A) Expression levels of miR-31 and *NIK* in individual ATL patients and normal controls using data set obtained by quantitative RT-PCR. Pearson's correlation coefficient within ATL samples was described in the graph.

adjacent to clusters of the *CDKN2* and *IFNA* families, and is a well-known hotspot of genomic loss in several types of human cancers. We performed genome-wide scans of genetic lesions in 168 ATL samples and demonstrated that 21 ATL cases (12.5%) had genomic deletion of 9p21.3 containing the *hsa-miR-31* coding region (Figure 4A; Figure S3A). All of these cases also have genomic defect in *CDKN2A* region. A major proportion of ATL cases that are without genetic deletion and somatic mutation in the *hsa-miR-31* region showed remarkable loss of miR-31 expression (Figure 4B). Detailed expression profiling revealed drastic downregulation of *Pri-miR-31* transcription in the primary ATL cells (Figure 4C). There was a strong correlation between the levels of mature miR-31 and primary transcript ($r = 0.9414$, $p = 5.45 \times 10^{-8}$). *hsa-miR-31* is located in intronic region of *LOC554202* gene. However, *LOC554202* mRNA levels were very low in primary T cells and there was no significant difference between ATL and normal cells, strongly suggesting that loss of miR-31 expression is due to specific transcriptional suppression in ATL cells. Using computational analysis, we identified a putative TATA box and transcriptional start site (TSS) 2500 bp upstream of the miR-31 coding region (Figure 4D). Although no CpG islands were found in this region, we unexpectedly discovered an assembly of YY1-binding motifs upstream of the miR-31 region in human and mouse (Figure 4D; Figure S3C). YY1 is a pivotal transcription factor and a recruiter of the Polycomb repressive complex (PRC) (Simon and Kingston, 2009). Convergence of the YY1 binding sequence, especially the repressive motif (Figure S3D), seems to be evolutionarily conserved, suggesting that YY1 is important in the regulation of miR-31 transcription. We further performed chromatin immunoprecipitation (ChIP) to evaluate repressive histone hallmarks, including di- and trimethylated H3K9 (H3K9me2 and H3K9me3) and trimethylated H3K27 (H3K27me3). The results showed higher levels of methylation at H3K9 and H3K27 in a broad area containing the miR-31 coding region (Figure 4E). As shown in Figures S3E–S3G, there was an inverse correlation between the levels of miR-31 expression and repressive histone methylation. These data allowed us to hypothesize that histone methylation, especially that of Polycomb family-dependent H3K27me3, may contribute to miR-31 repression. To confirm our hypothesis, we performed a YY1 knockdown experiment using a specific shRNA (Figures 4F–4I). As expected, knockdown of YY1 led to an increase in the levels of *Pri-miR-31* and mature miR-31 (Figures 4F and 4G). Furthermore, ChIP assays showed that

YY1 occupied the miR-31 region, especially in the upstream region of TSS, where there is an array of YY1 binding sites (Figures 4D and 4H). The results also demonstrated that decreased occupancy of YY1 and concomitant derecruitment of EZH2, a key component of PRC2, were induced by YY1 knockdown, indicating involvement of EZH2 in the repressive complex recruited to the miR-31 region (Figures 4H and 4I; Figure S3H). These results collectively suggest that YY1 regulates PRC2 localization and initiates miR-31 suppression. Indeed, we found significant escalation of methylated histone H3K9 and H3K27 at the miR-31 locus of peripheral blood lymphocytes of ATL patients (Figure 4J), indicating that aberrant abundance of suppressive histone methylation may be responsible for the loss of miR-31 in the primary ATL cells.

Overexpression of PRC2 Components Leads to miR-31 Repression

Given that Polycomb-mediated repressiveness affects miR-31 level, our findings imply that the amount of EZH2 is related to miR-31 expression (Figure 4I; Figures S3G and S4A). We found a significantly upregulated expression of PRC2 components, especially EZH2 and SUZ12, in the primary ATL cells (Figures 5A and 5B; Table S3). Quantitative RT-PCR revealed that miR-31 levels inversely correlated with both *EZH2* and *SUZ12*, respectively (Figure 5C). miR-101 and miR-26a, which are putative negative regulators of EZH2, seem to be associated with this relationship in ATL cells (Figures S4B–S4E). To further confirm our hypothetical mechanism linking the epigenetic machinery and miR-31 expression, we performed a “loss-of-PRC2-function” assay. Retroviral delivery of shSUZ12 and shEZH2 in the ATL cell lines resulted in a great increase in the levels of *Pri-miR-31* and its mature form (Figure 5D; Figure S4F). Knockdown of PRC2 induced histone demethylation at H3K27 in the miR-31 region, which is concomitant with the decrease in H3K9me3 levels, EZH2 occupancy, and HDAC1 recruitment (Figure 5E), suggesting that this multimeric complex leads to a completely closed chromatin architecture as a result of histone modifications in the miR-31 genomic region.

To further examine whether the proposed mechanism holds true in other human cancers, we analyzed a couple of carcinoma cell lines, including HeLa cells and nonmetastatic and metastatic breast carcinoma cell lines, MCF7 and MDA-MB-453 cells, respectively. qRT-PCR revealed that expression of *EZH2* and *SUZ12* inversely correlated with miR-31 levels (Figure S4G).

(B) miR-31 restoration by retroviral vector inhibits *NIK* RNA accumulation in TL-Om1 cells. The results of *NIK* and mature miR-31 quantifications are shown ($n = 3$, mean \pm SD).

(C) miR-31 or shNIK expression downregulates *NIK* protein expression and inhibits downstream pathway of noncanonical NF- κ B in TL-Om1 cells.

(D) Reduced nuclear translocation of RelA and RelB proteins in miR-31- or shNIK-expressing TL-Om1 cells.

(E) miR-31-dependent downregulation of NF- κ B activity in TL-Om1 cells examined by EMSA.

(F) NF- κ B-luciferase reporter assays ($n = 5$, mean \pm SD). FLAG-*NIK* plasmid was transiently introduced 48 hr prior to the assay.


(G) miR-31 level is relevant to proliferation of ATL cells. Cell proliferation curve of TL-Om1 cells were evaluated in two FCS conditions ($n = 3$, mean \pm SD).

(H) Apoptosis-related gene expression in TL-Om1 cells analyzed by qRT-PCR ($n = 3$, mean \pm SD) and western blots.

(I) Lentivirus-mediated *NIK* depletion promotes basal and Fas antibody-mediated apoptosis. Venus-positive population represented lentivirus-infected cells. Apoptotic cells were determined by PE-Annexin V / 7-AAD stainings ($n = 4$). Representative FACS analyses are shown.

(J) miR-31 activates Caspase 3/7 determined by FACS ($n = 3$).

(K) miR-31 expression and *NIK* depletion induce tumor cell death. Primary tumor cell from ATL patient and healthy CD3+ T cells were infected with lentivirus and analyzed by FACS. The apoptotic cells were defined by sequential gating beginning with FSC-SSC to select intact lymphocytes, subgating on the Venus-positive population, and calculating the PE-Annexin V and 7-AAD profilings. Representative result is shown and summarized data are presented in Figure 6J. See also Figure S2.



ChIP assays detected higher levels of trimethylated H3K27 and EZH2 occupancy in cells showing lower expression levels of miR-31 (Figure S4H). Furthermore, knockdown of EZH2 or SUZ12 restored miR-31 transcription in MDA-MB-453 and MCF7 cells (Figures 5F and 5G; Figure S4K, respectively), which are consistent with the results obtained with ATL cells. These results indicate a link between Polycomb-mediated epigenetic regulation and miR-31 transcription in ATL and breast cancer cell lines.

Polycomb Group Regulates NF- κ B Pathway by Controlling miR-31 Expression

Based on our findings, we considered an aspect of the biological communication between epigenetic silencing and the NF- κ B pathway through miR-31 regulation. The microarray data sets showed positive correlations between PRC2 components and miR-31 target gene, *NIK* expression (Figure 6A). The results also suggested that these factors tend to show higher levels in the aggressive subtype (acute type) than in the indolent subtypes (chronic and smoldering types), implying that these genes may play important roles in the clinical phenotype and prognosis of ATL. To examine this notion, we performed PRC2 knockdown in ATL cell lines. Western blots of these cells demonstrated decreased levels of NIK, p52, and phospho-I κ B α (Figure 6B; Figure S5A), suggesting suppression of both canonical and noncanonical NF- κ B cascade and activity (Figure 6C; Figures S5B and S5C). These results are consistent with those of miR-31 overexpression (Figures 3C–3F). Then, we tested whether exogenous manipulation of miR-31 could restore the effect of PRC2 loss. Anti-miR-31 treatment rescued impaired NF- κ B activity in PRC2-disrupted cells (Figure 6D). On the other hand, overexpression of EZH2 induced NF- κ B activation, which was partially canceled by the introduction of miR-31 precursor (Figure 6E; Figure S5D). These results suggest that Polycomb-mediated miR-31 suppression leads to NF- κ B activation. Indeed, knockdown of the PRC2 complex led to reduced levels of cell proliferation and greater sensitivity to serum deprivation in ATL cells (Figure 6F; Figure S5E). In addition, PRC2 disruption showed a reduction in cell migration (Figure S5F).

To gain further insight into this general network, we studied the functions of miR-31 and the PRC2 complex in breast cancer cell lines. NF- κ B activity was downregulated by knockdown of

PRC2 components in MDA-MB-453 cells (Figure 6G; Figures S5G and S5H), although no significant differences were observed in cell proliferation (data not shown). Repression of NF- κ B activity induced by knockdown of PRC2 components was partially restored by treatment with a miR-31 inhibitor, suggesting that PRC2 knockdown-mediated relief of NF- κ B repression is at least a part of the result of the miR-31 induction. In addition, knockdown of PRC2 components resulted in a reduced level of receptor-initiated accumulation of NIK in B cells (Figure 6H). Our findings indicate a common molecular mechanism comprising Polycomb-mediated epigenetic regulation, miR-31 expression and the NF- κ B signaling pathway.

Regulation of NF- κ B by Polycomb family may in turn control the cellular apoptosis responses. We found that lentivirus-mediated EZH2 knockdown led to increased apoptotic sensitivity in TL-Om1 cells (Figure 6I). Additional expression of NIK inhibited the cell death induced by EZH2 knockdown, suggesting the reciprocal relationship between Polycomb and NF- κ B cascades. By using primary tumor cells from patient, we tested the killing effect induced by miR-31, NIK knockdown, and EZH2 knockdown (Figure 6J; Figures S5I and S5J). All tested samples showed strong death response, demonstrating that survival of ATL cells was closely associated with miR-31, NIK, and EZH2, all of which show deregulated expression in ATL cells.

By qRT-PCR we finally examined the expression levels of some genes involved in the noncanonical NF- κ B pathway. As shown in Figure 6K, the results clearly demonstrated higher expression levels of positive regulators such as *NIK*, *CD40*, and *LTBR*, and lower expression levels of the negative regulators such as *BIRC2/3* (*cIAP1/2*), which are involved in proteasomal degradation of NIK (Zarnegar et al., 2008a). These observations are in line with a previous report on Multiple Myeloma cells (Annunziata et al., 2007). In addition to these data, we obtained convincing evidence for a molecular aspect of NIK accumulation in ATL cells. Polycomb-dependent epigenetic gene silencing may be associated with miR-31 loss, followed by NF- κ B activation and other signaling pathways (Figure 7).

DISCUSSION

Constitutive activation of NF- κ B contributes to abnormal proliferation and inhibition of apoptotic cell death in cancer cells,

Figure 4. Genetic and Epigenetic Abnormalities Cause miR-31 Loss in ATL Cells

(A) Genomic loss of chromosome 9p21.3 in primary ATL cells. Copy number analyses revealed tumor-associated deletion of miR-31 region (21/168) and *CDKN2* region (46/168). Recurrent genetic changes are depicted by horizontal lines based on CNAG output of the SNP array analysis.

(B) miR-31 expression in various sample sets. Expression levels were evaluated by real-time PCR.

Loss, samples with genomic loss of the miR-31 region; (–) samples without genomic loss of the miR-31 region.


(C) PCR-based miR-31 quantifications in primary ATL samples. ATL samples without genomic loss in miR-31 region ($n = 9$, Figure S3B), and normal CD4+ T cells ($n = 7$) were tested. p values (ATL versus normal) are shown.

(D) YY1 binding motif cluster around transcriptional start site (TSS) of miR-31 region. Arrows represent positions of the motifs. Regions of PCR amplification for ChIP assay are shown.

(E) Repression-associated histone methylation in miR-31 region determined by ChIP assay ($n = 3$, mean \pm SD). The results of relative enrichment against input control are presented and distance from miR-31 TSS is described. *MYT1* and *GAPDH* promoters are as positive or negative controls, respectively.

(F–I) YY1-dependent EZH2 occupancy in miR-31 locus. (F) YY1 knockdown in TL-Om1 cells. qRT-PCR (left, $n = 3$, mean \pm SD) and western blotting (right) showed decreased YY1 level. (G) YY1 knockdown led to both primary and mature miR-31 restoration in TL-Om1 cells ($n = 3$, mean \pm SD). (H) YY1 occupancy in miR-31 region analyzed by ChIP ($n = 3$, mean \pm SD). YY1 occupancy in miR-31 locus was reduced by YY1 knockdown. (I) EZH2 occupancy in miR-31 region analyzed by ChIP ($n = 3$, mean \pm SD). YY1 knockdown inhibited EZH2 recruitment in miR-31 region.

(J) Aberrant accumulation of repression-associated histone methylations widely in miR-31 region of primary ATL cells. PBMCs freshly isolated from ATL patients ($n = 6$) were analyzed by ChIP assay. PBMC from healthy adults were used for normal controls. See also Figure S3.



including ATL, diffuse large B cell lymphoma (DLBCL), Hodgkin lymphoma, breast cancer, prostate cancer and others (Prasad et al., 2010). NF- κ B is also essential for various cell functions, including inflammation, innate immunity, and lymphocytic development (Hayden and Ghosh, 2008). Identification of NF- κ B determinants will lead to marked progress in understanding molecular pathology.

Our global analyses demonstrated an interesting miRNA expression signature as well as an aberrant mRNA expression profile, which may be associated with leukemogenesis in the primary ATL cells (Figures 1 and 6A). We revealed downregulation of tumor-suppressive miRNA including Let-7 family, miR-125b, and miR-146b, which can contribute to aberrant tumor cell signaling. Recent studies have suggested unique expression profiles of miRNAs in ATL (Yeung et al., 2008; Bellon et al., 2009), but loss of miR-31 has not been focused. Cellular amount of miRNAs may be susceptible to various environments such as transcriptional activity, maturation processing, and also epigenetic regulation. The end results appear to be affected by methodology employed and conditions and types of samples used. Our integrated expression profiling of primary ATL cells are based on a significantly larger number of samples and fruitfully provides intriguing information that may be useful in improving the understanding of T cell biology as well as in the identification of biomarkers for diagnosis.

Pleiotropy of miR-31 was first reported by Valastyan et al. (2009). The authors elegantly demonstrated the function of miR-31 in vivo and also identified several target genes that contribute to cell migration and invasiveness. In the present study, we focused on the functional significance of miR-31 in the regulation of NF- κ B signaling that contributes to tumor cell survival.

Overexpression of NIK acts as an oncogenic driver in various cancers. In the present study, NIK was identified as a miR-31 target based on several lines of evidence. First, luciferase-3' UTR reporter assay showed that *NIK* 3' UTR sequence has a role for negative regulation (Figure S1B). By combining a specific inhibitor and mutations in miR-31-binding site, we demonstrated that miR-31 recognizes and negatively regulates the *NIK* 3' UTR (Figures 2A and 2D). Second, by introducing a miR-31 precursor or inhibitor, we showed that amount of miR-31 inversely correlates with levels of NIK expression and downstream signaling (Figures 2E–2K). Third, genetic evidence indicated strong base pairing and biological conservation (Bartel, 2009) (Figures S1L–S1O). Our experimental approach illustrated that mmu-miR-31 regulates mouse *Map3k14* gene. Fourth, individual assessments using gene expression data

clearly revealed an inverse correlation between the expression levels of miR-31 and *NIK* (Figure 3A). Collectively, we provide definitive evidence for the notion that miR-31 negatively regulates NIK expression and activity.

It is well known that the NIK level directly regulates NF- κ B activity in various cell types (Thu and Richmond, 2010). We experimentally showed that miR-31 regulates noncanonical NF- κ B activation stimulated by BAFF and CD40L, both of which are major B cell activating cytokines. Since signals from receptors are essential for the development and activity of B cells, the negative role of miR-31 in cytokines-induced NIK accumulation appears to be widely important in the noncanonical regulation of NF- κ B in B cells and other cell types (Figures 2H–2K). Again, our findings revealed the role of NIK in the regulation of canonical NF- κ B pathway. Strict regulation of NIK appears to be closely associated with the fate of lymphocytes.

The level of miR-31 was drastically suppressed in all tested primary ATL cells, and its magnitude is greater than that which has been reported in other cancers. Our results demonstrated a profound downregulation of miR-31 (fold change, 0.00403; Figure 1B) in all ATL cases, suggesting that miR-31 loss is a prerequisite for ATL development. Restoration of miR-31-repressed NF- κ B activity in ATL cells, resulting in impairment of the proliferative index and apoptosis resistance (Figure 3). Furthermore, our results demonstrate that inhibition of NF- κ B promotes tumor cell death in cell lines and also primary tumor cells from ATL patients (Figures 3 and 6), which are consistent with our previous observation (Watanabe et al., 2005). Since it is highly possible that miR-31 and relevant factors are pivotal in cancers, their expressions would have a great importance in view of biomarkers for the aberrant signaling and clinical outcomes.

By studying clinical samples and in vitro and ex vivo models, we obtained several biologically interesting results. First, we identified the Polycomb protein complex as a strong suppressor of miR-31. Generally, the Polycomb group constitutes a multimeric complex that negatively controls a large number of genes involved in cellular development, reproduction, and stemness (Sparmann and van Lohuizen et al., 2006). However, the key molecules involved in cancer development, progression, and prognosis are not yet fully understood. In breast and prostate cancers, oncogenic functions of EZH2 and NF- κ B activation were reported independently (Kleer et al., 2003; Varambally et al., 2002; Suh and Rabson, 2004). Interestingly, these tumors show low miR-31 levels (Valastyan et al., 2009; Schaefer et al., 2010). Recently, Min et al. (2010) reported that EZH2 activates NF- κ B by silencing the *DAB2IP* gene in prostate cancer cells.

Figure 5. Amount of PRC2 Components Epigenetically Links to miR-31 Expression in T Cells and Epithelial Cells

(A) Overexpression of PRC2 components in primary ATL cells measured by qRT-PCR (ATL, n = 28; normal, n = 7; mean \pm SD). These results were supported by the data of gene expression microarray (Table S3).
 (B) Escalation of EZH2 protein and trimethylated H3K27 levels in primary ATL cells illustrated by immunocytostaining (n = 4, a representative result is shown). Resting T cells were as normal control. Scale bars = 20 μ m.
 (C) Statistical correlation among the levels of miR-31, *EZH2*, and *SUZ12* in individual ATL samples. Correlation coefficients within ATL samples are shown in the graphs.
 (D and E) Loss of PRC2 function causes chromatin rearrangement and miR-31 upregulation. (D) TL-Om1 cells expressing shSUZ12, shEZH2, and scrambled RNA were established by retroviral vector. The levels of EZH2, SUZ12, *Pri-miR-31*, and mature miR-31 were measured by western blotting and qRT-PCR (n = 3, mean \pm SD). (E) Results of ChIP assays with indicated antibodies (n = 3, mean \pm SD). Amounts of immunoprecipitated DNA were analyzed by region-specific PCR. *GAPDH* promoter served as a region control.
 (F and G) Knockdown of Polycomb family proteins in MDA-MB-453 cells. (F) EZH2 and SUZ12 are shown by western blot. miR-31 level was examined by qRT-PCR (n = 3, mean \pm SD). (G) Histone methylation and EZH2 occupancy evaluated by ChIP assay (n = 3, mean \pm SD). See also Table S3 and Figure S4.




Figure 6. Epigenetic Change Driven by Polycomb Group Mediates NF- κ B Signaling through miR-31 Regulation

(A) Reciprocal relationship of mRNA expression between *NIK* and Polycomb group in primary samples. Pearson's correlation coefficients among ATL samples are shown.

(B) PRC2 knockdown negatively affects NF- κ B signaling in TL-Om1 cells. After establishment of PRC2 knockdown, the levels of *NIK* RNA ($n = 4$, mean \pm SD) and proteins of NIK, p52/p100, and phospho-I κ B α were examined.

(C) Downregulation of NF- κ B activity in PRC2-disrupted cells detected by EMSA.




Figure 7. Proposed Model for ATL and Other Tumor Cells
Polycomb repressive factors are linked to NIK-dependent NF-κB activation via miR-31 regulation.

In the present study, we found that the Polycomb group regulates miR-31 expression and that elevated expression of EZH2 leads to NF-κB activation via NIK-miR-31 regulation in ATL and breast cancer cells (Figure 6). We also showed that restoration of miR-31 partially impaired Polycomb-mediated NF-κB operation (Figures 6D, 6E, and 6G), suggesting that miR-31 is involved in this relationship. Furthermore, a connection between NIK and PRC2 was observed in B cells (Figure 6H). Polycomb group proteins are essential in lymphocyte development and activation (Su et al., 2003, 2005). Further, given the NF-κB is a pivotal transcription regulator in normal and oncogenic functions, practical participations of epigenetic regulators and miR-31 in NF-κB signaling will increase our understanding of the molecular mechanisms of T cell functions. For generalization of the molecular axis in other cancers and normal cells, further study will be needed.

Second, YY1 is a recruiter of PRC2 to the miR-31 region. In humans, the Polycomb response element (PRE) has not been precisely identified. A good candidate for a mammalian recruiter of PRC2 is YY1, the homolog of *D. melanogaster* PHO (Simon and Kingston, 2009). We found an assembly of the YY1 binding motif in the miR-31 locus and demonstrated that YY1 knockdown dislodged EZH2 in this region (Figure 4I), which supports previous findings (Carette et al., 2004). The detailed mechanism by which YY1 mediates recruitment of the Polycomb family may be important in the context of epigenetic regulation of orchestrated gene expression and T cell functions.

Third, Polycomb family proteins can control miRNA expression in an epigenetic fashion. The amount of PRC2 factors strongly influenced the degree of suppression of miR-31 expres-

sion (Figure 5). We speculate that, in addition to controlling the transcription, the Polycomb group can modulate translation via miRNA regulation. Furthermore, miR-101 and miR-26a are known to regulate EZH2 expression (Sander et al., 2008; Varambally et al., 2008), which is supported by our observation (Figure S4C). This signaling circuit will permit multiple gene regulation. Whereas genetic loss at the miR-31 locus is observed in some cases of ATL (Figure 4A), no genetic deletion in the miR-101-1 or miR-101-2 region was detected in ATL, which is not consistent with a previous finding in prostate cancer. Our results also suggested putative association between Let-7 family and EZH2 (Figure S4). Aberrant downregulations of these miRNAs in the primary ATL cells will be the next important questions to be addressed in efforts to improve understanding of the oncogenic signaling network.

By collaborative profiling of miRNA and mRNA expression, we identified a notable relationship between ATL subtypes and a gene cluster that contains miR-31, NIK, EZH2, and SUZ12. This finding suggests that an aberrant gene expression pattern correlates with the malignant phenotype, and this provides important clues about clinical manifestations and may help identify therapeutic targets against ATL cells (Figure 6A). Although HDAC inhibitors did not show effective responses (Figures S4I and S4J), emerging epigenetic drug such an EZH2 inhibitor (Fiskus et al., 2009) may pave a pathway leading to cures for various malignancies that involve constitutive activation of NF-κB.

In summary, we show that genetic and epigenetic loss of miR-31 is responsible for oncogenic NF-κB activation and malignant phenotypes in ATL. This provides evidence for the idea that miR-31 is an important tumor suppressor. An emerging pathway involving an epigenetic process, miR-31, and NF-κB will provide a conceptual advance in epigenetic reprogramming, inflammatory signaling, and oncogenic addiction.

EXPERIMENTAL PROCEDURES

Cell Lines and Primary ATL Cells

The primary peripheral blood mononuclear cells (PBMCs) from ATL patients and healthy volunteers used in the present work were a part of those collected with an informed consent as a collaborative project of the Joint Study on Prognostic Factors of ATL Development (JSPFAD). The project was approved by the Institute of Medical Sciences, the University of Tokyo (IMSUT) Human Genome Research Ethics Committee. Additional ATL clinical samples for copy number analysis were provided by Drs. Y. Yamada, Nagasaki University,

(D) NF-κB activity evaluated by reporter assays in the presence or absence of miR-31 inhibitor (n = 5, mean ± SD). Anti-miR-31 treatment partially rescued the NF-κB activity in PRC2 knockdowned TL-Om1 cells.
 (E) Overexpressed EZH2 activates NF-κB via miR-31. Jurkat cells were transfected with an EZH2 plasmid together with miR-31 precursor or control RNA (n = 5, mean ± SD).
 (F) PRC2 dysfunction changes TL-Om1 cell proliferation and response to serum starvation. Under conditions of 10% or 0.1% of FCS, cell growth curves were examined (n = 3, mean ± SD). PRC2 downregulation decreased cell growth with statistical significance.
 (G) NF-κB activity in PRC2-knockdowned MDA-MB-453 cells in the presence or absence of miR-31 inhibitor were examined (n = 5, mean ± SD).
 (H) PRC2 disruption inhibits BAFF-dependent NIK accumulation and IκBα phosphorylation in BJAB cells.
 (I) Apoptotic cell death induced by lentivirus-mediated EZH2 knockdown in TL-Om1. Venus-positive populations were analyzed by Annexin V/7-AAD stainings (n = 3) and representative of FACS data are shown.
 (J) Summary of primary tumor cell death. Lentivirus-based miR-31 expression, NIK knockdown, and EZH2 knockdown showed killing effects in six primary ATL samples. Statistical significances are shown in the graph. Results of FACS and qRT-PCR are shown in Figures S5I and S5J.
 (K) Expression levels of genes involved in noncanonical NF-κB pathway in primary ATL cells (ATL, n = 9; normal, n = 7; mean ± SD). Relative expression levels were tested by qRT-PCR (p < 0.05). See also Figure S5.

and K. Ohshima, Kurume University, where the projects were approved by the Research Ethics Committees of Nagasaki University and Kurume University, respectively. PBMC were isolated by Ficoll separation. ATL cells, primary lymphocytes, and all T cell lines were maintained in RPMI1640 supplemented with 10% of FCS and antibiotics. Clinical information of ATL samples is provided in Table S1.

Expression Analyses

Clinical samples for microarrays were collected by a collaborative study group, JSPFAD (Iwanaga et al., 2010). Gene expression microarray was used 4x44K Whole Human Genome Oligo Microarray (Agilent Technologies) and miRNA microarray was used Human miRNA microarray kit v2 (Agilent Technologies), respectively. Quantitative RT-PCR was performed with SYBRGreen (TAKARA). Mature miRNA assays were purchased from Applied Biosystems.

Copy Number Analyses

Genomic DNA from ATL patients was provided from the material bank of JSPFAD, Nagasaki University, and Kurume University, and was analyzed by Affymetrix GeneChip Human Mapping 250K Nsp Array (Affymetrix). Obtained data were analyzed by CNAG/AsCNAR program (Chen et al., 2008).

Oligonucleotides, Plasmids, and Retrovirus Vectors

All RT-PCR primers and oligonucleotides are described in Supplemental Experimental Procedures. miRNA precursor and inhibitor were from Applied Biosystems. Transfection of small RNA and other plasmid DNA were performed by Lipofectamine2000 (Invitrogen). For miRNA or shRNA expression, retroviral vectors (pSINsi-U6, TAKARA) were used.

3' UTR-Conjugated miR-31 Reporter Assay

HeLa cells were cotransfected with 3' UTR-inserted pMIR-REPORT firefly plasmid (Ambion), RSV-Renilla luciferase plasmid, and miRNA inhibitor. The cells were collected at 24 hr posttransfection, and Dual-luciferase reporter assay was performed (Promega).

Analysis of NF- κ B Pathway

NF- κ B activity was evaluated by EMSA and reporter assays as previously described (Horie et al., 2004). Antibodies for western blots are described in supplemental information. Cell proliferative assay was performed by Cell Counting Kit-8 (Dojindo).

Lentivirus Vectors and Apoptosis Analysis

A lentivirus vector (CS-H1-EVbSd) was provided from RIKEN, BRC, Japan. Lentivirus solution was produced by cotransfection with packaging plasmid (pCAG-HIVgp) and VSV-G- and Rev-expressing plasmid (pCMV-VSV-G-RSV-Rev) into 293FT cells. After infection of lentivirus, the apoptotic cell was evaluated by PE Annexin V / 7-AAD staining (BD Pharmingen) and analyzed by FACS Calibur (Becton, Dickinson). Collected data were analyzed by FlowJo software (Tree Star).

ChIP Assay

ChIP assay was previously described (Yamagishi et al., 2009). Briefly, cells were crosslinked with 1% of formaldehyde, sonicated, and subjected to chromatin-conjugated IP using specific antibodies. Precipitated DNA was purified and analyzed by real-time PCR with specific primers (see Supplemental Experimental Procedures).

Computational Prediction

To identify miR-31 target genes, we integrated the output results of multiple prediction programs; TargetScan, PicTar, miRanda, and PITA. RNAhybrid was for secondary structure of miRNA-3' UTR hybrid. TSSG program was for TATA box and TSS predictions. DNA methylation site was predicted by CpG island Searcher.

Statistical Analyses

Data were analyzed as follows: (1) Welch's t test for Gene Expression Microarray (p value cutoff at 10^{-6}) and miRNA Microarray (p value cutoff at 10^{-5}); (2) Pearson's correlation for two-dimensional hierarchical clustering analysis

and individual assessment of microarray data sets; (3) two-tailed paired Student's t test with $p < 0.05$ considered statistically significant for in vitro cell lines and primary cells experiments, including luciferase assay, RT-PCR, ChIP assay, cell growth assay, and migration assay. Data are presented as mean \pm SD.

ACCESSION NUMBERS

Coordinates have been deposited in Gene Expression Omnibus database with accession numbers, GSE31629 (miRNA microarray), GSE33615 (gene expression microarray), and GSE33602 (copy number analyses).

SUPPLEMENTAL INFORMATION

Supplemental Information includes three tables, five figures, and Supplemental Experimental Procedures and can be found with this article online at doi:10.1016/j.ccr.2011.12.015.

ACKNOWLEDGMENTS

We thank Dr. M. Iwanaga, Mr. M. Nakashima, and Ms. T. Akashi for support and maintenance of JSPFAD. We thank Drs. H. Miyoshi and A. Miyawaki for providing the Venus-encoding lentivirus vectors. We also thank Dr. R. Horie for experimental advices, and Drs. T. Kanno and T. Ishida for providing the MDA-MB-453. Grant support: Grants-in-Aid for Scientific Research from Ministry of Education, Culture, Sports, Science, and Technology of Japan to T.W. (No. 23390250) and by Grants-in-Aid from the Ministry of Health, Labour and Welfare to T.W. (H21-G-002 and H22-AIDS-I-002).

Received: November 3, 2010

Revised: August 12, 2011

Accepted: December 19, 2011

Published: January 17, 2012

REFERENCES

- Annunziata, C.M., Davis, R.E., Demchenko, Y., Bellamy, W., Gabrea, A., Zhan, F., Lenz, G., Hanamura, I., Wright, G., Xiao, W., et al. (2007). Frequent engagement of the classical and alternative NF-kappaB pathways by diverse genetic abnormalities in multiple myeloma. *Cancer Cell* 12, 115–130.
- Bartel, D.P. (2009). MicroRNAs: target recognition and regulatory functions. *Cell* 136, 215–233.
- Bellon, M., Lepelletier, Y., Hermine, O., and Nicot, C. (2009). Deregulation of microRNA involved in hematopoiesis and the immune response in HTLV-I adult T-cell leukemia. *Blood* 113, 4914–4917.
- Caretti, G., Di Padova, M., Micales, B., Lyons, G.E., and Sartorelli, V. (2004). The Polycomb Ezh2 methyltransferase regulates muscle gene expression and skeletal muscle differentiation. *Genes Dev.* 18, 2627–2638.
- Chen, Y., Takita, J., Choi, Y.L., Kato, M., Ohira, M., Sanada, M., Wang, L., Soda, M., Kikuchi, A., Igarashi, T., et al. (2008). Oncogenic mutations of ALK kinase in neuroblastoma. *Nature* 455, 971–974.
- Davis, B.N., Hilyard, A.C., Lagna, G., and Hata, A. (2008). SMAD proteins control DROSHA-mediated microRNA maturation. *Nature* 454, 56–61.
- Fiskus, W., Wang, Y., Sreekumar, A., Buckley, K.M., Shi, H., Jillella, A., Ustun, C., Rao, R., Fernandez, P., Chen, J., et al. (2009). Combined epigenetic therapy with the histone methyltransferase EZH2 inhibitor 3-deazaneplanocin A and the histone deacetylase inhibitor panobinostat against human AML cells. *Blood* 114, 2733–2743.
- Hayden, M.S., and Ghosh, S. (2008). Shared principles in NF-kappaB signaling. *Cell* 132, 344–362.
- Hironaka, N., Mochida, K., Mori, N., Maeda, M., Yamamoto, N., and Yamaoka, S. (2004). Tax-independent constitutive IkkappaB kinase activation in adult T-cell leukemia cells. *Neoplasia* 6, 266–278.
- Horie, R., Watanabe, M., Ishida, T., Koiwa, T., Aizawa, S., Itoh, K., Higashihara, M., Kadin, M.E., and Watanabe, T. (2004). The NPM-ALK oncoprotein



- abrogates CD30 signaling and constitutive NF- κ B activation in anaplastic large cell lymphoma. *Cancer Cell* 5, 353–364.
- Iwanaga, M., Watanabe, T., Utsunomiya, A., Okayama, A., Uchimaru, K., Koh, K.R., Ogata, M., Kikuchi, H., Sagara, Y., Uozumi, K., et al; Joint Study on Predisposing Factors of ATL Development investigators. (2010). Human T-cell leukemia virus type I (HTLV-1) proviral load and disease progression in asymptomatic HTLV-1 carriers: a nationwide prospective study in Japan. *Blood* 116, 1211–1219.
- Kleer, C.G., Cao, Q., Varambally, S., Shen, R., Ota, I., Tomlins, S.A., Ghosh, D., Sewalt, R.G., Otte, A.P., Hayes, D.F., et al. (2003). EZH2 is a marker of aggressive breast cancer and promotes neoplastic transformation of breast epithelial cells. *Proc. Natl. Acad. Sci. USA* 100, 11606–11611.
- Liao, G., Zhang, M., Harhaj, E.W., and Sun, S.C. (2004). Regulation of the NF- κ B-inducing kinase by tumor necrosis factor receptor-associated factor 3-induced degradation. *J. Biol. Chem.* 279, 26243–26250.
- Malinin, N.L., Boldin, M.P., Kovalenko, A.V., and Wallach, D. (1997). MAP3K-related kinase involved in NF- κ B induction by TNF, CD95 and IL-1. *Nature* 385, 540–544.
- Min, J., Zaslavsky, A., Fedele, G., McLaughlin, S.K., Reczek, E.E., De Raedt, T., Guney, I., Strohlic, D.E., Macconail, L.E., Beroukhim, R., et al. (2010). An oncogene-tumor suppressor cascade drives metastatic prostate cancer by coordinately activating Ras and nuclear factor- κ B. *Nat. Med.* 16, 286–294.
- Prasad, S., Ravindran, J., and Aggarwal, B.B. (2010). NF- κ B and cancer: how intimate is this relationship. *Mol. Cell. Biochem.* 336, 25–37.
- Ramakrishnan, P., Wang, W., and Wallach, D. (2004). Receptor-specific signaling for both the alternative and the canonical NF- κ B activation pathways by NF- κ B-inducing kinase. *Immunity* 21, 477–489.
- Saitoh, Y., Yamamoto, N., Dewan, M.Z., Sugimoto, H., Martinez Bruyn, V.J., Iwasaki, Y., Matsubara, K., Qi, X., Saitoh, T., Imoto, I., et al. (2008). Overexpressed NF- κ B-inducing kinase contributes to the tumorigenesis of adult T-cell leukemia and Hodgkin Reed-Sternberg cells. *Blood* 111, 5118–5129.
- Sander, S., Bullinger, L., Klapproth, K., Fiedler, K., Kestler, H.A., Barth, T.F., Möller, P., Stilgenbauer, S., Pollack, J.R., and Wirth, T. (2008). MYC stimulates EZH2 expression by repression of its negative regulator miR-26a. *Blood* 112, 4202–4212.
- Schaefer, A., Jung, M., Mollenkopf, H.J., Wagner, I., Stephan, C., Jentzmk, F., Miller, K., Lein, M., Kristiansen, G., and Jung, K. (2010). Diagnostic and prognostic implications of microRNA profiling in prostate carcinoma. *Int. J. Cancer* 126, 1166–1176.
- Simon, J.A., and Kingston, R.E. (2009). Mechanisms of polycomb gene silencing: knowns and unknowns. *Nat. Rev. Mol. Cell Biol.* 10, 697–708.
- Sparmann, A., and van Lohuizen, M. (2006). Polycomb silencers control cell fate, development and cancer. *Nat. Rev. Cancer* 6, 846–856.
- Su, I.H., Basavaraj, A., Krutchinsky, A.N., Hobert, O., Ullrich, A., Chait, B.T., and Tarakhovskiy, A. (2003). Ezh2 controls B cell development through histone H3 methylation and Igh rearrangement. *Nat. Immunol.* 4, 124–131.
- Su, I.H., Dobenecker, M.W., Dickinson, E., Oser, M., Basavaraj, A., Marqueron, R., Viale, A., Reinberg, D., Wülfing, C., and Tarakhovskiy, A. (2005). Polycomb group protein ezh2 controls actin polymerization and cell signaling. *Cell* 121, 425–436.
- Suh, J., and Rabson, A.B. (2004). NF- κ B activation in human prostate cancer: important mediator or epiphenomenon? *J. Cell. Biochem.* 91, 100–117.
- Thu, Y.M., and Richmond, A. (2010). NF- κ B inducing kinase: a key regulator in the immune system and in cancer. *Cytokine Growth Factor Rev.* 21, 213–226.
- Trabucchi, M., Briata, P., Garcia-Mayoral, M., Haase, A.D., Filipowicz, W., Ramos, A., Gherzi, R., and Rosenfeld, M.G. (2009). The RNA-binding protein KSRP promotes the biogenesis of a subset of microRNAs. *Nature* 459, 1010–1014.
- Valastyan, S., Reinhardt, F., Benaich, N., Calogrias, D., Szász, A.M., Wang, Z.C., Brock, J.E., Richardson, A.L., and Weinberg, R.A. (2009). A pleiotropically acting microRNA, miR-31, inhibits breast cancer metastasis. *Cell* 137, 1032–1046.
- Varambally, S., Dhanasekaran, S.M., Zhou, M., Barrette, T.R., Kumar-Sinha, C., Sanda, M.G., Ghosh, D., Pienta, K.J., Sewalt, R.G., Otte, A.P., et al. (2002). The polycomb group protein EZH2 is involved in progression of prostate cancer. *Nature* 419, 624–629.
- Varambally, S., Cao, Q., Mani, R.S., Shankar, S., Wang, X., Ateeq, B., Laxman, B., Cao, X., Jing, X., Ramnarayanan, K., et al. (2008). Genomic loss of microRNA-101 leads to overexpression of histone methyltransferase EZH2 in cancer. *Science* 322, 1695–1699.
- Ventura, A., and Jacks, T. (2009). MicroRNAs and cancer: short RNAs go a long way. *Cell* 136, 586–591.
- Watanabe, M., Ohsugi, T., Shoda, M., Ishida, T., Aizawa, S., Maruyama-Nagai, M., Utsunomiya, A., Koga, S., Yamada, Y., Kamihira, S., et al. (2005). Dual targeting of transformed and untransformed HTLV-1-infected T cells by DHMEQ, a potent and selective inhibitor of NF- κ B, as a strategy for chemoprevention and therapy of adult T-cell leukemia. *Blood* 106, 2462–2471.
- Yamagishi, M., Ishida, T., Miyake, A., Cooper, D.A., Kelleher, A.D., Suzuki, K., and Watanabe, T. (2009). Retroviral delivery of promoter-targeted shRNA induces long-term silencing of HIV-1 transcription. *Microbes Infect.* 11, 500–508.
- Yamaguchi, K., and Watanabe, T. (2002). Human T lymphotropic virus type-1 and adult T-cell leukemia in Japan. *Int. J. Hematol.* 76 (Suppl 2), 240–245.
- Yeung, M.L., Yasunaga, J., Bennasser, Y., Dusetti, N., Harris, D., Ahmad, N., Matsuoka, M., and Jeang, K.T. (2008). Roles for microRNAs, miR-93 and miR-130b, and tumor protein 53-induced nuclear protein 1 tumor suppressor in cell growth dysregulation by human T-cell lymphotropic virus 1. *Cancer Res.* 68, 8976–8985.
- Zarnegar, B.J., Wang, Y., Mahoney, D.J., Dempsey, P.W., Cheung, H.H., He, J., Shiba, T., Yang, X., Yeh, W.C., Mak, T.W., et al. (2008a). Noncanonical NF- κ B activation requires coordinated assembly of a regulatory complex of the adaptors cIAP1, cIAP2, TRAF2 and TRAF3 and the kinase NIK. *Nat. Immunol.* 9, 1371–1378.
- Zarnegar, B.J., Yamazaki, S., He, J.Q., and Cheng, G. (2008b). Control of canonical NF- κ B activation through the NIK-IKK complex pathway. *Proc. Natl. Acad. Sci. USA* 105, 3503–3508.

Carnosol, rosemary ingredient, induces apoptosis in adult T-cell leukemia/lymphoma cells via glutathione depletion: proteomic approach using fluorescent two-dimensional differential gel electrophoresis


Yo-ichi Ishida, Masao Yamasaki, Chizuko Yukizaki, Kazuo Nishiyama, Hirohito Tsubouchi, et al.

Human Cell

e-ISSN 1749-0774

Human Cell

DOI 10.1007/s13577-013-0083-6



 Springer

Your article is protected by copyright and all rights are held exclusively by Japan Human Cell Society and Springer Japan. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".