

Fig. 1 Clinical course during hospitalization. *DRV* darunavir, *RTV* ritonavir, *TDF* tenofovir, *FTC* emtricitabine, *HD-DXM* high-dose dexamethasone

bleeding). To rule out drug-induced thrombocytopenia, ART, together with clopidogrel and aspirin, which had been administered for years, were discontinued on admission. Although platelet transfusion was initiated for a couple of days, no change in platelet count was noted. Bone marrow examination on day 5 showed hypocellularity with a low number of megakaryocytes. No histopathological findings specific to myelodysplastic syndrome or leukemia were noted. On day 10, the patient developed intermittent epistaxis with a platelet count of 4,000/ μ l. On that day, a four-day course of orally administered HD-DXM of 40 mg/day was initiated, and ART with DRV/r plus TDF/FTC was reinitiated. The platelet count increased to 66,000/ μ l on day 10 after the above treatment, but it decreased to 12,000/ μ l on day 14. A second course of HD-DXM of 40 mg/day was initiated. The platelet count improved to 115,000/ μ l on day 10 after the second course, and 142,000/ μ l on day 15. Based on such improvement, no third course was considered necessary. The patient was discharged on day 39 from admission. No adverse event of dexamethasone was observed. The platelet count remained stable after discharge despite the re-initiation of clopidogrel and aspirin (Fig. 1). Three months after re-initiation of ART and thereafter, the HIV-1 viral load was suppressed to <100 copies/ml with good medication adherence. The patient experienced no relapses of HIV-ITP for 9 months.

Discussion

We reported here a patient with HIV-ITP who was treated successfully with a combination with two courses of HD-DXM and ART. The ITP likely relapsed when the platelet count diminished to <90,000/ μ l on day 10 after the first course of HD-DXM [3], thus justifying the second

course of HD-DXM. No additional courses were provided once the platelet count was above 90,000/ μ l on day 10 after the second course. That platelet count remained stable after re-initiating clopidogrel and aspirin negated any drug-induced thrombocytopenia. To our knowledge, this is the first case describing the use and effectiveness of the combination of HD-DXM and ART in the treatment of HIV-ITP.

In adult patients with non-HIV ITP, HD-DXM is preferred to conventional long-term prednisone [2], because of fewer adverse events, mainly due to the shorter term of steroid administration. The major side effect of steroid is immunosuppression, and it is important to avoid such complication, especially in immunocompromised hosts, such as HIV-1 infected patients. For the treatment of HIV-ITP, HD-DXM is probably as effective as in non-HIV ITP, because both diseases are considered to have a similar etiology [8]. Although about one-fifth of non-HIV ITP patients on HD-DXM treatment relapse by 8 months after treatment [5], the use of ART in patients with HIV-ITP can maintain long-term remission despite the lack of an immediate effect [9, 10]. In our patient, HD-DXM was applied when the clinical condition was severe with bleeding and thrombocytopenia, and it resulted in rapid improvement in platelet count. Thereafter, administration of ART resulted in suppression of viral load, which probably promoted long-term remission of HIV-ITP.

Notably, HD-DXM is cost-effective, compared to other treatments for ITP, such as immunoglobulin or rituximab. Although further studies are needed to confirm the efficacy and safety of the combination therapy, HD-DXM and ART is potentially suitable for treatment of HIV-ITP patients with severe thrombocytopenia or bleeding.

Acknowledgments The authors thank all the clinical staff at the AIDS Clinical Center, National Center for Global Health and Medicine. No financial support was received for this article.

Conflict of interest All authors declare no conflict of interest.

References

1. Ambler KL, Vickars LM, Leger CS, Foltz LM, Montaner JS, Harris M, et al. Clinical features, treatment, and outcome of HIV-associated immune thrombocytopenia in the HAART era. *Adv Hematol*. 2012;2012:910954.
2. Neunert C, Lim W, Crowther M, Cohen A, Solberg L Jr, Crowther MA. The American Society of Hematology 2011 evidence-based practice guideline for immune thrombocytopenia. *Blood*. 2011;117:4190–207.
3. Cheng Y, Wong RS, Soo YO, Chui CH, Lau FY, Chan NP, et al. Initial treatment of immune thrombocytopenic purpura with high-dose dexamethasone. *N Engl J Med*. 2003;349:831–6.
4. Godeau B, Bierling P. High-dose dexamethasone as initial treatment for immune thrombocytopenic purpura. *N Engl J Med*. 2003;349:2267–8. (author reply 2267–2268).

5. Mazzucconi MG, Fazi P, Bernasconi S, De Rossi G, Leone G, Gugliotta L, et al. Therapy with high-dose dexamethasone (HD-DXM) in previously untreated patients affected by idiopathic thrombocytopenic purpura: a GIMEMA experience. *Blood*. 2007; 109:1401–7.
6. Marroni M, Gresele P. Detrimental effects of high-dose dexamethasone in severe, refractory, HIV-related thrombocytopenia. *Ann Pharmacother*. 2000;34:1139–41.
7. Ramratnam B, Parameswaran J, Elliot B, Newstein M, Schiffman FJ, Rich JD, et al. Short course dexamethasone for thrombocytopenia in AIDS. *Am J Med*. 1996;100:117–8.
8. Bettaieb A, Fromont P, Louache F, Oksenhendler E, Vainchenker W, Duedari N, et al. Presence of cross-reactive antibody between human immunodeficiency virus (HIV) and platelet glycoproteins in HIV-related immune thrombocytopenic purpura. *Blood*. 1992; 80:162–9.
9. Arranz Caso JA, Sanchez Mingo C, Garcia Tena J. Effect of highly active antiretroviral therapy on thrombocytopenia in patients with HIV infection. *N Engl J Med*. 1999;341:1239–40.
10. Carbonara S, Fiorentino G, Serio G, Maggi P, Ingravallo G, Monno L, et al. Response of severe HIV-associated thrombocytopenia to highly active antiretroviral therapy including protease inhibitors. *J Infect*. 2001;42:251–6.

Prophylactic Effect of Antiretroviral Therapy on Hepatitis B Virus Infection

Hiroyuki Gatanaga,^{1,2} Tsunefusa Hayashida,^{1,2} Junko Tanuma,¹ and Shinichi Oka^{1,2}

¹AIDS Clinical Center, National Center for Global Health and Medicine, Tokyo, and ²Center for AIDS Research, Kumamoto University, Japan

Background. Hepatitis B virus (HBV) infection is common in individuals infected with human immunodeficiency virus, especially in men who have sex with men (MSM). Almost all currently used regimens of antiretroviral therapy (ART) contain lamivudine (LAM) or tenofovir disoproxil fumarate (TDF), both of which have significant anti-HBV activity. However, the prophylactic effect of ART on HBV infection has not been assessed previously.

Methods. Non-HBV-vaccinated HIV-infected MSM were serologically evaluated for HBV infection using stocked serum samples. Cases negative for HBV surface antigen (HBsAg), antibody to HBsAg (anti-HBs), and antibody to HBV core antigen (anti-HBc) in first serum samples were serologically followed until last available stocked samples. HBV genotype and LAM-resistant mutation (rtM204V/I) were analyzed in cases that became HBsAg-positive.

Results. The first stocked samples were negative for all analyzed HBV serological markers in 354 of 1434 evaluated patients. The analysis of their last samples indicated HBV incident infection in 43 of them during the follow-up period. The rate of incident infections was lower during LAM- or TDF-containing ART (0.669 incident infections in 100 person-years) than during no ART period (6.726 incident infections in 100 person-years) and other ART (5.263 incident infections in 100 person-years) ($P < .001$). Genotype A was most prevalent (76.5%), and LAM-resistant HBV was more frequent in incident infections during LAM-containing ART (50.0%) than in those during no ART and other ART (7.1%) ($P = .029$).

Conclusions. LAM- and TDF-containing ART regimens seem to provide prophylaxis against HBV infection, although drug-resistant strains seem to evade these effects.

Keywords. lamivudine; tenofovir disoproxil fumarate; resistant; chronic infection.

Patients with human immunodeficiency virus (HIV) infection are at high risk for both hepatitis B virus (HBV) infection and development of chronic infection [1–4]. Based on information from Western countries, the rate of coinfection varies according to risk categories; the highest rate is in men who have sex with men (MSM), with a slightly lower rate among intravenous drug users, and much lower in individuals infected through heterosexual contacts [5–8]. In Japan, HIV/

HBV coinfection is also significantly associated with MSM [9, 10]. The progression of chronic HBV infection to cirrhosis, end-stage liver diseases, and/or hepatocellular carcinoma is more rapid in HIV-infected persons than in those with chronic HBV infection alone [11, 12]. Vaccination of non-HBV-immunized HIV-infected individuals is recommended to prevent HBV infection [13]. However, all current recommended antiretroviral therapy (ART) regimens contain lamivudine (LAM) or tenofovir disoproxil fumarate (TDF), both of which have significant anti-HBV activity [14]. Do these ART regimens provide any prophylaxis against HBV infection? This is an important question, as a positive answer could influence the strategy applied to prevent HBV infection in HIV-infected individuals. To delineate the hepatitis B prophylactic effect of ART, we used stocked samples for serological evaluation of HBV infection in HIV-infected MSM. The present

Received 6 November 2012; accepted 25 February 2013; electronically published 13 March 2013.

Correspondence: Hiroyuki Gatanaga, MD, AIDS Clinical Center, National Center for Global Health and Medicine, 1-21-1 Toyama, Shinjuku-ku, Tokyo 162-8655, Japan (higatana@acc.ncgm.go.jp).

Clinical Infectious Diseases 2013;56(12):1812–9

© The Author 2013. Published by Oxford University Press on behalf of the Infectious Diseases Society of America. All rights reserved. For Permissions, please e-mail: journals.permissions@oup.com.

DOI: 10.1093/cid/cit145

study included those patients who had tested negative for hepatitis B surface antigen (HBsAg), antibody to HBsAg (anti-HBs), and antibody to hepatitis B core antigen (anti-HBc) using their first stocked blood samples, who were followed up serologically to identify new HBV incident infections among them. The other part of the study covered analysis of the relation between the frequency of incident infection and ART regimens.

METHODS

Patients

Since April 1997, we have stocked serum samples taken at routine clinical practice from HIV type 1 (HIV-1)-infected patients who visited the Outpatient Clinic of the AIDS Clinical Center, National Center for Global Health and Medicine, Tokyo, Japan, under signed informed consent for use in virologic research. Every patient had been interviewed at the first visit by clinical nurse specialists at the HIV outpatient clinic using a structured questionnaire that includes items on sexuality and history of HBV vaccination. Most of the patients regularly visited our clinic every 1–3 months, and we had collected and stored their sera at almost all visits. The ethics committee of the National Center for Global Health and Medicine approved the collection and analysis of the samples. First, we selected HIV-1-infected MSM who met the following inclusion criteria: (1) the first visit to our clinic was between April 1997 and December 2009, (2) they had not received HBV vaccination before the first visit, and (3) at least 2 serum samples were available and collected at least 6 months apart. The first sample was defined as the baseline serum sample, and baseline clinical data were defined as those recorded on the date of sampling of the first stocked serum. Patients' baseline characteristics, including age, race, hepatitis C virus antibody, results of *Treponema pallidum* hemagglutination assay, and CD4⁺ cell count were collected from the medical records.

HBV Analysis

In order to identify new HBV incident infection, we excluded patients with previously confirmed HBV infection. The baseline samples of the patients who met the inclusion criteria described above were serologically evaluated for HBsAg, anti-HBs, and anti-HBc using ARCHITECT HBsAg QT assay, anti-HBs assay, and anti-HBc assay, respectively (Abbott Laboratories, Chicago, Illinois) [15, 16]. Patients positive for any of HBsAg, anti-HBs, and anti-HBc at baseline were excluded from the serological follow-up. The remaining patients were considered to have never been infected with HBV before the baseline. Their last stocked sample taken before or in December 2010, or before HBV vaccination if performed during the follow-up period, was analyzed for HBsAg, anti-HBs, and anti-HBc. If the last sample was negative for all 3, the patient was

considered to have never been infected with HBV up to the sampling date of the last stocked serum. If HBsAg, anti-HBs, or anti-HBc was positive in the last stocked serum, the patient was considered to have HBV incident infection during the follow-up period. In the latter case, the baseline samples were subjected to polymerase chain reaction (PCR) analysis for HBV DNA [17, 18], and all the stocked samples during the follow-up period were serologically analyzed to determine the date of HBV incident infection. The date of incident infection was defined as the sampling date of the first positive serum for any HBV serological marker. The time from the baseline to HBV incident infection was analyzed by the Kaplan-Meier method. The data were censored at the sampling date of the last stocked sample if it was negative for all analyzed HBV serological markers. Patients' age and CD4⁺ cell count at the date of incident infection and alanine aminotransferase (ALT) values within 3 months of incident infection were collected. If an HBsAg-positive sample was available, HBV genotype and LAM-resistant mutation (rtM204V/I) were analyzed by PCR-invaser assay [17–19]. The diagnosis of chronic HBV infection was considered when HBsAg was still positive in sera taken at 6 months or longer after the incident infection.

Antiretroviral Therapy

To determine the type of ART under which HBV incident infection occurred, the regimen information of ART was collected from medical records over the period spanning from the baseline to the incidence infection or to the end of follow-up. The treatment status was divided into 4 categories: (1) No ART, no treatment with any antiretroviral agent; (2) Other-ART, ART with regimens that did not contain LAM, TDF, or emtricitabine (FTC); (3) LAM-ART, ART with LAM-containing regimens that did not contain TDF or FTC; and (4) TDF-ART, ART with TDF-containing regimens with or without LAM or FTC. Data were censored on the sampling date of the last stocked sample if it was negative for all analyzed HBV serological markers. When the treatment category was modified, the data were censored on the date of category change for the previous treatment category and a new follow-up as a different case was initiated for the replacement treatment category.

Statistical Analysis

The time from the baseline to HBV incident infection was analyzed by the Kaplan-Meier method. The Cox proportional hazards regression analysis was used to assess the risk of HBV incident infections. The impact of patients' baseline characteristics, year of entry, the use of antiretroviral agents (any antiretroviral, and any of LAM, TDF, or FTC), and the frequency of changing ART regimen during the follow-up period was estimated with univariate analysis, and those with statistical significance were incorporated into multivariate analysis. The




Figure 1. Patient selection process: 1434 patients met the inclusion criteria. Of these patients, 1080 were excluded because of positive hepatitis B virus serology in the first samples. The results of various serological tests are shown. The remaining 354 were enrolled for serological follow-up. Of these, 43 were positive in the last sample analysis. Their stocked samples were analyzed serologically and the results of HBV serology using the first positive samples are indicated. Abbreviations: anti-HBc, antibody to HBV core antigen; anti-HBs, antibody to HBsAg; HBsAg, hepatitis B surface antigen; HBV, hepatitis B virus; MSM, men who have sex with men.

frequency and risk of HBV incident infection during each treatment category was also assessed by univariate Cox proportional hazards regression analysis. We used hazard ratios and 95% confidence intervals to estimate the impact of each variable on incident infection. Patients' age and CD4⁺ cell count on the date of incident infection, and peak value of ALT within 3 months of incident infection were compared between transient infection and chronic infection with Wilcoxon rank-sum test. The differences in rates of HBV genotype A and rtM204V/I mutation were compared with χ^2 test (ie, the Fisher exact test).

Statistical significance of difference was defined as a 2-sided *P* value of <.05. All statistical analyses were performed with the Statistical Package for Social Sciences version 17.0 (SPSS, Chicago, Illinois).

RESULTS

Figure 1 shows the patient selection procedure. A total of 1434 HIV-1-infected MSM met the inclusion criteria described in the Methods section. Of these, 146 patients (10.2%) were positive for HBsAg, 737 (51.4%) were positive for anti-HBs, and 197 (13.7%) were solely positive for anti-HBc using baseline samples. The remaining 354 patients (24.7%; negative for HBsAg, anti-HBs, and anti-HBc at baseline), who were considered to have never been infected with HBV, were enrolled for serological follow-up. Table 1 lists their baseline characteristics. Serological analysis of the last sample of each of these patients showed HBV incident infection during follow-up in 43 (12.1%). Their baseline samples were found to be PCR-negative for HBV DNA, confirming that the incident infection in these patients occurred during the follow-up period. All stocked samples of the 43 patients were analyzed serologically to determine the date of HBV incident infection. HBV incident infections occurred every year between 1997 and 2010 except in 1998. The median time period from the baseline to HBV incident infection was 1.6 years (interquartile range [IQR], 192–1151 days; range, 28–4068 days). The total observation period was 1607 person-years (median, 3.7 years [IQR], 1.9–6.5 years). Figure 2 shows the Kaplan-Meier curve for the HBV incident infection for the whole cohort of enrolled patients.

In order to assess the risk of HBV incident infections, patients' baseline characteristics, year of entry, the use of any anti-retroviral agents, the use of any of LAM, TDF, or FTC, and the frequency of changing ART regimen during the follow-up

Table 1. Baseline Characteristics of the 354 Enrolled Patients

Characteristic	Total (n = 354)	Year of Entry			
		1997–2000 (n = 61)	2001–2003 (n = 79)	2004–2006 (n = 112)	2007–2009 (n = 102)
Age, y, median (IQR)	32.0 (27.0–38.0)	32.0 (27.8–37.3)	31.0 (27.0–37.8)	32.0 (27.0–38.0)	35.0 (27.0–42.0)
Race/ethnicity					
Japanese	340 (96.0)	59 (96.7)	78 (98.7)	109 (97.3)	94 (92.2)
Asian other than Japanese	4 (1.1)	0 (0.0)	0 (0.0)	1 (0.9)	3 (2.9)
Caucasian	10 (2.8)	2 (3.3)	1 (1.3)	2 (1.8)	5 (4.9)
HCV antibody, positive	8 (2.3)	1 (1.6)	2 (2.5)	1 (0.9)	4 (3.9)
TPHA positive	101 (28.5)	23 (37.7)	20 (25.3)	30 (26.8)	28 (27.5)
CD4 ⁺ cell count, cells/mm ³ , median (IQR)	277 (151–404)	277 (169–417)	313 (97–443)	316 (176–413)	252 (129–359)
HIV RNA, log ₁₀ copies/mL, median (IQR)	4.6 (3.8–5.2)	4.5 (3.6–5.2)	4.8 (3.9–5.4)	4.4 (3.8–4.9)	4.7 (3.9–5.2)

Data are No. (%) unless otherwise specified.

Abbreviations: HCV, hepatitis C virus; HIV, human immunodeficiency virus; IQR, interquartile range; TPHA, *Treponema pallidum* hemagglutination assay.




Figure 2. Kaplan-Meier curve showing the time to hepatitis B virus incident infection.

period were estimated using a proportional hazards model (Table 2). Younger age and higher CD4⁺ cell count correlated positively, and use of any antiretroviral, use of LAM, TDF, or FTC, and the frequency of changing ART regimen correlated negatively with HBV incident infection, with statistical significance in univariate analysis. However, in multivariate analysis, the use of LAM, TDF, or FTC continued to show significant relation. Then, we focused on the relation between treatment status and HBV incident infection. The observation period in each patient was divided into 4 categories by treatment status: No ART, no treatment with any antiretroviral agent; Other-ART, ART with regimens that did not contain LAM, TDF, or FTC; LAM-ART, ART with LAM-containing regimens that did not contain TDF or FTC; or TDF-ART, ART with TDF-containing regimens with or without LAM or FTC. No

participant received FTC single tablet (Emtriva). All the participants who took FTC received the combination tablet of TDF/FTC (Truvada), and therefore, such treatment status was categorized as TDF-ART. The total categorized observation period of No ART, Other-ART, LAM-ART, and TDF-ART was 446, 114, 814, and 233 person-years, respectively. The number of the HBV incident infections was 30 during the No ART period, 6 during Other-ART period, 7 during LAM-ART period, and 0 during TDF-ART period. No incident infection occurred at the time of changing ART regimen. The proportional hazards model showed a significantly lower frequency of HBV incident infection during LAM- or TDF-ART (0.669 incident infections per 100 person-years) compared with that during No ART (6.726 incident infections per 100 person-years), although there was no significant difference between Other-ART (5.263 incident infections per 100 person-years) and No ART, suggesting that ART regimens with anti-HBV activity can reduce HBV incident infections by 90% (Table 3). During LAM-ART, the HIV-1 load around the period of incident infection remained below the detection limit in all the 7 infected patients, indicating excellent adherence to ART.

Figure 3 shows peak ALT levels for the 43 HBV incident infections. Among the 36 incident infections observed the No ART and Other-ART groups, 16 infections (44.4%) were asymptomatic and not associated with significant increases in ALT (peak ALT, <60 IU/L). We were able to serologically follow 33 of the 36 cases for 6 months after the date of incident infection (TDF-ART was introduced within 6 months of incident infection in the other 3 cases). Among the 33 patients, 13 (39.4%) developed chronic infection (HBsAg was still positive 6 months after the date of incident infection). The median CD4⁺

Table 2. Cox Proportional Hazards Regression Analysis for the Risk of Hepatitis B Virus Incident Infection

Factors	Univariate Analysis		Multivariate Analysis	
	Hazard Ratio (95% CI)	P Value	Hazard Ratio (95% CI)	P Value
Year of entry, per 1 y increase	.942 (.860–1.033)	.207		
Baseline characteristics				
Age, per 1 y increase	.921 (.879–.965)	.001	.958 (.917–1.001)	.054
Race (Japanese)	21.243 (.010–45 657.613)	.435		
HCV antibody	.048 (<.001–346.311)	.503		
TPHA	1.475 (.792–2.747)	.220		
CD4 ⁺ cell count, per 100 cells/mm ³ increase	1.121 (1.008–1.246)	.035	.882 (.752–1.034)	.121
HIV RNA, per 1 log ₁₀ copies/mL increase	1.387 (.999–1.924)	.051		
Antiretroviral use during follow-up period				
Any antiretroviral	.097 (.052–.184)	<.001	.927 (.305–2.818)	.893
LAM, TDF, or FTC	.075 (.039–.146)	<.001	.110 (.031–.390)	.001
Frequency of changing regimen	.245 (.145–.414)	<.001	.700 (.385–1.270)	.240

Abbreviations: CI, confidence interval; FTC, emtricitabine; HCV, hepatitis C virus; HIV, human immunodeficiency virus; LAM, lamivudine; TDF, tenofovir disoproxil fumarate; TPHA, *Treponema pallidum* hemagglutination assay.

Table 3. Frequency and Hazard Ratio of Hepatitis B Virus Incident Infection in Each Treatment Status Category

ART	Observation Period (Person-Years)	Incident Infection	Hazard Ratio (95% CI)	P Value
No ART	446	30	1	...
Other-ART	114	6	.924 (.381–2.239)	.861
ART containing at least 1 of LAM, TDF, and FTC ^a	1047	7	.113 (.049–.261)	<.001
LAM-ART	814	7		
TDF-ART	233	0		

Abbreviations: ART, antiretroviral therapy; CI, confidence interval; FTC, emtricitabine; LAM, lamivudine; TDF, tenofovir disoproxil fumarate; LAM-ART, ART with LAM-containing regimens that did not contain TDF or FTC; Other-ART, ART with regimens that did not contain LAM, TDF, or FTC; TDF-ART, ART with TDF-containing regimens with or without LAM or FTC.

^a No participant received FTC single tablet (Emtriva) during the observation period. All the participants who took FTC received the combination tablet of TDF/FTC (Truvada), and therefore, such treatment status was categorized into TDF-ART.

cell count was lower in the patients who developed chronic infection than in those with transient infection, although the difference was not significant ($P = .068$; Table 4), indicating that HIV-related immunodeficiency may play a role in the induction of chronic HBV infection. Among the 7 incident infections observed during LAM-ART, only 2 patients (28.6%) were symptomatic, had significant rise in ALT, and developed chronic HBV infection, and both of these infections were caused by LAM-resistant HBV (Table 5). The other 5 cases were asymptomatic and transient. Three of them were caused by LAM-sensitive strains and 1 was by LAM-resistant strain. HBsAg-positive serum sample was not available in the last case. LAM-resistant HBV was more frequently identified in analyzed incident infections during LAM-containing ART (50.0%) than in those during no ART and other ART (7.1%) ($P = .029$). Considered together, LAM seems to prevent acquisition of HBV infection, progression to symptomatic hepatitis, and development of chronic infection even after the development of infection, although these effects may be less pronounced in patients with LAM-resistant strains.

Among the 43 infection cases observed during total serological follow-up, HBsAg-positive samples were available in 34 cases and their HBV genotype was determined. Genotype A was the most frequent, as reported previously [10, 20–22], and genotypes B, G, and H were also identified. The rate of development of chronic infection was higher in genotype A than in other genotypes as previously reported [23], although the difference was not significant in our study. In the remaining 9 cases, only anti-HBc with (7 cases) or without (2 cases) anti-HBs were detected, although their samples were available and




Figure 3. Peak alanine aminotransferase (ALT) values in hepatitis B virus (HBV) incident infections according to treatment regimen. Thirty, 6, and 7 HBV incident infections were observed during No antiretroviral therapy (ART), Other-ART, and lamivudine (LAM)-ART, respectively. No incident infection was identified during tenofovir disoproxil fumarate (TDF)-ART. No participant received emtricitabine (FTC) single tablet (Emtriva) during the observation period. All the participants who took FTC received the combination tablet of TDF/FTC (Truvada), and therefore, such treatment status was categorized into TDF-ART. Data are peak ALT values measured within 3 months of the date of incident infections. LAM-resistant mutation (rtM204V/I) was analyzed in 34 cases using the available hepatitis B surface antigen (HBsAg)-positive samples. Open squares: patients infected with LAM-resistant HBV. Closed circles and squares: patients who developed chronic infection (HBsAg-positive 6 months after the date of incident infection). Checked circles and squares: patients who received TDF-containing ART within 6 months of incident infection. Abbreviations: ALT, alanine aminotransferase; ART, antiretroviral therapy; LAM, lamivudine.

serologically analyzed at least every 3 months around the incident infection.

DISCUSSION

The results of this serological follow-up study indicated that LAM- and TDF-containing ART regimens protect against HBV incident infection. Furthermore, the results also suggested that LAM prevents progression to symptomatic hepatitis and development of chronic infection even after the development of HBV incident infection, provided such infection is caused by LAM-sensitive strains. However, it seems that LAM-resistant strains may evade this protective effect. One previous study that estimated the incidence of acute HBV infection among HIV-infected patients reported similar frequencies in patients receiving ART with and without LAM [5]. However, the authors defined immunoglobulin M anti-HBs positivity as a marker of HBV incident infection and did not exclude anti-HBc-positive patients at study entry. This probably made it difficult to distinguish incident infection from reactivation of chronic infection, as discussed in the report. In this study, we identified a

Table 4. Patient Characteristics and Clinical Features of Hepatitis B Virus Incident Infections in the No Antiretroviral Therapy (ART) and Other-ART Treatment Categories

Factors	Transient (n = 20)	Chronic ^a (n = 13)	Treated ^b (n = 3)	P Value ^c
Age, y, median (IQR)	31.0 (28.0–33.0)	29.0 (25.0–38.3)	25.0 (21.0–35.0) ^d	.406
CD4 ⁺ cell count, cells/mm ³ , median (IQR)	371 (308–518)	320 (235–383)	674 (206–1935) ^d	.068
Peak ALT level ^e , U/L, median (IQR)	65 (30–573)	264 (115–774)	31 (15–314) ^d	.162
HBV genotype, No. (%)				.645
Genotype A	9 (45.0)	11 (84.6)	2 (66.7)	
Other genotypes	3 (15.0)	2 (15.4)	1 (33.3)	
Genotype unknown	8 (40.0)	0 (0.0)	0 (0.0)	
HBV rtM204V/I mutation, No. (%)				.480
Positive	1 (5.0)	0 (0.0)	1 (33.3)	
Negative	11 (55.0)	13 (100.0)	2 (66.7)	
Unknown	8 (40.0)	0 (0.0)	0 (0.0)	

Abbreviations: ALT, alanine aminotransferase; HBV, hepatitis B virus; IQR, interquartile range.

^a Hepatitis B surface antigen–positive 6 months after the date of incident infection.

^b Treated cases with tenofovir disoproxil fumarate–containing antiretroviral therapy within 6 months of incident infection.

^c P values between transient and chronic cases calculated with Wilcoxon rank-sum tests for continuous variables and χ^2 tests for proportions.

^d Minimum and maximum values.

^e Peak ALT level within 3 months of incident infection.

significant number of isolated anti-HBc–positive patients, a finding in agreement with previous reports [24–27], and

Table 5. Patient Characteristics and Clinical Features of Hepatitis B Virus Incident Infections During LAM-ART Treatment

Factors	Transient (n = 5)	Chronic ^a (n = 2)	P Value ^b
Age, y, median (IQR)	33.0 (30.3–36.5)	38.0 (33.0–43.0) ^c	.329
CD4 ⁺ cell count, cells/mm ³ , median (IQR)	430 (267–648)	362 (360–364) ^c	.699
Peak ALT level ^d , U/L, median (IQR)	22 (14–51)	1133 (941–1325) ^c	.051
HBV genotype, No. (%)			>.999
Genotype A	3 (60.0)	1 (50.0)	
Other genotypes	1 (20.0)	1 (50.0)	
Genotype unknown	1 (20.0)	0 (0.0)	
HBV rtM204V/I mutation, No. (%)			.400
Positive	1 (20.0)	2 (100.0)	
Negative	3 (60.0)	0 (0.0)	
Unknown	1 (20.0)	0 (0.0)	

Abbreviations: ALT, alanine aminotransferase; HBV, hepatitis B virus; IQR, interquartile range; LAM-ART, ART with LAM-containing regimens that did not contain TDF or FTC.

^a Hepatitis B surface antigen–positive 6 months after the date of incident infection.

^b P values calculated with Wilcoxon rank-sum tests for continuous variables and χ^2 tests for proportions.

^c Minimum and maximum values.

^d Peak ALT level within 3 months of incident infection.

excluded them from the serological follow-up to avoid improper inclusion of isolated anti-HBc–positive ones as HBV-naïve [28, 29].

HBV vaccination is recommended for individuals seeking evaluation or treatment for sexually transmitted diseases, HIV-infected patients, sexually active persons with >1 partner, and MSM [13]. However, the response and durability of adequate titers of anti-HBs are often reduced in HIV-infected patients [30–34]. Modified regimens of vaccination have been reported to improve anti-HBs response in HIV-infected patients, although the response rate was still low in those with low CD4⁺ cell counts [35–37]. Our study demonstrated the HBV prophylactic effects of LAM- and TDF-containing ART regimens, suggesting that ART should be initiated before HBV vaccination, especially in those with low CD4⁺ cell counts. Early introduction of ART was recommended recently not only for HIV-infected individuals, but also for prevention of transmission to others [38, 39]. Early introduction of treatment may also be recommended to prevent HBV infection to the patients themselves if they are HBV-naïve. One randomized clinical trial reported the prophylactic effect of TDF combined with FTC in HIV prevention in seronegative MSM [40]. However, in that trial, HBV vaccination was offered to all susceptible participants, which made it impossible to estimate the prophylactic effect of the treatment on HBV prevention.

Our study carries certain limitations related to its retrospective nature. Patients on ART might have more opportunities to improve their behavior to prevent transmission of HIV to others, which could reduce HBV infection in themselves but

introduce bias in our analysis. However, the results suggest prophylaxis against potential HBV infection by oral medications, which could be useful for nonimmunized medical care providers.

Notes

Acknowledgments. We thank Y. Takahashi and F. Negishi for the helpful assistance in sample processing, and A. Nakano for the excellent project coordination. We also thank all physicians and nurses at the AIDS Clinical Center, National Center for Global Health and Medicine, for the dedicated clinical practice and patient care.

Financial support. This work was supported in part by Grants-in Aid for AIDS research from the Ministry of Health, Labor, and Welfare (H23-AIDS-001), and the Global COE Program (Global Education and Research Center Aiming at the Control of AIDS), MEXT, Japan.

Potential conflicts of interest. H. G. has received honoraria from MSD K.K., Abbott Japan, Janssen Pharmaceutical K.K., Torii Pharmaceutical, and ViiV Healthcare. S. O. has received honoraria and research funding from MSD K.K., Abbott Japan, Janssen Pharmaceutical K.K., Pfizer, Roche Diagnostics K.K., and ViiV Healthcare, and has received honoraria from Astellas Pharmaceutical K.K., Bristol-Myers K.K., Daiichisankyo, Dainippon Sumitomo Pharma, GlaxoSmithKline, K.K., Taisho Toyama Pharmaceutical, and Torii Pharmaceutical. All other authors report no potential conflicts.

All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

- Colin JF, Cazals-Hatem D, Lioriot MA, et al. Influence of human immunodeficiency virus infection on chronic hepatitis B in homosexual men. *Hepatology* **1999**; 29:1306–10.
- Lacombe K, Bottero J, Lemoine M, et al. HIV/hepatitis B virus coinfection: current challenges and new strategies. *J Antimicrob Chemother* **2010**; 65:10–7.
- Soriano V, Vispo E, Labarga P, et al. Viral hepatitis and HIV coinfection. *Antiviral Res* **2010**; 85:303–15.
- Kourtis AP, Bulters M, Hu DJ, et al. HIB-HBV coinfection—a global challenge. *N Engl J Med* **2012**; 366:1749–52.
- Kellerman SE, Hanson DL, McNaghten AD, et al. Prevalence of chronic hepatitis B and incidence of acute hepatitis B infection in human immunodeficiency virus-infected subjects. *J Infect Dis* **2003**; 188:571–7.
- Lincoln D, Petoumenos K, Dore GJ, et al. HIV/HBV and HIV/HCV coinfection, and outcomes following highly active antiretroviral therapy. *HIV Med* **2003**; 4:241–9.
- Nunez M, Soriano V. Management of hepatitis co-infected with hepatitis B virus and HIV. *Lancet Infect Dis* **2005**; 6:374–82.
- Koziel MJ, Peters MG. Viral hepatitis in HIV infection. *N Engl J Med* **2007**; 356:1445–54.
- Hattori J, Shiino T, Gatanaga H, et al. Trends in transmitted drug-resistant HIV-1 and demographic characteristics of newly diagnosed patients: nationwide surveillance from 2003 to 2008 in Japan. *Antivir Res* **2010**; 88:72–9.
- Fujisaki S, Yokomaku Y, Shiino T, et al. Outbreak of infections by hepatitis B virus genotype A and transmission of genetic drug resistance in patients coinfecting with HIV-1 in Japan. *J Clin Microbiol* **2011**; 49:1017–24.
- Thio CL, Seaberg EC, Skolasky RL, et al. HIV-1, hepatitis B virus, and risk of liver-related mortality in the Multicenter AIDS Cohort Study (MACS). *Lancet* **2002**; 360:1921–6.
- Thio CL. Hepatitis B and human immunodeficiency virus coinfection. *Hepatology* **2009**; 49:S138–45.
- Brook G, Main J, Nelson M, et al. British HIV association guidelines for the management of coinfection with HIV-1 and hepatitis B or C virus 2010. *HIV Med* **2010**; 11:1–30.
- Thompson MA, Aberg JA, Hoy JF, et al. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International AIDS Society–USA panel. *JAMA* **2012**; 308:387–402.
- Kinn S, Akhavan S, Agut H, et al. Performance of the DiaSorin LIAISON anti-HBs II for the detection of hepatitis B surface antibodies: comparison with the Abbott ARCHITECT anti-HBs assay. *J Clin Virol* **2011**; 50:297–302.
- Ollier L, Laffont C, Kechkekian A, et al. Detection of antibodies to hepatitis B core antigen using the Abbott ARCHITECT anti-HBc assay: analysis of borderline reactive sera. *J Virol Methods* **2008**; 154:206–9.
- Tadokoro K, Kobayashi M, Yamaguchi T, et al. Classification of hepatitis B virus genotypes by the PCR-invader method with genotype-specific probes. *J Virol Methods* **2006**; 138:30–9.
- Tadokoro K, Suzuki F, Kobayashi M, et al. Rapid detection of drug-resistant mutations in hepatitis B virus by the PCR-Invader assay. *J Virol Methods* **2011**; 171:67–73.
- Zoulim F, Locarnini S. Hepatitis B virus resistance to nucleos(t)ide analogues. *Gastroenterology* **2009**; 137:1593–608.
- Koibuchi T, Hitani A, Nakamura T, et al. Predominance of genotype A HBV in an HBV-HIV-1 dually positive population compared with an HIV-1-negative counterpart in Japan. *J Med Virol* **2001**; 64:435–40.
- Matsuura K, Tanaka Y, Hige S, et al. Distribution of hepatitis B virus genotypes with chronic infection in Japan shifting toward an increase of genotype A. *J Clin Microbiol* **2009**; 47:1476–83.
- Kobayashi M, Ikeda K, Arase Y, et al. Change of hepatitis B virus genotypes in acute and chronic infections in Japan. *J Med Virol* **2008**; 80:1880–4.
- Lin CL, Kao JH. The clinical implications of hepatitis B virus genotype: recent advances. *J Gastroenterol Hepatol* **2011**; 26:S123–30.
- Gandhi RT, Wurcel A, Lee H, et al. Isolated antibody to hepatitis B core antigen in human immunodeficiency virus type-1-infected individuals. *Clin Infect Dis* **2003**; 36:1602–5.
- French AL, Operskalski E, Peters M, et al. Isolated hepatitis B core antibody is associated with HIV and ongoing but not resolve hepatitis C virus infection in a cohort of US women. *J Infect Dis* **2007**; 195:1437–42.
- Perez-Rodriguez MT, Sopena B, Crespo M, et al. Clinical significance of “anti-HBc alone” in human immunodeficiency virus-positive patients. *World J Gastroenterol* **2009**; 15:1237–41.
- Koo YX, Tan DS, Tan IB, et al. “Anti-HBc alone” in human immunodeficiency virus-positive and immune-suppressed lymphoma patients. *World J Gastroenterol* **2009**; 15:3834–5.
- Gandhi RT, Wurcel A, Lee H, et al. Response to hepatitis B vaccine in HIV-1-positive subjects who test positive for isolated antibody to hepatitis B core antigen: implications for hepatitis B vaccine strategies. *J Infect Dis* **2005**; 191:1435–41.
- French AL, Lin MY, Evans CT, et al. Long-term serologic follow-up of isolated hepatitis B core antibody in HIV-infected and HIV-uninfected women. *Clin Infect Dis* **2009**; 49:148–54.
- Biggar RJ, Goedert JJ, Hoofnagle J. Accelerated loss of antibody to hepatitis B surface antigen among immunodeficient homosexual men infected with HIV. *N Engl J Med* **1987**; 316:630–1.
- Collier AC, Corey L, Murphy VL, et al. Antibody to human immunodeficiency virus (HIV) and suboptimal response to hepatitis B vaccination. *Ann Intern Med* **1988**; 109:101–5.
- Loke RH, Murray-Lyon IM, Coleman JC, et al. Diminished response to recombinant hepatitis B vaccine in homosexual men with HIV antibody: an indicator of poor prognosis. *J Med Virol* **1990**; 31:109–11.
- Tayal SC, Sankar KN. Impaired response to recombinant hepatitis B vaccine in asymptomatic HIV-infected individuals. *AIDS* **1994**; 8:558–9.
- del Pozo Balado Mdel M, Leal M, Mendez Lagares G, et al. Increased regulatory T cell counts in HIV-infected nonresponders to hepatitis B virus vaccine. *J Infect Dis* **2010**; 202:362–9.
- Fonseca MO, Pang LW, de Paula Cavalheiro N, et al. Randomized trial of recombinant hepatitis B vaccine in HIV-infected adult

- patients comparing a standard dose to a double dose. *Vaccine* **2005**; *23*: 2902–8.
36. Launay O, van der Vliet D, Rosenberg AR, et al. Safety and immunogenicity of 4 intramuscular double doses and 4 intradermal low doses vs standard hepatitis vaccine regimen in adults with HIV-1: a randomized controlled trial. *JAMA* **2011**; *305*:1432–40.
37. Flynn PM, Cunningham CK, Rudy B, et al. Hepatitis B vaccination in HIV-infected youth: a randomized trial of three regimens. *J Acquir Immune Defic Syndr* **2011**; *56*:325–32.
38. Wilson DP, Law MG, Grulich AE, et al. Relation between HIV viral load and infectiousness: a model-based analysis. *Lancet* **2008**; *372*: 314–20.
39. Cohen MS, Chen YQ, McCauley M, et al. Prevention of HIV-1 infection with early antiretroviral therapy. *N Engl J Med* **2011**; *365*: 493–505.
40. Grant RM, Lama JR, Anderson PL, et al. Preexposure chemoprophylaxis for HIV prevention in men who have sex with men. *N Engl J Med* **2010**; *363*:2587–99.

Restriction fragment mass polymorphism (RFMP) analysis based on MALDI-TOF mass spectrometry for detecting antiretroviral resistance in HIV-1 infected patients

J.-H. Lee^{1*}, A. Hachiya^{2,3*}, S.-K. Shin¹, J. Lee¹, H. Gatanaga³, S. Oka³, K. A. Kirby², Y. T. Ong², S. G. Sarafianos^{2,4}, W. R. Folk⁴, W. Yoo¹, S. P. Hong¹ and S.-O. Kim¹

1) Research and Development Center, GeneMatrix Inc., Seongnam, South Korea, 2) Christopher Bond Life Science Center, Department of Molecular Microbiology and Immunology, University of Missouri School of Medicine, Columbia, MO, USA, 3) AIDS Clinical Center, National Center for Global Health and Medicine, Tokyo, Japan and 4) Department of Biochemistry, University of Missouri, Columbia, MO, USA

Abstract

Viral genotype assessment is important for effective clinical management of HIV-1 infected patients, especially when access and/or adherence to antiretroviral treatment is reduced. In this study, we describe development of a matrix-assisted laser desorption/ionization-time of flight mass spectrometry-based viral genotyping assay, termed restriction fragment mass polymorphism (RFMP). This assay is suitable for sensitive, specific and high-throughput detection of multiple drug-resistant HIV-1 variants. One hundred serum samples from 60 HIV-1-infected patients previously exposed to nucleoside reverse transcriptase inhibitors (NRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs) and protease inhibitors (PIs) were analysed for the presence of drug-resistant viruses using the RFMP and direct sequencing assays. Probit analysis predicted a detection limit of 223.02 copies/mL for the RFMP assay and 1268.11 copies/mL for the direct sequencing assays using HIV-1 RNA Positive Quality Control Series. The concordance rates between the RFMP and direct sequencing assays for the examined codons were 97% (K65R), 97% (T69Ins/D), 97% (L74VI), 97% (K103N), 96% (V106AM), 97% (Q151M), 97% (Y181C), 97% (M184VI) and 94% (T215YF) in the reverse transcriptase coding region, and 100% (D30N), 100% (M46I), 100% (G48V), 100% (I50V), 100% (I54LS), 99% (V82A), 99% (I84V) and 100% (L90M) in the protease coding region. Defined mixtures were consistently and accurately identified by RFMP at 5% relative concentration of mutant to wild-type virus while at 20% or greater by direct sequencing. The RFMP assay based on mass spectrometry proved to be sensitive, accurate and reliable for monitoring the emergence and early detection of HIV-1 genotypic variants that lead to drug resistance.

Keywords: Drug, HIV-1, MALDI-TOF, resistance mutation, RFMP

Original Submission: 28 August 2012; **Revised Submission:** 4 January 2013; **Accepted:** 15 January 2013

Editor: G. Antonelli

Article published online: 11 March 2013

Clin Microbiol Infect 2013; **19**: E263–E270

10.1111/1469-0691.12167

Corresponding authors: S.-O. Kim or S. P. Hong, Research and Development Center, GeneMatrix Inc., Korea Bio Park, 694-1, Sampyung-dong, Bundang-gu, Seongnam-si, Gyeonggi-do, South Korea
E-mails: sookim@genematrix.net; sunphong@genematrix.net

*These authors contributed equally

Introduction

Highly active antiretroviral therapy (HAART) can dramatically suppress HIV-1 replication, improve immunological

response and extend a patient's lifespan. However, less than excellent adherence to HAART or conditions that result in reduced treatment efficacy leads to a higher risk of the emergence of antiretroviral (ARV) drug-resistant viral strains, which eventually leads to increased viral loads, poor immunological response and eventually treatment failure [1]. Especially, women who have received single-dose nevirapine to prevent mother-to-child HIV-1 transmission are at increased risk of virological failure as a result of the replication of low-abundance nevirapine-resistant variants when treated with a subsequent nevirapine-containing regimen [2]. Of importance in the effective management of HIV-1 infections is the timely and efficient detection of

drug-resistant viral strains and their specific mutations in a patient's clinical samples.

The guidelines for use of ARV drugs in HIV-1-infected adults and adolescents established by the US Department of Health and Human Services (DHHS) (<http://www.aidsinfo.nih.gov/guidelines>) recommend monitoring viral genotypic changes in patient samples and use of this information to determine which therapeutic regimens are most appropriate for the specific patient [3].

The restriction fragment mass polymorphism (RFMP) method is based on amplification and mass detection of oligonucleotides excised by type-III restriction enzyme digestion, using matrix-assisted laser desorption and ionization time of flight mass spectrometry (MALDI-TOF MS). RFMP-based drug-resistance testing and genotyping has been shown to be a sensitive, accurate and reliable method for clinical utility in many fields [4–12]. Especially important is that RFMP enables sensitive detection of mutations without population-based cloning and subsequent sequencing analysis [6].

In this study, we applied the RFMP assay for detection of mutations in the coding sequences for reverse transcriptase (RT) and protease (PR) of HIV-1 that engender resistance to nucleoside reverse transcriptase inhibitors (NRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs) and protease inhibitors (PIs). Compared with direct sequencing, RFMP is shown to be a sensitive and reliable method for genotypic testing of drug-resistance mutations in HIV-1 infected patients.

Materials and Methods

Specimens

A total of 100 plasma samples were collected from 60 HIV-1 infected patients who had received HAART (including NRTIs, NNRTIs and PIs) at the AIDS Clinical Center, National Center for Global Health and Medicine, Japan, between 1999 and 2009. Written informed consent was obtained from each participant, and the experimental protocol conformed to the ethical guidelines of the 1975 Declaration of Helsinki, as reflected in *a priori* approval (NCGM-H22-938) by the Ethics Committee of the AIDS Clinical Center, National Center for Global Health and Medicine, Japan. The demographic characteristics are summarized in Table 1.

HIV-1 performance panels

To assess the limit of detection ability of the RFMP assay, the HIV-1 RNA Positive Quality Control Series (ACCURUN® 315) obtained from SeraCare Life Sciences (Milford, MA, USA) was used to measure viral load in HIV-1 performance panels.

TABLE 1. Demographic characteristics of 60 HIV-1-infected patients

Characteristic	Value
Mean age in years (range)	42 (22–67)
No. male (%)	53 (88)
No. female (%)	7 (12)
Race (%)	
Asian	58 (97)
African	2 (3)
Risk factor for HIV (% of patients)	
Heterosexual	14 (23)
Homosexual	27 (45)
Haemophilia (infected blood products)	19 (32)
CDC clinical stage (%)	
A1/A2/A3	0/0/0
B1/B2/B3	17/23/13
C1/C2/C3	5/12/20
Unknown	10
Mean CD4 cell count (No. of cells/ μ L [range])	320 (12–759)
No. of unknown (%)	3 (5)
Mean HIV-1 RNA (No. of RNA copies/mL [range])	43 000 (50–1 200 000)
No. of unknown (%)	6 (10)
History of actual treatment (No. of patients)	
With NRTI	2
With NRTI plus NNRTI	23
With NRTI plus PI	21
With NNRTI plus PI	1
With NRTI plus NNRTI plus PI	33
With NRTI plus PI plus INI	1
With NRTI plus INI	2
With NRTI plus PI plus INI plus FI	1
With NRTI plus NNRTI plus PI plus INI plus FI	1
Interruption	15

NRTI, nucleoside reverse transcriptase inhibitor; lamivudine, abacavir, emtricitabine, tenofovir, stavudine, didanosine, zidovudine; NNRTI, non-nucleoside reverse transcriptase inhibitor; efavirenz, nevirapine; PI, protease inhibitor; atazanavir, ritonavir, lopinavir, darunavir, fosamprenavir, amprenavir, nelfinavir; INI, integrase inhibitor; raltegravir; FI, fusion inhibitor; enfuvirtide.

Construction of recombinant HIV-1 clones

Recombinant infectious HIV-1 clones with various mutations in the RT region were constructed using site-directed mutagenesis. Briefly, the mutations were introduced into the *Xma*I-*Nhe*I fragment (759 bp) of pTZNX1, which encodes Gly-15 to Ala-267 of HIV-1 RT (strain BH 10), by oligonucleotide-based mutagenesis [13]. The *Xma*I-*Nhe*I fragment was inserted into a pNL4-3-based plasmid, generating various molecular clones with the desired mutations. Each molecular clone (10^7 /mL as DNA) was transfected into human 293T cells (4×10^5 cells/100-mm-diameter dish) with Fugene transfection reagent (Roche Diagnosis, Basel, Switzerland). After 48 h, culture supernatants were harvested and stored at -80°C until use. Viral loads were determined using the COBAS® Amplicor HIV-1 Monitor Test, v1.5.

RNA extraction and cDNA amplification

HIV-1 RNA was extracted from 200 μ L of plasma using the High Pure Viral RNA Kit (Roche Diagnostics, Mannheim, Germany) according to manufacturer's instructions. Purified viral RNA was dissolved in 50 μ L elution buffer (nuclease-free, sterile, double distilled water). cDNA was synthesized using only the reverse transcription step component of the RNA PCR kit (TaKaRa, Otsu, Japan).

RFMP assay

PCR reactions were performed in 25 μ L reaction mixtures containing 20 mM Tris-HCl (pH 8.4), 50 mM KCl, 0.2 mM deoxynucleoside triphosphates (dNTPs), 10 pmol of primers and 0.4 U of Platinum Taq DNA polymerase (Invitrogen, Carlsbad, CA, USA). The initial denaturing phase of 5 min at 94°C was followed by a 35-cycle amplification phase containing a denaturation step at 94°C for 15 s, annealing step at 50°C for 15 s and elongation step at 72°C for 15 s, and completed with a final extension phase at 72°C for 5 min. For the RFMP analysis of codons 65, 69, 74, 103, 106, 151, 181, 184 and 215 in the HIV-1 RT region and codons 30, 46, 48, 50, 54, 82, 84 and 90 in the HIV-1 PR region, each of the forward and reverse primers contained the viral target sequence and the *FokI* recognition sequence *ggatg* (Table S1). Restriction enzyme digests were performed by mixing the PCR reaction with 10 μ L of buffer (50 mM potassium acetate, 20 mM Tris-acetate, 10 mM magnesium acetate and 1 mM dithiothreitol) and 1 U of *FokI* enzyme (New England Biolabs, Beverly, MA, USA). The reaction mixtures were incubated at 37°C for 2 h. Subsequently, the digest desalting and mass analysis were performed as described previously [14].

MALDI-TOF instrumentation and calibration

Mass spectra were acquired on a Biflex IV linear MALDI-TOF MS (Bruker Daltonics) workstation equipped with a 337 nm nitrogen laser and a nominal ion flight path length of 1.25 m. The samples were analyzed in the negative-ion mode with a total acceleration voltage of 20 kV, extraction voltage of 18.25 kV, laser attenuation of 55, and delayed extraction of long time delay mode. Typically, time-of-flight data from 20 to 50 individual laser pulses were recorded and averaged on a transient digitizer with a time base of 2 ns and delay of 24 000 ns, after which the averaged spectra were automatically converted to mass by the accompanying data-processing software. Using these settings the instrument typically provided mass accuracy of 40–80 ppm (10^{-6}), mass resolution of 1500–2000 and sensitivity of 10–50 fmol in the 2- to 6-kDa mass range for oligonucleotides. Oligonucleotide standards of 6mer (5'-ACGTAC-3'; 1762.2 Da) and 16mer (5'-ACGTACGTACGTACGT-3'; 4881.2 Da) with no terminal phosphate were used for mass calibration of the instrument. The presence of metal cations produces salt adducts, leading to reduced resolution and low sensitivity, so C18 reverse phase micro-column chromatography was used for desalting oligonucleotides. Non-homogeneous crystallization is obtained with the classic dried droplet preparation, and a search for a 'sweet spot' is required. Re-crystallization of sample DNAs on matrix-spotted anchorchip plates allowed robust formation of small single crystals.

Direct sequencing assay

To amplify the HIV-1 RT and PR regions for analysis by direct sequencing, PCR was performed with the following primers: 5'-AACAAATGGCCATTGACAGAAGAAA-3' (2614–2637 bp of HXB2), 5'-CTGTATGTCATTGACAGTCCAGCT-3' (3299–3323 bp of HXB2) for the RT region and 5'-CTTCCCTCAGATCACTCTTTGGCAA-3' (2248–2273 bp of HXB2), 5'-AGGGCTAATGGGAAAATTTAAAGT-3' (2238–2561 bp of HXB2) for the PR region. PCR products were sequenced using the BigDye Terminator (version 3.1) Cycle Sequencing kit and an ABI PRISM 310 Analyzer (Applied Biosystems, Foster City, CA, USA).

Statistics

A limit of detection test was performed by probit analysis to compare sensitivity between the RFMP and direct sequencing assays using the statistical package SAS (version 8; SAS Institute Inc., Cary, NC, USA).

Results

RFMP assay strategy

The RFMP assay is based on mass spectrometric analysis of small DNA fragments that include sites of mutation (Fig. S1). The first step requires PCR amplification with forward and reverse primers that introduce the *FokI* enzyme site, *ggatg* (Table S1). The diagnostic fragments released by enzymatic digestion consist of various sizes from 8 nt oligomers to 14 nt oligomers for nine codons in the RT region and eight codons in the PR region, leading to facile identification of sequence variation by mass spectrum analysis. Genotypic analysis of mutations at codons 65, 69, 74, 103, 106, 151, 181, 184 and 215 in the RT region and codons 30, 46, 48, 50, 54, 82, 84 and 90 in the PR region, as assessed by the RFMP assay, was determined for 100 plasma samples. The RFMP results showed distinct peaks relevant to each codon, with the mass values for each diagnostic fragment being exactly as predicted (Supplementary Material Table S2).

Estimation of limit of detection and ability to detect mixed genotype populations

The detection limit was estimated using replicates of each of nine dilutions of HIV-1 RNA Positive Quality Control Series (ACCURUN[®] 315) material ranging between 10 and 5000 copies/ml. Analysis of various calibrated HIV-1 RNA dilution series determined the lower detection limit to be 223.02 copies/mL for the RFMP and 1268.11 copies/mL for the direct sequencing assays by probit analysis. The probit analysis predicts a 95% CI: 132.64–693.00 for the RFMP and 863.09–3656.80 for the direct sequencing assays (Table 2).

TABLE 2. Limit of detection of the RFMP and direct sequencing assays

HIV-1 RNA copies/ml	No. tested	RFMP		Direct sequencing	
		No. detected	Per cent detected	No. detected	Per cent detected
5000	10	10	100%	10	100%
2500	10	10	100%	10	100%
1000	10	10	100%	9	90%
500	10	10	100%	4	40%
250	10	10	100%	1	10%
100	10	8	80%	0	0%
50	10	3	30%	0	0%
25	10	1	10%	0	0%
10	10	1	10%	0	0%
Limit of detection		223.02 copies/mL (95% CI, 132.64–693.00)		1268.11 copies/mL (95% CI, 863.09–3656.80)	

Defined dilutions of HIV-1 RNA Positive Quality Control Series were made from 10 copies to 5000 copies/mL and limit of detection abilities were calculated by probit analysis at a 95% detection level.

To evaluate the ability of the RFMP assay to determine small amounts of mutant virus in mixed populations, assays were performed with recombinant HIV-1 clones composed of different ratios of wild-type (K103 in the RT region) and mutant genotypes (N103 in the RT region). Defined mixtures were prepared with the following percentages of K103N mutant virus in the total virus population: 100%, 50%, 20%, 10%, 5% and 1%. The K103N mutant virus could be detected in concentrations as low as 5% of the total virus by RFMP, whereas direct sequencing assays were able to detect mutant virus only when present in 20% or more of the total virus population (Fig. 1).

Comparison of RFMP with direct sequencing analyses

All 100 clinical samples from 60 patients were analysed by the RFMP and direct sequencing assays for the presence of drug resistance-related mutations: nine codons in the RT region and eight codons in the PR region of the HIV-1 *pol* gene (a total of 17 codons).

The overall concordance rates between RFMP and sequencing assays were excellent, irrespective of PR and RT regions (Table 3). Concordance rates in the RT region were 97% (97/100) at codons 65, 69, 74, 103, 151, 181 and 184, 96% (96/100) at codon 106, and 94% (94/100) at codon 215 (Fig. 2a).




FIG. 1. Evaluation of the sensitivity of the RFMP assay for detection of minor amounts of virus with a defined mixture of K103N. The MALDI-TOF MS spectra and direct sequencing chromatograms shown are representative of experiments repeated three times using mixed populations of wild-type (K103) and NNRTI mutant (N103). The wild-type plasmids were mixed with mutant type at various ratios as follows: (a) 100%, (b) 50%, (c) 20%, (d) 10%, (e) 5% and (f) 1%. Molecular masses of 3133.0 and 3157.0 correspond to N103 and K103, respectively. AI is absolute intensity; m/z is mass-to-charge ratio.

TABLE 3. Comparison of the results obtained by the RFMP and direct sequencing assays in 100 clinical specimens

pol gene	Codon	Discordant (n)	Concordant (n)	Compatible (n)		Amino acid	
				RFMP only	Direct sequencing only	RFMP	Direct sequencing
RT	65	3	97	—	—		
	69	3	97	—	—		
	74	3	97	—	—		
	103	3	97	—	—		
	106	3	96	1	—	V/A	V
	151	3	97	—	—		
	181	3	97	—	—		
	184	3	97	—	—		
	215	3	94	3	—	T/F T/Y T/Y	T/F or I/S or T/I/F T/Y or N/S or T/S/Y T/Y or N/S or T/S/Y
	PR	30	—	100	—	—	
46		—	100	—	—		
48		—	100	—	—		
50		—	100	—	—		
54		—	100	—	—		
82		—	99	1	—	V/A	V
84		—	99	1	—	I/V	V
90		—	100	—	—		

RT, reverse transcriptase; PR, protease.

Similarly, concordance rates in the PR region were 100% (100/100) at codons 30, 46, 48, 50, 54 and 90, and 99% (99/100) at codons 82 and 84 in the PR region (Fig. 2b). Both assays showed identical base substitution and amino acid composition in these positions. Rate of compatible cases were observed 1% (1/100) at codons 106, and 3% (3/100) at codon 215 in the RT region and 1% (1/100) at codons 82 and 84 in the PR region, respectively. Three samples (mixed-type) at codon 215 containing double mutations in a single codon were identified only by RFMP, as a result of the inability of the direct sequencing assay to determine the variants present in the clinical samples. As shown in Fig 2(c), 215T/Y mixtures detected by RFMP could be scored as 215T (ACC) plus 215Y (TAC), or 215N (AAC) plus 215S (TCC), or 215T (ACC) plus 215S (TCC) plus 215Y (TAC), by direct sequencing. A compatible single nucleotide mixture at one position was observed in three samples at three codons (codon 106 in the RT; codon 82, 84 in the PR), respectively. Of these, the RFMP assay detected more mixed samples than the direct sequencing assay. The details of mixtures detected by both assays are shown in Table 3. Discordances between the two assays occurred for three samples at RT region codons, which had undetectable viral loads; the correct viral genotypes were identified only by RFMP assay.

Discussion

ARV drug-resistance is a major obstacle in the effective clinical management of HIV-1-infected patients [3, 15] and therapeutic strategies must maximize the early detection of drug resistance mutations. Having a sensitive genotyping assay that can detect with high accuracy and reliability, drug-resistance

mutations that emerge during HAART can be very important for the optimization of ARV regimens, improvement of patient treatment and outcome of therapy. The RFMP assay has been demonstrated to be a sensitive, accurate and reliable method for genotyping and detecting drug-resistance mutations in several viruses, including hepatitis and papillomavirus [4, 6, 8–12].

In the present study, we validated use of the MALDI-TOF MS-based RFMP assay to detect oligonucleotides containing 8–14 nucleotides for codons implicated with ARV drug-resistance mutations. Specifically, we established successful detection at codons 65, 69, 74, 103, 106, 151, 181, 184 and 215 in the HIV-1 RT coding region, and 30, 46, 48, 50, 54, 82, 84 and 90 in the HIV-1 PR coding region. These codons address resistance to all approved NRTI and NNRTI inhibitors [16–18]: mutation at RT codon 65 (K65R) confers resistance to tenofovir, didanosine and abacavir; RT mutation L74V confers resistance to didanosine and abacavir; the K103N RT mutation engenders resistance to the NNRTIs efavirenz and nevirapine; the Q151M RT mutation causes resistance to AZT, D4T, didanosine and abacavir through the decreased incorporation mechanism; Y181C causes resistance to nevirapine, etravirine and rilpivirine; M184V/I confers resistance to 3TC and FTC, and also affects resistance to rilpivirine; and finally, T215Y causes resistance to AZT and D4T through the excision mechanism. In addition, mutations at the 30, 46, 48, 50, 54, 82, 84 and 90 sites of the HIV-1 PR coding region cause resistance to all known protease inhibitors: specifically, D30N causes resistance to nelfinavir; M46I/L causes resistance to nelfinavir and indinavir; G48V/M causes resistance to atazanavir, nelfinavir and saquinavir; I50L/V causes resistance to atazanavir, darunavir, fosamprenavir and lopinavir; and mutations at




FIG. 2. Comparison of the RFMP and direct sequencing assays for detection of mixed genotypes. Sera were taken from patients infected with HIV-1 carrying ARV drug-resistant mutations and examined by the RFMP and sequencing assays. (a) For codon 103, molecular masses of 3088.0/3157.0 and 3113.0/3133.0 represent Lys (AAA) and Asn (AAC), respectively. (b) For codon 46, molecular masses of 2988.0/3164.0 and 3003.0/3148.0 represent Met (ATG) and Ile (ATA), respectively. (c) For codon 215, molecular masses of 3740.4/3796.4 and 3724.4/3811.4 represent Thr (ACC) and Tyr (TAC), respectively. Each codon was indicated by a red box in the sequencing chromatogram. AI is absolute intensity; m/z is mass-to-charge ratio.

residues 54, 82, 84 and 90 cause resistance to all protease inhibitors to a varying extent (<http://hivdb.stanford.edu/DR/PIResiNote.html>).

The detection limit of the RFMP assay was determined to be 223.02 copies/mL and able to identify a minority mutant at a concentration as low as 5% of the circulating mixed populations, whereas the detection limit of the direct sequencing method was 1268.11 copies/mL and able to detect variants only when present at >20% of the total population (Table 2 and Fig. 1). A clear correlation was observed between peak ratios and relative genotype concentration of mixed populations.

The performance of the RFMP assay in detecting mutations related to ARV drug-resistance was compared with direct sequencing assays for 100 clinical samples from 60 HIV-1-infected patients who experienced HAART therapy. The RFMP assay successfully identified genotypic changes at all 17 codons tested in clinical samples. Compared with direct sequencing, the RFMP assay exhibited 96.6% concordance in the RT region and 99.8% concordance in the PR region (Table 3 and Fig 2). The PR data were not significantly more concordant than the RT data (the difference was only two cases). The reason for the slight difference is not clear, thus further investigation is required with more samples. Notably, the RFMP assay

outperformed direct sequencing for the detection of single and double mutations in compatible samples. The RFMP assay detected 1% (1/100) more mutant viruses in codons 106, 82 and 84 and 3% (3/100) more mutant viruses in codon 215. All discordances between the two assays were due to the inability of the direct sequencing assay to identify the residues at nine RT codons in three patients. Hence, the discrepancy among the two assays may be due to a lower sensitivity of direct sequencing.

HAART can clearly extend the life expectancy of HIV-1 patients. However, as adherence is usually imperfect, continuous ARV drug-resistance testing can be an important management tool. There are two major methods of assessing ARV drug-resistance: phenotypic assays and genotypic assays that provide complementary information and be preferable to the other [19,20]. Current treatment guidelines define as treatment failure the detection of more than 200 copies/mL of HIV-1 RNA [3]. However, existing genotypic assays, such as direct sequencing and the reverse hybridization-based assay, do not reliably detect fewer than 1000 copies/ml of HIV-1 RNA, nor do they enable detection of sequence variants present at <20% of minority variants in mixed populations [21,22]. This performance does not allow facile interpretation of ratios between multiple virus mixtures, especially when a double mutation is present in a single codon. While the reverse hybridization-based assay has somewhat higher sensitivity than direct sequencing it gives rise to false-positive and false-negative results more frequently than direct sequencing [21–26].

With the advent of Next Generation Sequencing methods it is possible to detect, by 454 pyrosequencing or an Illumina Genome Analyzer, minor viral variants whose prevalence is <1%. 454 pyrosequencing, also called ultra-deep pyrosequencing (UDPS), relies on fixing nebulized and adapter-ligated DNA fragments to small DNA-capture beads in a water-in-oil emulsion. The DNA fixed to these beads is then amplified by PCR. An advantage of UDPS in the case of viral genomic analysis is that it yields long sequence information for each sample (average ~800 bases). However, UDPS has some technical challenges. A major limitation of the UDPS relates to resolution of homopolymer-containing DNA segments, such as AAA and GGG [27]. Because there is no terminating moiety preventing multiple consecutive incorporations at a given cycle, pyrosequencing relies on the magnitude of light emitted to determine the number of repetitive bases. This is prone to greater error than misincorporation. Hence, the dominant error type for the 454 platform is insertion-deletion, rather than substitution. The decrease in single read accuracy from 99.4% for test fragments to 96% for genomic libraries is primarily due to a lack of clonality in a fraction of the genomic templates in the emulsion [28]. Moreover, based on current list prices for the UDPS, the

current cost for all the reagents, including the picotiter plate, library preparation kits and emulsion PCR kits, to perform a single experiment is \$1000–7000 [29].

By combining the merits of unique assay chemistry and the mature nature of MALDI-TOF mass spectrometry, the RFMP assay can be used to screen for HIV drug-resistance mutations in a robust high-throughput manner (e.g. 96 samples can be analysed in 3 h with Bruker software (flexcontrol 3.0), which is faster than existing hybridization or sequencing-based methods). In terms of cost-effectiveness, the direct cost per test (reagents and labour) of the RFMP assay can be <\$30, including viral DNA extraction, PCR, restriction digestion, desalting and matrix for mass analysis, which is cheaper than the sequencing or hybridization assays that are c. \$50–100 per test. These costs do not include the equipment costs, which are slightly greater for the RFMP method. However, with the advent of many diagnostic assays operating on a mass spectrometer platform, such as clinical genotyping and microorganism identification, and the gradual spread of compact mass spectrometers into laboratories as medical devices (i.e. Bruker Microflex), the burden of the amortization cost should be substantially reduced [30]. A limitation of the RFMP assay is that it determines only molecular mass and is thus applicable only to known resistance mutations and may fail to detect DNA sequence changes that do not affect molecular mass. Moreover, the occurrence of a novel resistance mutation with sequence alterations that cause a deviation from the predicted molecular mass pattern should be addressed by periodic updating of mass patterns for unambiguous result interpretation from up-to-date HIV databases.

We demonstrate here that the mass spectrometry-based RFMP assay is highly sensitive and able to successfully detect HIV-1 carrying drug-resistant mutations that are present in <5% of the total virus population. Hence, this assay can be used for the efficient assessment of genotype dynamics of viral quasi-species. Therefore, this simple procedure can be easily adapted to a high-throughput format, should enable earlier detection of drug-resistant viruses and help elucidate mechanisms of HIV-1 resistance, as well as guide and optimize treatment decisions.

Acknowledgements

We thank Doctors Yukiko Takahashi and Fujie Negishi for clinical sample preparation.

Transparency Declaration

This work was supported by grants from the Ministry of Knowledge and Economy, Bilateral International Collaborative

R&D Program, Republic of Korea, and in part by NIH grants AI076119, AI094715, AI074389, AI073975 and AI077424.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Figure S1. Schematic representation of the RFMP strategy.

Table S1. Primers used in PCR amplification for the RFMP assay.

Table S2. Expected masses of oligonucleotides resulting from restriction enzyme digestion of PCR products.

References

- Chen TK, Aldrovandi GM. Review of hiv antiretroviral drug resistance. *Pediatr Infect Dis J* 2008; 27: 749–752.
- Boltz VF, Zheng Y, Lockman S et al. Role of low-frequency hiv-1 variants in failure of nevirapine-containing antiviral therapy in women previously exposed to single-dose nevirapine. *Proc Natl Acad Sci USA* 2011; 108: 9202–9207.
- Thompson MA, Aberg JA, Hoy JF et al. Antiretroviral treatment of adult hiv infection: 2012 recommendations of the international antiviral society-USA panel. *JAMA* 2012; 308: 387–402.
- Hong SP, Shin SK, Lee EH et al. High-resolution human papillomavirus genotyping by maldi-tof mass spectrometry. *Nat Protoc* 2008; 3: 1476–1484.
- Oh HB, Kim SO, Cha CH et al. Identification of hepatitis c virus genotype 6 in korean patients by analysis of 5' untranslated region using a matrix assisted laser desorption/ionization time of flight-based assay, restriction fragment mass polymorphism. *J Med Virol* 2008; 80: 1712–1719.
- Cha CK, Kwon HC, Cheong JY et al. Association of lamivudine-resistant mutational patterns with the antiviral effect of adefovir in patients with chronic hepatitis b. *J Med Virol* 2009; 81: 417–424.
- Cha YS, Choi SH, Lee JH et al. Analysis of tpox short tandem repeat locus with matrix-associated laser desorption/ionization time-of-flight-based restriction fragment mass polymorphism assay. *Anal Biochem* 2011; 412: 79–84.
- Cho SW, Koh KH, Cheong JY et al. Low efficacy of entecavir therapy in adefovir-refractory hepatitis b patients with prior lamivudine resistance. *J Viral Hepatitis* 2010; 17: 171–177.
- Choe WH, Hong SP, Kim BK et al. Evolution of hepatitis b virus mutation during entecavir rescue therapy in patients with antiviral resistance to lamivudine and adefovir. *Antivir Ther* 2009; 14: 985–993.
- Han KH, Hong SP, Choi SH et al. Comparison of multiplex restriction fragment mass polymorphism and sequencing analyses for detecting entecavir resistance in chronic hepatitis b. *Antivir Ther* 2011; 16: 77–87.
- Kim SS, Cheong JY, Lee D et al. Adefovir-based combination therapy with entecavir or lamivudine for patients with entecavir-refractory chronic hepatitis b. *J Med Virol* 2012; 84: 18–25.
- Lee JM, Kim HJ, Park JY et al. Rescue monotherapy in lamivudine-resistant hepatitis b e antigen-positive chronic hepatitis b: Adefovir versus entecavir. *Antivir Ther* 2009; 14: 705–712.
- Kodama EI, Kohgo S, Kitano K et al. 4'-ethynyl nucleoside analogs: potent inhibitors of multidrug-resistant human immunodeficiency virus variants in vitro. *Antimicrob Agents Chemother* 2001; 45: 1539–1546.
- Kim HS, Han KH, Ahn SH et al. Evaluation of methods for monitoring drug resistance in chronic hepatitis b patients during lamivudine therapy based on mass spectrometry and reverse hybridization. *Antivir Ther* 2005; 10: 441–449.
- Tsibris AM, Hirsch MS. Antiretroviral therapy in the clinic. *J Virol* 2010; 84: 5458–5464.
- Singh K, Marchand B, Kirby KA, Michailidis E, Sarafianos SG. Structural aspects of drug resistance and inhibition of hiv-1 reverse transcriptase. *Viruses* 2010; 2: 606–638.
- Sarafianos SG, Marchand B, Das K et al. Structure and function of hiv-1 reverse transcriptase: molecular mechanisms of polymerization and inhibition. *J Mol Biol* 2009; 385: 693–713.
- Tang MW, Shafer RW. Hiv-1 antiretroviral resistance: scientific principles and clinical applications. *Drugs* 2012; 72: e1–e25.
- Underwood MR, Ross LL, Irlbeck DM et al. Sensitivity of phenotypic susceptibility analyses for nonthymidine nucleoside analogues conferred by k65r or m184v in mixtures with wild-type hiv-1. *J Infect Dis* 2009; 199: 84–88.
- Qari SH, Winters M, Vandamme AM, Merigan T, Heneine W. A rapid phenotypic assay for detecting multiple nucleoside analogue reverse transcriptase inhibitor-resistant hiv-1 in plasma. *Antivir Ther* 2002; 7: 131–139.
- Sen S, Tripathy SP, Paranjape RS. Antiretroviral drug resistance testing. *J Postgrad Med* 2006; 52: 187–193.
- Garcia-Gonzalez C, Garcia-Bujalance S, Ruiz-Carrascoso G et al. Detection and quantification of the k103n mutation in hiv reverse transcriptase by pyrosequencing. *Diagn Microbiol Infect Dis* 2012; 72: 90–96.
- Clevenbergh P, Durant J, Halfon P et al. Persisting long-term benefit of genotype-guided treatment for hiv-infected patients failing haart. The viradapt study: week 48 follow-up. *Antivir Ther* 2000; 5: 65–70.
- Cortez KJ, Maldarelli F. Clinical management of hiv drug resistance. *Viruses* 2011; 3: 347–378.
- Erali M, Page S, Reimer LG, Hillyard DR. Human immunodeficiency virus type 1 drug resistance testing: a comparison of three sequence-based methods. *J Clin Microbiol* 2001; 39: 2157–2165.
- Garcia-Bujalance S, de Guevara CL, Gonzalez-Garcia J, Arribas JR, Gutierrez A. Comparison between sequence analysis and a line probe assay for testing genotypic resistance of human immunodeficiency virus type 1 to antiretroviral drugs. *J Clin Microbiol* 2005; 43: 4186–4188.
- Rothberg JM, Leamon JH. The development and impact of 454 sequencing. *Nat Biotechnol* 2008; 26: 1117–1124.
- Margulies M, Egholm M, Altman WE et al. Genome sequencing in microfabricated high-density picolitre reactors. *Nature* 2005; 437: 376–380.
- Hert DG, Fredlake CP, Barron AE. Advantages and limitations of next-generation sequencing technologies: a comparison of electrophoresis and non-electrophoresis methods. *Electrophoresis* 2008; 29: 4618–4626.
- Nagy E, Maier T, Urban E, Terhes G, Kostrzewa M. Species identification of clinical isolates of bacteroides by matrix-assisted laser-desorption/ionization time-of-flight mass spectrometry. *Clin Microbiol Infect* 2009; 15: 796–802.

Urinary beta-2 microglobulin and alpha-1 microglobulin are useful screening markers for tenofovir-induced kidney tubulopathy in patients with HIV-1 infection: a diagnostic accuracy study

Takeshi Nishijima · Takuro Shimbo · Hirokazu Komatsu · Misao Takano · Junko Tanuma · Kunihisa Tsukada · Katsuji Teruya · Hiroyuki Gatanaga · Yoshimi Kikuchi · Shinichi Oka

Received: 26 November 2012 / Accepted: 14 February 2013 / Published online: 7 March 2013
© Japanese Society of Chemotherapy and The Japanese Association for Infectious Diseases 2013

Abstract Kidney tubulopathy is a well-known adverse event of antiretroviral agent tenofovir. A cross-sectional study was conducted to compare the diagnostic accuracy of five tubular markers, with a collection of abnormalities in these markers as the reference standard. The study subjects were patients with HIV-1 infection on ritonavir-boosted darunavir plus tenofovir/emtricitabine with suppressed viral load. Kidney tubular dysfunction (KTD) was predefined as the presence of at least three abnormalities in the following five parameters: β 2-microglobulinuria (β 2M), α 1-microglobulinuria (α 1M), high urinary *N*-acetyl- β -D-glucosaminidase (NAG), fractional excretion of phosphate (FE_{IP}), and fractional excretion of uric acid (FE_{UA}). Receiver operating characteristic curves and areas under the curves (AUC) were estimated, and the differences between the largest AUC and each of the other AUCs were tested using a nonparametric method. The cutoff value of each tubular marker was determined using raw data of 100 % sensitivity

with maximal specificity. KTD was diagnosed in 19 of the 190 (10 %) patients. The AUCs (95 % CIs) of each tubular marker were β 2M, 0.970 (0.947–0.992); α 1M, 0.968 (0.944–0.992); NAG, 0.901 (0.828–0.974); FE_{IP} , 0.757 (0.607–0.907), and FE_{UA} , 0.762 (0.653–0.872). The AUCs of β 2M and α 1M were not significantly different, whereas those of the other three markers were smaller. The optimal cutoff values with 100 % sensitivity were 1,123 μ g/gCr (β 2M, specificity 89 %), 15.4 mg/gCr (α 1M, specificity 87 %), 3.58 U/gCr (NAG, specificity 46 %), 1.02 % (FE_{IP} , specificity 0 %), and 3.92 % (FE_{UA} , specificity 12 %). Urinary β 2M and α 1M are potentially suitable screening tools for tenofovir-induced KTD. Monitoring either urinary β 2M or α 1M should be useful in early detection of tenofovir nephrotoxicity.

Keywords Tenofovir · Kidney tubular dysfunction · Screening · Urinary beta-2 microglobulin · Urinary alpha-1 microglobulin · HIV-1 infection

T. Nishijima · M. Takano · J. Tanuma · K. Tsukada · K. Teruya · H. Gatanaga (✉) · Y. Kikuchi · S. Oka
AIDS Clinical Center, National Center for Global Health and Medicine, 1-21-1, Toyama, Shinjuku,
Tokyo 162-0052, Japan
e-mail: hingatana@acc.ncgm.go.jp

T. Nishijima · H. Gatanaga · S. Oka
Center for AIDS Research, Kumamoto University,
Kumamoto, Japan

T. Shimbo
Department of Clinical Research and Informatics, International
Clinical Research Center, National Center for Global Health
and Medicine, 1-21-1, Toyama, Shinjuku, Tokyo, Japan

H. Komatsu
Department of Community Care, Saku Central Hospital,
Nagano, Japan

Introduction

Tenofovir disoproxil fumarate (TDF), a prodrug of tenofovir, is a widely used nucleotide reverse transcriptase inhibitor (NRTI) as part of the initial antiretroviral therapy for patients with human immunodeficiency (HIV) infection, as well as in treatment of hepatitis B virus infection [1–4]. Tenofovir is excreted through the kidney by glomerular filtration and active tubular secretion. Renal proximal tubular damage is a well-known adverse effect of tenofovir, which sometimes leads to diabetes insipidus, Fanconi syndrome, and acute renal failure [5–7], and patients on TDF are more likely to develop renal dysfunction than those on other NRTIs [8, 9]. Of note, low

bone mineral density, another adverse effect of TDF, sometimes occurs as a result of kidney proximal tubulopathy associated with phosphate wasting and increased bone turnover [10, 11].

The risk of clinically relevant tenofovir-induced nephrotoxicity is considered to be relatively low [9, 12, 13]. However, tenofovir renal safety has not been confirmed in the long term. Importantly, both HIV infection and hepatitis B infection require long-term treatment, especially for HIV infection, where the treatment is lifelong. In tenofovir-induced nephrotoxicity, tubular dysfunction usually precedes the decline in glomerular filtration rate (GFR), suggesting that tubular markers are more sensitive than estimated GFR (eGFR) calculated with serum creatinine in screening for tenofovir nephrotoxicity [9, 14–16]. Thus, it is important to identify a renal tubular marker with high sensitivity to screen for tenofovir-induced KTD.

There is no gold standard for the diagnosis of tenofovir-induced kidney tubular dysfunction (KTD). Previous reports usually applied a collection of abnormalities in tubular function parameters as diagnostic criteria [17, 18]. However, the criteria used in each study differ, and it is often cumbersome and costly to monitor multiple renal tubular markers in practice. To our knowledge, there are no studies that have compared the diagnostic accuracy of various tubular markers for tenofovir-induced KTD. Furthermore, several pathological conditions are associated with KTD, such as active infection, diabetic nephropathy, and preexisting renal impairment [19, 20], making it difficult to evaluate KTD induced exclusively by tenofovir.

Based on the foregoing background, we designed the present study to identify the best screening marker(s) of tubular dysfunction in tenofovir-induced KTD, using a collection of abnormalities in kidney tubular markers as the reference standard.

Patients and methods

Ethics statement

This study was approved by the Human Research Ethics Committee of the National Center for Global Health and Medicine, Tokyo, Japan. Each patient in this study provided a written informed consent for publication of clinical data for research purpose. The study was conducted according to the principles expressed in the Declaration of Helsinki.

Study design

This study was designed and reported according to the recommendations of the standards for reporting of diagnostic accuracy (STARD) statement [21]. We performed a single-

center cross-sectional study to compare the diagnostic accuracies of various kidney tubular markers for tenofovir-induced KTD, with a collection of abnormalities in these markers as the reference standard, to identify the best screening marker in tenofovir-treated Japanese patients with HIV infection.

Study subjects

The study population has been previously reported [22]. The study enrolled consecutive Japanese patients with HIV infection, aged >17 years, with HIV-1 viral load <200 copies/ml, and on at least 4-week treatment with once-daily ritonavir (100 mg)-boosted darunavir (800 mg) plus fixed-dose tenofovir (300 mg)/emtricitabine (200 mg), seen at our clinic between October 1, 2011 and March 31, 2012. The exclusion criteria were (1) active infection, (2) malignancy, (3) diabetes mellitus, defined by the use of glucose-lowering agents or fasting plasma glucose >126 mg/dl or plasma glucose >200 mg/dl on two different days, (4) alanine aminotransferase 2.5 times more than the upper limit of normal, (5) estimated glomerular filtration rate (eGFR) calculated by Cockcroft–Gault equation of <50 ml/min [creatinine clearance = [(140 – age) × weight (kg)] / (serum creatinine × 72)(×0.85 for females)] [23], and (6) patients who did not sign the consent form.

Measurements

Blood and spot urine samples were collected on the same day (either on the day of enrollment or on the next visit), together with body weight measurement. The blood samples were used to measure serum creatinine, serum uric acid, serum phosphate, CD4 count, and C-reactive protein (CRP); urine samples were used to measure phosphate, uric acid, creatinine, β 2-microglobulin (β 2M), α 1-microglobulin (α 1M), and *N*-acetyl- β -D-glucosaminidase (NAG). The values of β 2M, α 1M, and NAG measured in the urine samples were expressed relative to urinary creatinine of 1 g/l (g Cr) [24].

Urinary concentrations of β 2M and α 1M were measured with a latex aggregation assay kit (β 2M, BMG-Latex X1 “Seiken”; Denka Seiken, Niigata, Japan; α 1M, Eiken α 1M-III; Eiken Chemical, Tokyo, Japan), and those of NAG by colorimetric assay of enzyme activity with 6-methyl-2-pyridyl-*N*-acetyl-1-thio- β -D-glucosaminide as substrate (Nittobo Medical, Tokyo, Japan).

Definition of renal proximal tubular dysfunction

KTD was predefined as the presence of at least three abnormalities in the following five parameters: fractional excretion of phosphate (FE_{IP}) {[urine phosphate × serum creatinine] / (serum phosphate × urine creatinine)} × 100} > 18 %,