

杉浦 互

1. Yotsumoto M, Shinozawa K, Yamamoto Y, Sugiura W, Miura T, and Fukutake K. Mutations to the probe of Cobas TaqMan HIV-1 ver. 1.0 assay causing undetectable viral load in a patient with acute HIV-1 infection. *J Infect Chemother* 17(6): 863-865, 2011.
2. Yoshida I, Sugiura W, Shibata J, Ren F, Yang Z, and Tanaka H. Change of positive selection pressure on HIV-1 envelope gene inferred by early and recent samples. *PLoS One* 6(4): e18630, 2011.
3. Ibe S, and Sugiura W. Clinical significance of HIV reverse transcriptase inhibitor-resistant mutations. *Future Microbiol* 6(3): 295-315, 2011.
4. Shibata J, Sugiura W, Odee H, Iwatani Y, Sato H, Tsang H, Matsuda M, Hasegawa N, Ren F, and Tanaka H. Within-host co-evolution of Gag P453L and protease D30 N/N88D demonstrates virological advantage in a highly protease inhibitor-exposed HIV-1 case. *Antivir Res* 90(1): 33-41, 2011.
5. Fujisaki S, Yokomaku Y, Shiino T, Koibuchi T, Hattori J, Ibe S, Iwatani Y, Iwamoto A, Shirasaka T, Hamaguchi M, and Sugiura W. Outbreak of hepatitis B virus genotype A and transmission of genetic drug resistance in cases coinfecting with HIV-1 in Japan. *J Clin Microbiol* 49(3): 1017-1024, 2011.
6. Bunupuradah T, Imahashi M, Iampornsin T, Matsuoka K, Iwatani Y, Puthanakit T, Ananworanich J, Sophonphan J, Mahanontharit A, Naoe T, Vonthanak S, Phanuphak P, Sugiura W, and On Behalf Of The Predict Study Team. Association of APOBEC3G genotypes and CD4 decline in Thai and Cambodian HIV-infected children with moderate immune deficiency. *AIDS Res Ther* 9(1): 34, 2012.
7. Tsuzuki T, Iwase H, Shimada M, Hirashima N, Hibino Y, Ryuge N, Saito M, Tamaki D, Kamiya A, Yokoi M, Yokomaku Y, Fujisaki S, Sugiura W, and Goto H. Clinical evaluation of peginterferon alpha plus ribavirin for patients co-infected with HIV and HCV at Nagoya Medical Center. *The Jpn J Gastroenterol* 109(7): 1186-1196, 2012.
8. Ode H, Nakashima M, Kitamura S, Sugiura W, and Sato H. Molecular dynamics simulation in virus research. *Front Microbiol* 3: 258, 2012.
9. Miyamoto T, Nakayama EE, Yokoyama M, Ibe S, Takehara S, Kono K, Yokomaku Y, Pizzato M, Luban J, Sugiura W, Sato H, and Shioda T. The Carboxyl-Terminus of Human Immunodeficiency Virus Type 2 Circulating Recombinant form 01_AB Capsid Protein Affects Sensitivity to Human TRIM5 α . *PloS one* 7(10): e47757, 2012.
10. Matsunaga S, Sawasaki T, Ode H, Morishita R, Furukawa A, Sakuma R, Sugiura W, Sato H, Katahira M, Takaori-Kondo A, Yamamoto N, Ryo A. Molecular and enzymatic characterization of XMRV protease by a cell-free proteolytic analysis. *J Proteomics* 75(15): 4863-4873, 2012.

11. Kitamura S, Ode H, Nakashima M, Imahashi M, Naganawa Y, Kurosawa T, Yokomaku Y, Yamane T, Watanabe N, Suzuki A, Sugiura W, and Iwatani Y. The APOBEC3C crystal structure and the interface for HIV-1 Vif binding. *Nat Struct Mol Biol* 19(10): 1005-1010, 2012.
12. Hirano A, Ikemura K, Takahashi M, Shibata M, Amioka K, Nomura T, Yokomaku Y, and Sugiura W. Short communication: lack of correlation between UGT1A1*6, *28 genotypes, and plasma raltegravir concentrations in Japanese HIV type 1-infected patients. *AIDS Res Hum Retroviruses* 28(8): 776-779, 2012.
13. Jahanbakhsh F, Ibe S, Hattori J, Monavari SHR, Matsuda M, Maejima M, Iwatani Y, Memarnejadian A, Keyvani H, Azadmanesh K, and Sugiura W. Molecular epidemiology of HIV-1 infection in Iran: genomic evidence of CRF35_AD predominance and CRF01_AE infection among individuals associated with injection drug use. *AIDS Res Hum Retroviruses* 29: 198-203, 2013.
14. Nishizawa M, Hattori J, Shiino T, Matano T, Heneine W, Johnson JA, and Sugiura W. Highly-Sensitive Allele-Specific PCR Testing Identifies a Greater Prevalence of Transmitted HIV Drug Resistance in Japan. *PLoS One* 8(12): e83150, 2013.
15. Shibata M, Takahashi M, Yoshino M, Kuwahara T, Nomura T, Yokomaku Y, and Sugiura W. Development and application of a simple LC-MS method for the determination of plasma rilpivirine (TMC-278) concentrations. *JMI* 60(1-2): 35-40, 2013.
16. Saito A, Nomaguchi M, Kono K, Iwatani Y, Yokoyama M, Yasutomi Y, Sato H, Shioda T, Sugiura W, Matano T, Adachi A, Nakayama EE, and Akari H. TRIM5 genotypes in cynomolgus monkeys primarily influence inter-individual diversity in susceptibility to monkey-tropic human immunodeficiency virus type 1. *J Gen Virol* 94(Pt 6): 1318-1324, 2013.
17. Nii-Trebi NI, Ibe S, Barnor JS, Ishikawa K, Brandful JA, Ofori SB, Yamaoka S, Ampofo WK, and Sugiura W. HIV-1 Drug-Resistance Surveillance among Treatment-Experienced and -Naive Patients after the Implementation of Antiretroviral Therapy in Ghana. *PloS one* 8(8): e71972, 2013.
18. Katano H, Yokomaku Y, Fukumoto H, Kanno T, Nakayama T, Shingae A, Sugiura W, Ichikawa S, and Yasuoka A. Seroprevalence of Kaposi's sarcoma-associated herpesvirus among men who have sex with men in Japan. *J Medical Virol* 85(6): 1046-1052, 2013.
19. Jahanbakhsh F, Hattori J, Matsuda M, Ibe S, Monavari SH, Memarnejadian A, Aghasadeghi MR, Mostafavi E, Mohraz M, Jabbari H, Kamali K, Keyvani H, Azadmanesh K, and Sugiura W. Prevalence of transmitted HIV drug resistance in Iran between 2010 and 2011. *PloS one* 8(4): e61864, 2013.
20. Gatanaga H, Murakoshi H, Hachiya A, Hayashida T, Chikata T, Ode H, Tsuchiya K, Sugiura W, Takiguchi M, and Oka S. Naturally Selected Rilpivirine-Resistant HIV-1 Variants by Host Cellular Immunity. *Clin Infect Dis* 57(7): 1051-1055, 2013.

21. Kudoh A, Takahama S, Sawasaki T, Ode H, Yokoyama M, Okayama A, Ishikawa A, Miyakawa K, Matsunaga S, Kimura H, Sugiura W, Sato H, Hirano H, Ohno S, Yamamoto N and Ryo A. The phosphorylation of HIV-1 Gag by atypical protein kinase C facilitates viral infectivity by promoting Vpr incorporation into virions. *Retrovirology* 11(1): 9, 2014.

吉村 和久

1. Narumi T, Arai H, Yoshimura K, Harada S, Nomura W, Matsushita S, and Tamamura H. Small Molecular CD4 Mimics as HIV Entry Inhibitors. *Bioorg Med Chem*, 19: 6735-6742, 2011.
2. Maeda Y, Yoshimura K, Miyamoto F, Kodama E, Harada S, Yuan Y, Harada S, and Yusa K. *In vitro* and *In vivo* Resistance to Human Immunodeficiency Virus Type 1 Entry Inhibitors. *J AIDS Clin Res* S2:004. doi: 10.4172/2155-6113.S2-004, 2012.
3. Ong YT, Kirby KA, Hachiya A, Chiang LA, Marchand B, Yoshimura K, Murakami T, Singh K, Matsushita S, and Sarafianos SG. Preparation of biologically active single-chain variable antibody fragments that target the HIV-1 gp120 V3 loop. *Cell Mol Biol* 58:71-79, 2012.
4. Yokoyama M, Naganawa S, Yoshimura K, Matsushita S, and Sato H. Structural Dynamics of HIV-1 Envelope Gp120 Outer Domain with V3 Loop. *PLoS ONE* 7: e37530, 2012.
5. Harada S, Yoshimura K, Yamaguchi A, Yusa K, and Matsushita S. Impact of antiretroviral pressure on selection of primary HIV-1 envelope sequences in vitro. *J Gen Virol* 94: 933-943, 2013.
6. Kuwata T, Takaki K, Yoshimura K, Enomoto I, Wu F, Ourmanov KI, Hirsch VM, Yokoyama M, Sato H, and Matsushita S. Conformational Epitope Consisting of the V3 and V4 Loops as a Target for Potent and Broad Neutralization of Simian Immunodeficiency Viruses. *J Virol* 87: 5424-5436, 2013.
7. Narumi T, Arai H, Yoshimura K, Harada S, Hirota Y, Ohashi N, Hashimoto C, Nomura W, Matsushita S, and Tamamura H. CD4 mimics as HIV entry inhibitors: lead optimization studies of the aromatic substituents. *Bioorg Med Chem* 21: 2518-2526, 2013.
8. Kuwata T, Takaki K, Enomoto I, Yoshimura K, and Matsushita S. Increased infectivity in human cells and resistance to antibody-mediated neutralization by truncation of the SIV gp41 cytoplasmic tail. *Front Microbiol Virol* 4: 1-7, 2013.
9. Hashimoto C, Narumi T, Otsuki H, Hirota Y, Arai H, Yoshimura K, Harada S, Ohashi N, Nomura W, Miura T, Igarashi T, Matsushita S, and Tamamura H. A CD4 mimic as an HIV entry inhibitor: Pharmacokinetics. *Bioorg Med Chem* 21: 7884-7889, 2013.
10. Otsuki H, Hishiki T, Miura T, Hashimoto C, Narumi T, Tamamura H, Yoshimura K, Matsushita S, and Igarashi T. Generation of a replication-competent simian-human immunodeficiency virus, the neutralisation sensitivity of which can be enhanced in the presence of a small molecule CD4 mimic. *J Gen Virol* 94: 2710-2716, 2013.

川村 龍吉

1. Kawamura T, Ogawa Y, Nakamura Y, Nakamizo S, Ohta Y, Nakano H, Kabashima K, Katayama I, Koizumi S, Kodama T, Nakao A, and Shimada S. Severe dermatitis with loss of epidermal Langerhans cells in human and mouse zinc deficiency. *J Clin Invest* 122(2): 722-732, 2012.
2. Ogawa Y, Kawamura T, Matsuzawa T, Aoki R, Gee P, Yamashita A, Moriishi K, Yamasaki K, Koyanagi Y, Blauvelt A, Shimada S. Antimicrobial peptide LL-37 produced by HSV-2-infected keratinocytes enhances HIV infection of Langerhans cells. *Cell Host Microbe* 16;13(1): 77-86, 2013.
3. Matsuzawa T, Kawamura T, Ogawa Y, Takahashi M, Aoki R, Moriishi K, Koyanagi Y, Gatanaga H, Blauvelt A, and Shimada S. Oral administration of the CCR5 inhibitor, maraviroc, blocks HIV ex vivo infection of Langerhans cells within the epithelium. *J Invest Dermatol* 133(12): 2803-2805, 2013.
4. Aoki R, Kawamura T, Goshima F, Ogawa Y, Nakae S, Nakao A, Moriishi K, Nishiyama Y, and Shimada S. Mast cells play a key role in host defense against herpes simplex virus infection through TNF- α and IL-6 production. *J Invest Dermatol* 133(9): 2170-9, 2013.
5. Matsuzawa T, Kawamura T, Ogawa Y, Maeda K, Nakata H, Moriishi K, Koyanagi Y, Gatanaga H, Shimada S, and Mitsuya H. EFdA, a Reverse Transcriptase Inhibitor, Potently Blocks HIV-1 Ex Vivo Infection of Langerhans Cells within Epithelium. *J Invest Dermatol* 2014 in press

児玉 栄一

1. Hachiya A, Kodama EN, Schuckmann MM, Kirby KA, Michailidis E, Sakagami Y, Oka S, Singh K, and Sarafianos SG. K70Q adds high-level tenofovir resistance to “Q151M complex” HIV reverse transcriptase through the enhanced discrimination mechanism. *PLoS One* 6:e16242, 2011
2. Kirby KA, Singh K, Michailidis E, Marchand B, Kodama EN, Ashida N, Mitsuya H, Parniak MA, and Sarafianos SG. The sugar ring conformation of 4'-ethynyl-2'-fluoro- 2'-deoxyadenosine and its recognition by the polymerase active site of HIV reverse transcriptase. *Cellular and Molecular Biology* 57: 39-44, 2011.
3. Li S, Hattori T, and Kodama EN. Epigallocatechin gallate inhibits HIV reverse transcription step. *Antiviral Chem Chemother* 21: 239-243, 2011.
4. Izumi K, Watanabe K, Oishi S, Fujii N, Matsuoka M, Sarafianos SG and Kodama EN. Potent anti-HIV-1 activity of N-HR-derived peptides including a deep pocket-forming region without antagonistic effect on T-20. *Antiviral Chem Chemother* 22: 51-55, 2011.
5. Tanyaradzwa P, Ndongwe TP, Adedeji AO, Michailidis E, Ong YT, Hachiya A, Marchand B, Ryan EM, Rai DK, Kirby KA, Whatley AS, Burke D, Johnson M, Ding S, Zheng YM, Liu SL, Kodama E, Delviks-Frankenberry KA, Pathak VK, Mitsuya H, Parniak MA, Singh K, and Sarafianos SG. Biochemical, inhibition, and inhibitor resistance studies of xenotropic murine leukemia virus-related virus reverse transcriptase. *Nucleic Acids Research* 40: 345-359, 2012.

6. Miyamoto F, and Kodama EN. Novel Human Immunodeficiency Virus Type 1 (HIV-1) Fusion Inhibition Peptides: Designing the Next Generation of Drugs. *Antiviral Chem Chemother* 22:151-158, 2012.
7. Maeda Y, Yoshimura K, Miyamoto F, Kodama E, Harada S, Yuan Y, Harada S, and Yusa K. *In vitro* and *In vivo* Resistance to Human Immunodeficiency Virus Type 1 Entry Inhibitors. *J AIDS Clin Res* S2:004. doi: 10.4172/2155-6113.S2-004, 2012.
8. Masuda R, Oishi S, Tanahara N, Ohno H, Hirasawa A, Tsujimoto G, Kodama E, Matsuoka M, Fuji N. Development and application of fluorescent SDF-1 derivatives. *Future Med Chem* 4: 837-844, 2012.
9. Li X, Qian H, Miyamoto F, Naito T, Kawaji K, Kajiwara K, Hattori T, Matsuoka M, Watanabe K, Oishi S, Fujii N, and Kodama EN. A simple, rapid, and sensitive system for the evaluation of anti-viral drugs in rats. *Biochem Biophys Res Commun* 424: 257-261, 2012.
10. Hachiya A, Marchand B, Kirby KA, Michailidis E, Tu X, Krzystof Palczewski, Ong YT, Griffin DT, Schuckmann MM, Junko Tanuma, Oka S, Singh K, Kodam ENa, and Sarafianos SG. HIV-1 reverse transcriptase (RT) polymorphism 172K, suppresses the effect of clinically relevant drug resistance mutations to both nucleoside and nonnucleoside RT inhibitors. *J Bio Chem* 287: 29988-29999, 2012.
11. Michailidis E, Singh K, Ryan EM, Hachiya A, Ong YT, Kirby KA, Marchand B, Kodama EN, Mitsuya H, Parniak MA, and Sarafianos SG. Effect of translocation defective reverse transcriptase inhibitors on the activity of n348i, a connection subdomain drug resistant hiv-1 reverse transcriptase mutant. *Cell Mol Biol* 58: 187-195, 2012.
12. Izumi K, Kawaji K, Miyamoto F, Shimane K, Shimura K, Sakagami Y, Hattori T, Watanabe K, Oishi S, Fujii N, Matsuoka M, Kaku M, Sarafianos SG, and Kodama EN. Mechanism of Resistance to S138A Substituted Enfuvirtide and its Application to Peptide Design. *Int J Biochem Cell Biol* 45: 908-915, 2013.
13. Shimane K, Kawaji K, Miyamoto F, Oishi S, Watanabe K, Sakagami Y, Fujii N, Shimura K, Matsuoka M, Kaku M, Sarafianos SG, and Kodama E. HIV-1 resistance mechanism to an electrostatically constrained peptide fusion inhibitor that is active against T-20-resistant strains. *Antimicrob Agent Chemother* 57:4035-4038, 2013.
14. Michailidis E, Ryan EM, Hachiya A, Kirby KA, Marchand B, Leslie MD, Huber AD, Ong YT, Jackson JC, Singh K, Kodama EN, Mitsuya H, Parniak MA and Sarafianos SG. Hypersusceptibility Mechanism of Tenofovir-Resistant HIV to EFdA. *Retrovirology* 10: 65 doi:10.1186/1742-4690-10-65, 2013
15. Hachiya A, Reeve A, Marchand B, Michailidis E, Ong Y, Kirby K, Leslie M, Oka S, Kodama E, Rohan L, Mitsuya H, Parniak M, and Sarafianos SG. Evaluation of combinations of 4'-ethynyl-2'-fluoro-2'-deoxyadenosine with clinically used antiretroviral drugs. *Antimicrob Agent Chemother* 57: 4554-4558, 2013.

16. Kirby KA, Michailidis E, Fetterly TL, Steinbach MA, Singh K, Marchand B, Leslie MD, Hagedorn AN, **Kodama EN**, Marquez VE, Hughes SH, Mitsuya H, Parniak MA, Sarafianos SG. Effects of substitutions at the 4' and 2' positions on the bioactivity of 4'-ethynyl-2'-fluoro-2'-deoxyadenosine. *Antimicrob Agent Chemother* 57:6254-6264, 2013.
17. Maeda K, Desai DV, Aoki M, Nakata H, **Kodama EN**, and Mitsuya H. Delayed emergence of HIV-1 variants resistant to 4'-ethynyl-2'-fluoro-2'-deoxyadenosine: comparative sequential passage study with lamivudine, tenofovir, emtricitabine and BMS-986001. *Antivir Ther* 19:179-89, 2014.
18. Miyamoto F and **Kodama EN**. Development of small molecule HIV-1 fusion inhibitors: linking Biology to Chemistry. *Curr Pharm Des* 19:1827-1834, 2013
19. Fujiwara S, Kimura H, Imadome K, Arai A, **Kodama E**, Morio T, Shimizu N, and Wakiguchi H. Current Studies on Chronic Active Epstein-Barr virus Infection in Japan. *Pediatr Int* 54:159-166, 2014

横幕 能行

1. Uzu T, Yokoyama H, Itoh H, Koya D, Nakagawa A, Nishizawa M, Maegawa H, **Yokomaku Y**, Araki S, Abiko A, and Haneda M. Elevated serum levels of interleukin-18 in patients with overt diabetic nephropathy: effects of miglitol. *Clin Exp Nephrol* 15: 58-63, 2011.
2. Fujisaki S, **Yokomaku Y**, Shiino T, Koibuchi T, Hattori J, Ibe S, Iwatani Y, Iwamoto A, Shirasaka T, Hamaguchi M, and **Sugiura W**. Outbreak of Infections by Hepatitis B Virus Genotype A and Transmission of Genetic Drug Resistance in Patients Coinfected with HIV-1 in Japan. *J Clin Microbiol* 49(3): 1017-1024, 2011.
3. Tsuzuki T, Iwase H, Shimada M, Hirashima N, Hibino Y, Ryuge N, Saito M, Tamaki D, Kamiya A, Yokoi M, **Yokomaku Y**, Fujisaki S, **Sugiura W**, and Goto H. Clinical evaluation of peginterferon alpha plus ribavirin for patients co-infected with HIV and HCV at Nagoya Medical Center. *Jpn J Gastroenterol* 109(7):1186-1196, 2012.
4. Miyamoto T, Nakayama EE, Yokoyama M, Ibe S, Takehara S, Kono K, **Yokomaku Y**, Pizzato M, Luban J, **Sugiura W**, Sato H, and Shioda T. The Carboxyl-Terminus of Human Immunodeficiency Virus Type 2 Circulating Recombinant form 01_AB Capsid Protein Affects Sensitivity to Human TRIM5 α . *PLoS one* 7(10):e47757, 2012.
5. Kitamura S, Ode H, Nakashima M, Imahashi M, Naganawa Y, Kurosawa T, **Yokomaku Y**, Yamane T, Watanabe N, Suzuki A, **Sugiura W**, and Iwatani Y. The APOBEC3C crystal structure and the interface for HIV-1 Vif binding. *Nat Struct Mol Biol* 19(10): 1005-1010, 2012.
6. Hirano A, Ikemura K, Takahashi M, Shibata M, Amioka K, Nomura T, **Yokomaku Y**, and **Sugiura W**. Short communication: lack of correlation between UGT1A1*6, *28 genotypes, and plasma raltegravir concentrations in Japanese HIV type 1-infected patients. *AIDS Res Hum Retroviruses* 28(8):776-779, 2012.

7. Nishijima T, Takano M, Ishisaka M, Komatsu H, **Gatanaga H**, Kikuchi Y, Endo T, Horiba M, Kaneda S, Uchiyama H, Koibuchi T, Naito T, Yoshida M, Tachikawa N, Ueda M, **Yokomaku Y**, Fujii T, Higasa S, Takada K, Yamamoto M, Matsushita S, Tateyama M, Tanabe Y, Mitsuya H, and Oka S. Abacavir/lamivudine versus tenofovir/emtricitabine with atazanavir/ritonavir for treatment-naive Japanese patients with HIV-1 infection: a randomized multicenter trial. *Intern Med* 52(7): 735-744, 2013.
8. Shibata M, Takahashi M, Yoshino M, Kuwahara T, Nomura T, **Yokomaku Y**, and **Sugiura W**. Development and application of a simple LC-MS method for the determination of plasma rilpivirine (TMC-278) concentrations. *JMI* 60(1-2): 35-40, 2013.
9. Katano H, **Yokomaku Y**, Fukumoto H, Kanno T, Nakayama T, Shingae A, **Sugiura W**, Ichikawa S, and Yasuoka A. Seroprevalence of Kaposi's sarcoma-associated herpesvirus among men who have sex with men in Japan. *J Med Virol* 85(6): 1046-1052, 2013.

Impact of Small Body Weight on Tenofovir-Associated Renal Dysfunction in HIV-Infected Patients: A Retrospective Cohort Study of Japanese Patients

Takeshi Nishijima^{1,3*}, Hirokazu Komatsu², Hiroyuki Gatanaga^{1,3}, Takahiro Aoki¹, Koji Watanabe^{1,3}, Ei Kinai¹, Haruhito Honda¹, Junko Tanuma¹, Hirohisa Yazaki¹, Kunihisa Tsukada¹, Miwako Honda¹, Katsuji Teruya¹, Yoshimi Kikuchi¹, Shinichi Oka^{1,3}

1 AIDS Clinical Center, National Center for Global Health and Medicine, Tokyo, Japan, **2** Department of Community Care, Saku Central Hospital, Nagano, Japan, **3** Center for AIDS Research, Kumamoto University, Kumamoto, Japan

Abstract

Background: Treatment with tenofovir is sometimes associated with renal dysfunction. Limited information is available on this side effect in patients with small body weight, although the use of tenofovir will spread rapidly in Asia and Africa, where patients are likely to be of smaller body weight.

Methods: In a single-center cohort, Japanese patients with HIV infection who started tenofovir-containing antiretroviral therapy were retrospectively analyzed. The incidence of tenofovir-associated renal dysfunction, defined as more than 25% decrement of estimated glomerular filtration rate (eGFR) from the baseline, was determined. The effects of small body weight and body mass index (BMI) on tenofovir-associated renal dysfunction, respectively, were estimated in univariate and multivariate Cox hazards models as the primary exposure. Other possible risk factors were evaluated by univariate analysis and those found significant were entered into the multivariate analysis.

Results: The median weight of 495 patients was 63 kg. Tenofovir-related renal dysfunction occurred in 97 (19.6%) patients (incidence: 10.5 per 100 person-years). Univariate analysis showed that the incidence of tenofovir-related renal dysfunction was significantly associated with smaller body weight and BMI, respectively (per 5 kg decrement, HR=1.23; 95% CI, 1.10–1.37; $p<0.001$) (per 1 kg/m² decrement, HR=1.14; 95% CI, 1.05–1.23; $p=0.001$). Old age, high baseline eGFR, low serum creatinine, low CD4 count, high HIV viral load, concurrent nephrotoxic drugs, hepatitis C infection, and current smoking were also associated with tenofovir-related renal dysfunction. Multivariate analysis identified small body weight as a significant risk (adjusted HR=1.13; 95% CI, 1.01–1.27; $p=0.039$), while small BMI had marginal significance (adjusted HR=1.07; 95% CI 1.00–1.16; $p=0.058$).

Conclusion: The incidence of tenofovir-associated renal dysfunction in Japanese patients was high. Small body weight was identified as an independent risk factor for tenofovir-associated renal dysfunction. Close monitoring of renal function is advocated for patients with small body weight treated with tenofovir.

Citation: Nishijima T, Komatsu H, Gatanaga H, Aoki T, Watanabe K, et al. (2011) Impact of Small Body Weight on Tenofovir-Associated Renal Dysfunction in HIV-Infected Patients: A Retrospective Cohort Study of Japanese Patients. PLoS ONE 6(7): e22661. doi:10.1371/journal.pone.0022661

Editor: Gary Maartens, University of Cape Town, South Africa

Received: February 24, 2011; **Accepted:** June 28, 2011; **Published:** July 25, 2011

Copyright: © 2011 Nishijima et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by a Grant-in-Aid for AIDS Research from the Ministry of Health, Labor, and Welfare of Japan (H20-AIDS-002). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: tnishiji@acc.ncgm.go.jp

Introduction

Tenofovir disoproxil fumarate (TDF) is one of the most widely used nucleotide reverse transcriptase inhibitors (NRTI) for patients with HIV infection, with proven efficacy and safety [1–6]. However, TDF is known to cause renal proximal tubular dysfunction, and several case reports have been published with TDF-related Fanconi syndrome, diabetes insipidus, and acute tubular necrosis, which sometimes lead to acute renal failure [7–10]. Long-term TDF use also reduces glomerular filtration rate more than other NRTIs [11–14]. To date, the nephrotoxic effect of TDF is regarded as mild and tolerable. A recently published

meta-analysis has reported that the use of TDF is associated with a statistically significant but only modest renal dysfunction, and recommended that TDF use should not be restricted even when regular monitoring of renal function and serum phosphate levels is impractical [15]. However, the TDF-related renal dysfunction has hardly been evaluated in patients with small body weight, who are potentially at higher risk for larger drug exposure and thus, more severe toxicity [16–19].

The 2010 WHO guideline on antiretroviral therapy for HIV infection in adults and adolescents, usually applied to resource-constrained settings, recommends TDF as one of the components of first line therapies (URL:<http://whqlibdoc.who.int/publications/>

2010/9789241599764_eng.pdf). It is expected that the use of TDF will spread rapidly in Asia and Africa in the near future, where patients are more likely to be of small body weight. Thus, at this stage, it is important to establish the relationship between TDF-associated renal dysfunction and body weight. A small body weight is considered a risk factor for TDF-associated renal dysfunction, in addition to old age, high baseline serum creatinine level, low CD4 count, concurrent use of ritonavir-boosted protease inhibitor, and concurrent use of nephrotoxic drugs [4,17,19–21]. To our knowledge, there is almost no report that primarily analyzed the influence of body weight on TDF-associated renal dysfunction. Since Japanese are generally of smaller stature and have a lower median body weight than Whites and African Americans, who mostly comprise the cohorts of studies published to date, it is important to investigate the impact of TDF-associated renal dysfunction in Japanese patients.

Based on the above background, the present study was designed to determine the incidence of TDF-associated renal dysfunction in Japanese patients and analyze the impact of small body weight on TDF-associated renal dysfunction.

Methods

Ethics Statement

This study was approved by the Human Research Ethics Committee of National Center for Global Health and Medicine (Text S1). All patients included in this study provided a written informed consent for their clinical and laboratory data to be used and be published for research purposes. This study has been conducted according to the principles expressed in the Declaration of Helsinki.

Study Design and Settings

We performed a single-center, retrospective cohort study of HIV-infected Japanese patients using medical records at the National Center for Global Health and Medicine, Tokyo, Japan. Our facility is one of the largest clinics for patients with HIV infection in Japan with more than 2,700 registered patients.

Study Subjects

The study population were patients >17 years of age who commenced treatment with standard 300 mg/day of TDF-containing antiretroviral regimen at our clinic between January 1, 2002 to March 31, 2009. Both treatment-naïve and patients with experience in antiretroviral treatment but not TDF, with an estimated glomerular filtration rate (eGFR) of >60 ml/min/1.73 m² were enrolled. Patients were followed up until September 31, 2009. Patients were excluded if their follow-up period at our facility was less than 24 weeks after commencement of TDF-based therapy, if they had started TDF at other facilities, or if there was evidence of prior TDF use. We only included Japanese patients in order to examine a population with comparatively homogenous basic demographics and background.

Measurements

Outcome measure: TDF-associated renal dysfunction. We defined TDF-associated renal dysfunction as more than 25% decrease in eGFR relative to the baseline [17,22,23]. Baseline eGFR was estimated for each patient from the average of two successive serum creatinine measurements made closest to and preceding the commencement of TDF by no more than 90 days. Changes in eGFR were plotted from the baseline measurement until the value diminished to less than 75% of the baseline or at the end of the follow-up period. The eGFR values at occurrence of TDF-

associated renal dysfunction, at censoring, and closest to and preceding 24, 48, and 96 weeks to the diagnosis were collected. Patients generally visited our clinic between every month to every 3 months, and measurement of eGFR was usually conducted on every visit. eGFR was calculated using the equation from the 4-variable Modification of Diet in Renal Disease (MDRD) study [24].

Primary exposure variable. Our primary exposure variables were body weight and body mass index (BMI) at the time of commencement of TDF-containing antiretroviral therapy (ART). BMI was calculated by the equation: BMI = [body weight (kg)/height (m)²].

Other variables: potential risk factors. Potential risk factors for TDF-associated renal dysfunction were determined according to previous studies and collected together with the basic demographics from the medical charts [4,19,20,25]. They included sex, age, baseline laboratory data: CD4 cell count, HIV viral load, and serum creatinine, and other medical conditions (antiretroviral treatment-naïve or experienced, concurrent ritonavir-boosted protease inhibitors, concurrent nephrotoxic drugs such as ganciclovir, sulfamethoxazole/trimethoprim, ciprofloxacin, and NSAIDs, diabetes mellitus, co-infection with hepatitis B defined by positive hepatitis B surface antigen, co-infection with hepatitis C defined by positive HCV viral load, hypertension defined by current treatment with antihypertensive agents, dyslipidemia defined by current treatment with lipid-lowering agents, and current smoking) [26]. We used the data on or closest to and preceding the day of starting TDF-containing ART by no more than 90 days. The data on weight change from the baseline to the end of follow-up period and the frequency of eGFR monitoring for each patient were collected.

Statistical analysis

The time to 25% decline in eGFR from the baseline was calculated from the date of treatment initiation to the date of occurrence of TDF-associated renal dysfunction. Censored cases represented those who discontinued TDF, dropped out, referred to other facilities, or at the end of follow-up period. The time from TDF initiation to 25% decrease in eGFR was analyzed by the Kaplan Meier method for the whole cohort. To estimate the impact of body weight on the incidence of TDF-associated renal dysfunction, we calculated the impact of every 5 kg decrement from the median weight using Cox proportional hazards regression analysis. The impact of every 1 kg/m² decrement in BMI on the incidence of TDF-associated renal dysfunction was estimated by the same method. The impact of each basic demographics, baseline laboratory data, and other medical conditions listed above was also estimated with univariate Cox proportional hazards regression.

To estimate the unbiased prognostic impact of weight on TDF-associated renal dysfunction, we conducted three models using multivariate Cox proportional hazards regression analysis. Model 1 was the aforementioned univariate analysis for every 5 kg decrement. Model 2 included sex, age plus model 1 in order to adjust for basic characteristics. In model 3, we added variables with P values <0.05 in univariate analysis for adjustment (these included age per 10 years, serum creatinine >0.8 mg/dl, CD4 count <200/μl, HIV viral load per log₁₀/ml, concurrent nephrotoxic drugs, co-infection with hepatitis C, and current smoking). Concurrent ritonavir-boosted protease inhibitors were also added in Model 3 although their p value was 0.116 in the univariate analysis. This was based on the results of several studies suggesting that concurrent use of ritonavir-boosted protease inhibitors is a risk factor for TDF-associated renal dysfunction

[19,20]. The eGFR was excluded from multivariate analysis because of its multicollinearity with sex, age, and serum creatinine, since eGFR was gained by the equation of those variables. The impact of every 1 kg/m² decrement in BMI on the incidence of TDF-associated renal dysfunction was estimated by the same method with Model 1 to Model 3.

Four other analyses were conducted to further examine the relationship between low body weight and TDF-associated renal dysfunction. First, the time from initiation of TDF therapy to 25% decrease in eGFR was analyzed by the Kaplan Meier method for intertertile baseline body weight categories: <59, 59–67, and >67 kg. The log-rank test was used to determine statistical significance. Second, to investigate the impact of changes in muscle mass on changes in the eGFR as calculated by MDRD, we compared weight changes with one-way ANOVA among intertertile baseline weight categories. We also conducted the sensitivity analysis by adding the variable “weight change” in multivariate analysis. Third, the median and interquartile value for the actual fall in eGFR from the baseline to 24, 48, and 96 weeks for the whole cohort and three baseline weight categories, respectively, were calculated. The eGFR value at 24, 48, and 96 weeks included those that were censored before reaching 24, 48, and 96 weeks, respectively, so that we could interpret the data for actual fall in eGFR, including not only survived cases but also censored cases. Fourth, we counted the number of patients whose eGFR decreased to <60 and <10 ml/min/1.73 m², and who discontinued TDF with the clinical diagnosis of renal dysfunction due to TDF. Chi-square test was used to determine whether the difference among the weight categories was statistically significant.

Statistical significance was defined at two-sided p values < 0.05. We used hazard ratios (HRs) and 95% confidence intervals (CIs) to estimate the impact of each variable on TDF-associated renal dysfunction. All statistical analyses were performed with The Statistical Package for Social Sciences ver. 17.0 (SPSS, Chicago, IL).

Results

Between January 1, 2002 to March 31, 2009, 599 patients started TDF-containing ART (Figure 1). Of these, 104 patients were excluded based on the abovementioned criteria. Thus 495 patients were included in the present study (Dataset S1). Table 1 shows the demographics, laboratory data, and medical conditions of the study population at baseline. Two patients received ART with 3 NRTIs, 3 patients received ART with one protease inhibitor (PI), one non-NRTI (NNRTI), and tenofovir/emtricitabine, and the remaining patients had a standard ART with 2 NRTIs and either PI, NNRTI, or integrase inhibitor (INI). The median body weight and BMI were 63 kg and 21.9 kg/m², respectively. The median age of the patients was 38 years and 95.2% were males. The eGFR was well maintained (median: 120.9 ml/min/1.73 m²), and the median baseline CD4 count was 247/μl. Of the total, 208 patients (42%) were antiretroviral treatment naïve, while 287 were treatment-experienced patients. Viral load was suppressed to <50 copies/ml in 162 (32.7%) patients. 403 (81.4%) were on concurrent PIs as the key drug, 367 (74.1%) were on ritonavir-boosted PIs, and only 83 (16.8%) had NNRTIs as the key drug. Smoking was prevalent among the study population, as 240 (48.5%) were identified as a current smoker.

TDF-associated renal dysfunction defined by more than 25% decrease of eGFR from baseline occurred in 97 patients (19.6%), with an estimated incidence of 10.5 per 100 person-years. The median time from commencement of TDF to occurrence of TDF-




Figure 1. Flow diagram of patient selection.
doi:10.1371/journal.pone.0022661.g001

associated renal dysfunction was 39 weeks (IQR 13.5–99.4 weeks) (range: 1–1,841 days). The total observation period was 924.7 patient-years (median 72 weeks, IQR 38.6–139.3 weeks). Figure 2 shows the Kaplan-Meier survival curve for the occurrence of TDF-associated renal dysfunction for the whole cohort.

Univariate analysis showed a significant relationship between TDF-associated renal dysfunction and every 5 kg less than the median body weight (HR = 1.23; 95% CI, 1.10–1.37; $p < 0.001$), and 1 kg/m² less BMI than the median BMI (HR = 1.14; 95% CI, 1.05–1.23; $p = 0.001$) (Table 2). Furthermore, old age, high eGFR, low serum creatinine, low CD4 counts, high HIV viral load, concurrent use of nephrotoxic drugs, presence of chronic hepatitis C, and smoking were associated with TDF-related renal dysfunction. On the other hand, concurrent use of PIs, ritonavir boosted PIs, and LPV/r tended to be associated with TDF-related renal dysfunction, albeit statistically insignificant. Treatment-naïve or Treatment-experienced was not associated with TDF-related renal dysfunction.

Multivariate analysis showed that every 5 kg less than the median body weight was a significant risk for TDF-associated renal dysfunction after adjustment for sex and age (adjusted HR = 1.21; 95% CI, 1.07–1.36; $p = 0.002$) (Table 3, Model 2), and also after adjustment for other risk factors (adjusted HR = 1.13; 95% CI, 1.01–1.27; $p = 0.039$) (Table 3, Model 3). Similarly, every 1 kg/m² less than the median BMI was also a significant risk factor for TDF-associated renal dysfunction even after adjustment for sex and age (adjusted HR = 1.13; 95% CI 1.05–1.22; $p = 0.002$) (Table 4, Model 2), and tended to be a significant factor after adjustment for other variables (adjusted HR = 1.07; 95% CI 1.00–1.16; $p = 0.058$) (Table 4, Model 3). Old age and current smoking were also independent risk factors in both multivariate analysis for body weight and BMI (Table 3, Model 3 and Table 4, Model 3).

In complementary analyses, First, Figure 3 shows the relation between probability of TDF-associated nephrotoxicity and time from initiation of TDF therapy to 25% decrease in eGFR analyzed by the Kaplan Meier method for intertertile baseline weight categories. Compared to patients with baseline body weight >67 kg, patients with baseline weight <59 kg were significantly more likely to develop >25% decline in eGFR ($p = 0.002$). On the other hand, the difference in this probability between patients with baseline weight 59–67 kg and those >67 kg was only marginally significant ($p = 0.073$, log-rank test). Secondly, one-way ANOVA

Table 1. Baseline demographics and laboratory data.


Characteristics		
Median (IQR) weight (kg)	63	(57–69)
Median (IQR) BMI (kg/m ²)	21.9	(20.3–23.8)
Male, n (%)	471	(95.2)
Median (IQR) age	38	(33–46)
Median (IQR) eGFR (ml/min/1.73 m ²)	120.9	(104.8–138.2)
Median (IQR) serum creatinine (mg/dl)	0.72	(0.64–0.81)
Median (IQR) CD4 count (/μl)	247	(159–371)
Median (IQR) HIV viral load (log ₁₀ /ml)	3.73	(1.60–4.81)
HIV viral load <50 copies/ml, n (%)	162	(32.7)
Antiretroviral therapy naïve, n (%)	208	(42.0)
Key drugs, n (%)*		
PIs	403	(81.4)
Ritonavir-boosted PIs	367	(74.1)
LPV/r	175	(35.4)
ATV/r	131	(26.5)
FPV/r	52	(10.5)
DRV/r	9	(1.8)
FPV	14	(2.8)
ATV	4	(0.8)
NFV	15	(3)
SQV	2	(0.4)
IDV	1	(0.2)
NNRTIs	83	(16.8)
EFV	65	(13.1)
NVP	17	(3.4)
ETR	1	(0.2)
INI		
RAL	10	(2.0)
Concurrent use of nephrotoxic drug, n (%)	131	(26.5)
Diabetes mellitus, n (%)	30	(6.1)
Hepatitis B, n (%)	75	(15.2)
Hepatitis C, n (%)	52	(10.5)
Hypertension, n (%)	28	(5.7)
Dyslipidemia, n (%)	40	(8.1)
Smoking, n (%)	240	(48.5)
Median (IQR) weight change (kg)	0.0	(–2.0–2.25)
Median (IQR) frequency of eGFR monitoring	16	(9.0–27)

(n = 495).

*Two patients did not take any key drugs. Three patients took both PI and NNRTI.

IQR: interquartile range, BMI: body mass index, eGFR: estimated glomerular filtration rate, PI: protease inhibitor, LPV/r: lopinavir/ritonavir, ATV: atazanavir, FPV: fosamprenavir, DRV: darunavir, NFV: nelfinavir, SQV: saquinavir, IDV: indinavir, NNRTI: non-nucleos(t)ide reverse transcriptase inhibitor, EFV: efavirenz, NVP: nevirapine, ETR: etravirine, INI: integrase inhibitor, RAL: raltegravir.

doi:10.1371/journal.pone.0022661.t001

**Figure 2.** Kaplan-Meier curve showing the time to 25% reduction in eGFR for the whole cohort. eGFR: estimated glomerular filtration rate.

doi:10.1371/journal.pone.0022661.g002

the median and interquartile values for the actual falls in eGFR from the baseline to 24, 48, and 96 weeks. The eGFR decreased gradually in all categories, except for patients with baseline weight

Table 2. Univariate analysis for TDF-associated renal dysfunction.

	HR	95%CI	P value
Weight per 5 kg decrement	1.23	1.10–1.37	<0.001
BMI per 1 kg/m ² decrement	1.14	1.05–1.23	0.001
Male gender	0.54	0.26–1.11	0.094
Age per 10 years	1.22	1.02–1.45	0.027
eGFR per 10 ml/min/1.73 m ²	1.10	1.05–1.15	<0.001
Serum creatinine >0.8 mg/dl	0.51	0.30–0.88	0.014
CD4 count <200/μl	1.97	1.32–2.93	0.001
HIV viral load per log ₁₀ /ml	1.15	1.01–1.30	0.037
Antiretroviral therapy naïve	0.98	0.63–1.52	0.927
Concurrent key drugs			
Any PIs	1.52	0.89–2.59	0.124
Ritonavir boosted PIs	1.46	0.91–2.33	0.116
LPV/r	1.45	0.97–2.17	0.072
ATV/r	1.05	0.66–1.68	0.826
Concurrent nephrotoxic drug	1.59	1.04–2.42	0.031
Diabetes mellitus	1.57	0.76–3.24	0.220
Hepatitis B	1.36	0.82–2.24	0.231
Hepatitis C	1.80	1.07–3.04	0.028
Hypertension	1.18	0.51–2.69	0.702
Dyslipidemia	0.97	0.47–2.00	0.932
Smoking	1.57	1.05–2.36	0.028

TDF: tenofovir, HR: hazard ratio, CI: confidence interval, BMI: body mass index, eGFR: estimated glomerular filtration rate, PI: protease inhibitor, LPV/r: lopinavir/ritonavir, ATV: atazanavir.

doi:10.1371/journal.pone.0022661.t002

showed that weight changes among the three baseline weight categories were not significantly different ($p = 0.206$). Sensitivity analysis after adding the variable “weight change” in Model 3 multivariate analysis (Table 3) showed that adjusted hazard ratio for weight per 5 kg decrement hardly changed (adjusted HR 1.131; 95% CI, 1.007–1.271; $p = 0.038$). Thirdly, Table 5 shows

Table 3. Multivariate analysis to estimate the effect of lower body weight on TDF-associated renal dysfunction.

	Model 1 Crude		Model 2 Adjusted		Model 3 Adjusted	
	HR	95%CI	HR	95%CI	HR	95%CI
Weight per 5 kg decrement [†]	1.23	1.10–1.37	1.21	1.07–1.36	1.13	1.01–1.27
Male gender			0.88	0.41–1.89	0.57	0.26–1.26
Age per 10 years [†]			1.16	0.98–1.38	1.24	1.04–1.49
Serum creatinine >0.8 mg/dl					0.62	0.35–1.07
CD4 count <200/ μ l					1.65	0.97–2.79
HIV viral load per log ₁₀ /ml					1.05	0.90–1.23
Boosted PIs					1.54	0.93–2.54
Concurrent use of nephrotoxic drug					1.23	0.77–1.97
Hepatitis C					1.57	0.92–2.69
Smoking [†]					1.65	1.09–2.48

[†]P<0.05 in Model 3.

TDF: tenofovir, HR: hazard ratio, CI: confidence interval, PI: protease inhibitor.
doi:10.1371/journal.pone.0022661.t003

>67 kg. Fourthly, the number (percentage) of patients whose eGFR decreased to <60 ml/min/1.73 m² was not different among the baseline weight categories (p=0.229), whereas the number of patients who discontinued TDF with a clinical diagnosis of renal dysfunction due to TDF varied significantly according to body weight (p=0.001, chi-square test, Table 6). None of the patients showed reduction of eGFR to <10 ml/min/1.73 m².

Discussion

In this Japanese cohort, 19.6% of the patients experienced eGFR decline of more than 25% from the baseline after commencement of TDF. The incidence of TDF-associated renal dysfunction was 10.5 per 100 person-years. Multivariate analysis identified smaller body weight and smaller body mass index as significant and almost significant factors, respectively, for TDF-associated renal dysfunction.

The incidence of TDF-associated renal dysfunction in patients with small body weight might be higher than previously

reported in studies of patients with larger statures. Studies from North America, Europe, and Australia reported an incidence of <1% to 4.3% for TDF-related renal dysfunction, although the definition used for the diagnosis of renal impairment was different among the studies and varied from an increase in serum creatinine from >0.5 to >2 mg/dL from baseline [1–3,5,13]. Several studies conducted in these regions indicated that the range of patients' mean body weight was 69–74 kg, indicating that their patients were heavier than those of the present study with a median weight of 63 kg [2,6,12,14]. The impact of the comparatively lower body weight seems stronger in our patients probably because they do not appear to have many of the other established risk factors for TDF-associated renal dysfunction despite the high incidence of 10.5 per 100 person-years. For example, they were comparatively young with a median age of 38 years, CD4 count was relatively maintained, and approximately 30% had suppressed HIV viral load at baseline (Table 1). Furthermore, they were less likely to have hypertension, dyslipidemia, and diabetes mellitus.

Table 4. Multivariate analysis to estimate the impact of BMI decrement on TDF-associated renal dysfunction.

	Model 1 Crude		Model 2 Adjusted		Model 3 Adjusted	
	HR	95%CI	HR	95%CI	HR	95%CI
BMI per 1 kg/m ² decrement	1.14	1.05–1.23	1.13	1.05–1.22	1.07	1.00–1.16
Male gender			0.67	0.32–1.38	0.48	0.23–1.03
Age per 10 years [†]			1.20	1.01–1.43	1.27	1.06–1.52
Serum creatinine >0.8 mg/dl					0.60	0.35–1.04
CD4 count <200/ μ l					1.64	0.97–2.79
HIV viral load per log ₁₀ /ml					1.05	0.90–1.23
Boosted PIs					1.49	0.90–2.45
Concurrent use of nephrotoxic drugs					1.22	0.76–1.94
Hepatitis C					1.62	0.94–2.76
Smoking [†]					1.63	1.08–2.46

[†]P<0.05 in Model 3.

BMI: body mass index, TDF: tenofovir, HR: hazard ratio, CI: confidence interval, PI: protease inhibitor.
doi:10.1371/journal.pone.0022661.t004




Figure 3. Kaplan-Meier curve showing the time to 25% reduction in eGFR according to baseline weight categories. Compared to patients with body weight >67 kg, those with weight <59 kg were more likely to develop $>25\%$ decline in eGFR ($P=0.002$), whereas those with weight $59\text{--}67$ kg showed only a marginal significance ($P=0.073$, log-rank test). eGFR: estimated glomerular filtration rate.
doi:10.1371/journal.pone.0022661.g003

The results of multivariate analysis that each 5 kg decrement in body weight was significantly associated with TDF-associated renal dysfunction but not each 1 kg/m^2 decrement in BMI suggests that weight might be more useful and handy information to estimate the risk for TDF-associated renal dysfunction than BMI. Thus, patient's body weight is an important risk factor to consider at the time of TDF prescription.

Our study is one of a few that have examined the impact of TDF-associated renal dysfunction in patients with small body weight, but is the first to examine the impact of small body weight as a primary exposure by creating the model used for multivariate analysis [16–18]. One study from Thailand that included patients with a median weight of 56.5 kg reported a similar incidence of 16.2 per 100 person-years for developing TDF-associated renal dysfunction [17]. They concluded that the small body weight of their patients was probably associated with the high incidence of TDF-associated renal dysfunction. Our study confirmed that conclusion and provided statistically-backed evidence that small body weight is a significant risk factor of TDF-associated renal dysfunction by using a multivariate model with least multicollinearity to evaluate the impact of small body weight. The results of the present study could be applied to many countries in

Asia and Africa, where stature and body weight of the population are comparatively smaller.

This study adopted a decrease in eGFR of $>25\%$ as a definition for TDF-associated renal dysfunction. This criterion is one of common methods in evaluating renal function [22,23]. Using this definition, however, does not mean that all patients with $>25\%$ fall in eGFR have severe renal dysfunction. However, the definition of renal dysfunction based on a fall in eGFR of $>25\%$ is probably more sensitive than that based on eGFR $<60\text{ ml/min/1.73 m}^2$, in patients with comparatively good baseline renal function, such as patients of our study. Adopting this definition could be useful in detecting early renal dysfunction and in the clinical decision making regarding the need for certain interventions, for example, discontinuation of TDF. Early detection of renal dysfunction is particularly important in patients with HIV infection, because kidney disease may be associated with AIDS and death, and TDF-associated renal dysfunction might be irreversible [27,28].

Since the calculation of eGFR using the MDRD formula is based on serum creatinine, age, race and gender, any fall in eGFR is influenced by hypercreatininemia caused by increased muscle mass [29]. It is possible that the muscle mass increases in patients on ART, especially those with low weight at baseline compared to those with higher weight, reflecting reversal of wasting in those patients who were most malnourished. Such increase in muscle mass could then result in a fall in eGFR despite no change in actual renal elimination of creatinine. However, complementary analysis showed that weight change throughout the follow-up period was not significantly different among patients with different baseline weight, and the sensitivity analysis demonstrated that weight change did not alter the significance of every 5 kg decrement.

In the present study, high eGFR and low serum creatinine levels at baseline were identified as risk factors for falls in eGFR of more than 25%, in contrast to several previous studies that showed high serum creatinine and low eGFR were risk factors [4,10,25]. While the exact reason for this discrepancy is unknown at present, it could be related to differences in the definition of TDF-associated renal dysfunction. The aforementioned Thai study used the same definition applied in the present study and a Canadian study that used the definition of 1.5-fold increase in serum creatinine from baseline also reported high eGFR and low serum creatinine level at baseline as risk factors [17,30]. Thus, it is plausible to observe a fall in eGFR when the baseline value is high, since Horberg et al. reported that patients with baseline eGFR of $>80\text{ ml/min/1.73 m}^2$ were likely to show a pronounced fall in eGFR with TDF use [31].

Multivariate analysis also suggested that old age and current smoking are significant risks for TDF-associated renal dysfunction

Table 5. Median and interquartile range of the actual fall in eGFR from the baseline to 24, 48, and 96 weeks, according to body weight.

	Total (n = 495)		<59 kg (n = 167)		59–67 kg (n = 168)		>67 kg (n = 160)	
	fall in eGFR(ml/min/1.73 m ²)		fall in eGFR		fall in eGFR		fall in eGFR	
	median	IQR	median	IQR	median	IQR	median	IQR
to 24 weeks	7.8	(–1.7–18.1)	9.8	(–3.6–22.6)	6.8	(–1.5–17.3)	7.3	(–1.8–15.4)
to 48 weeks	9.0	(–0.7–21.9)	13.0	(–0.2–29.3)	7.2	(–1.2–20.0)	8.1	(–0.6–18.6)
to 96 weeks	9.3	(–0.5–23.1)	13.4	(1.2–33.2)	8.6	(–0.2–21.7)	7.5	(–2.4–19.8)

eGFR: estimated glomerular filtration rate, IQR: interquartile range.
doi:10.1371/journal.pone.0022661.t005

Table 6. Number of patients whose eGFR decreased to <60 ml/min/1.73 m² and who discontinued tenofovir with clinical diagnosis of renal dysfunction due to tenofovir.

	<59 kg (n = 167)	59–67 kg (n = 168)	>67 kg (n = 160)	p value
eGFR <60 ml/min/1.73 m ²	4 (2.4%)	1 (0.6%)	1 (0.6%)	0.229
Discontinued tenofovir	16 (9.6%)	8 (4.8%)	1 (0.6%)	0.001
Reasons for discontinuation				
>25% eGFR decrement	8 (4.8%)	4 (2.4%)	0 (0%)	
Urine $\beta 2$ microglobulin >5000 $\mu\text{g/l}$	11 (6.6%)	4 (2.4%)	1 (0.6%)	

Among the patients who discontinued tenofovir, both $>25\%$ fall in eGFR and urine $\beta 2$ microglobulin >5000 $\mu\text{g/l}$ were registered in six patients with body weight <59 kg, and in three patients with body weight 59–67 kg.
eGFR: estimated glomerular filtration rate.

doi:10.1371/journal.pone.0022661.t006

(Table 3, Model 3 and Table 4, Model 3). However, these results have to be interpreted with caution, because these multivariate analyses were formulated to primarily evaluate weight decrement, not age or smoking.

The mechanism of TDF-associated renal dysfunction is not fully understood. TDF-associated renal dysfunction probably develops as a result of complex interaction of pharmacological, environmental, and genetic factors, rather than small body weight only [32]. It should be noted, however, that small body weight has been identified as a risk factor for TDF-associated renal dysfunction not only in clinical trials, but also in *in vitro* and pharmacokinetic studies [33–36]. TDF is the prodrug of acyclic nucleotide analog tenofovir, which is excreted by both glomerular filtration and active tubular secretion. *In vitro* studies showed that tenofovir exhibits mitochondrial toxicity in renal proximal tubular cells, and animal studies demonstrated that renal tubular dysfunction was associated with the dose and plasma drug concentrations of TDF [34,35]. Furthermore, pharmacokinetic studies showed that small body weight is associated with reduced plasma TDF clearance and thus high plasma TDF concentrations, which could result in renal tubular dysfunction. [33,36].

There are several limitations to our study. First, because of the retrospective nature of the study, patients with possible risks for TDF-associated renal dysfunction could have not been prescribed TDF. Because of this selection bias, the incidence of TDF-associated renal dysfunction might be underestimated. Second, the study did not compare the incidence of renal dysfunction in a control group (TDF-free ART). Due to the small body weight in Japanese or other factors such as genetics, the use of ART without TDF might cause higher incidence of renal dysfunction as well. Third, as discussed above, the definition of TDF-associated renal dysfunction, especially the criteria used to evaluate proximal renal tubular damage, is not uniformly established in the field and is different in the published studies. Accordingly, we decided to adopt changes in eGFR, instead of parameters for proximal renal

tubular damage. Using the eGFR as a marker for TDF-associated renal dysfunction, our results might have underestimated the incidence of TDF-associated renal dysfunction.

In conclusion, the present study demonstrated a high incidence of TDF-associated renal dysfunction among Japanese patients, a potentially high-risk group due to the low median body weight. The results also identified small body weight as a risk for TDF-associated renal dysfunction in a statistical model that included small body weight as a primary exposure. TDF is certainly a drug of choice for one of the components of the first line therapies for HIV infection. However, the importance of close monitoring for renal function in patients with small body weight should be emphasized for early detection of TDF-associated renal dysfunction.

Supporting Information

Text S1 Letter of Approval from Human Research Ethics Committee of National Center for Global Health and Medicine.

(PDF)

Dataset S1 Raw data of the target population.

(XLS)

Acknowledgments

The authors thank all the clinical staff at the AIDS Clinical Center for their help in completion of this study.

Author Contributions

Conceived and designed the experiments: AN TH KU TN MKS. Performed the experiments: AN H. Sakai. Analyzed the data: AN TH KU KO IK H. Sakai. TN MKS. Contributed reagents/materials/analysis tools: H. Suemori H. Sakai. Wrote the manuscript: AN TN MKS.

References

- Gallant JE, DeJesus E, Arribas JR, Pozniak AL, Gazzard B, et al. (2006) Tenofovir DF, emtricitabine, and efavirenz vs. zidovudine, lamivudine, and efavirenz for HIV. *N Engl J Med* 354: 251–260.
- Gallant JE, Staszewski S, Pozniak AL, DeJesus E, Suleiman JM, et al. (2004) Efficacy and safety of tenofovir DF vs stavudine in combination therapy in antiretroviral-naïve patients: a 3-year randomized trial. *JAMA* 292: 191–201.
- Izzedine H, Hulot JS, Vittecoq D, Gallant JE, Staszewski S, et al. (2005) Long-term renal safety of tenofovir disoproxil fumarate in antiretroviral-naïve HIV-1-infected patients. Data from a double-blind randomized active-controlled multicentre study. *Nephrol Dial Transplant* 20: 743–746.
- Nelson MR, Katlama C, Montaner JS, Cooper DA, Gazzard B, et al. (2007) The safety of tenofovir disoproxil fumarate for the treatment of HIV infection in adults: the first 4 years. *AIDS* 21: 1273–1281.
- Arribas JR, Pozniak AL, Gallant JE, DeJesus E, Gazzard B, et al. (2008) Tenofovir disoproxil fumarate, emtricitabine, and efavirenz compared with zidovudine/lamivudine and efavirenz in treatment-naïve patients: 144-week analysis. *J Acquir Immune Defic Syndr* 47: 74–78.
- Post FA, Moyle GJ, Stellbrink HJ, Domingo P, Podzamczar D, et al. (2010) Randomized comparison of renal effects, efficacy, and safety with once-daily abacavir/lamivudine versus tenofovir/emtricitabine, administered with efavir-

- enz, in antiretroviral-naive, HIV-1-infected adults: 48-week results from the ASSERT study. *J Acquir Immune Defic Syndr* 55: 49–57.
7. Verhelst D, Monge M, Meynard JL, Fouquerey B, Mougenot B, et al. (2002) Fanconi syndrome and renal failure induced by tenofovir: a first case report. *Am J Kidney Dis* 40: 1331–1333.
 8. Schaaf B, Aries SP, Kramme E, Steinhoff J, Dalhoff K (2003) Acute renal failure associated with tenofovir treatment in a patient with acquired immunodeficiency syndrome. *Clin Infect Dis* 37: e41–43.
 9. Rollet F, Nazal EM, Chauvelot-Moachon L, Kelaidi C, Daniel N, et al. (2003) Tenofovir-related Fanconi syndrome with nephrogenic diabetes insipidus in a patient with acquired immunodeficiency syndrome: the role of lopinavir-ritonavir-didanosine. *Clin Infect Dis* 37: e174–176.
 10. Peyriere H, Reynes J, Rouanet I, Daniel N, de Boever CM, et al. (2004) Renal tubular dysfunction associated with tenofovir therapy: report of 7 cases. *J Acquir Immune Defic Syndr* 35: 269–273.
 11. Kinai E, Hanabusa H (2009) Progressive renal tubular dysfunction associated with long-term use of tenofovir DF. *AIDS Res Hum Retroviruses* 25: 387–394.
 12. Winston A, Amin J, Mallon P, Marriott D, Carr A, et al. (2006) Minor changes in calculated creatinine clearance and anion-gap are associated with tenofovir disoproxil fumarate-containing highly active antiretroviral therapy. *HIV Med* 7: 105–111.
 13. Gallant JE, Winston JA, DeJesus E, Pozniak AL, Chen SS, et al. (2008) The 3-year renal safety of a tenofovir disoproxil fumarate vs. a thymidine analogue-containing regimen in antiretroviral-naive patients. *AIDS* 22: 2155–2163.
 14. Fux CA, Simcock M, Wolbers M, Bucher HC, Hirschel B, et al. (2007) Tenofovir use is associated with a reduction in calculated glomerular filtration rates in the Swiss HIV Cohort Study. *Antivir Ther* 12: 1165–1173.
 15. Cooper RD, Wiebe N, Smith N, Keiser P, Naicker S, et al. (2010) Systematic review and meta-analysis: renal safety of tenofovir disoproxil fumarate in HIV-infected patients. *Clin Infect Dis* 51: 496–505.
 16. Gayet-Ageron A, Ananworanich J, Jupimai T, Chetchotisakd P, Prasithsirikul W, et al. (2007) No change in calculated creatinine clearance after tenofovir initiation among Thai patients. *J Antimicrob Chemother* 59: 1034–1037.
 17. Chaisiri K, Bowonwatanuwong C, Kasetratat N, Kiertiburanakul S (2010) Incidence and risk factors for tenofovir-associated renal function decline among Thai HIV-infected patients with low-body weight. *Curr HIV Res* 8: 504–509.
 18. Reid A, Stohr W, Walker AS, Williams IG, Kityo C, et al. (2008) Severe renal dysfunction and risk factors associated with renal impairment in HIV-infected adults in Africa initiating antiretroviral therapy. *Clin Infect Dis* 46: 1271–1281.
 19. Gatanaga H, Tachikawa N, Kikuchi Y, Teruya K, Genka I, et al. (2006) Urinary beta2-microglobulin as a possible sensitive marker for renal injury caused by tenofovir disoproxil fumarate. *AIDS Res Hum Retroviruses* 22: 744–748.
 20. Goicoechea M, Liu S, Best B, Sun S, Jain S, et al. (2008) Greater tenofovir-associated renal function decline with protease inhibitor-based versus nonnucleoside reverse-transcriptase inhibitor-based therapy. *J Infect Dis* 197: 102–108.
 21. Rodriguez-Novoa S, Labarga P, Soriano V, Egan D, Albalater M, et al. (2009) Predictors of kidney tubular dysfunction in HIV-infected patients treated with tenofovir: a pharmacogenetic study. *Clin Infect Dis* 48: e108–116.
 22. Bash LD, Coresh J, Kottgen A, Parekh RS, Fulop T, et al. (2009) Defining incident chronic kidney disease in the research setting: The ARIC Study. *Am J Epidemiol* 170: 414–424.
 23. Chue CD, Edwards NC, Davis LJ, Steeds RP, Townend JN, et al. (2011) Serum phosphate but not pulse wave velocity predicts decline in renal function in patients with early chronic kidney disease. *Nephrol Dial Transplant*;doi:10.1093/ndt/gfq78.
 24. Levey AS, Coresh J, Greene T, Stevens LA, Zhang YL, et al. (2006) Using standardized serum creatinine values in the modification of diet in renal disease study equation for estimating glomerular filtration rate. *Ann Intern Med* 145: 247–254.
 25. Gupta SK, Eustace JA, Winston JA, Boydstun II, Ahuja TS, et al. (2005) Guidelines for the management of chronic kidney disease in HIV-infected patients: recommendations of the HIV Medicine Association of the Infectious Diseases Society of America. *Clin Infect Dis* 40: 1559–1585.
 26. Lomaestro BM (2000) Fluoroquinolone-induced renal failure. *Drug Saf* 22: 479–485.
 27. Wever K, van Agtmael MA, Carr A (2010) Incomplete reversibility of tenofovir-related renal toxicity in HIV-infected men. *J Acquir Immune Defic Syndr* 55: 78–81.
 28. Szczech LA, Hoover DR, Feldman JG, Cohen MH, Gange SJ, et al. (2004) Association between renal disease and outcomes among HIV-infected women receiving or not receiving antiretroviral therapy. *Clin Infect Dis* 39: 1199–1206.
 29. Baxmann AC, Ahmed MS, Marques NC, Menon VB, Pereira AB, et al. (2008) Influence of muscle mass and physical activity on serum and urinary creatinine and serum cystatin C. *Clin J Am Soc Nephrol* 3: 348–354.
 30. Antoniou T, Raboud J, Chirhin S, Yoong D, Govan V, et al. (2005) Incidence of and risk factors for tenofovir-induced nephrotoxicity: a retrospective cohort study. *HIV Med* 6: 284–290.
 31. Horberg M, Tang B, Towner W, Silverberg M, Bersoff-Matcha S, et al. (2010) Impact of tenofovir on renal function in HIV-infected, antiretroviral-naive patients. *J Acquir Immune Defic Syndr* 53: 62–69.
 32. Post FA, Holt SG (2009) Recent developments in HIV and the kidney. *Curr Opin Infect Dis* 22: 43–48.
 33. Jullien V, Treluyer JM, Rey E, Jaffray P, Krivine A, et al. (2005) Population pharmacokinetics of tenofovir in human immunodeficiency virus-infected patients taking highly active antiretroviral therapy. *Antimicrob Agents Chemother* 49: 3361–3366.
 34. Van Rompay KK, Durand-Gasselín L, Brignolo LL, Ray AS, Abel K, et al. (2008) Chronic administration of tenofovir to rhesus macaques from infancy through adulthood and pregnancy: summary of pharmacokinetics and biological and virological effects. *Antimicrob Agents Chemother* 52: 3144–3160.
 35. Kohler JJ, Hosseini SH, Hoying-Brandt A, Green E, Johnson DM, et al. (2009) Tenofovir renal toxicity targets mitochondria of renal proximal tubules. *Lab Invest* 89: 513–519.
 36. Kiser JJ, Fletcher CV, Flynn PM, Cunningham CK, Wilson CM, et al. (2008) Pharmacokinetics of antiretroviral regimens containing tenofovir disoproxil fumarate and atazanavir-ritonavir in adolescents and young adults with human immunodeficiency virus infection. *Antimicrob Agents Chemother* 52: 631–637.

Renal Function Declines More in Tenofovir- than Abacavir-Based Antiretroviral Therapy in Low-Body Weight Treatment-Naïve Patients with HIV Infection

Takeshi Nishijima^{1,4}, Hiroyuki Gatanaga^{1,4*}, Hirokazu Komatsu³, Kuniyoshi Tsukada¹, Takuro Shimbo², Takahiro Aoki¹, Koji Watanabe^{1,4}, Ei Kinai¹, Haruhito Honda¹, Junko Tanuma¹, Hirohisa Yazaki¹, Miwako Honda¹, Katsuji Teruya¹, Yoshimi Kikuchi¹, Shinichi Oka^{1,4}

1 AIDS Clinical Center, National Center for Global Health and Medicine, Tokyo, Japan, **2** Department of Clinical Research and Informatics, International Clinical Research Center, National Center for Global Health and Medicine, Tokyo, Japan, **3** Department of Community Care, Saku Central Hospital, Nagano, Japan, **4** Center for AIDS Research, Kumamoto University, Kumamoto, Japan

Abstract

Objective: To compare the rate of decline of renal function in tenofovir- and abacavir-based antiretroviral therapy (ART) in low-body weight treatment-naïve patients with HIV infection.

Design: We conducted a single-center retrospective cohort study of 503 Japanese patients who commenced on either tenofovir- or abacavir-based initial ART.

Methods: The incidence of renal dysfunction, defined as more than 25% fall in estimated glomerular filtration rate (eGFR) from the baseline, was determined in each group. The effect of tenofovir on renal dysfunction was estimated by univariate and multivariate Cox hazards models as the primary exposure. Changes in eGFR until 96 weeks were estimated in both groups with a repeated measures mixed model.

Results: The median body weight of the cohort was 64 kg. The estimated incidence of renal dysfunction in the tenofovir and the abacavir arm was 9.84 per 100 and 4.55 per 100 person-years, respectively. Tenofovir was significantly associated with renal dysfunction by univariate and multivariate analysis (HR=1.747; 95% CI, 1.152–2.648; p=0.009) (adjusted HR=2.080; 95% CI, 1.339–3.232; p<0.001). In subgroup analysis of the patients stratified by intertertile baseline body weight, the effect of tenofovir on renal dysfunction was more evident in patients with lower baseline body weight by multivariate analysis (≤ 60 kg: adjusted HR=2.771; 95%CI, 1.494–5.139; p=0.001) (61–68 kg: adjusted HR=1.908; 95%CI, 0.764–4.768; p=0.167) (> 68 kg: adjusted HR=0.997; 95%CI, 0.318–3.121; p=0.995). The fall in eGFR was significantly greater in the tenofovir arm than the abacavir arm after starting ART (p=0.003).

Conclusion: The incidence of renal dysfunction in low body weight patients treated with tenofovir was twice as high as those treated with abacavir. Close monitoring of renal function is recommended for patients with small body weight especially those with baseline body weight < 60 kg treated with tenofovir.

Citation: Nishijima T, Gatanaga H, Komatsu H, Tsukada K, Shimbo T, et al. (2012) Renal Function Declines More in Tenofovir- than Abacavir-Based Antiretroviral Therapy in Low-Body Weight Treatment-Naïve Patients with HIV Infection. PLoS ONE 7(1): e29977. doi:10.1371/journal.pone.0029977

Editor: Claire Thorne, UCL Institute of Child Health, University College London, United States of America

Received: September 21, 2011; **Accepted:** December 7, 2011; **Published:** January 5, 2012

Copyright: © 2012 Nishijima et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by a Grant-in Aid for AIDS research from the Japanese Ministry of Health, Labour, and Welfare (H20-AIDS-002), and the Global Center of Excellence Program (Global Education and Research Center Aiming at the Control of AIDS) from the Japanese Ministry of Education, Science, Sports and Culture. No additional external funding was received for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: higtana@acc.ncgm.go.jp

Introduction

Tenofovir disoproxil fumarate (TDF) and abacavir sulfate (ABC) are widely used nucleot(s)ide reverse transcriptase inhibitors (NRTIs) as part of the initial antiretroviral therapy for patients with HIV infection in the developed countries (URL:<http://www.aidsinfo.nih.gov/ContentFiles/AdultandAdolescentGL.pdf>) (URL:http://www.europeanaidscinicalsociety.org/images/stories/EACS-Pdf/1_treatment_of_hiv_infected_adults.pdf). TDF is generally preferred to ABC, since ABC is reported to cause serious hypersensitivity

reaction in 5–8% of the patients and its efficacy in viral suppression is reported to be inferior to TDF among patients with baseline HIV viral load of $> 100,000$ copies/ml [1,2]. On the other hand, renal proximal tubular damage and renal dysfunction are well-known adverse effects of TDF [3–9]. A meta-analysis study that compared TDF and other NRTIs concluded that the decline in renal function with TDF use is significant but modest, and the ASSERT study conducted in Europe compared randomly-selected treatment naïve patients who commenced treatment with either TDF or ABC with efavirenz and showed no difference in estimated glomerular filtration

rate (eGFR) between the two groups at 48 weeks [9,10]. To date, the nephrotoxicity of TDF have been regarded as mild and tolerable [2,5–7,9–11].

However, the TDF-related nephrotoxicity has hardly been evaluated in patients with small body weight, who are potentially at higher risk for larger drug exposure and thus, more severe toxicity [12–15]. Indeed, some recent studies including ours reported a higher incidence of TDF-related renal dysfunction among Asian patients with low body weight compared with previous studies on mostly Whites and African Americans with larger body weight [13,16]. Thus, it is important to provide more evidence in support of TDF-associated nephrotoxicity in patients with low body weight since such data can elucidate whether TDF-related nephrotoxicity is as mild in low-body-weighted patients as previously reported in Europe and the USA. This is also important because there is increasing use of TDF in resource-limited settings, where patients are often of relatively small body weight, following the revised 2010 WHO guidelines that recommend TDF as one of the components of first line therapies (URL:http://whqlibdoc.who.int/publications/2010/9789241599764_eng.pdf) [13,16–19]. To our knowledge, there are no studies that compared renal function in treatment naïve Asian patients who commenced treatment with TDF or ABC.

Based on the above background, the present study was designed to compare the incidence of renal dysfunction and change in eGFR between treatment-naïve Japanese patients with low body weight who started either TDF or ABC as part of the antiretroviral regimen.

Methods

Ethics Statement

This study was approved by the Human Research Ethics Committee of National Center for Global Health and Medicine, Tokyo. All patients included in this study provided a written informed consent for their clinical and laboratory data to be used and published for research purposes. This study has been conducted according to the principles expressed in the Declaration of Helsinki.

Study Subjects

We performed a retrospective, single-center cohort study of HIV-infected Japanese patients using the medical records at the National Center for Global Health and Medicine, Tokyo, Japan. Our facility is one of the largest clinics for patients with HIV infection in Japan with more than 2,700 registered patients. The study population was treatment-naïve patients with HIV infection, aged >17 years, who commenced treatment with either the recommended 300 mg/day dose of TDF or 600 mg/day dose of ABC-containing antiretroviral regimen at our clinic between January 1, 2004 and March 31, 2009. During this inclusion period, all except two patients at our clinic started ART with either ABC or TDF. Patients with an eGFR of >60 ml/min/1.73 m² were enrolled. Patients were followed up until March 31, 2011. They were excluded if they started ART with both TDF and ABC, their follow-up period at our facility was less than 24 weeks after commencement of ART, or if they had started ART at other facilities. Only Japanese patients were included in order to examine a population with comparatively homogenous basic demographics and background. The attending physician selected either TDF or ABC at baseline, and the use of these two drugs was based on the Japanese guidelines, which place both ABC and TDF as the preferred NRTIs (<http://www.haart-support.jp/guideline2011.pdf>, in Japanese). The attending physician also selected

the key drug [non-nucleoside reverse transcriptase inhibitor (NNRTI), protease inhibitor (PI), or integrase inhibitor (INI)]. All patients received standard ART with two NRTIs combined with either PI, NNRTI, or INI.

Measurements

We defined renal dysfunction as more than 25% decrease in eGFR relative to the baseline [13,16,20,21]. The baseline eGFR was estimated for each patient from the average of two successive serum creatinine measurements made closest to and preceding the commencement of antiretroviral therapy by no more than 90 days. Changes in eGFR were plotted from the baseline measurement until the average value of two successive measurements diminished to less than 75% of the baseline, discontinuation of TDF or ABC, or at the end of the follow-up period. Discontinuation of TDF and ABC was the choice of the attending physician, and was based on virologic failure or ART-related side effects other than renal dysfunction. Before the initiation of ART and until suppression of HIV-1 viral load, patients visited our clinic every month. However, after viral load suppression, the visit interval was extended up to every three months. Serum creatinine and eGFR were measured in every visit, and the frequency of measurements was similar in patients on TDF and ABC. eGFR was calculated using the equation from the 4-variable Modification of Diet in Renal Disease (MDRD) study, $eGFR = 186 \times [\text{serum creatinine}]^{-1.154} \times [\text{age}]^{-0.203} \times [0.742 \text{ if patient is female}] \times [1.212 \text{ if patient is African American}]$ [22]. In this study, the primary exposure variable was TDF use over ABC as part of the initial ART.

The potential risk factors for renal dysfunction were determined according to previous studies and collected together with the basic demographics from the medical records [15,23–25]. They included age, sex, body weight, body mass index, (BMI) = {body weight (kg) / [(height (m))²]}, baseline laboratory data (CD4 cell count, HIV viral load, and serum creatinine), and presence or absence of other medical conditions (concurrent use of ritonavir-boosted protease inhibitors, concurrent nephrotoxic drugs such as ganciclovir, sulfamethoxazole/trimethoprim, and non-steroidal anti-inflammatory agents, diabetes mellitus defined by using anti-diabetic agents or fasting plasma glucose >126 mg/dl or plasma glucose >200 mg/dl on two different days, co-infection with hepatitis B defined by positive hepatitis B surface antigen, co-infection with hepatitis C defined by positive HCV viral load, hypertension defined by current treatment with antihypertensive agents or two successive measurements of systolic blood pressure >140 mmHg or diastolic blood pressure >90 mmHg at the clinic, dyslipidemia defined by current treatment with lipid-lowering agents, and current smoking). At our clinic, weight and blood pressure were measured on every visit whereas other variables were measured in the first visit and at least once annually. We used the data on or closest to and preceding the day of starting ART by no more than 90 days.

Statistical analysis

The time to 25% decline in eGFR from the baseline was calculated from the date of commencement of treatment to the date of diagnosis of the above-defined renal dysfunction. Censored cases represented those who discontinued ABC or TDF, dropped out, were referred to other facilities, or at the end of follow-up period. The time from the start of ART to >25% decrease in eGFR was analyzed by the Kaplan Meier method for patients who started TDF (TDF arm) and ABC (ABC arm), and the log-rank test was used to determine the statistical significance. The Cox proportional hazards regression analysis was used to estimate the

impact of TDF use over ABC on the incidence of more than 25% decrease in eGFR relative to the baseline. The impact of each basic demographics, baseline laboratory data, and other medical conditions listed above was also estimated with univariate Cox proportional hazards regression.

To estimate the unbiased prognostic impact of TDF use over ABC for renal dysfunction, we conducted three models using multivariate Cox proportional hazards regression analysis. Model 1 was the aforementioned univariate analysis for TDF use over ABC. Model 2 included age and weight plus model 1 in order to adjust for basic characteristics. In model 3, we added variables with P values <0.05 in univariate analysis for adjustment (these included age per 1 year, weight per 1 kg decrement, CD4 count per 1 μl decrement, HIV viral load per \log_{10}/ml , serum creatinine per 1 mg/dl , concurrent use of nephrotoxic drug(s), hepatitis B infection, and diabetes mellitus). The eGFR and the BMI were excluded from multivariate analysis because of their multicollinearity with age and serum creatinine, and weight, respectively, since eGFR and BMI are gained by the equation of those variables [22,26]. We chose to add weight instead of BMI because our previous work showed that weight was more useful and handy information to estimate the risk for TDF-related nephrotoxicity than BMI [16].

As a sensitivity analysis, creatinine clearance was similarly calculated with Cockcroft-Gault equation for each patient, $\text{creatinine clearance} = [(140 - \text{age}) \times \text{weight (kg)}] / (\text{serum creatinine} \times 72) (\times 0.85 \text{ for females})$ [27]. Actual body weight was used for the calculation. The impact of TDF use over ABC for $>25\%$ decrement of creatinine clearance from the baseline was estimated in univariate analysis and multivariate analysis adjusted with the before mentioned variables with Cox proportional hazards model.

To estimate the impact of weight on TDF-related nephrotoxicity, we did subgroup analysis for intertertile baseline body weight categories: ≤ 60 , 61–68, and >68 kg. Then, the abovementioned multivariate analysis with eGFR was conducted for each subgroup.

We also used a repeated measures mixed model to estimate and compare changes in eGFR between ABC and TDF from baseline to 2 years after initiation of ART by 6-month intervals adjusted for baseline eGFR and weight [10]. For each patient, the eGFR values at closest to and preceding 24, 48, 72 and 96 weeks after commencement of ART were collected. In this analysis, censoring occurred at discontinuation of TDF or ABC, leaving care, or reaching the end of the observation period before 96 weeks. Sensitivity analysis with creatinine clearance calculated by Cockcroft-Gault equation was similarly conducted.

Statistical significance was defined at two-sided p values <0.05 . We used hazard ratios (HRs) and 95% confidence intervals (95% CIs) to estimate the impact of each variable on renal dysfunction. All statistical analyses were performed with The Statistical Package for Social Sciences ver. 17.0 (SPSS, Chicago, IL).

Results

The study subjects were 199 patients in the TDF arm and 304 patients in the ABC arm who fulfilled the abovementioned criteria. Table 1 shows the demographics, laboratory data, and medical conditions of the study population at baseline. The majority of the study population was males, comparatively young and had a small stature (median weight, 64 kg, median BMI, 22.2 kg/m^2). More than 80% of the patients in the two arms had ritonavir-boosted PI. In the ABC arm, patients had significantly lower CD4 count ($p=0.006$), were significantly more likely to have hypertension

($p<0.001$), and tended to use more nephrotoxic drugs ($p=0.109$). On the other hand, in the TDF arm, patients had marginally higher baseline eGFR ($p=0.098$) and were significantly more likely to have hepatitis B virus infection ($P<0.001$). However, all other major background parameters were similar in the two groups (Table 1).

More than 25% decrement in eGFR from baseline occurred in 44 patients (22.1%) in the TDF arm and 41 (13.5%) in the ABC arm, with an estimated incidence of 9.84 and 4.55 per 100 person-years, respectively. Figure 1 shows the time from ART initiation to $>25\%$ decrease in eGFR by the Kaplan Meier method in the two groups. Patients who started TDF-containing ART were significantly more likely to develop renal dysfunction, compared to the ABC group ($p=0.001$, Log-rank test). The median time from commencement of ART to occurrence of $>25\%$ decrement in eGFR was 246 days (range, 1–1,339 days) for the TDF arm and 501 days (range, 7–2,022) for ABC arm. The total observation period was 447.2 patient-years [median, 839 days, interquartile range (IQR), 357–1137 days] for the TDF arm and 901.7 patient-years (median, 1,119 days, IQR, 660.5–1509 days) for the ABC arm.

Univariate analysis showed a significant relationship between TDF use and $>25\%$ decrement in eGFR (HR = 1.747; 95%CI, 1.152–2.648; $p=0.009$) (Table 2). Furthermore, old age, small body weight, low baseline CD4 count, high HIV viral load, high eGFR, low serum creatinine, concurrent use of nephrotoxic drugs, hepatitis B infection, and diabetes mellitus were associated with renal dysfunction. On the other hand, concurrent use of ritonavir boosted PIs was not associated with renal dysfunction (HR = 1.220; 95%CI, 0.663–2.244; $p=0.523$). Multivariate analysis identified TDF use as a significant risk for $>25\%$ decrement in eGFR after adjustment for age and weight (adjusted HR = 1.893; 95%CI, 1.243–2.881; $p<0.003$) (Table 3, Model 2), and also after adjustment for other risk factors (adjusted HR = 2.080; 95%CI, 1.339–3.232; $p<0.001$) (Table 3, Model 3). We also conducted a sensitivity analysis using BMI decrement instead of weight as a variable in Table 3, Model 3. The results were almost identical; TDF use over ABC use was a risk for renal dysfunction (adjusted HR 1.957, 95% CI 1.262–3.036, $p=0.003$).

Sensitivity analysis with creatinine clearance confirmed the abovementioned findings: both univariate and multivariate analyses showed that TDF use was significantly associated with $>25\%$ decrement in eGFR (univariate analysis: HR = 2.212; 95%CI, 1.340–3.653; $p=0.002$) (multivariate analysis: adjusted HR = 2.544; 95%CI, 1.493–4.335; $p=0.001$).

Subgroup analysis of the patients stratified by intertertile baseline body weight showed that the lower the baseline body weight, the more evident the impact of TDF on renal dysfunction (≤ 60 kg: adjusted HR = 2.771; 95%CI, 1.494–5.139; $p=0.001$) (61–68 kg: adjusted HR = 1.908; 95%CI, 0.764–4.768; $p=0.167$) (>68 kg: adjusted HR = 0.997; 95%CI, 0.318–3.121; $p=0.995$) (Table 4). These findings suggest that there is the effect modification by baseline body weight on TDF-associated renal dysfunction.

Data analysis by repeated measures mixed models showed a significant decrease in adjusted mean eGFR from the baseline to 96 weeks in both groups (TDF: $-9.984 \text{ ml}/\text{min}/1.73\text{m}^2$, 95%CI -12.05 to $-7.914 \text{ ml}/\text{min}/1.73\text{m}^2$, $p<0.001$; ABC: $-5.393 \text{ ml}/\text{min}/1.73\text{m}^2$, 95%CI -7.087 to $-3.699 \text{ ml}/\text{min}/1.73\text{m}^2$, $p<0.001$) (Figure 2). There was a statistically significant interaction between the two arms over time ($p=0.003$), indicating that adjusted mean eGFR decreased more significantly in the TDF group than in the ABC group after initiation of ART. Analysis of eGFR in each group demonstrated a rapid decrease during the first 24 weeks,

Table 1. Baseline demographics and laboratory data of patients who received tenofovir- and abacavir-based antiretroviral therapy (n = 503).

	TDF (n = 199)	ABC (n = 304)	P value
Sex (male), n (%)	196 (98.5)	296 (97.4)	0.539
Median (IQR) age	36 (31–44)	37 (31–43)	0.436
Median (IQR) weight (kg)	64 (58–69)	64 (58.0–70.9)	0.426
Median (IQR) BMI (kg/m ²)	22.1 (20.4–23.9)	22.2 (20.3–24.6)	0.321
Median (IQR) eGFR (ml/min/1.73m ²)	119.4 (103.0–135.0)	115.6 (102.4–132.2)	0.098
Median (IQR) serum creatinine (mg/dl)	0.74 (0.67–0.84)	0.75 (0.68–0.83)	0.250
Median (IQR) CD4 count (/μl)	199 (109–272)	178.5 (75.3–234.8)	0.006
Median (IQR) HIV RNA viral load (log ₁₀ /ml)	4.63 (4.20–5.20)	4.74 (4.23–5.20)	0.731
Ritonavir-boosted protease inhibitors, n (%)	173 (86.9)	256 (84.2)	0.441
Protease inhibitors (unboosted), n (%)	5 (2.5)	20 (6.6)	0.038
NNRTIs, n (%)	16 (8.0)	26 (8.6)	0.848
INIs, n (%)	5 (2.5)	2 (0.7)	0.119
Hypertension, n (%)	5 (2.5)	53 (17.4)	<0.001
Dyslipidemia, n (%)	4 (2.0)	4 (1.3)	0.718
Diabetes mellitus, n (%)	8 (4.0)	12 (3.9)	1.000
Concurrent use of nephrotoxic drugs, n (%)	65 (32.7)	121 (39.8)	0.109
Hepatitis B, n (%)	35 (17.6)	9 (3.0)	<0.001
Hepatitis C, n (%)	7 (3.5)	7 (2.3)	0.421
Current smoker, n (%)	93 (46.7)	149 (49.3)	0.585

TDF: tenofovir, ABC: abacavir, IQR: interquartile range, BMI: body mass index, eGFR: estimated glomerular filtration rate, NNRTI: non- nucleoside reverse transcriptase inhibitor, INI: integrase inhibitor.

doi:10.1371/journal.pone.0029977.t001




Figure 1. Kaplan-Meier curve showing the time to renal dysfunction in patients treated with TDF or ABC. Compared to treatment-naïve patients who commenced treatment with ABC, those on TDF were more likely to develop >25% fall in eGFR ($p = 0.001$, Log-rank test). TDF: tenofovir, ABC: abacavir, ART: antiretroviral therapy, eGFR: estimated glomerular filtration rate.

doi:10.1371/journal.pone.0029977.g001

Table 2. Univariate analysis to estimate the risk of various factors in inducing more than 25% fall in eGFR.

	Hazard ratio	95% CI	P value
TDF vs. ABC use	1.747	1.152–2.648	0.009
Female gender	0.048	0.000–16.93	0.310
Age per 1 year	1.031	1.011–1.051	0.002
Weight per 1 kg decrement	1.047	1.023–1.072	<0.001
BMI per 1 kg/m ² decrement	1.152	1.066–1.244	<0.001
CD4 count per 1 / μ l decrement	1.006	1.004–1.008	<0.001
HIV viral load per log ₁₀ /ml	1.562	1.179–2.071	0.002
Ritonavir-boosted protease inhibitors	1.220	0.663–2.244	0.523
Baseline eGFR per 1 ml/min/1.73m ²	1.009	1.005–1.014	<0.001
Baseline serum creatinine per 1mg/dl	0.016	0.003–0.086	<0.001
Concurrent nephrotoxic drug	2.134	1.417–3.214	<0.001
Hepatitis B	1.866	1.038–3.356	0.037
Hepatitis C	1.721	0.631–4.695	0.289
Diabetes mellitus	2.558	1.181–5.540	0.017
Hypertension	0.865	0.448–1.669	0.664
Current smoking	0.989	0.657–1.489	0.958

eGFR: estimated glomerular filtration rate, CI: confidence interval, TDF: tenofovir, ABC: abacavir, BMI: body mass index.
doi:10.1371/journal.pone.0029977.t002

followed by a plateau until 96 weeks. In sensitivity analysis with creatinine clearance calculated by Cockcroft-Gault equation, the result was the same; a significant decrease from the baseline to 96 weeks in both groups (TDF: -10.62 ml/min, 95%CI -13.78 to -7.458 ml/min; ABC: -4.325 ml/min, 95%CI -6.893 to -1.756 ml/min) and significantly more eGFR decrement in the TDF group ($p = 0.019$).

Discussion

In this observational Japanese cohort, treatment-naïve patients who started TDF-containing ART experienced eGFR decline of >25% approximately twice as likely compared to those treated with ABC-containing regimen. Univariate and multivariate analyses identified TDF use as an independent risk factor for

Table 4. Multivariate analysis to estimate the risk of TDF- over ABC-based antiretroviral therapy in the induction of more than 25% fall in eGFR according to baseline body weight.

	Adjusted HR	95% CI	P value
Baseline body weight ≤ 60 kg (n = 171)			
TDF vs. ABC use	2.771	1.494–5.139	0.001
Baseline body weight 61–68 kg (n = 167)			
TDF vs. ABC use	1.908	0.764–4.768	0.168
Baseline body weight >68 kg (n = 165)			
TDF vs. ABC use	0.997	0.318–3.121	0.995

TDF use was adjusted with the same variables indicated in Model 3, Table 3: age per 1 year, weight per 1 kg decrement, CD4 count per 1 / μ l decrement, HIV viral load per log₁₀/ml, serum creatinine per 1 mg/dl, concurrent use of nephrotoxic drugs, hepatitis B infection, and diabetes mellitus.
doi:10.1371/journal.pone.0029977.t004

renal dysfunction. Subgroup analysis showed that the effect of TDF on renal dysfunction was more evident in patients with lower body weight. Furthermore, eGFR decrement was significantly larger in the TDF group than in ABC group over the 2-year observation period.

In our previous study, we demonstrated a high incidence of TDF-associated nephrotoxicity in patients with low body weight, and the use of a robust statistical model indicated a greater decline in renal function in patients of low body weight treated with TDF [16]. The results of the present study further emphasize the importance of low body weight as a risk factor for TDF-related nephrotoxicity by showing that in a cohort of patients with low body weight, the incidence of renal dysfunction was twice higher with TDF use than with ABC use.

Among the studies designed to compare renal function after the commencement of TDF and ABC-containing ART for treatment-naïve patients, our cohort had the lowest median body weight (64 kg). This is lower than the median body weight of patients of the ASSERT study conducted in European countries (72 kg) [10]. The

Table 3. Multivariate analysis to estimate the risk of TDF- over ABC-based antiretroviral therapy in inducing more than 25% fall in eGFR.

	Model 1 Crude		Model 2 Adjusted		Model 3 Adjusted	
	HR	95% CI	HR	95%CI	HR	95%CI
TDF vs. ABC use [†]	1.747	1.152–2.648	1.893	1.243–2.881	2.080	1.339–3.232
Age per 1 year			1.029	1.010–1.048	1.020	1.000–1.040
Weight per 1 kg decrement [†]			1.046	1.022–1.071	1.028	1.005–1.052
CD4 count per 1 / μ l decrement [†]					1.004	1.002–1.007
HIV viral load per log ₁₀ /ml					1.048	0.749–1.466
Serum creatinine per 1 mg/dl [†]					0.053	0.009–0.304
Use of nephrotoxic drug					1.309	0.825–2.077
Hepatitis B					1.070	0.573–2.000
Diabetes mellitus					1.565	0.684–3.582

[†]P<0.05 in Model 3.

TDF: tenofovir, ABC: abacavir, eGFR: estimated glomerular filtration rate, HR: hazard ratio, CI: confidence interval.
doi:10.1371/journal.pone.0029977.t003