

Table 1. List of 132 Human Proteins Interacting with the HCV NSSA Protein

gene ID	symbol	description	refs
47	ACLY	ATP citrate lyase	22
60	ACTB	actin, beta	101
79026	AHNAK	AHNAK nucleoprotein	22
10598	AHSA1	AHA1, activator of heat shock 90 kDa protein ATPase homologue 1 (yeast)	102
207	AKT1	v-akt murine thymoma viral oncogene homologue 1	22
302	ANXA2	annexin A2	105
335	APOA1	apolipoprotein A-I	22
348	APOE	apolipoprotein E	22
116985	ARAP1	ArfGAP with RhoGAP domain, ankyrin repeat and PH domain 1	22
27236	ARFIP1	ADP-ribosylation factor interacting protein 1	22
23204	ARL6IP1	ADP-ribosylation factor-like 6 interacting protein 1	this study
4508	ATP6	ATP synthase F0 subunit 6	this study
8312	AXIN1	axin 1	22
581	BAX	BCL2-associated X protein	22
222389	BEND7	BEN domain containing 7	22
274	BIN1	bridging integrator 1	this study; 22,47
89927	C16orf45	chromosome 16 open reading frame 45	this study
8618	CADPS	Ca ²⁺ -dependent secretion activator	22
93664	CADPS2	Ca ²⁺ -dependent secretion activator 2	22
79080	CCDC86	coiled-coil domain containing 86	22
983	CDK1	cyclin-dependent kinase 1	22
1021	CDK6	cyclin-dependent kinase 6	22
1060	CENPC1	centromere protein C 1	22
153241	CEP120	centrosomal protein 120 kDa	22
11190	CEP250	centrosomal protein 250 kDa	22
9702	CEPS7	centrosomal protein 57 kDa	22
80254	CEP63	centrosomal protein 63 kDa	22
1381	CRABP1	cellular retinoic acid binding protein 1	22
1445	CSK	c-src tyrosine kinase	22
1452	CSNK1A1	casein kinase 1, alpha 1	104
1457	CSNK2A1	casein kinase 2, alpha 1 polypeptide	63,105
1499	CTNNB1	catenin (cadherin-associated protein), beta 1, 88 kDa	84,106
9093	DNAJA3	DnaJ (Hsp40) homologue, subfamily A, member 3	22
2202	EFEMP1	EGF containing fibulin-like extracellular matrix protein 1	22
5610	EIF2AK2	eukaryotic translation initiation factor 2-alpha kinase 2	22
2051	EPHB6	EPH receptor B6	this study
54942	FAM206A	family with sequence similarity 206, member A	22
25827	FBXL2	F-box and leucine-rich repeat protein 2	22
2274	FHL2	four and a half LIM domains 2	22
23770	FKBP8	FK506 binding protein 8, 38 kDa	this study; 43,45
2316	FLNA	filamin A, alpha	12
2495	FTH1	ferritin, heavy polypeptide 1	22
8880	FUBP1	far upstream element (FUSE) binding protein 1	107
2534	FYN	FYN oncogene related to SRC, FGR, YES	22
11345	GABARAPL2	GABA(A) receptor-associated protein-like 2	this study
54826	GIN1	gypsy retrotransposon integrase 1	22
2801	GOLGA2	golgin A2	22
2874	GPS2	G protein pathway suppressor 2	22
2885	GRB2	growth factor receptor-bound protein 2	22
2931	GSK3A	glycogen synthase kinase 3 alpha	22
2932	GSK3B	glycogen synthase kinase 3 beta	22
3055	HCK	hemopoietic cell kinase	22
3320	HSP90AA1	heat shock protein 90 kDa alpha (cytosolic), class A member 1	22
3303	HSPA1A	heat shock 70 kDa protein 1A	108
3315	HSPB1	heat shock 27 kDa protein 1	109
3537	IGLC1	immunoglobulin lambda constant 1 (Mcg marker)	22
79711	IPO4	importin 4	22
3843	IPOS	importin 5	22
3683	ITGAL	integrin, alpha L (antigen CD11A (p180), lymphocyte function-associated antigen 1; alpha polypeptide)	22
6453	ITSN1	intersectin 1	this study
3716	JAK1	Janus kinase 1	22

Table 1. continued

gene ID	symbol	description	refs
3932	LCK	lymphocyte-specific protein tyrosine kinase	22
55679	LIMS2	LIM and senescent cell antigen-like domains 2	22
4067	LYN	v-yes-1 Yamaguchi sarcoma viral related oncogene homologue	22
9448	MAP4K4	mitogen-activated protein kinase kinase kinase kinase 4	this study
6300	MAPK12	mitogen-activated protein kinase 12	22
4155	MBP	myelin basic protein	22
4256	MGP	matrix Gla protein	110
55233	MOB1A	MOB kinase activator 1A	22
4673	NAP1L1	nucleosome assembly protein 1-like 1	22
4674	NAP1L2	nucleosome assembly protein 1-like 2	22
10397	NDRG1	N-myc downstream regulated 1	22
4778	NFE2	nuclear factor (erythroid-derived 2), 45 kDa	22
11188	NISCH	nischarin	this study
4924	NUCB1	nucleobindin 1	22
4938	OAS1	2'-5'-oligoadenylate synthetase 1, 40/46 kDa	22
5007	OSBP	oxysterol binding protein	111
64098	PARVG	parvin, gamma	22
5170	PDPK1	3-phosphoinositide dependent protein kinase-1	22
5297	PI4KA	phosphatidylinositol 4-kinase, catalytic, alpha	22
5291	PIK3CB	phosphoinositide-3-kinase, catalytic, beta polypeptide	22
5295	PIK3R1	phosphoinositide-3-kinase, regulatory subunit 1 (alpha)	55,84,106
5300	PIN1	peptidylprolyl cis/trans isomerase, NIMA-interacting 1	112
5307	PITX1	paired-like homeodomain 1	22
5347	PLK1	polo-like kinase 1	113
10654	PMVK	phosphomevalonate kinase	22
5478	PPIA	peptidylprolyl isomerase A (cyclophilin A)	114,115
10848	PPP1R13L	protein phosphatase 1, regulatory subunit 13 like	22
5515	PPP2CA	protein phosphatase 2, catalytic subunit, alpha isozyme	116
5518	PPP2R1A	protein phosphatase 2, regulatory subunit A, alpha	116
5698	PSMB9	proteasome (prosome, macropain) subunit, beta type, 9 (large multifunctional peptidase 2)	22
5757	PTMA	prothymosin, alpha	22
5894	RAF1	v-raf-1 murine leukemia viral oncogene homologue 1	22
6142	RPL18A	ribosomal protein L18a	22
6167	RPL37	ribosomal protein L37	this study
6238	RRBP1	ribosome binding protein 1 homologue 180 kDa (dog)	22
91543	RSAD2	radical S-adenosyl methionine domain containing 2	117
6252	RTN1	reticulum 1	this study
10313	RTN3	reticulum 3	this study
6424	SFRP4	secreted frizzled-related protein 4	22
81858	SHARPIN	SHANK-associated RH domain interactor	22
64754	SMYD3	SET and MYND domain containing 3	22
8470	SORBS2	sorbin and SH3 domain containing 2	22
10174	SORBS3	sorbin and SH3 domain containing 3	22
6714	SRC	v-src sarcoma (Schmidt-Ruppin A-2) viral oncogene homologue (avian)	22
10847	SRCAP	Snf2-related CREBBP activator protein	22
6741	SSB	Sjogren syndrome antigen B (autoantigen La)	22
284297	SSCS5D	scavenger receptor cysteine rich domain containing (5 domains)	110
6772	STAT1	signal transducer and activator of transcription 1	118
25777	SUN2	Sad1 and UNC84 domain containing 2	this study
6850	SYK	spleen tyrosine kinase	119
4070	TACSTD2	tumor-associated calcium signal transducer 2	22
6880	TAF9	TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32 kDa	22
6908	TBP	TATA box binding protein	22
7046	TGFBR1	transforming growth factor, beta receptor 1	22
7057	THBS1	thrombospondin 1	22
374395	TMEM179B	transmembrane protein 179B	22
7110	TMF1	TATA element modulatory factor 1	22
7157	TP53	tumor protein p53	22
7159	TP53BP2	tumor protein p53 binding protein, 2	22
7186	TRAF2	TNF receptor-associated factor 2	22
11078	TRIOBP	TRIO and F-actin binding protein	22

Table 1. continued

gene ID	symbol	description	refs
51061	TXNDC11	thioredoxin domain containing 11	22
53347	UBASH3A	ubiquitin associated and SH3 domain containing A	22
10869	USP19	ubiquitin specific peptidase 19	22
9218	VAPA	VAMP (vesicle-associated membrane protein)-associated protein A, 33 kDa	22
9217	VAPB	VAMP (vesicle-associated membrane protein)-associated protein B and C	this study; ^{22,28,46}
10493	VAT1	vesicle amine transport protein 1 homologue (<i>T. californica</i>)	this study
55737	VPS35	vacuolar protein sorting 35 homologue (<i>S. cerevisiae</i>)	22
6293	VPS52	vacuolar protein sorting 52 homologue (<i>S. cerevisiae</i>)	22
140612	ZFP28	zinc finger protein 28 homologue (mouse)	this study
9726	ZNF646	zinc finger protein 646	22

Quantitative Reverse-Transcription PCR (qRT-PCR)

Total RNA was prepared from the cell and culture supernatant using the RNeasy mini kit (QIAGEN, Hilden, Germany) and QIAamp Viral RNA Mini Kit (QIAGEN), respectively. First-strand cDNA was synthesized using high capacity cDNA reverse transcription kit (Applied biosystems, Carlsbad, CA, USA) with random primers. Each cDNA was estimated by Platinum SYBR Green qPCR Super Mix UDG (Invitrogen) as per the manufacturer's protocol. Fluorescent signals of SYBR Green were analyzed with ABI PRISM 7000 (Applied Biosystems). The HCV internal ribosomal entry site (IRES) region and human glyceraldehyde-3-phosphate dehydrogenase (GAPDH) gene were amplified with the primer pairs 5'-GAGTGTCTGTCAGCCTCCA-3' and 5'-CACTCGCAAGCACCTATCA-3', and 5'-GAAGTCTGGAGTCAACGGATT-3' and 5'-GATGACAAGCTTCCCCTTCTC-3', respectively.⁴² The quantities of the HCV genome and the other host mRNAs were normalized with that of GAPDH mRNA. RTN1 and RTN3 genes were amplified using the primer pairs purchased from QIAGEN.

Cell Lines and Virus Infection

Cells from the Huh7OK1 cell line are highly permissive to HCV JFH1 strain (genotype 2a) infection compared to Huh 7.5.1 and exhibit the highest propagation efficiency for JFH1.⁴³ These cells were maintained at 37 °C in a humidified atmosphere and 5% CO₂ in the Dulbecco's modified Eagle's medium (DMEM) (Sigma, St. Louis, MO, USA) supplemented with nonessential amino acids (NEAA) and 10% fetal calf serum (FCS). The viral RNA of JFH1 was introduced into Huh7OK1 as described by Wakita et al.⁴⁴ The viral RNA of JFH1 derived from the plasmid pJFH1 was prepared as described by Wakita et al.⁴⁴

Statistical Analysis

Experiments for RNAi transfection and qRT-PCR were performed two times. The estimated values were represented as the mean \pm standard deviation ($n = 2$). The significance of differences in the means was determined by the Student's *t*-test.

RESULTS AND DISCUSSION

Identifying Host Proteins That Interact with HCV NSSA Protein

We employed an integrated approach that combined an experimental Y2H assay and comprehensive literature mining to identify human host proteins interacting with NSSA.

First, we performed an Y2H screening to characterize the interactions between NSSA and host proteins. The analysis of positive colonies revealed 17 host factors as interacting partners

of NSSA (Tables 1, S1, Supporting Information), 14 of which are novel. The other three interactions have been characterized previously; vesicle-associated membrane protein (VAMP)-associated protein B (VAPB), a membrane trafficking factor, and FK506-binding protein 8 (FKBP8), an immunoregulation protein, independently regulate HCV replication via interactions with NSSA;^{28,43,45,46} Bridging integrator 1 (BIN1), a tumor suppressor protein, interacts with NSSA and significantly contributes to HCC.⁴⁷ Among the newly discovered interactors, MAP4K4 is overexpressed in HCC, and knock-down of MAP4K4 expression inhibits HCC progression;⁴⁸ RTN1 and VAT1 were previously observed to be elevated in HCV infected cells,⁴⁹ and ARL6IP1, EPHB6, GABARAPL2, ITSN1 and NISCH were differentially expressed in HCV infection *in vitro*.⁵⁰ Furthermore, five (ARL6IP1, FKBP8, RTN1, RTN3, VAPB) of the 17 interactors (29.4%) localize to the endoplasmic reticulum (ER; GO:0005783; $p = 0.0028$), which is consistent with the role of NSSA as a crucial constituent of the HCV replication complex associated with the ER.⁵¹ These results suggest that the PPIs detected by our Y2H assay may closely reflect NSSA interactions *in vivo*.

We next scanned the biomedical literature to expand the repertoire of NSSA–host interactions. Because of an ever increasing volume of biomedical literature describing the pathogenesis of infectious diseases, the identification of specific host–pathogen interactions and their roles in pathogenicity is a nontrivial task, and therefore, recent years have witnessed a rapid development of computational tools for biomedical literature mining. We performed extensive literature mining using computational tools that facilitate the retrieval and extraction of relevant information from the biomedical literature (Pubmed, EBIMed, Protein Coral) and followed it up with a careful manual inspection to identify additional host factors, which directly interact with NSSA and which were not present in the Y2H data set. One hundred and fifteen pairwise interactions between NSSA and human proteins (consisting of 93 catalogued by a high throughput study of binary HCV–host interactions²² and 22 from assorted reports; see Supporting Information, Table S2) were extracted from the literature in this manner and were added to the existing interactors. The resulting NSSA–human interactome thus comprised 132 human host proteins directly interacting with NSSA (Table 1), all of which are expressed in the liver (see Supporting Information, Table S3).

Network Topological Analysis of the NSSA–host Interactions: NSSA Preferentially Targets Hubs and Bottlenecks in the Host Protein Interactome

To further understand the biological significance of the NSSA–host interactions, we retrieved PPIs for the nodes targeted by




Figure 1. Topological analysis of the NSSA infection network. (A) The node degree distributions of the NSSA interactors in the HPI, NSSA infection network, and HPI are represented as box plots. The average degree of the NSSA interactors in HPI (19.02) was higher than those of the NSSA infection network (8.24) and HPI (5.96). Median node degrees (indicated by thick horizontal lines) of the NSSA interactors in HPI, NSSA infection network, and HPI are 8, 5, and 3, respectively. (B) The average degree of the nodes targeted by NSSA in HPI was much higher than mean average degree of 1000 sets of the randomly selected 108 nodes in HPI. (C) The shortest path length distributions of the NSSA infection network and HPI. The path length is represented on the *x*-axis while the *y*-axis describes the frequency, i.e., the percentage of node (protein) pairs within the PPI network with a given shortest path length. For simplicity, only the node frequencies for path lengths 1–5 in the HPI are displayed. (D) The number of bottlenecks among the nodes targeted by NSSA in HPI was much higher than mean of the number of bottlenecks among 1000 sets of the randomly selected 108 nodes in HPI. *: $p < 0.001$.

NSSA in the HPI and incorporated them with the initial interactions to infer an extended NSSA infection network. PPIs for 108 of 132 NSSA interactors were retrieved in this manner; 24 of 132 NSSA interactors had no PPIs in the HPI (Supporting Information, Tables S4, S5a, S5b). For the NSSA infection network and the HPI, we computed the node degree distribution and the characteristic/average path length measures to capture the topologies of the two networks. The degree of a protein, which corresponds to the number of its interacting partners, may often reflect its biological relevance since a better connected protein is likely to have a higher ability to influence biological networks via PPIs. Average path lengths provide an approximate measure of the relative ease and speed of dissemination of information between the proteins in a network.

The NSSA infection network consisted of 1442 entities (nearly all of which are expressed in the liver; see Supporting Information) with 6263 interactions between them (Supporting Information, Tables S4, S5a). The average degree (defined as the number of interactions for a given protein) of the NSSA infection network (8.24) was notably higher than the degree inferred for the HPI (5.96) (Figure 1A). Furthermore, the

average degree of the nodes targeted by NSSA in the HPI (19.02) was even higher; this number is significantly greater than the average degree obtained from a sample of randomly selected nodes (6.17 ± 1.08 with $p < 0.001$; Figure 1B; see Supporting Information). Also the degrees inferred for the majority of the NSSA interactors in the HPI (65 of 108; 60.18%) were higher than the mean degree of the HPI (5.96) (Figure 1A). Our observations therefore suggest that NSSA preferentially targets several highly connected cellular proteins (hubs) with an ability to influence a large number of host factors in HCV infection. The average (shortest) path length of the NSSA infection network (3.26) was significantly shorter than the HPI (4.54), and also the distribution of shortest path lengths was shifted toward the left (Figure 1C), thereby suggesting that the NSSA influenced cellular network is more compact and inclined toward faster communication between the constituents relative to the host cellular network.

Next, we examined the betweenness measures of the NSSA interactors in the HPI to assess their significance in the HPI and the NSSA infection network. The betweenness of a node, determined by the number of shortest paths passing through it, reflects the importance of that node in the network; the nodes

with the highest betweenness prominently regulate the flow of signaling information and are therefore “bottlenecks”, representing central points for communication in an interaction network.⁵² Previously, proteins with high betweenness have been implicated in crucial roles in HCV infection and pathogenesis.^{53,54} To investigate if NSSA preferentially targets bottlenecks (defined as the top 10% of the nodes in the HPI ranked by betweenness), we estimated the fraction of NSSA interactors that were bottlenecks in the HPI. A significant proportion (39 of 108; 36.1%) of the NSSA interactors were identified as bottlenecks in the HPI (Supporting Information, Table S6); this number is significantly higher than the number of bottlenecks among randomly selected nodes (10.72 ± 3.17 with $p < 0.001$; Figure 1D; see Supporting Information). These include growth factor receptor-binding protein 2 (GRB2), which plays an important role in the subversion of host signaling pathways by NSSA;⁵⁵ tumor protein 53 (TP53), a key mediator of the oncogenic effect of NSSA in HCV-induced HCC;⁵⁶ and tyrosine kinase SRC, which regulates the formation of NSSA-containing HCV replication complex.⁵⁷ Among the NSSA interacting proteins identified by our Y2H screening, ITSN1, an endocytic traffic associated protein, and GABARAPL2, an autophagy associated protein, were identified as network bottlenecks.

Our observations therefore suggest that NSSA preferentially interacts with highly central proteins in the host protein interactome; these interactions may help the virus to regulate efficiently the flow of the infection-related information in the host cellular network and manipulate the host metabolic machinery for its own survival and pathogenesis. Our observations are consistent with studies that suggested that viral pathogens tend to interact with well-connected host proteins that are central to the host cellular networks, thus enabling them to appropriate essential cellular functions.^{21,22,26,58,59}

Functional Analysis of NSSA Interaction Network

Next, we investigated the NSSA infection network for the enrichment of specific biological associations (KEGG pathways, CATH structural domains; GO terms and Reactome Pathways; Supporting Information, Tables S7a, S7b, S7c and S7d). Notably, a significant proportion of the proteins in the NSSA infection network were mapped to the CATH Phosphorylase Kinase; domain 1, domain (CATH:3.30.200.20; 138 out of 1442, $p = 2.61 \times 10^{-45}$) including 23 of the 132 NSSA interacting host proteins ($p = 3.38 \times 10^{-14}$) (13 of which are bottlenecks in the HPI), based on the Gene3D protein domain assignments (Supporting Information, Table S7b). These include two novel interactions between EPHB6 (a kinase deficient receptor) and MAP4K4 and NSSA, identified by our Y2H assay (Table 1). The significant representation of cellular kinases in the NSSA infection network is consistent with the key roles played by reversible phosphorylation of NSSA in modulating various NSSA functions in HCV pathogenesis. Impairing NSSA hyperphosphorylation has been shown to inhibit HCV replication, and thus, the cellular kinases that regulate NSSA phosphorylation are important targets for anti-HCV therapy.^{9,60–63}

The analysis of NSSA infection network revealed an enrichment of 79 KEGG pathways (Supporting Information, Table S7a). Furthermore, 31 of the 39 NSSA interacting bottlenecks (hereafter referred to as bottlenecks) were mapped to 75 of the 79 enriched KEGG pathways (Supporting

Information, Table S5). Among the 75 bottleneck-associated enriched KEGG pathways, the highest numbers were associated with various cancers and infectious diseases (31 enriched KEGG pathways; 27 bottlenecks), followed by immune system, signal transduction and endocrine system (23 enriched KEGG pathways; 27 bottlenecks), cell growth and death (4 enriched KEGG pathways; 9 bottlenecks), nervous system (4 enriched KEGG pathways; 8 bottlenecks) and cellular communication (3 enriched KEGG pathways; 14 bottlenecks) among others (Tables 2, S8a, Supporting Information). Below we describe our observations on the most prominent enriched biological themes of interest that were associated with the NSSA infection network, with a specific focus on the bottlenecks.

Cancers and Infectious Diseases

The analysis of the NSSA interaction network revealed that NSSA specifically targets host factors that participate in various complex human diseases. Thirty-four NSSA interactors including 24 bottlenecks were mapped to one or more of the 17 enriched KEGG pathways associated with different infectious diseases (Supporting Information, Tables S7a, S8a). Among the most prominent associations, 12 bottlenecks were mapped to “Epstein–Barr virus infection” ($p = 1.36 \times 10^{-27}$); 10 to “Hepatitis C” ($p = 3.47 \times 10^{-24}$); 10 to “HTLV-I infection” ($p = 1.39 \times 10^{-20}$); 9 to “Hepatitis B” ($p = 3.33 \times 10^{-26}$); 8 to “Measles” ($p = 5.69 \times 10^{-17}$); 7 bottlenecks were mapped to “Influenza A” ($p = 5.01 \times 10^{-12}$); 7 to “Herpes simplex infection” ($p = 1.47 \times 10^{-13}$) and 6 to “Tuberculosis” ($p = 3.02 \times 10^{-6}$) (Supporting Information, Tables S7a, S8a). These associations include infectious diseases induced by various bacterial and viral pathogens thereby suggesting that HCV and other pathogens may systematically target specific host factors, the perturbation of which may contribute to the onset of various human diseases.

Also, 19 bottlenecks were mapped to one or more of the 16 enriched KEGG pathways associated with various cancers. Among the most prominent associations, 10 bottlenecks were mapped to “Viral carcinogenesis” ($p = 1.3 \times 10^{-30}$); 8 each were mapped to “Prostate cancer” ($p = 4.27 \times 10^{-25}$), “Endometrial cancer” ($p = 5.52 \times 10^{-21}$) and “Colorectal cancer” ($p = 4.22 \times 10^{-18}$); 7 to “Pancreatic cancer” ($p = 1.94 \times 10^{-18}$); 6 to “Chronic myeloid leukemia” ($p = 1.61 \times 10^{-30}$) and 5 each to “Non-small cell lung cancer” ($p = 8.66 \times 10^{-15}$) and “Glioma” ($p = 2.38 \times 10^{-14}$) (Supporting Information, Tables S7a, S8a). The significant association of HCV with host factors central to various cancer pathways (including tumor suppressors such as TP53) is consistent with previous observations that viral pathogens significantly targeted host proteins associated with cancer pathways,^{59,64,65} which likely plays major roles in tumorigenesis.

Immune System and Signal Transduction

HCV infection induces various active and passive host immune responses including the recognition of viral RNA by host cell receptors. These events lead to the production of Type I interferons (IFN- α/β) and inflammatory cytokines in the infected hepatocytes, initiating the antiviral response. HCV persistence in the host is determined by the virus’s ability to impair host immune responses.^{66–69}

The analysis of the NSSA interaction network revealed that 21 of the 132 NSSA interacting proteins, including 16 bottlenecks and their interacting partners, were mapped to one or more enriched KEGG pathways associated with the immune system (Supporting Information, Tables S7a, S8a).

Table 2. KEGG Pathway Functional Categories (Subclasses) Sorted by the Number of Enriched Pathways (≥ 3) Associated with One or More NSSA Interacting Bottlenecks

category	no. of enriched pathways	no. of bottlenecks	associated bottlenecks	KEGG pathways in the given category associated with most number of bottlenecks
infectious diseases	16	24	ACTB, AKT1, CDKL1, CSNK2A1, CTNNB1, FLNA, FYN, GRS2, GRB2, GSK3B, HSPB1, JAK1, LCK, LYN, PIK3R1, PPP2CA, RAF1, SRC, STAT1, SYK, TBP, TGFBR1, TP53, TRAF2	"Epstein-Barr virus infection"; "H1N1v infection"; "Hepatitis C"; "Hepatitis B"; "Measles"; "Influenza A"; "Herpes simplex infection"; "Tuberculosis"; "Toxoplasmosis"; "Chagas disease (American trypanosomiasis)"; "Bacterial invasion of epithelial cells"
cancers	16	19	AKT1, AXIN1, CDK1, CTNNB1, GRB2, GSK3B, HSP90AA1, JAK1, LYN, RAF1, SRC, STAT1, SYK, TBP, TGFBR1, THBS1, TP53, TRAF2	"Pathways in cancer"; "Viral carcinogenesis"; "Prostate cancer"; "Endometrial cancer"; "Colorectal cancer"; "Pancreatic cancer"; "Chronic myeloid leukemia"; "Non-small cell lung cancer"; "Glioma"; "Small cell lung cancer"; "Renal cell carcinoma"; "Melanoma"; "Acute myeloid leukemia"
immune system	10	16	ACTB, AKT1, CTNNB1, FYN, GRB2, GSK3B, HSP90AA1, LCK, LYN, PIK3R1, PIN1, RAF1, SRC, STAT1, SYK, TRAF	"Chemokine signaling pathway"; "T cell receptor signaling pathway"; "Fc epsilon RI signaling pathway"; "B cell receptor signaling pathway"; "Natural killer cell mediated cytotoxicity"; "Fc gamma R-mediated phagocytosis"
signal transduction	9	22	AKT1, AXIN1, CSNK2A1, CTNNB1, FLN, GRB2, GSK3B, HSP90AA1, HSPB1, JAK1, LCK, LYN, PIK3R1, PPP2CA, RAF1, SRC, STAT1, SYK, TGFBR1, THBS1, TP53, TRAF2	"PI3K-Akt signaling pathway"; "MAPK signaling pathway"; "Wnt signaling pathway"; "ErbB signaling pathway"; "VEGF signaling pathway"; "NF-kappa B signaling pathway"; "Jak-STAT signaling pathway"
nervous system	5	8	AKT1, GRB2, GSK3B, LYN, PIK3R1, PPP2CA, RAF1, TP53	"Neurotrophin signaling pathway"; "Long-term depression"; "Dopaminergic synapse"; "Long-term potentiation"
endocrine system	4	10	AKT1, CDK1, GRB2, GSK3B, HSP90AA1, PIK3R1, PLK1, RAF1, SRC, TRAF2	"Progesterone-mediated oocyte maturation"; "Insulin signaling pathway"; "GnRH signaling pathway"; "Adipocytokine signaling pathway"
cell growth and death	4	9	AKT1, CDK1, GSK3B, PIK3R1, PLK1, PPP2CA, THBS1, TP53, TRAF2	"Cell cycle"; "Apoptosis"; "p53 signaling pathway"; "Oocyte meiosis"
cell communication	3	14	ACTB, AKT1, CSNK2A1, CTNNB1, FLNA, FYN, GRB2, GSK3B, PIK3R1, PPP2CA, RAF1, SRC, TGFBR1, THBS1	"Focal adhesion"; "Tight junction"; "Adherens junction"
development	3	12	AKT1, FHL2, FYN, GRB2, GSK3B, JAK1, LCK, PIK3R1, STAT1, SYK, TGFBR1, THBS1	"Osteoclast differentiation"; "Axon guidance"; "Dorso-ventral axis formation"

Eight bottlenecks were mapped to the enriched KEGG pathway "Chemokine signaling pathway" ($p = 2.27 \times 10^{-10}$), which is consistent with the modulation of host interferon signaling by NSSA in HCV infection.⁷⁰ In addition, 7 bottlenecks each were mapped to "T cell receptor signaling pathway" ($p = 4.6 \times 10^{-24}$), "Fc epsilon RI signaling pathway" ($p = 2.86 \times 10^{-14}$) and "B cell receptor signaling pathway" ($p = 1.8 \times 10^{-14}$) and 6 bottlenecks were mapped to "Natural killer cell mediated cytotoxicity" ($p = 1.92 \times 10^{-12}$). Three bottlenecks (AKT1, PIK3R1 and STAT1) were also mapped to the enriched KEGG pathway "Toll-like receptor signaling pathway" ($p = 3.23 \times 10^{-7}$; Supporting Information, Tables S7a, S8a). Toll-like receptor 3 mediated chemokine and cytokine signaling plays an important role in the host immune response in HCV infection.⁷¹ Therefore, NSSA interaction with bottlenecks, which function in various aspects of the host immune response, may significantly contribute to the perturbation of the host immune system in HCV pathogenesis.

Additionally, 32 of 132 NSSA interacting proteins examined in the present study, including 24 bottlenecks, were mapped to various pathways associated with the signal transduction and the endocrine system (Supporting Information, Tables S7a, S8a), many of which are implicated in HCV infection and HCC progression and are targets for molecular therapy in HCC.^{22,72-74}

Eleven bottlenecks were mapped to the enriched KEGG pathway "PI3K-Akt signaling pathway" ($p = 2.2 \times 10^{-24}$; Supporting Information, Tables S7a, S8a), which is consistent with a previous study that NSSA stimulates the activation of PI3K-Akt pathway, which contributes to HCC in HCV infection.⁷⁵ Eight bottlenecks were mapped to the enriched KEGG pathway "MAPK signaling pathway" ($p = 2.4 \times 10^{-19}$; Supporting Information, Tables S7a, S8a). Elements of the MAPK signaling cascades are directly involved in the progression of HCV infection, particularly in association with HCV Core and E2 proteins,^{22,24,76,77} thereby suggesting that NSSA interactions with the key facilitators of MAPK signaling in the host interactome may play an important role in regulating the reversible phosphorylation of NSSA and may contribute to the progression of HCV pathogenesis.

Bottlenecks AKT1, GRB2, GSK3B, PIK3R1 and RAF1 and many of their interactors were mapped to the enriched KEGG pathway "Insulin signaling pathway" ($p = 2.42 \times 10^{-13}$; Supporting Information, Tables S7a, S8a); these proteins are highlighted in Figure 2. Insulin signaling plays an important role in regulating glucose and lipid metabolism, and the disruption of this process may contribute to insulin resistance (IR). IR is linked with steatosis, fibrosis progression and poor interferon- α response in HCV infection.⁷⁸⁻⁸⁰ Suppression of AKT1 and GSK3B activity in HCV infection disrupts glucose metabolism and contributes to IR.^{81,82} Furthermore, PIK3R1 and NSSA interactor PIK3CB (Figure 2) are subunits of phosphatidylinositol 3-kinase (PI3K), which controls insulin secretion;⁸³ PI3K also facilitates the activation of the proto-oncogene beta-catenin (CTNNB1) by NSSA, which contributes to the development of HCC in HCV pathogenesis.⁸⁴ Previously, HCV Core protein has been directly implicated in the induction of IR in HCV infection,⁸⁵ while there is little evidence suggesting definitive links between NSSA and IR. Our observations, however, suggest that NSSA directly interacts with key regulators of insulin metabolism and may, therefore, play a major role in modulating HCV-induced IR and eventually HCC.




Figure 2. NS5A interacting bottlenecks and their interacting partners associated with the enriched KEGG pathway hsa04910: "Insulin signaling pathway".

Cell Adhesion and Communication

The perturbation of adherens and tight junction associated proteins has been implicated in HCV entry, cell–cell transmission and hepatoma migration in HCV infection.^{86–88} In the NS5A infection network, eight bottlenecks (ACTB, AKT1, CSNK2A1, CTNNB1, FYN, PPP2CA, SRC and TGFBR1) were mapped to either or both of the enriched KEGG pathways "Adherens Junction" ($p = 1.03 \times 10^{-15}$) and "Tight junction" ($p = 1.19 \times 10^{-5}$), which are associated with cell adhesion junctions and cellular communication (Supporting Information, Tables S7a, S8a). CSNK2A1 is the catalytic (alpha) subunit of Casein Kinase II (CK2), which phosphorylates NS5A and regulates the production of infectious viral particles.⁶³ CTNNB1, a key component of cell-adhesion complexes, is positively regulated by CK2.⁸⁹ Furthermore, the activation of CTNNB1 by NS5A significantly contributes to HCC.⁸⁴ Taken together, our observations suggest that NS5A interactions with bottlenecks, which regulate cell–cell adhesion (CSNK2A1, CTNNB1) and cytoskeletal organization (ACTB), may significantly contribute to the progression of HCV life cycle and tumorigenesis in HCV pathogenesis.

Eleven bottlenecks were mapped to the enriched KEGG pathway "Focal Adhesion" ($p = 1.02 \times 10^{-17}$; Supporting Information, Tables S7a, S8a), thereby reiterating that focal adhesion is a major target of NS5A.²² Focal adhesion regulates cell migration and adhesion, and some of its components were directly implicated in the regulation of HCV replication and propagation in our earlier study.²⁴ Our observations thus suggest that NS5A interactions with key components of the focal adhesion machinery may play important roles in the HCV lifecycle. For instance, NS5A interacts with bottleneck THBS1 (Thrombospondin-1), a glycoprotein, which was mapped to the KEGG "Focal Adhesion" pathway. THBS1 plays a key role in NS5A-mediated activation of the cytokine TGF- β 1, which facilitates HCV replication and progressive liver fibrosis in HCV infection.⁹⁰ Our observations suggest that direct NS5A interactions with the bottlenecks THBS1 and TGFBR1 (TGF- β receptor 1; KEGG Pathway "Adherens Junction"), a key facilitator of TGF- β downstream signaling, may be crucial in facilitating HCV replication and tumorigenesis in HCV pathogenesis.




Figure 3. ER-localized host factors RTN1 and RTN3 were found to interact (blue edges) with NS5A in an Y2H screening of human liver cDNA library using NS5A as bait.




Figure 4. Effects of knockdown of RTN1 and RTN3 on HCV propagation and replication. Host factors RTN1 and RTN3 were suppressed by RNAi (A) in Huh7OK1 cells infected with HCV JFH1 strain (genotype 2a). The amounts of viral titer (B) and intracellular viral RNA (C) were estimated. Each value was represented as percentage of the cells transfected with the control siRNA. FFU: Focus-forming units; *: $p < 0.05$, **: $p < 0.01$.

Cellular Transport

Cellular factors associated with endocytic trafficking are key facilitators of the HCV life cycle, particularly HCV entry into the hepatic cells.^{91–93} Endocytosis of the extracellular growth factor receptor (EGFR) in association with the cell surface glycoprotein CD81 plays a crucial role in HCV internalization and entry and is, therefore, an attractive target of anti-HCV strategies.⁹⁴ In the NS5A infection network, NS5A interactors ARAP1 and HSPA1A together with two bottlenecks (SRC, TGFBR1) were mapped to the enriched KEGG pathway “Endocytosis” ($p = 2.97 \times 10^{-8}$; Supporting Information, Tables S7a, S8a). ARAP1, a Golgi associated protein, negatively regulates EGFR trafficking, and decreased ARAP1 expression contributes to enhanced EGFR endocytosis.⁹⁵ Therefore, NS5A

interaction with ARAP1 may facilitate EGFR internalization and thus viral entry in HCV infection.

NS5A Interacting Host Proteins RTN1 and RTN3 Function in HCV Propagation but Not Replication

Traditionally, viral and host proteins associated with the HCV lifecycle (internalization, replication, assembly and release) have been preferred targets in the anti-HCV studies. During infection, HCV localizes to the detergent-resistant membrane fraction (DRM) derived from the ER, where the viral replication and assembly take place.⁴ Thus, of the novel interactions identified in our Y2H assay, we focused on two ER-localized host factors RTN1 and RTN3 (Figure 3). RTN1 and RTN3 belong to a group of proteins named Reticulons, which are integral to maintaining the shape and organization of the

ER and have been implicated in facilitating the replication of various positive-strand RNA viruses.^{96–98} Furthermore, both RTN1 and RTN3 have been specifically detected in the very low density lipoprotein (VLDL) transport vesicle (VTV);⁹⁹ VTV is a key component of the VLDL secretory pathway, which plays an essential role in the production and the release of the infectious HCV particles.¹⁰⁰ Therefore, NSSA interactions with RTN1 and RTN3 suggested novel and potentially crucial roles of the two host proteins in the replication and/or release stages of the HCV lifecycle.

We performed cellular assays to assess the impact of RTN1 and RTN3 siRNA knockdowns on HCV replication and release. Since the HCV-production systems using the HCV JFH1 infectious strain (genotype 2a) isolates alone are capable of both efficient replication and the production of the infectious HCV particles, JFH1 was used to infect the Huh7OK1 cell line 24h after transfection with each siRNA (see Materials and Methods). The infected cells were harvested after 72 h postinfection, and the expression of each host protein was assessed by qRT-PCR (Figure 4A). The viral titer was significantly decreased by individual and double knockdowns of RTN1 and RTN3 (Figure 4B). However, RTN1 and RTN3 knockdowns had no effect on the intracellular viral RNA levels in the HCV infected cells (Figure 4C), suggesting that RTN1 and RTN3 regulate HCV propagation but not HCV replication.

CONCLUSIONS

We describe here our observations of PPIs between HCV NSSA and host proteins. By employing a multifold approach involving an experimental Y2H assay and literature mining, we derived a comprehensive set of experimentally determined binary interactions between NSSA and host proteins. We proceeded to map the combined NSSA–host interactions onto an overall interaction network, which comprised a repertoire of connections, which potentially enable NSSA to link up with and modulate the components of the host cellular networks. We then employed a network-based approach to understand the biological context of these connections in HCV pathogenesis with the help of the TargetMine data warehouse.

A functional analysis of the PPI networks highlighted NSSA interactions with several well connected host factors (hubs) and centrally located “bottlenecks” in the host cellular networks that function in cellular pathways associated with immune system and cell signaling, cellular adhesion and cell transport, cell growth and cell death and ER homeostasis among others. The “bottlenecks” include several proteins that were previously implicated in HCV pathogenesis, thereby suggesting that NSSA interactions with centrally connected host factors may enable the virus to influence strongly the host cellular processes in HCV infection. Notably, many bottlenecks were mapped to pathways associated with the infectious diseases induced by diverse bacterial and viral pathogens of the human host. These observations thus suggest the presence of some common themes underlying the onset of various human diseases associated with pathogenic infection in humans, a better understanding of which may be helpful in optimizing broad spectrum approaches to counteracting a wide range of pathogenic infections.

Cellular assays based on siRNA knockdowns in the HCV infected and replicon cells demonstrated RTN1 and RTN3, ER-localized NSSA interacting proteins, to be novel regulators of HCV propagation, but not replication, and thus promising novel candidates for anti-HCV therapy.

Our analysis therefore provides further insights into the role of NSSA–host interactions in HCV infection, a deeper understanding of which may aid in the identification of new clinically relevant targets for optimizing the therapeutic strategies to manipulate HCV–host interactions and thus more effectively combating HCV infection. Our analysis also emphasizes the importance of elaborate network-based computational approaches that integrate diverse biological data types in investigating host–pathogen interactions.

ASSOCIATED CONTENT

Supporting Information

Supporting methods, figures, and tables. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*E-mail: kenji@nibio.go.jp (K.M.); lokesh@nibio.go.jp (L.P.T.). Tel: +81-72-641-9890. Fax: +81-72-641-9881.

Author Contributions

†L. P. Tripathi and H. Kambara contributed equally to this work.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This study was supported by the Industrial Technology Research Grant Program in 2007 from New Energy and Industrial Technology Development Organization (NEDO) of Japan and also by grants-in-aid from the Ministry of Health, Labor, and Welfare; the Ministry of Education, Culture, Sports, Science, and Technology; the Osaka University Global Center of Excellence Program; and the Foundation for Biomedical Research and Innovation.

REFERENCES

- (1) Dubuisson, J. Hepatitis C virus proteins. *World J. Gastroenterol.* **2007**, *13* (17), 2406–15.
- (2) Moriishi, K.; Matsuura, Y. Host factors involved in the replication of hepatitis C virus. *Rev. Med. Virol.* **2007**, *17* (5), 343–54.
- (3) Myrmel, H.; Ulvestad, E.; Asjo, B. The hepatitis C virus enigma. *APMIS* **2009**, *117* (5–6), 427–39.
- (4) Tang, H.; Grise, H. Cellular and molecular biology of HCV infection and hepatitis. *Clin. Sci.* **2009**, *117* (2), 49–65.
- (5) Pol, S.; Vallet-Pichard, A.; Corouge, M.; Mallet, V. O. Hepatitis C: epidemiology, diagnosis, natural history and therapy. *Contrib. Nephrol.* **2012**, *176*, 1–9.
- (6) Kuiken, C.; Simmonds, P. Nomenclature and numbering of the hepatitis C virus. *Methods Mol. Biol.* **2009**, *510*, 33–53.
- (7) Moradpour, D.; Penin, F.; Rice, C. M. Replication of hepatitis C virus. *Nat. Rev. Microbiol.* **2007**, *5* (6), 453–63.
- (8) Love, R. A.; Brodsky, O.; Hickey, M. J.; Wells, P. A.; Cronin, C. N. Crystal structure of a novel dimeric form of NSSA domain I protein from hepatitis C virus. *J. Virol.* **2009**, *83* (9), 4395–403.
- (9) Yamasaki, L. H.; Arcuri, H. A.; Jardim, A. C.; Bittar, C.; de Carvalho-Mello, I. M.; Rahal, P. New insights regarding HCV-NSSA structure/function and indication of genotypic differences. *Virol. J.* **2012**, *9*, 14.
- (10) Appel, N.; Zayas, M.; Miller, S.; Krijnse-Locker, J.; Schaller, T.; Friebe, P.; Kallis, S.; Engel, U.; Bartenschlager, R. Essential role of domain III of nonstructural protein 5A for hepatitis C virus infectious particle assembly. *PLoS Pathog.* **2008**, *4* (3), e1000035.

- (11) Gale, M. J., Jr.; Korth, M. J.; Tang, N. M.; Tan, S. L.; Hopkins, D. A.; Dever, T. E.; Polyak, S. J.; Gretch, D. R.; Katze, M. G. Evidence that hepatitis C virus resistance to interferon is mediated through repression of the PKR protein kinase by the nonstructural 5A protein. *Virology* **1997**, *230* (2), 217–27.
- (12) Ghosh, S.; Ahrens, W. A.; Phatak, S. U.; Hwang, S.; Schrum, L. W.; Bonkovsky, H. L. Association of filamin A and vimentin with hepatitis C virus proteins in infected human hepatocytes. *J. Viral Hepatitis* **2011**, *18* (10), e568–77.
- (13) Gao, M.; Nettles, R. E.; Belema, M.; Snyder, L. B.; Nguyen, V. N.; Fridell, R. A.; Serrano-Wu, M. H.; Langley, D. R.; Sun, J. H.; O'Boyle, D. R., 2nd; Lemm, J. A.; Wang, C.; Knipe, J. O.; Chien, C.; Colonna, R. J.; Grasela, D. M.; Meanwell, N. A.; Hamann, L. G. Chemical genetics strategy identifies an HCV NSSA inhibitor with a potent clinical effect. *Nature* **2010**, *465* (7294), 96–100.
- (14) Lee, C. Discovery of hepatitis C virus NSSA inhibitors as a new class of anti-HCV therapy. *Arch. Pharmacol. Res.* **2011**, *34* (9), 1403–7.
- (15) Lemm, J. A.; O'Boyle, D., 2nd; Liu, M.; Nower, P. T.; Colonna, R.; Deshpande, M. S.; Snyder, L. B.; Martin, S. W.; St Laurent, D. R.; Serrano-Wu, M. H.; Romine, J. L.; Meanwell, N. A.; Gao, M. Identification of hepatitis C virus NSSA inhibitors. *J. Virol.* **2010**, *84* (1), 482–91.
- (16) Lemon, S. M.; McKeating, J. A.; Pietschmann, T.; Frick, D. N.; Glenn, J. S.; Tellinghuisen, T. L.; Symons, J.; Furman, P. A. Development of novel therapies for hepatitis C. *Antiviral Res.* **2010**, *86* (1), 79–92.
- (17) Fusco, D. N.; Chung, R. T. Novel therapies for hepatitis C: insights from the structure of the virus. *Annu. Rev. Med.* **2012**, *63*, 373–87.
- (18) Buhler, S.; Bartenschlager, R. New targets for antiviral therapy of chronic hepatitis C. *Liver Int.* **2012**, *32* (Suppl 1), 9–16.
- (19) Sarrazin, C.; Hezode, C.; Zeuzem, S.; Pawlotsky, J. M. Antiviral strategies in hepatitis C virus infection. *J. Hepatol.* **2012**, *56* (Suppl), S88–S100.
- (20) Wang, S.; Wu, X.; Pan, T.; Song, W.; Wang, Y.; Zhang, F.; Yuan, Z. Viperin inhibits hepatitis C virus replication by interfering with binding of NSSA to host protein hVAP-33. *J. Gen. Virol.* **2012**, *93* (Pt1), 83–92.
- (21) Durmus Tekir, S.; Cakir, T.; Ulgen, K. O. Infection strategies of bacterial and viral pathogens through pathogen-human protein-protein interactions. *Front. Microbiol.* **2012**, *3*, 46.
- (22) de Chasse, B.; Navratil, V.; Tafforeau, L.; Hiet, M. S.; Aublin-Gex, A.; Agaugue, S.; Meiffren, G.; Pradezynski, F.; Faria, B. F.; Chantier, T.; Le Breton, M.; Pellet, J.; Davoust, N.; Mangeot, P. E.; Chaboud, A.; Penin, F.; Jacob, Y.; Vidalain, P. O.; Vidal, M.; Andre, P.; Rabourdin-Combe, C.; Lotteau, V. Hepatitis C virus infection protein network. *Mol. Syst. Biol.* **2008**, *4*, 230.
- (23) Tan, S. L.; Ganji, G.; Paepfer, B.; Proll, S.; Katze, M. G. Systems biology and the host response to viral infection. *Nat. Biotechnol.* **2007**, *25* (12), 1383–9.
- (24) Tripathi, L. P.; Kataoka, C.; Taguwa, S.; Moriishi, K.; Mori, Y.; Matsuura, Y.; Mizuguchi, K. Network based analysis of hepatitis C virus Core and NS4B protein interactions. *Mol. BioSyst.* **2010**, *6* (12), 2539–53.
- (25) Friedel, C. C.; Haas, J. Virus-host interactomes and global models of virus-infected cells. *Trends Microbiol.* **2011**, *19* (10), 501–8.
- (26) Tafforeau, L.; Rabourdin-Combe, C.; Lotteau, V. Virus-human cell interactomes. *Methods Mol. Biol.* **2012**, *812*, 103–20.
- (27) Aizaki, H.; Aoki, Y.; Harada, T.; Ishii, K.; Suzuki, T.; Nagamori, S.; Toda, G.; Matsuura, Y.; Miyamura, T. Full-length complementary DNA of hepatitis C virus genome from an infectious blood sample. *Hepatology* **1998**, *27* (2), 621–7.
- (28) Hamamoto, I.; Nishimura, Y.; Okamoto, T.; Aizaki, H.; Liu, M.; Mori, Y.; Abe, T.; Suzuki, T.; Lai, M. M.; Miyamura, T.; Moriishi, K.; Matsuura, Y. Human VAP-B is involved in hepatitis C virus replication through interaction with NSSA and NSSB. *J. Virol.* **2005**, *79* (21), 13473–82.
- (29) Rebholz-Schuhmann, D.; Kirsch, H.; Arregui, M.; Gaudan, S.; Riethoven, M.; Stoehr, P. EBI Med—text crunching to gather facts for proteins from Medline. *Bioinformatics* **2007**, *23* (2), e237–44.
- (30) Rebholz-Schuhmann, D.; Arregui, M.; Gaudan, S.; Kirsch, H.; Jimeno, A. Text processing through Web services: calling Whatizit. *Bioinformatics* **2008**, *24* (2), 296–8.
- (31) Stark, C.; Breitkreutz, B. J.; Reguly, T.; Boucher, L.; Breitkreutz, A.; Tyers, M. BioGRID: a general repository for interaction datasets. *Nucleic Acids Res.* **2006**, *34* (Database issue), D535–9.
- (32) Turner, B.; Razick, S.; Turinsky, A. L.; Vlasblom, J.; Crowdy, E. K.; Cho, E.; Morrison, K.; Donaldson, I. M.; Wodak, S. J. iRefWeb: interactive analysis of consolidated protein interaction data and their supporting evidence. *Databse* **2010**, *2010*, baq023.
- (33) Chen, Y. A.; Tripathi, L. P.; Mizuguchi, K. TargetMine, an integrated data warehouse for candidate gene prioritisation and target discovery. *PLoS One* **2011**, *6* (3), e17844.
- (34) Cline, M. S.; Smoot, M.; Cerami, E.; Kuchinsky, A.; Landys, N.; Workman, C.; Christmas, R.; Avila-Campilo, I.; Creech, M.; Gross, B.; Hanspers, K.; Isserlin, R.; Kelley, R.; Killcoyne, S.; Lotia, S.; Maere, S.; Morris, J.; Ono, K.; Pavlovic, V.; Pico, A. R.; Vailaya, A.; Wang, P. L.; Adler, A.; Conklin, B. R.; Hood, L.; Kuiper, M.; Sander, C.; Schmulevich, I.; Schwikowski, B.; Warner, G. J.; Ideker, T.; Bader, G. D. Integration of biological networks and gene expression data using Cytoscape. *Nat. Protoc.* **2007**, *2* (10), 2366–82.
- (35) Smoot, M. E.; Ono, K.; Ruschinski, J.; Wang, P. L.; Ideker, T. Cytoscape 2.8: new features for data integration and network visualization. *Bioinformatics* **2011**, *27* (3), 431–2.
- (36) Assenov, Y.; Ramirez, F.; Schelhorn, S. E.; Lengauer, T.; Albrecht, M. Computing topological parameters of biological networks. *Bioinformatics* **2008**, *24* (2), 282–4.
- (37) Lees, J.; Yeats, C.; Perkins, J.; Sillito, L.; Rentsch, R.; Dessailly, B. H.; Orengo, C. Gene3D: a domain-based resource for comparative genomics, functional annotation and protein network analysis. *Nucleic Acids Res.* **2012**, *40* (Database issue), D465–71.
- (38) Ashburner, M.; Ball, C. A.; Blake, J. A.; Botstein, D.; Butler, H.; Cherry, J. M.; Davis, A. P.; Dolinski, K.; Dwight, S. S.; Eppig, J. T.; Harris, M. A.; Hill, D. P.; Issel-Tarver, L.; Kasarskis, A.; Lewis, S.; Matese, J. C.; Richardson, J. E.; Ringwald, M.; Rubin, G. M.; Sherlock, G. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat. Genet.* **2000**, *25* (1), 25–9.
- (39) Aoki-Kinoshita, K. F.; Kanehisa, M. Gene annotation and pathway mapping in KEGG. *Methods Mol. Biol.* **2007**, *396*, 71–91.
- (40) Benjamini, Y.; Hochberg, Y. Controlling the false discovery rate—A practical and powerful approach to multiple testing. *J. R. Stat. Soc. B* **1995**, *57* (1), 289–300.
- (41) Noble, W. S. How does multiple testing correction work? *Nat. Biotechnol.* **2009**, *27* (12), 1135–7.
- (42) Okamoto, K.; Mori, Y.; Komoda, Y.; Okamoto, T.; Okochi, M.; Takeda, M.; Suzuki, T.; Moriishi, K.; Matsuura, Y. Intramembrane processing by signal peptide peptidase regulates the membrane localization of hepatitis C virus core protein and viral propagation. *J. Virol.* **2008**, *82* (17), 8349–61.
- (43) Okamoto, T.; Omori, H.; Kaname, Y.; Abe, T.; Nishimura, Y.; Suzuki, T.; Miyamura, T.; Yoshimori, T.; Moriishi, K.; Matsuura, Y. A single-amino-acid mutation in hepatitis C virus NSSA disrupting FKBP8 interaction impairs viral replication. *J. Virol.* **2008**, *82* (7), 3480–9.
- (44) Wakita, T.; Pietschmann, T.; Kato, T.; Date, T.; Miyamoto, M.; Zhao, Z.; Murthy, K.; Habermann, A.; Krausslich, H. G.; Mizokami, M.; Bartenschlager, R.; Liang, T. J. Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat. Med.* **2005**, *11* (7), 791–6.
- (45) Taguwa, S.; Kambara, H.; Omori, H.; Tani, H.; Abe, T.; Mori, Y.; Suzuki, T.; Yoshimori, T.; Moriishi, K.; Matsuura, Y. Co-chaperone activity of human butyrate-induced transcript 1 facilitates hepatitis C virus replication through an Hsp90-dependent pathway. *J. Virol.* **2009**, *83* (20), 10427–36.
- (46) Kukihara, H.; Moriishi, K.; Taguwa, S.; Tani, H.; Abe, T.; Mori, Y.; Suzuki, T.; Fukuhara, T.; Taketomi, A.; Maehara, Y.; Matsuura, Y.