

研究成果の刊行に関する一覧表

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Enooku K, Nakagawa H, Soroida Y, Ohkawa R, Kageyama Y, Uranbileg B, Watanabe N, Tateishi R, Yoshida H, <u>Koike K</u> , Yatomi Y, Ikeda H.	Increased serum mitochondrial creatine kinase activity as a risk for hepatocarcinogenesis in chronic hepatitis C patients.	<b>Int J Cancer</b>			[Epub ahead of print]
Nakagawa H, Hikiba Y, Hirata Y, Font-Burgada J, Sakamoto K, Hayakawa Y, Taniguchi K, Umemura A, Kinoshita H, Sakitani K, Nishikawa Y, Hirano K, Ikenoue T, Ijichi H, Dhar D, Shibata W, Akanuma M, <u>Koike K</u> , Karin M, Maeda S.	Loss of liver E-cadherin induces sclerosing cholangitis and promotes carcinogenesis.	<b>Proc Natl Acad Sci U S A</b>			[Epub ahead of print]
Otsuka M, Kishikawa T, Yoshikawa T, Ohno M, Takata A, Shibata C, <u>Koike K</u> .	The role of microRNAs in hepatocarcinogenesis: current knowledge and future prospects.	<b>J Gastroenterol</b>			[Epub ahead of print]
Uchino R, Isayama H, Tsujino T, Sasahira N, Ito Y, Matsubara S, Takahara N, Arizumi T, Toda N, Mohri D, Togawa O, Yagioka H, Yanagihara Y, Nakajima K, Akiyama D, Hamada T, Miyabayashi K, Mizuno S, Kawakubo K, Kogure H, Sasaki T, Yamamoto N, Nakai Y, Hirano K, Tada M, <u>Koike K</u> .	Results of the Tokyo Trial of Prevention of Post-ERCP Pancreatitis with Risperidone-2: a multicenter, randomized, placebo-controlled, double-blind clinical trial.	<b>Gastrointest Endosc</b>			[Epub ahead of print]

Ikeda K, Izumi N, Tanaka E, Yotsuyanagi H, Takahashi Y, Fukushima J, Kondo F, Fukusato T, <u>Koike K</u> , Hayashi N, Tsubouchi H, Kumada H.	Discrimination of fibrotic staging of chronic hepatitis C using multiple fibrosis markers.	<b>Hepatol Res</b>			[Epub ahead of print]
Shibata C, Kishikawa T, Otsuka M, Ohno M, Yoshikawa T, Takata A, Yoshida H, <u>Koike K</u> .	Inhibition of microRNA122 decreases SREBP1 expression by modulating suppressor of cytokine signaling 3 expression.	<b>Biochem Biophys Res Commun</b>			[Epub ahead of print]
Minami T, Tateishi R, Shiina S, Fujiwara N, Mikami S, Sato M, Uchino K, Enooku K, Asaoka Y, Kondo Y, Yoshida H, <u>Koike K</u> .	Spontaneous clearance of serum hepatitis C virus RNA during the clinical course of hepatocellular carcinoma in patients with chronic hepatitis C.	<b>Hepatol Res</b>			[Epub ahead of print]
Sato M, Tateishi R, Yasunaga H, Horiguchi H, Yoshida H, Matsuda S, Fushimi K, <u>Koike K</u> .	Acute liver disease in Japan: a nationwide analysis of the Japanese Diagnosis Procedure Combination database.	<b>J Gastroenterol</b>			[Epub ahead of print]
Sato M, Kato N, Tateishi R, Muroyama R, Kowatari N, Li W, Goto K, Otsuka M, Shiina S, Yoshida H, Omata M, <u>Koike K</u> .	IL28B minor allele is associated with a younger age of onset of hepatocellular carcinoma in patients with chronic hepatitis C virus infection.	<b>J Gastroenterol</b>			[Epub ahead of print]
Mikoshiba N, Miyashita M, Sakai T, Tateishi R, <u>Koike K</u> .	Depressive symptoms after treatment in hepatocellular carcinoma survivors: prevalence, determinants, and impact on health-related quality of life.	<b>Psychooncology</b>			[Epub ahead of print]

Ohki T, Tateishi R, Akahane M, Mikami S, Sato M, Uchino K, Arano T, Enooku K, Kondo Y, Yamashiki N, Goto T, Shiina S, Yoshida H, Matsuyama Y, Omata M, Ohtomo K, <u>Koike K.</u>	CT with hepatic arterioportography as a pretreatment examination for hepatocellular carcinoma patients: a randomized controlled trial.	<b>Am J Gastroenterol</b>	108	1305-1313	2013
Inoue Y, Tomiya T, Nishikawa T, Ohtomo N, Tanoue Y, Ikeda H, <u>Koike K.</u>	Induction of p53-Dependent p21 Limits Proliferative Activity of Rat Hepatocytes in the Presence of Hepatocyte Growth Factor.	<b>PLoS One</b>	8(11)	e78346	2013
Hikita H, Enooku K, Satoh Y, Yoshida H, Nakagawa H, Masuzaki R, Tateishi R, Soroida Y, Sato M, Suzuki A, Gotoh H, Iwai T, Yokota H, <u>Koike K.</u> , Yatomi Y, Ikeda H.	Perihepatic lymph node enlargement is a negative predictor for sustained responses to pegylated interferon- $\alpha$ and ribavirin therapy for Japanese patients infected with hepatitis C virus genotype 1.	<b>Hepatol Res</b>	43(10)	1005-1012	2013
Kishikawa T, Otsuka M, Yoshikawa T, Ohno M, Takata A, Shibata C, Kondo Y, Akanuma M, Yoshida H, <u>Koike K.</u>	Regulation of the expression of the liver cancer susceptibility gene MICA by microRNAs.	<b>Sci Rep</b>	3	2739	2013
Liu Y, Higashitsuji H, Higashitsuji H, Itoh K, Sakurai T, <u>Koike K.</u> , Hirota K, Fukumoto M, Fujita J.	Overexpression of gankyrin in mouse hepatocytes induces hemangioma by suppressing factor inhibiting hypoxia-inducible factor-1 (FIH-1) and activating	<b>Biochem Biophys Res Commun</b>	432(1)	22-27	2013

	hypoxia-inducible factor-1.				
<u>Koike K.</u>	The oncogenic role of hepatitis C virus.	<b>Recent Results Cancer Res</b>	193	97-111	2014
Uchino K, Tateishi R, Nakagawa H, Shindoh J, Sugawara Y, Akahane M, Shibahara J, Yoshida H, <u>Koike K.</u>	Uninodular combined hepatocellular and cholangiocarcinoma with multiple non-neoplastic hypervascular lesions appearing in the liver of a patient with HIV and HCV coinfection.	<b>J Clin Virol</b>	57(2)	173-177	2013
Ohno M, Shibata C, Kishikawa T, Yoshikawa T, Takata A, Kojima K, Akanuma M, Kang YJ, Yoshida H, Otsuka M, <u>Koike K.</u>	The flavonoid apigenin improves glucose tolerance through inhibition of microRNA maturation in miRNA103 transgenic mice.	<b>Sci Rep</b>	3	2553	2013
Ikeda K, Izumi N, Tanaka E, Yotsuyanagi H, Takahashi Y, Fukushima J, Kondo F, Fukusato T, <u>Koike K.</u> , Hayashi N, Kumada H.	Fibrosis score consisting of four serum markers successfully predicts pathological fibrotic stages of chronic hepatitis B.	<b>Hepatol Res</b>	43(6)	596-604	2013
Urabe Y, Ochi H, Kato N, Kumar V, Takahashi A, Muroyama R, Hosono N, Otsuka M, Tateishi R, Lo PH, Tanikawa C, Omata M, <u>Koike K.</u> , Miki D, Abe H, Kamatani N, Toyota J, Kumada H, Kubo M, Chayama K, Nakamura Y, Matsuda K.	A genome-wide association study of HCV induced liver cirrhosis in the Japanese population identifies novel susceptibility loci at MHC region.	<b>J Hepatol</b>	58(5)	875-882	2013

Hikita H, Nakagawa H, Tateishi R, Masuzaki R, Enooku K, Yoshida H, Omata M, Soroida Y, Sato M, Gotoh H, Suzuki A, Iwai T, Yokota H, <u>Koike K</u> , Yatomi Y, Ikeda H.	Perihepatic lymph node enlargement is a negative predictor of liver cancer development in chronic hepatitis C patients.	<b>J Gastroenterol</b>	48(3)	366-373	2013
Lo PH, Urabe Y, Kumar V, Tanikawa C, <u>Koike K</u> , Kato N, Miki D, Chayama K, Kubo M, Nakamura Y, Matsuda K.	Identification of a functional variant in the mica promoter which regulates mica expression and increases HCV-related hepatocellular carcinoma risk.	<b>PLoS One</b>	8(4)	e61279	2013
Minami T, Kishikawa T, Sato M, Tateishi R, Yoshida H, <u>Koike K</u> .	Meta-analysis: mortality and serious adverse events of peginterferon plus ribavirin therapy for chronic hepatitis C.	<b>J Gastroenterol</b>	48(2)	254-268	2013
Ikeda H, Enooku K, Ohkawa R, <u>Koike K</u> , Yatomi Y.	Plasma lysophosphatidic acid levels and hepatocellular carcinoma.	<b>Hepatology</b>	57	417-418	2013
Takata A, Otsuka M, Yoshikawa T, Kishikawa T, Hikiba Y, Obi S, Goto T, Kang YJ, Maeda S, Yoshida H, Omata M, Asahara H, <u>Koike K</u> .	MiRNA-140 acts as a liver tumor suppressor by controlling NF-κB activity via directly targeting Dnmt1 expression.	<b>Hepatology</b>	57	162-170	2013
Nishimura T, Yamaguchi K, Fujii H, Okada Y, Yokomizo C, Niimi T, Sumida Y, Yasui K, Mitsuyoshi H, Minami M, Umemura A, Shima T, Okanoue T, Itoh Y	Prediction of favorable clinical course in hepatitis C virus carriers with persistently normal serum alanine aminotransferase levels: A long-term follow-up study	Hepatol Res	43	557-562	2013

Okada Y, Yamaguchi K, Nakajima T, Nishikawa T, Jo M, Mitsumoto Y, Kimura H, Nishimura T, Tochiki N, Yasui K, Mitsuyoshi H, Minami M, Kagawa K, <u>Okanoue T</u> , Itoh Y	Rosuvastatin ameliorates high-fat and high-cholesterol diet-induced nonalcoholic steatohepatitis in rats	Liver Int	33	301-311	2013
Yoneda M, Imajo K, Eguchi Y, Fujii H, Sumida Y, Hyogo H, Ono M, Suzuki Y, Kawaguchi T, Aoki N, Sata M, Kanemasa K, Kohgo Y, Saibara T, Chayama K, Itoh Y, Yoshikawa T, Anza I K, Fujimoto K, <u>Okanoue T</u> , Nakajima A	Noninvasive scoring systems in patients with nonalcoholic fatty liver disease with normal alanine aminotransferase levels	J Gastroenterol	48	1051-1060	2013
Sumida Y, Naito Y, Tanaka S, Sakai K, Inada Y, Taketani H, Kanemasa K, Yasui K, Itoh Y, <u>Okanoue T</u> , Yoshikawa T	Long-term (>2 Yr) efficacy of vitamin E for non-alcoholic steatohepatitis	Hepato-Gastroenterology	60	1445-1450	2013
Nakahara T, Hyogo H, Yoneda M, Sumida Y, Eguchi Y, Fujii H, Ono M, Kawaguchi T, Imajo K, Aikata H, Tanaka S, Kanemasa K, Fujimoto K, Anzai K, Saibara T, Sata M, Nakajima A, Itoh Y, Chayama K, <u>Okanoue T</u>	Type 2 diabetes mellitus is associated with the fibrosis severity in patients with nonalcoholic fatty liver disease in a large retrospective cohort of Japanese patients	J Gastricenterol	online 26 November		2013
Kondo Y, Hasegawa G, Okada H, Senmaru T, Fukui M, Nakamura N, Sawada M, Kitawaki J, <u>Okanoue T</u> , Kishimoto Y, Amano A, Maruyama N,	<i>Leprdb/db</i> mice with senescence marker protein-30 knockout ( <i>Leprdb/db Smp30Y/-</i> ) exhibit increase in small dense –LDL severe fatty	PLoS One	8(6)	e65698	2013

Obayashi H, Ishigami A	liver despite being fed a standard diet				
Shima T, Uto H, Ueki K, Takamura T, Kohgo Y, Kawata S, Yasui K, Park H, Nakamura N, Nakatou T, Tanaka N, Umemura A, Mizuno M, Tanaka J, <u>Okanoue T</u>	Clinicopathological features of liver injury in patients with type 2 diabetes mellitus and comparative study of histologically proven nonalcoholic fatty liver disease with or without type 2 diabetes mellitus	J Gastroenterol	48	515-525	2013
Ue M, Ikebe N, Munekage K, Ochi T, Ono M, <u>Saibara T.</u>	Hepatocyte destruction with enhanced collagen synthesis in the liver of chronic hepatitis C patients on hemodialysis.	J Viral Hepat	20	350-357	2013
Osawa Y, Hoshi M, Yasuda I, <u>Saibara T.</u> , Moriwaki H, Kozawa O.	Tumor necrosis factor- $\alpha$ promotes cholestasis-induced liver fibrosis in the mouse through tissue inhibitor of metalloproteinase-1 production in hepatic stellate cells.	PLoS ONE	8	e65251	2013
Yoneda M, Imajo K, Eguchi Y, Fujii H, Sumida Y, Hyogo H, Ono M, Suzuki Y, Kawaguchi T, Aoki N, Sata M, Kanemasa K, Kohgo Y, <u>Saibara T</u> , Chayama K, Itoh Y, Yoshikawa T, Anzai K, Fujimoto K, Okanoue T, Nakajima A; Japan Study Group of Nonalcoholic Fatty Liver Disease.	Noninvasive scoring systems in patients with nonalcoholic fatty liver disease with normal alanine aminotransferase levels.	J Gastroenterol	48	1051-1060.	2013

Saibara T, Enomoto N, Kaneko S, Chayama K, Sata M, Imawari M, Onishi S, Okita K.	Clinical efficacy of combination therapy with ME3738 and pegylated interferon alpha-2a in patients with hepatitis C virus genotype 1.	Hepatol Res		doi: 10.1111/hepr .12139.	2013
Ono M, Ogasawara M, Hirose A, Mogami S, Ootake N, Aritake K, Higuchi T, Okamoto N, Sakamoto S, Yamamoto M, Urade Y, <u>Saibara T</u> , Oben JA.	Bofutsushosan, a Japanese herbal (Kampo) medicine, attenuates progression of nonalcoholic steatohepatitis in mice.	J Gastroenterol		DOI 10.1007/s00 535-013-085 2-8	
Taketani H, Sumida Y, Tanaka S, Imajo K, Yoneda M, Hyogo H, Ono M, Fujii H, Eguchi Y, Kanemasa K, Chayama K, Itoh Y, Yoshikawa T, <u>Saibara T</u> , Fujimoto K, Nakajima A; Japan Study Group of NAFLD	The association of insomnia with gastroesophageal reflux symptoms in biopsy-proven nonalcoholic fatty liver disease.	J Gastroenterol.		Aug 22	2013
Hashiba M, Ono M, Hyogo H, Ikeda Y, Masuda K, Yoshioka R, Ishikawa Y, Nagata Y, Munekage K, Ochi T, Hirose A, Nozaki-Fujimura Y, Noguchi S, Okamoto N, Chayama K, Suganuma N, <u>Saibara T</u> .	Glycemic variability is an independent predictive factor for development of hepatic fibrosis in nonalcoholic Fatty liver disease.	PLoS One.	8	e76161	2013
Toda K, Hayashi Y, Yamashita A, Okabe M, Ono M, <u>Saibara T</u> .	Aromatase-null mice expressing enhanced green fluorescent protein in germ cells provide a model system to assess estrogen-dependent	Transgenic Res		Nov 22.	2013

	ovulatory responses.				
Nakahara T, Hyogo H, Yoneda M, Sumida Y, Eguchi Y, Fujii H, Ono M, Kawaguchi T, Imajo K, Aikata H, Tanaka S, Kanemasa K, Fujimoto K, Anzai K, <u>Saibara T</u> , Sata M, Nakajima A, Itoh Y, Chayama K, Okanoue T; Japan Study Group of Nonalcoholic Fatty Liver Disease.	Type 2 diabetes mellitus is associated with the fibrosis severity in patients with nonalcoholic fatty liver disease in a large retrospective cohort of Japanese patients.	J Gastroenterol.		Nov 26.	2013
Tanaka K, Hyogo H, Ono M, Takahashi H, Kitajima Y, Ono N, Eguchi T, Fujimoto K, Chayama K, <u>Saibara T</u> , Anzai K, Eguchi Y, Japan Study Group of Nonalcoholic Fatty Liver Disease.	The upper limit of normal serum alanine aminotransferase levels in Japanese subjects.	Hepatology Research		DOI: 10.1111/hepr .12293	2013
<u>Hashimoto E</u> , Taniai M, Tokushige K.	Characteristics and diagnosis of NAFLD/NASH.	J Gastroenterol Hepatol	28 Suppl 4	64-70	2013
TokushigeK, <u>HashimotoE</u> , Kodama K.	Hepatocarcinogenesis in non-alcoholic fatty liver disease in Japan.	J Gastroenterol Hepatol	28 Suppl 4	88-92	2013
Sato K, <u>Tanaka S</u> , Mitsunori Y, Mogushi K, Yasen M, Aihara A, Ban D, Ochiai T, Irie T, Kudo A, Nakamura N, Tanaka H, Arii S.	Contrast-enhanced intraoperative ultrasonography for vascular imaging of hepatocellular carcinoma; clinical and biological significance.	Hepatology	57(4)	1436-1447	2013

Muramatsu S, <u>Tanaka S</u> , Mogushi K, Adikrisna R, Aihara A, Ban D, Ochiai T, Irie T, Kudo A, Nakamura N, Tanaka H, Nakayama K, Tanaka H, Yamaoka S, Arii S.	Visualization of stem cell features in human hepatocellular carcinoma enlightened in vivo significance of tumor-host interaction and clinical implication.	Hepatology	58(1)	218-228	2013
Furuta M, Kozaki KI, Tanimoto K, <u>Tanaka S</u> , Arii S, Shimamura T, Niida A, Miyano S, Inazawa J.	The Tumor-suppressive mir-497-195 cluster targets multiple cell-cycle regulators in hepatocellular carcinoma.	PLoS One	8(3)	e60155	2013
Katsuta E, <u>Tanaka S</u> , Mogushi K, Matsumura S, Ban D, Ochiai T, Irie T, Kudo A, Nakamura N, Tanaka H, Tanabe M, Arii S.	Age-related clinicopathological and molecular features of patients received curative hepatectomy for hepatocellular carcinoma.	American Journal of Surgery			in press
Kudo A, <u>Tanaka S</u> , Ban D, Matsumura S, Irie T, Ochiai T, Nakamura N, Arii S, Tanabe M.	Alcohol consumption and recurrence of non-B or non-C hepatocellular carcinoma after hepatectomy: a propensity score analysis.	Journal of Gastroenterolo gy			in press
Ogawa K, <u>Tanaka S</u> , Matsumura S, Murakata A, Ban D, Ochiai T, Irie T, Kudo A, Nakamura N, Tanabe M, Arii S.	EpCAM-Targeted therapy for human hepatocellular carcinoma.	Annals of Surgical Oncology			in press
Kudo A, Mogushi K, Takayama T, Matsumura S, Ban D, Irie T, Ochiai T, Tanaka H, Anzai N, Sakamoto M, <u>Tanaka S</u> , Arii S.	Mitochondrial metabolism in the Noncancerous Liver Determine the Occurrence of Hepatocellular Carcinoma: A Prospective Study.	Journal of Gastroenterolo gy			in press

Okada-Iwabu M, Yamauchi T, Iwabu M, Honma T, Hamagami K, Matsuda K, Yamaguchi M, Tanabe H, Kimura-Someya T, Shirouzu M, Ogata H, Tokuyama K, <u>Ueki K</u> , Nagano T, Tanaka A, Yokoyama S, Kadowaki T.	A small-molecule AdipoR agonist for type 2 diabetes and short life in obesity.	Nature	503	493-499	2013
Moller JB, Pedersen M, Tanaka H, Ohsugi M, Overgaard RV, Lyngé J, Almind K, Vasconcelos NM, Poulsen P, Keller C, <u>Ueki K</u> , Ingwersen SH, Pedersen BK, Kadowaki T	Body Composition is Main Determinant for the Difference in Type 2 Diabetes Pathophysiology between Japanese and Caucasians.	Diabetes Care		Oct 15 (Epub ahead of print)	2013
Nakaya K, Kubota N, Takamoto I, Kubota T, Katsuyama H, Sato H, Tokuyama K, Hashimoto S, Goto M, Jomori T, <u>Ueki K</u> , Kadowaki T.	Dipeptidyl peptidase-4 inhibitor anagliptin ameliorates diabetes in mice with haploinsufficiency of glucokinase on a high-fat diet.	Metabolism	62	933-951	2013
<u>Akuta N</u> , Suzuki F, Kawamura Y, Sezaki H, Hosaka T, Kobayashi M, Hara T, Kobayashi M, Saitoh S, Arase Y, Ikeda K, Kumada H	Utility of telaprevir-resistant variant detection for prediction of treatment efficacy in HCV genotype 1.	J Clin Microbiol.	52	193-200	2014
Arase Y, Kobayashi M, Kawamura Y, Suzuki F, Suzuki Y, <u>Akuta N</u> , Kobayashi M, Sezaki H, Saitoh S, Hosaka T, Ikeda K, Kumada H, Kobayashi T	Impact of virus clearance for the development of hemorrhagic stroke in chronic hepatitis C.	J Med Virol.	86	169-175	2014
<u>Akuta N</u> , Suzuki F, Kobayashi M, Hara T, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Saitoh S, Ikeda K,	Correlation between hepatitis B virus surface antigen level and alpha-fetoprotein in	J Med Virol.	86	131-138	2014

Kumada H	patients free of hepatocellular carcinoma or severe hepatitis			
Sezaki H, Suzuki F, Hosaka T, <u>Akuta N</u> , Fukushima T, Hara T, Kawamura Y, Kobayashi M, Suzuki Y, Saitoh S, Arase Y, Ikeda K, Kumada H	Effectiveness and safety of reduced-dose telaprevir-based triple therapy in chronic hepatitis C patients	Hepatol Res.		Epub ahead of print
Tanaka M, Suzuki F, Seko Y, Hara T, Kawamura Y, Sezaki H, Hosaka T, Akuta N, Kobayashi M, Suzuki Y, Saitoh S, Arase Y, Ikeda K, Kobayashi M, Kumada H.	Renal dysfunction and hypophosphatemia during long-term lamivudine plus adefovir dipivoxil therapy in patients with chronic hepatitis B.	J Gastroenterol.		Epub ahead of print
Kobayashi M, Hosaka T, Suzuki F, <u>Akuta N</u> , Sezaki H, Suzuki Y, Kawamura Y, Kobayashi M, Saitoh S, Arase Y, Ikeda K, Miyakawa Y, Kumada H	Seroclearance rate of hepatitis B surface antigen in 2,112 patients with chronic hepatitis in Japan during long-term follow-up	J Gastroenterol.		Epub ahead of print
Kawamura Y, Ikeda K, Fukushima T, Hara T, Kobayashi M, Saitoh S, Sezaki H, <u>Akuta N</u> , Suzuki F, Suzuki Y, Arase Y, Kumada H	Potential of no-touch pincer ablation procedure for small hepatocellular carcinoma that uses a multipolar radiofrequency ablation system: an experimental animal study.	Hepatol Res.		Epub ahead of print
Suzuki F, Hosaka T, Suzuki Y, <u>Akuta N</u> , Sezaki H, Hara T, Kawamura Y, Kobayashi M, Saitoh S, Arase Y, Ikeda K, Kobayashi M, Watahiki S, Mineta R, Kumada H	Long-term efficacy and emergence of multidrug resistance in patients with lamivudine-refractory chronic hepatitis B treated by combination therapy with adefovir plus	J Gastroenterol.		Epub ahead of print

	lamivudine				
Arase Y, Kobayashi M, Suzuki F, Suzuki Y, Kawamura Y, <u>Akuta N</u> , Kobayashi M, Sezaki H, Saito S, Hosaka T, Ikeda K, Kumada H, Kobayashi T.	Effect of type 2 diabetes on risk for malignancies included hepatocellular carcinoma in chronic hepatitis C	Hepatology.	3	964-973	2013
Hirakawa M, Ikeda K, Kobayashi M, Kawamura Y, Hosaka T, Sezaki H, <u>Akuta N</u> , Suzuki F, Suzuki Y, Saitoh S, Arase Y, Kumada H	Randomized controlled trial of a new procedure of radiofrequency ablation using an expandable needle hepatocellular carcinoma	Hepatol Res.	43	846-852	2013
Suzuki F, Suzuki Y, Sezaki H, <u>Akuta N</u> , Seko Y, Kawamura Y, Hosaka T, Kobayashi M, Saito S, Arase Y, Ikeda K, Mineta R, Watahiki S, Kobayashi M, Nakayasu Y, Tsuda H, Aoki K, Yamada I, Kumada H	Exploratory Study on telaprevir given every 8h at 500mg or 750mg with peginterferon-alpha-2b and rivavirin in hepatitis C patients	Hepatol Res.	43	691-701	2013
Hosaka T, Suzuki F, Kobayashi M, Seko Y, Kawamura Y, Sezaki H, <u>Akuta N</u> , Suzuki Y, Saitoh S, Arase Y, Ikeda K, Kobayashi M, Kumada H	Long-term entecavir treatment reduces hepatocellular carcinoma incidence in patients with hepatitis B virus infection	Hepatology.	58	98-107	2013
<u>Akuta N</u> , Suzuki F, Seko Y, Kawamura Y, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Hara T, Kobayashi M, Saitoh S, Arase Y, Ikeda K, Kumada H	Emergence of telaprevir-resistant variants detected by ultra-deep sequencing after triple therapy in patients infected with HCV genotype 1	J Med Virol.	85	1028-1036	2013

Kawamura Y, Saitoh S, Arase Y, Ikeda K, Fukushima T, Hara T, Seko Y, Hosaka T, Kobayashi M, Sezaki H, <u>Akuta N</u> , Suzuki F, Suzuki Y, Fukuzawa K, Hamada Y, Takahashi J, Kobayashi M, Kumada H	Three-dimensional magnetic resonance imaging for stringent diagnosis of advanced fibrosis associated with nonalcoholic steatohepatitis	Hepatol Int.	7	850-858	2013
Seko Y, Ikeda K, Kawamura Y, Fukushima T, Hara T, Sezaki H, Hosaka T, <u>Akuta N</u> , Suzuki F, Kobayashi M, Suzuki Y, Saitoh S, Arase Y, Kumada H	Antiumor efficacy of transcatheter arterial chemoembolization with warmed miriplatin in hepatocellular carcinoma	Hepatol Res.	43	942-949	2013
Hosaka T, Suzuki F, Kobayashi M, Seko Y, Kawamura Y, Sezaki H, <u>Akuta N</u> , Suzuki Y, Saitoh S, Arase Y, Ikeda K, Kobayashi M, Kumada H	Clearance of hepatitis B surface antigen during long-term nucleot(s)ide analog treatment in chronic hepatitis B: results from a nine-year longitudinal study	J Gastroenterol.	48	930-941	2013
<u>Akuta N</u> , Suzuki F, Fukushima T, Kawamura Y, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Hara T, Kobayashi M, Saitoh S, Arase Y, Ikeda K, Kumada H	Prediction of treatment efficacy and telaprevir-resistant variants after triple therapy in patients infected with HCV genotype 1	J Clin Microbiol.	51	2862-2868	2013
Kawamura Y, Ikeda K, Fukushima T, Seko Y, Hara T, Sezaki H, Hosaka T, <u>Akuta N</u> , Kobayashi M, Saitoh S, Suzuki F, Suzuki Y, Arase Y, Kumada H	What Is the most effective drug delivery system for cisplatin during the treatment of hepatic tumors with single-session transcatheter chemotherapy? A pilot study.	Gut Liver.	5	576-584	2013

Hara T, <u>Akuta N</u> , Suzuki F, Sezaki H, Susuki Y, Hosaka, Kobayashi M, Saitoh, Kumada H	A pilot study of triple therapy with telaprevir, peginterferon and ribavirin for elderly patients with genotype 1 chronic Hepatitis C.	J Med Virol.	85	1746-1753	2013
Imai N, Ikeda K, Seko Y, Kawamura Y, Sezaki H, Hosaka T, <u>Akuta N</u> , Kobayashi M, Saitoh S, Suzuki F, Susuki Y, Arase Y, Kumada H	Transcatheter arterial chemotherapy with miriplatin for hepatocellular carcinoma patients with chronic renal failure: report of three cases.	Gut Liver.	7	246-251	2013
Oze T, Hiramatsu N, Mita E, <u>Akuta N</u> , Sakamoto N, Nagano H, Itoh Y, Kaneko S, Izumi N, Nomura H, Hayashi N, Takehara T.	A multicenter survey of re-treatment with pegylated interferon plus ribavirin combination therapy for patients with chronic hepatitis C in Japan.	Hepatol Res	43	35-43	2013
Ogura S, Saitoh S, Kawamura Y, Sezaki H, Hosaka T, <u>Akuta N</u> , Kobayashi M, Suzuki F, Suzuki Y, Arase Y, Ikeda K, Kumada H	Magnetic resonance laparoscopy: A new non-invasive technique for the assessment of chronic viral disease.	Hepatol Res.	43	836-845	2013
<u>Akuta N</u> , Suzuki F, Seko Y, Kawamura Y, Sezaki H, Hosaka T, Kobayashi M, Kobayashi M, Saitoh S, Arase Y, Ikeda K, Kumada H	Efficacy and anticarcinogenic activity of ribavirin combination therapy for hepatitis C virus-related compensated cirrhosis	Intervirology.	56	37-45	2013
Seko Y, <u>Akuta N</u> , Suzuki F, Kawamura Y, Sezaki H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Saitoh S, Arase Y, Ikeda K, Kumada H	Amino acid substitutions in the hepatitis C Virus core region and lipid metabolism are associated with hepatocarcinogenesis in nonresponders to	Intervirology.	56	13-21	2013

	interferon plus ribavirin combination therapy.				
Yamada S, Kawaguchi A, <u>Kawaguchi T</u> , Fukushima N, Kuromatsu R, Sumie S, Takata A, Nakano M, Satani M, Tonan T, Fujimoto K, Shima H, Kakuma T, Torimura T, Charlton MR, Sata M	Serum albumin level is a notable profiling factor for non-B, non-C hepatitis virus-related hepatocellular carcinoma: A data-mining analysis	Hepatol Res.			in press
<u>Kawaguchi T</u> , Nagao Y, Sata M	Independent factors associated with altered plasma active ghrelin levels in HCV-infected patients	Liver Int.	33	1510-6	2013
Taniguchi E, <u>Kawaguchi T</u> , Sakata M, Itou M, Oriishi T, Sata M	Lipid profile is associated with the incidence of cognitive dysfunction in viral cirrhotic patients: A data-mining analysis	Hepatol Res.	43	418-24	2013
Sakata M, Kawahara A, <u>Kawaguchi T</u> , Akiba J, Taira T, Taniguchi E, Abe M, Koga H, Kage M, Sata M	Decreased expression of insulin and increased expression of pancreatic transcription factor PDX-1 in islets in patients with liver cirrhosis: a comparative investigation using human autopsy specimens	J Gastroenterol.	48	277-85	2013
Katoh H, Okamoto T, Fukuhara T, Kambara H, Morita E, Mori Y, Kamitani W, <u>Matsuura Y</u> .	Japanese Encephalitis Virus Core Protein Inhibits Stress Granule Formation through an Interaction with Caprin-1 and Facilitates Viral Propagation	J Virol	87	489-502	2013

Suzuki R, Matsuda M, Watashi K, Aizaki H, <u>Matsuura Y</u> , Wakita T, Suzuki T.	Signal peptidase complex subunit 1 participates in the assembly of hepatitis C virus through an interaction with E2 and NS2	PLoS Pathog		doi:10.1371/journal.ppat.1003589	2013
Lee H, Komano J, Saitoh Y, Yamaoka S, Kozaki T, Misawa T, Takahama M, Satoh T, Takeuchi O, Yamamoto N, <u>Matsuura Y</u> , Saitoh T, Akira S.	Zinc-finger antiviral protein mediates retinoic acid inducible gene I-like receptor-independent antiviral response to murine leukemia virus	Proc Natl Acad Sci U S A	110	12379-12384	2013
Yoshio S, Kanto T, Kuroda S, Matsubara T, Higashitani K, Kakita N, Ishida H, Hiramatsu N, Nagano H, Sugiyama M, Murata K, Fukuhara T, <u>Matsuura Y</u> , Hayashi N, Mizokami M, Takehara T.	Human BDCA3 (+) dendritic cells are a potent producer of IFN-λ in response to hepatitis C virus	Hepatology	57	1705-1715	2013
Kimura T, Katoh H, Kayama H, Saiga H, Okuyama M, Okamoto T, Umemoto E, <u>Matsuura Y</u> , Yamamoto M, Takeda K.	Ifit1 inhibits Japanese encephalitis virus replication through binding to 5' capped 2'-O unmethylated RNA	J Virol	87	9997-10003	2013
Tripathi LP, Kambara H, Chen YA, Nishimura Y, Moriishi K, Okamoto T, Morita E, Abe T, Mori Y, <u>Matsuura Y</u> , Mizuguchi K.	Understanding the biological context of NS5A-host interactions in HCV infection: a network-based approach	J Proteome Res	12	2537-2551	2013
Ratnoglik SL., Aoki C., Sudarmono P., Komoto M., Deng L., <u>Shoji I.</u> , Fuchino H., Kawahara N., and Hotta H.	Antiviral activity of extracts from <i>Morinda citrifolia</i> leaves and chlorophyll catabolites pheophorbide a and pyropheophorbide a, against hepatitis C virus.	Microbiol. Immunol.		doi: 10.1111/1348-0421.12133	2014

Adianti M., Aoki C., Komoto M., Deng L., <u>Shoji I.</u> , Wahyuni T., Lusida M., Soetjipto S., Fuchino H., Kawahara N., and Hotta H.	Anti-hepatitis C virus compounds obtained from Glycyrrhiza uralensis and other Glycyrrhiza species.	Microbiol. Immunol.		doi: 10.1111/1348-0421.12127	2014
Tao RR, Huang JY, Lu YM, Hong LJ, Wang H, Masood MA, Ye WF, Zhu DY, Huang Q, Fukunaga K, Lou YJ, <u>Shoji I.</u> , Wilcox CS, Lai EY, Han F.	Nitrosative stress induces peroxiredoxin 1 ubiquitination during ischemic insult via E6AP activation in endothelial cells both in vitro and in vivo.	Antioxid & Redox Sign.		doi: 10.1089/ars.2013.5381	2013
Mawatari S., Uto H., Ido A., Nakashima K., Suzuki T., Kanmura S., Kumagai K., Oda K., Tabu K., Tamai T., Moriuchi A., Oketani M., Shimada Y., Sudoh M., <u>Shoji I.</u> , and Tsubouchi H.	Hepatitis C virus NS3/4A protease inhibits complement activation by cleaving complement component 4.	PLoS One	8	e82094, 1-7	2013
Wahyuni TS., Tumewu L., Permanasari AA., Apriani E., Adianti M., Rahman A., Widyawaruyanti A., Lusida MI., Fuad A., Soetjipto, Nasronudin, Fuchino H., Kawahara N., <u>Shoji I.</u> , Deng L., Aoki C., and Hotta H.	Antiviral activities of Indonesian medicinal plants in the East Java region against hepatitis C virus.	Virol. J.	10	259, 1-9	2013
Ichimura T, Taoka M, <u>Shoji I.</u> , Kato H, Hatakeyama S, Isobe T, and Hachiya N.	14-3-3 Proteins sequester a pool of soluble TRIM32 ubiquitin ligase to repress autoubiquitination and cytoplasmic body formation.	J. Cell. Sci.	126	2014-2026	2013
El-Shamy A, Shindo M, <u>Shoji I.</u> , Deng L, Okuno T, and Hotta H.	Polymorphisms of the Core, NS3 and NS5A proteins of hepatitis C virus genotype 1b associate with development of hepatocellular carcinoma.	Hepatology	58	555-563	2013

Horiuchi Y, Takagi A, Kobayashi N, Moriya O, Nagai T, <u>Moriya K</u> , Tsutsumi T, Koike K, Akatsuka T.	The effect of the infectious dose and the presence of HCV core gene on mouse intrahepatic CD8 T-cells.	<b>Hepatol Res</b>			[Epub ahead of print]
Urbnileg B, Enooku K, Soroida Y, Ohkawa R, Kudo Y, Nakagawa H, Tateishi R, Yoshida H, Shinzawa S, <u>Moriya K</u> , Ohtomo N, Nishikawa T, Inoue Y, Tomiya T, Kojima S, Matsuura T, Koike K, Yatomi Y, Ikeda H.	High ubiquitous mitochondrial creatine kinase expression in hepatocellular carcinoma denotes a poor prognosis with highly malignant potential.	Int J Cancer.	134	2189-98.	2014
Yotsuyanagi H, Ito K, Yamada N, Takahashi H, Okuse C, Yasuda K, <u>Suzuki M, Moriya K</u> , Mizokami M, Miyakawa Y, Koike K	High levels of hepatitis B virus after the onset of disease lead to chronic infection in patients with acute hepatitis B.	Clin Infect Dis.	57(7)	935-42	2013
He G, Dhar D, Nakagawa H, Font-Burgada J, Ogata H, Jiang Y, Shalapour S, Seki E, Yost SE, Jepsen K, Frazer KA, Harismendy O, Hatziapostolou M, Iliopoulos D, Suetsugu A, Hoffman RM, <u>Tateishi R, Koike K</u> , Karin M.	Identification of Liver Cancer Progenitors Whose Malignant Progression Depends on Autocrine IL-6 Signaling	Cell	155(2)	284-96	2013
Sato M, Kato N, <u>Tateishi R, Muroyama</u> R, Kowatari N, Li W, Goto K, Otsuka M, Shiina S, Yoshida H, Omata M, Koike K.	The Impact of Pnpla3 Polymorphisms on the Development of Hepatocellular Carcinoma in Patients with Chronic Hepatitis C Virus Infection.	Hepatol Res			2014