

Perihepatic lymph node enlargement is a negative predictor of liver cancer development in chronic hepatitis C patients

Hiromi Hikita · Hayato Nakagawa · Ryosuke Tateishi · Ryota Masuzaki · Kenichiro Enooku · Haruhiko Yoshida · Masao Omata · Yoko Soroida · Mamiko Sato · Hiroaki Gotoh · Atsushi Suzuki · Tomomi Iwai · Hiromitsu Yokota · Kazuhiko Koike · Yutaka Yatomi · Hitoshi Ikeda

Received: 5 January 2012 / Accepted: 19 June 2012 / Published online: 12 July 2012
© Springer 2012

Abstract

Background Perihepatic lymph node enlargement (PLNE) is a common ultrasound finding in chronic hepatitis C patients. Although PLNE is considered to reflect the inflammatory response to hepatitis C virus (HCV), its clinical significance remains unclear.

Methods Between December 2004 and June 2005, we enrolled 846 chronic hepatitis C patients in whom adequate ultrasound examinations had been performed. PLNE was defined as a perihepatic lymph node that was at least 1 cm in the longest axis by ultrasonography. We analyzed the clinical features of patients with PLNE and prospectively investigated the association between PLNE and hepatocellular carcinoma (HCC) development.

Results We detected PLNE in 169 (20.0 %) patients. Female sex, lower body mass index (BMI), and HCV serotype 1 were independently associated with the presence of

PLNE. However, there were no significant differences in liver function tests, liver stiffness, and hepatitis C viral loads between patients with and without PLNE. During the follow-up period (mean 4.8 years), HCC developed in 121 patients. Unexpectedly, patients with PLNE revealed a significantly lower risk of HCC development than those without PLNE ($p = 0.019$, log rank test). Multivariate analysis revealed that the presence of PLNE was an independent negative predictor of HCC development (hazard ratio 0.551, $p = 0.042$). In addition, the sustained viral response rate in patients who received interferon (IFN) therapy was significantly lower in patients with PLNE than in patients without PLNE.

Conclusions Patients with PLNE had a lower risk of HCC development than those without PLNE. This study may provide new insights into daily clinical practice and the pathophysiology of HCV-induced hepatitis and hepatocarcinogenesis.

Keywords Perihepatic lymph node enlargement · Chronic hepatitis C · Hepatocarcinogenesis

H. Hikita and H. Nakagawa contributed equally to this work.

H. Hikita · H. Nakagawa · K. Enooku · Y. Soroida · M. Sato · H. Gotoh · A. Suzuki · T. Iwai · H. Yokota · Y. Yatomi · H. Ikeda

Department of Clinical Laboratory Medicine,
Graduate School of Medicine, University of Tokyo,
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan

H. Nakagawa (✉) · R. Tateishi · R. Masuzaki · K. Enooku · H. Yoshida · K. Koike · H. Ikeda
Department of Gastroenterology, Graduate School of Medicine,
University of Tokyo, 7-3-1 Hongo, Bunkyo-ku,
Tokyo 113-8655, Japan
e-mail: n-hayato@y17.so-net.ne.jp

M. Omata
Yamanashi Prefectural Hospital Organization,
1-1-1 Fujimi, Kofu, Yamanashi 400-8506, Japan

Abbreviations

ALT	Alanine aminotransferase
AFP	α -Fetoprotein
PLNE	Perihepatic lymph node enlargement
HCV	Hepatitis C virus
HCC	Hepatocellular carcinoma
IFN	Interferon
LN	Lymph node
US	Ultrasonography

Introduction

Hepatocellular carcinoma (HCC) is the fifth most common cancer worldwide and chronic hepatitis C virus (HCV)

infection is a major cause of HCC in the United States, southern European countries, and Japan [1, 2]. The host immune responses to HCV are often not strong enough to completely clear the infection, resulting in chronic stimulation of the antigen-specific immune response. Accumulating basic and clinical lines of evidence indicate that a sustained inflammatory reaction in the liver is the major contributing factor to HCC development [3–5].

Inflammatory processes in organs frequently lead to hyperplasia of regional lymph nodes (LNs). Perihepatic LN enlargement (PLNE) is a common finding in patients with chronic hepatitis, especially in patients with hepatitis C [6, 7]. Some studies have revealed that PLNE in chronic hepatitis C patients was associated with a higher HCV viral load, [7] higher histological grade of hepatic inflammation and fibrosis [8, 9], and higher CD8 lymphocyte level in the blood [10]; therefore, PLNE is considered to reflect an inflammatory response to HCV. However, such associations as those noted above are inconsistent among studies [7–15], and the precise mechanism and clinical relevance of PLNE are not fully understood. Furthermore, to our knowledge, there have been no studies designed to investigate the association between PLNE and hepatocarcinogenesis. To clarify whether we should pay attention to the risk of HCC development in patients with PLNE is very important, because we encounter such situations very often.

The purpose of this study was to reevaluate the clinical relevance of PLNE and to carry out a prospective assessment of the association between PLNE and HCC development. To elucidate these matters, we investigated a well-characterized chronic hepatitis C cohort in which we previously reported the utility of performing transient elastography for risk assessment of HCC development [16]. In the present study, we prospectively assessed the association between PLNE and HCC development in chronic hepatitis C patients in that cohort.

Patients and methods

Patients and screening for perihepatic LN enlargement

As described previously [16], we enrolled 866 chronic hepatitis C patients, excluding those with HCC or a past history of it, who visited the University of Tokyo Hospital between December 2004 and June 2005. All patients were positive for HCV-RNA and showed at least a transiently elevated serum alanine aminotransferase (ALT) level. Patients with concomitant hepatitis B virus surface antigen positivity, patients with uncontrollable ascites, patients on interferon (IFN) therapy, and patients who visited only for consultation purposes were excluded from this study.

Each subject was screened for HCC with ultrasonography (US) at or immediately after the first visit. At the same time, we surveyed the presence of PLNE with US. The US examination was performed using the SSD-2000 (Aloka, Tokyo, Japan). To identify LNs the following criteria were used, according to a previous report [17]: one or more masses with an ovoid shape and less echogenic than liver parenchyma, separated from adjacent organs and vessels by a clear-cut cleavage on repeated transverse, sagittal, and oblique scans. LNs were searched for near the trunk of the portal vein, hepatic artery, celiac axis, superior mesenteric vein, and pancreas head. Furthermore, we used Doppler US to differentiate LNs from vessels.

We defined PLNE as an LN that was at least 1 cm in the longest axis. There were two reasons for this definition. First, we preliminarily investigated the prevalence of PLNE with US in 465 healthy subjects who had had medical check-ups and did not have liver disease or other underlying causes of LN enlargement. We found that only 15 (3.2 %) of these subjects had a perihepatic LN larger than 1 cm in the longest axis (unpublished data). Second, when an LN was smaller than 1 cm, it was sometimes difficult to distinguish the LN from other structures. If two or more LNs were detected with US, we determined PLNE to be present based on the length of the largest LN. In 20 of the 866 patients in the present study, adequate visualization of the liver hilum was not achieved with US because of severe obesity or excessive meteorism. Therefore, we analyzed the association between PLNE and the subsequent incidence of HCC in 846 patients.

HCV RNA was measured using Amplicore HCV version 2.0 (Roche, Tokyo, Japan), and the HCV serotype was examined using a serotyping assay (SRL, Tokyo, Japan). We also monitored IFN therapy and responses during the follow-up period. A sustained virological response (SVR) was defined as undetectable HCV-RNA at least 24 weeks after the end of therapy. All blood tests were performed at the time of US examination.

The study protocol conformed to the ethical guidelines of the 1975 Declaration of Helsinki.

Patient follow up

Patients were followed up every 3–6 months at the outpatient clinic, when blood tests (including tumor markers) and US were carried out. Contrast-enhanced computed tomography (CT) was performed when HCC was suspected, based on US, and/or if the serum α -fetoprotein (AFP) level showed an abnormal increase. HCC was diagnosed by dynamic CT, and hyperattenuation in the arterial phase with washout in the late phase was considered a definite sign of HCC [18]. When a diagnosis of HCC was ambiguous, ultrasound-guided tumor biopsy was

performed and a pathologic diagnosis was made based on the Edmondson and Steiner criteria. Time to HCC occurrence was defined as the interval between the date of the first US screening and the diagnosis of HCC. Patients were censored at the time of death without HCC development, the last visit when lost to follow up, or the end of the study period. The last observation in this study was made on December 31, 2010. Thus, the time of observation was extended from that of our previous study, which was censored on May 31, 2008 [16].

Transient elastography

Transient elastography was performed using Fibroscan (Echosens, Paris, France) as described previously [16].

Statistical analysis

Data were expressed as means \pm standard deviation (SD) unless otherwise indicated. Categorical variables were compared by χ^2 tests, whereas continuous variables were compared by the unpaired Student's *t*-test (parametric) or the Mann–Whitney *U*-test (non-parametric). Multivariate logistic regression analysis was used to identify factors that were independently associated with the presence of PLNE. Cumulative HCC incidence was estimated using the Kaplan–Meier method, and the difference between groups was assessed with the log-rank test. In the analysis of risk factors for hepatocarcinogenesis, we tested the following variables in univariate analysis and multivariate Cox proportional hazard regression analysis: age, sex, platelet count, serum albumin concentration, total bilirubin concentration, ALT and aspartate aminotransferase (AST) levels, higher AFP concentration (>10 ng/ml), prothrombin activity, heavy alcohol drinking (alcohol intake >80 g/day), BMI, higher liver stiffness measurement (LSM) (>10 kPa), HCV serotype, HCV viral load (>100 kIU/ml), IFN treatment, achievement of SVR, and presence of PLNE. A *p* value of less than 0.05 on a two-tailed test was considered significant. Data processing and analysis were performed using StatView (ver. 5.0; SAS Institute, Cary, NC, USA) and SPSS (ver. 14.0; SPSS, Chicago, IL, USA) software.

Results

Patient profiles

We detected PLNE in 169 of 846 (20.0 %) patients with chronic hepatitis C. A representative ultrasound image is shown in Fig. 1. The mean (\pm SD) length of the longest axis was $1.7 (\pm 0.5)$ cm (range 1.0–3.5 cm). The clinical

features of patients with and without PLNE are summarized in Table 1. The proportion of females was significantly higher in the PLNE-positive group than in the PLNE-negative group (63.3 vs. 52.9 %), and BMI was slightly but significantly lower in the PLNE-positive group than in the PLNE-negative group (21.9 ± 2.6 vs. 22.5 ± 2.9). The proportion of HCV serotype 1 patients was higher in the PLNE-positive group than in the PLNE-negative group, with borderline significance. There was a tendency of a higher serum ALT level in the PLNE-positive group, but the difference was without statistical significance. There were no significant differences in other liver function test results, or in liver stiffness and hepatitis C viral load between the two groups. Multivariate logistic regression analysis using the factors of sex, serum ALT, BMI, and HCV serotype revealed that female sex, lower BMI, and HCV serotype 1 were independently associated with the presence of PLNE (Table 2).

Incidence of HCC

The mean follow-up period was 4.8 years, constituting a total observation of 4,021 person-years. During the observation period, 70 (8.3 %) patients were lost to follow up: 15 (8.8 %) patients in the PLNE-positive group and 55 (8.1 %) patients in the PLNE-negative group. There were no patients in whom an enlarged perihepatic LN turned out to be caused by other underlying diseases including metastasis of HCC. The SVR rate in patients who received IFN therapy during the follow-up period was significantly lower in the PLNE-positive group compared with that in the PLNE-negative group [7/34 (20.6 %) vs. 93/172




Fig. 1 Representative ultrasound image of enlarged perihepatic lymph node (LN) in a patient with chronic hepatitis C

Table 1 Clinical features of patients with and without PLNE

Variable	PLNE-positive group (n = 169)	PLNE-negative group (n = 677)	p value
Age (years)	62.4 ± 10.1 (29–83)	62.4 ± 11.5 (17–89)	0.58
Male, n (%)	62 (36.7)	319 (47.1)	0.018
Serum albumin (g/dl)	4.0 ± 0.4 (2.8–4.8)	4.0 ± 0.4 (2.5–5.0)	0.93
Total bilirubin (mg/dl)	0.8 ± 0.3 (0.3–2.1)	0.9 ± 0.5 (0.3–4.6)	0.23
AST (IU/l)	52 ± 34.1 (17–223)	50 ± 33.8 (9–286)	0.16
ALT (IU/l)	57 ± 48.7 (4–374)	53 ± 45.2 (2–503)	0.10
Platelet count (×10 ³ /μl)	16.1 ± 6.6 (2.1–42.2)	16.1 ± 6.7 (3.2–43.6)	0.89
Prothrombin time (%)	86.0 ± 12.1 (50.3–100.0)	85.7 ± 12.4 (38.9–100.0)	0.88
AFP (ng/ml)	22.0 ± 67.9 (1–592)	13.4 ± 37.1 (1–563)	0.61
BMI (kg/m ²)	21.9 ± 2.6 (14.4–28.7)	22.5 ± 2.9 (15.1–29.8)	0.007
Liver stiffness (kPa)	10.9 ± 7.8 (2.8–42.2)	12.0 ± 10.0 (2.5–75.0)	0.59
Alcohol consumption >80 g/day, n (%)	6 (3.6)	25 (3.7)	0.82
HCV viral load (kIU/ml)	549 ± 646 (5–5000)	651 ± 842 (5–5000)	0.48
HCV serotype 1, n (%)	146 (86.3)	538 (79.5)	0.053
Patients who received IFN, n (%)	34 (20.1)	172 (25.4)	0.18
Patients who achieved SVR, n (%)	7 (4.1)	93 (13.7)	0.0009

PLNE perihepatic lymph node enlargement, AST aspartate aminotransferase, ALT alanine aminotransferase, AFP α-fetoprotein, BMI body mass index, HCV hepatitis C virus

Table 2 Factors associated with the presence of PLNE: multivariate analysis

Variable	Odds ratio (95 % confidence interval [CI])	p value
Male sex	0.667 (0.464–0.936)	0.024
ALT level (per 1 IU/l)	1.003 (0.999–1.006)	0.10
BMI (per 1 kg/m ²)	0.919 (0.864–0.978)	0.017
HCV serotype 1	1.64 (1.02–2.66)	0.043

(54.1 %), p = 0.0005]. This finding was consistent with previous reports [12, 19].

By the end of the follow-up period, HCC had developed in 121 patients (3.0 % per 1 person-year). The cumulative incidence rates of HCC at 3 and 5 years estimated by the Kaplan–Meier method were 8.9 and 13.7 %, respectively. We then assessed the incidence of HCC stratified by the presence of PLNE. Unexpectedly, the PLNE-positive group revealed a significantly lower incidence of HCC than the PLNE-negative group (p = 0.019, log-rank test) (Fig. 2). The cumulative incidence rates at 3 and 5 years were 3.6 and 8.2 %, respectively, in the PLNE-positive group, and 10.1 and 15.1 % in the PLNE-negative group. These results indicate that patients with PLNE have a lower risk of HCC development despite having a lower SVR rate with IFN therapy.

Risk analyses

We analyzed the risk factors for HCC development. In the univariate analyses, older age, male sex, lower serum




Fig. 2 Cumulative incidence of hepatocellular carcinoma (HCC) development stratified by the presence of perihepatic lymph node enlargement (PLNE)

albumin concentration, higher total bilirubin concentration, higher AST level, higher ALT level, lower prothrombin activity, lower platelet count, heavy alcohol drinking, higher BMI, LSM greater than 10 kPa, AFP level greater than 10 ng/ml, HCV serotype 1, not receiving IFN, not achieving SVR, and absence of PLNE were significant risk factors for HCC (Table 3). As we had reported previously, a higher LSM (i.e., greater than 10 kPa) was a strong predictor of HCC development [hazard ratio (HR) 15.4, 95 % confidence interval (CI) 8.6–27.0, p < 0.0001]. Multivariate proportional hazard regression analyses

Table 3 Risk factors for HCC development: univariate and multivariate analyses

Variable	Univariate analysis		Multivariate analysis	
	Hazard ratio (95 % CI)	<i>p</i> value	Hazard ratio (95 % CI)	<i>p</i> value
Age (per 1 year age)	1.07 (1.05–1.09)	<0.0001	1.04 (1.01–1.06)	0.002
Male sex	1.45 (1.02–2.08)	0.041	1.49 (1.02–2.17)	0.039
Platelet count (per 10 ³ /μl)	0.852 (0.823–0.882)	<0.0001	0.965 (0.926–1.005)	0.089
Total bilirubin (per 1 mg/dl)	1.88 (1.45–2.45)	<0.0001	0.825 (0.567–1.2)	0.32
Serum albumin level (per 1 g/dl)	0.12 (0.084–0.17)	<0.0001	0.441 (0.263–0.739)	0.002
AST level (per 1 IU/l)	1.01 (1.007–1.014)	<0.0001	1.002 (0.991–1.013)	0.71
ALT level (per 1 IU/l)	1.004 (1.002–1.007)	0.002	1.0 (0.991–1.013)	0.94
AFP level >10 ng/ml	6.76 (4.69–9.8)	<0.0001	1.9 (1.22–2.97)	0.005
Prothrombin time (per 1 %)	0.973 (0.966–0.979)	<0.0001	0.989 (0.976–1.001)	0.072
Alcohol consumption >80 g/day	2.73 (1.43–5.24)	0.002	3.53 (1.76–7.09)	0.0004
BMI (per 1 kg/m ²)	1.09 (1.03–1.16)	0.006	1.09 (1.01–1.17)	0.025
Liver stiffness >10 kPa	15.4 (8.6–27.0)	<0.0001	4.41 (2.24–8.7)	<0.0001
HCV serotype 1	1.76 (1.03–3.03)	0.04	1.36 (0.774–2.38)	0.29
HCV-RNA >100 kIU/ml	1.36 (0.87–2.13)	0.18	1.24 (0.781–1.97)	0.36
Patients treated with IFN	0.44 (0.262–0.75)	0.002	0.59 (0.315–1.11)	0.1
Patients with SVR	0.175 (0.055–0.549)	0.003	0.621 (0.169–2.28)	0.47
Presence of PLNE	0.53 (0.31–0.91)	0.02	0.551 (0.31–0.978)	0.042

HCC hepatocellular carcinoma, IFN interferon, SVR sustained viral response

revealed that older age, male sex, lower serum albumin concentration, AFP level greater than 10 ng/ml, heavy alcohol drinking, higher BMI, LSM greater than 10 kPa, and absence of PLNE were independent risk factors for HCC (Table 3). These results suggest that the presence of PLNE is an independent negative predictor of HCC development in chronic hepatitis C patients.

Subgroup analysis of non-obese patients

To further rule out the possibility that obesity acted as a confounder in the association between the presence of PLNE and HCC development, we reanalyzed the contribution of PLNE to HCC development in a subgroup of non-obese patients, defined as those with BMI <25 kg/m² (*n* = 695), because we could clearly visualize the liver hilum in such individuals. As shown in Fig. 3, the PLNE-positive group had a significantly lower incidence of HCC than the PLNE-negative group even in the non-obese subgroup (*p* = 0.02). Thus, we further confirmed that the presence of PLNE was negatively associated with HCC development independently of obesity.

Significance of the size of perihepatic LNs

To examine the significance of the size of perihepatic LNs, we divided patients with PLNE into two groups: a smaller LN group (longest axis of LN 1 cm to 2 cm, *n* = 122) and a larger LN group (longest axis of LN ≥2 cm, *n* = 47).




Fig. 3 Cumulative incidence of HCC development stratified by the presence of PLNE: subgroup analysis of non-obese patients (body mass index [BMI] <25 kg/m²)

The characteristics of each group are summarized in Table 4. The proportion of male patients and the LSM value tended to be higher in the larger LN group than in the smaller LN group, but the difference was not statistically significant for either factor. There were no significant differences in other factors between the two groups. Furthermore, there was no significant difference in HCC incidence rates between the two groups (Fig. 4), although the larger LN group revealed a slightly higher incidence of

Table 4 Comparison of clinical features between patients with small perihepatic lymph nodes (LNs) and those with large perihepatic LNs

Variable	LN size 1 cm to <2 cm, (n = 122)	LN size ≥2 cm (n = 47)	p value
Age (years)	62.7 ± 10.4 (29–83)	61.5 ± 9.2 (32–77)	0.32
Male, n (%)	40 (32.8)	22 (46.8)	0.092
Serum albumin (g/dl)	4.0 ± 0.3 (3.0–4.8)	4.0 ± 0.4 (2.8–4.8)	0.96
Total bilirubin (mg/dl)	0.8 ± 0.3 (0.3–1.6)	0.8 ± 0.4 (0.3–2.1)	0.98
AST (IU/l)	53 ± 36.0 (17–223)	50 ± 28.7 (17–181)	0.83
ALT (IU/l)	56 ± 49.0 (4–374)	59 ± 48.2 (6–308)	0.48
Platelet count (×10 ³ /μl)	16.3 ± 6.2 (2.1–36.4)	15.4 ± 7.4 (4.8–42.2)	0.25
Prothrombin time (%)	86.7 ± 11.7 (57.4–100.0)	84.2 ± 13.0 (50.3–100.0)	0.27
AFP (ng/ml)	13.0 ± 27.7 (1–168)	45.3 ± 118.6 (1–592)	0.19
BMI (kg/m ²)	21.8 ± 2.6 (16.8–28.0)	22.0 ± 2.6 (14.4–28.7)	0.75
Liver stiffness (kPa)	10.2 ± 7.1 (2.8–37.4)	12.5 ± 9.3 (4.2–42.2)	0.064
Alcohol consumption >80 g/day, n (%)	5 (4.1)	1 (2.1)	0.54
HCV viral load (kIU/ml)	658 ± 788	504 ± 582	0.17
HCV serotype 1, n (%)	107 (87.7)	39 (83.0)	0.58
Patients who received IFN, n (%)	24 (19.6)	10 (21.2)	0.98
Patients who achieved SVR, n (%)	5 (4.1)	2 (4.3)	0.99




Fig. 4 Cumulative incidence of HCC development in patients with PLNE stratified by the size of perihepatic LNs: i.e., smaller (longest axis of LN 1 to <2 cm) and larger (longest axis of LN ≥2 cm)

HCC. These results suggest that the size of perihepatic LNs in chronic hepatitis C patients may not be clinically as important as the presence of PLNE itself.

Discussion

Although PLNE is a common finding in patients with chronic hepatitis C, its clinical significance has remained unclear. In the present study, we reevaluated the clinical relevance of PLNE in a large cohort of chronic hepatitis C patients. We found, by prospective analysis, that patients with PLNE had a lower risk of HCC development than

those without PLNE. To our knowledge, this is the first study reporting a negative association between the presence of PLNE and HCC development.

Before we started this study, we expected that patients with PLNE would have a higher risk of HCC development, based on previous reports showing positive associations between PLNE and liver inflammation and fibrosis [8, 13–15]. However, in our study, neither inflammatory markers, such as serum AST and ALT levels, nor fibrosis markers, such as the platelet count and LSM, had statistically significant associations with the presence of PLNE. On the contrary, patients with PLNE revealed a significantly lower risk of HCC development. One possible explanation for this result is that obesity may affect the ability of US to detect perihepatic LNs, although patients with severe obesity were excluded from the study. To rule out the effect of confounders, especially obesity, we performed multivariate analysis and subgroup analysis of non-obese patients, and the results showed that the presence of PLNE was an independent negative predictor of HCC development. Additionally, of the 846 patients enrolled in this study, 175 patients underwent abdominal computed tomography (CT) within one year from the date of the US examination. The concordance rate for the diagnosis of PLNE between CT and US in these patients was 91.4 % (160/175). Therefore, we consider that the diagnostic accuracy of US for PLNE was acceptable in this study.

Although the mechanism of PLNE in patients with hepatitis C is still unknown, hyperplasia of regional LNs is generally considered to reflect inflammatory responses in the adjacent organs. The volume of perihepatic LNs has been reported to significantly decrease after antiviral

therapy, especially in patients with an SVR, supporting the hypothesis that PLNE reflects the inflammatory response to HCV [19–21]. In fact, PLNE was reported to be associated with CD8 lymphocyte counts in the peripheral blood [10]. Furthermore, HCV-specific IFN- γ production and proliferative responses of T cells were found most commonly in perihepatic LNs rather than in liver tissue or in the peripheral blood, indicating that there was ongoing T-cell activation in perihepatic LNs [22]. Our results, taken together with these previous reports, suggest that the presence of PLNE may reflect an adequate host immune response to HCV. T-cell immunity is very important in the control of HCV infection and in the prevention of hepatocarcinogenesis [23–25], and a T-cell response that is too weak may accelerate hepatocarcinogenesis, as seen in patients co-infected with HCV and human immunodeficiency virus [26, 27]. Thus, a weak T-cell response may be one explanation of the higher risk of HCC development in patients without PLNE. On the other hand, too strong an anti-HCV T-cell response may induce hepatocellular damage and lead to subsequent hepatocarcinogenesis [28], so patients with larger perihepatic LNs may have a slightly higher tendency to develop HCC. However, from the present type of observational study, we cannot evaluate a causal relationship between PLNE and hepatocarcinogenesis, so further studies are needed to clarify this point.

As mentioned above, several studies have shown that PLNE was positively associated with the degree of liver inflammation or fibrosis [8, 13–15], but, in the present study we could not find such associations, except for slight serum ALT elevation. However, because of ethical concerns regarding the performance of liver biopsy, we did not assess liver histology, so we cannot conclude whether or not PLNE is really associated with liver inflammation and fibrosis. Of note, the reported relationships of PLNE to liver function tests and liver inflammation and fibrosis are inconsistent among studies [7–15]. One reason may be that these findings were based on relatively small samples. Another reason is that there is a lack of established criteria for the diagnosis of PLNE. The lack of definite criteria may also contribute to the wide variation in the prevalence of PLNE among studies (from 20 to 100 %) [8–10, 21]. We defined PLNE as an LN that was at least 1 cm in the longest axis, and this definition was based on the report by Grier et al. [21] and our preliminary investigation in healthy subjects. Some studies have used more detailed measurements of LNs with calculations of node volume and shape [8, 10, 20]. These methods are certainly more accurate in terms of the assessment of nodal volume, but may be too complicated in the clinical setting, as discussed by Grier et al. [21]. We used a simpler method, because our study included a large number of patients and was conducted to examine the significance of PLNE in daily clinical practice. Admittedly, a

more detailed method would be appropriate to elucidate more clearly the involvement of PLNE in the pathophysiology of hepatitis and hepatocarcinogenesis.

In the present study, female sex, lower BMI, and HCV serotype 1 were independently associated with the presence of PLNE. Soresi et al. [29] also reported that PLNE was observed significantly more often in female patients than in male patients with chronic hepatitis C. Although we cannot clarify the mechanism underlying this association, this finding may be interesting from the point of view of gender differences in immune systems and hepatocarcinogenesis. In the study by Soresi et al., BMI in patients with PLNE tended to be lower than that in patients without PLNE, although the difference was not statistically significant [29], and this finding may be in line with our present results. Regarding BMI in patients with chronic HCV infection, an anti-HCV specific immune response was reportedly associated with lower BMI through the expression of adiponectin, one of the major adipokines [30]. Thus, the active immune response to HCV in patients with lower BMI might cause PLNE. Recent studies have reported that obesity and obesity-induced dysregulation of adipokines play important roles in hepatocarcinogenesis [31–33], so the examination of adipokine expression may help to explain the relationship of PLNE to BMI and hepatocarcinogenesis.

Another important finding in our study was that the SVR rate in patients who received IFN therapy was significantly lower in patients with PLNE than in patients without PLNE. This finding is consistent with previous reports [12, 19]. Although the proportion of individuals with HCV serotype 1 was higher in our patients with PLNE than in patients without PLNE, subgroup analysis of the patients with HCV serotype 1 also revealed a significantly lower SVR rate in patients with PLNE than in patients without PLNE (data not shown). Therefore, further analyses are planned to clarify the relationship of PLNE to HCV serotype and response to IFN therapy.

In conclusion, the presence of PLNE is an independent negative predictor of HCC development in chronic hepatitis C patients. This study may provide new insights into daily clinical practice and the pathophysiology of HCV-induced hepatitis and hepatocarcinogenesis.

Acknowledgments This study was not supported by any grants.

Conflict of interest The authors have no conflicts of interest regarding this study.

References

1. Bosch FX, Ribes J, Diaz M, Cleries R. Primary liver cancer: worldwide incidence and trends. *Gastroenterology*. 2004;127: S5–16.

2. El-Serag HB, Rudolph KL. Hepatocellular carcinoma: epidemiology and molecular carcinogenesis. *Gastroenterology*. 2007;132:2557–76.
3. Levvero M. Viral hepatitis and liver cancer: the case of hepatitis C. *Oncogene*. 2006;25:3834–47.
4. Berasain C, Castillo J, Perugorria MJ, Latasa MU, Prieto J, Avila MA. Inflammation and liver cancer: new molecular links. *Ann N Y Acad Sci*. 2009;1155:206–21.
5. Nakagawa H, Maeda S, Yoshida H, Tateishi R, Masuzaki R, Ohki T, et al. Serum IL-6 levels and the risk for hepatocarcinogenesis in chronic hepatitis C patients: an analysis based on gender differences. *Int J Cancer*. 2009;125:2264–9.
6. Kuo HT, Lin CY, Chen JJ, Tsai SL. Enlarged lymph nodes in porta hepatis: sonographic sign of chronic hepatitis B and C infections. *J Clin Ultrasound*. 2006;34:211–6.
7. Watanabe T, Sassa T, Hiratsuka H, Hattori S, Abe A. Clinical significance of enlarged perihepatic lymph node on ultrasonography. *Eur J Gastroenterol Hepatol*. 2005;17:185–90.
8. Dietrich CF, Lee JH, Herrmann G, Teuber G, Roth WK, Caspary WF, et al. Enlargement of perihepatic lymph nodes in relation to liver histology and viremia in patients with chronic hepatitis C. *Hepatology*. 1997;26:467–72.
9. Soresi M, Carroccio A, Bonfissuto G, Agate V, Magliarisi C, Aragona F, et al. Ultrasound detection of abdominal lymphadenomegaly in subjects with hepatitis C virus infection and persistently normal transaminases: a predictive index of liver histology severity. *J Hepatol*. 1998;28:544–9.
10. Muller P, Renou C, Harafa A, Jouve E, Kaplanski G, Ville E, et al. Lymph node enlargement within the hepatoduodenal ligament in patients with chronic hepatitis C reflects the immunological cellular response of the host. *J Hepatol*. 2003;39:807–13.
11. Cassani F, Valentini P, Cataleta M, Manotti P, Francesconi R, Giostra F, et al. Ultrasound-detected abdominal lymphadenopathy in chronic hepatitis C: high frequency and relationship with viremia. *J Hepatol*. 1997;26:479–83.
12. del Olmo JA, Esteban JM, Maldonado L, Rodriguez F, Escudero A, Serra MA, et al. Clinical significance of abdominal lymphadenopathy in chronic liver disease. *Ultrasound Med Biol*. 2002;28:297–301.
13. Tavakoli-Tabasi S, Ninan S. Clinical significance of perihepatic lymphadenopathy in patients with chronic hepatitis C infection. *Dig Dis Sci*. 2011;56:2137–44.
14. Soresi M, Carroccio A, Agate V, Bonfissuto GD, Magliarisi C, Fulco M, et al. Evaluation by ultrasound of abdominal lymphadenopathy in chronic hepatitis C. *Am J Gastroenterol*. 1999;94:497–501.
15. Soresi M, Bonfissuto G, Magliarisi C, Riili A, Terranova A, Di Giovanni G, et al. Ultrasound detection of abdominal lymph nodes in chronic liver diseases. A retrospective analysis. *Clin Radiol*. 2003;58:372–7.
16. Masuzaki R, Tateishi R, Yoshida H, Goto E, Sato T, Ohki T, et al. Prospective risk assessment for hepatocellular carcinoma development in patients with chronic hepatitis C by transient elastography. *Hepatology*. 2009;49:1954–61.
17. Cassani F, Zoli M, Baffoni L, Cordiani MR, Brunori A, Bianchi FB, et al. Prevalence and significance of abdominal lymphadenopathy in patients with chronic liver disease: an ultrasound study. *J Clin Gastroenterol*. 1990;12:42–6.
18. Bruix J, Sherman M. Management of hepatocellular carcinoma. *Hepatology*. 2005;42:1208–36.
19. Friedrich-Rust M, Forestier N, Sarrazin C, Reesink HW, Herrmann E, Zeuzem S. Ultrasound evaluation of perihepatic lymph nodes during antiviral therapy with the protease inhibitor telaprevir (VX-950) in patients with chronic hepatitis C infection. *Ultrasound Med Biol*. 2007;33:1362–7.
20. Dietrich CF, Stryjek-Kaminska D, Teuber G, Lee JH, Caspary WF, Zeuzem S. Perihepatic lymph nodes as a marker of antiviral response in patients with chronic hepatitis C infection. *AJR Am J Roentgenol*. 2000;174:699–704.
21. Grier S, Patel N, Kuo YT, Cosgrove DO, Goldin RC, Thomas HC, et al. Perihepatic lymph nodes as markers of disease response in patients with hepatitis C-related liver disease: a prospective clinical evaluation. *Eur J Gastroenterol Hepatol*. 2010;22:257–63.
22. Moonka D, Milkovich KA, Rodriguez B, Abouljoud M, Lederman MM, Anthony DD. Hepatitis C virus-specific T-cell gamma interferon and proliferative responses are more common in perihepatic lymph nodes than in peripheral blood or liver. *J Virol*. 2008;82:11742–8.
23. Bowen DG, Walker CM. Adaptive immune responses in acute and chronic hepatitis C virus infection. *Nature*. 2005;436:946–52.
24. Nakagawa H, Hirata Y, Takeda K, Hayakawa Y, Sato T, Kinoshita H, et al. Apoptosis signal-regulating kinase 1 inhibits hepatocarcinogenesis by controlling the tumor-suppressing function of stress-activated mitogen-activated protein kinase. *Hepatology*. 2011;54:185–95.
25. Kang TW, Yevsa T, Woller N, Hoenicke L, Wuestefeld T, Dauch D, et al. Senescence surveillance of pre-malignant hepatocytes limits liver cancer development. *Nature*. 2011;479:547–51.
26. Garcia-Samaniego J, Rodriguez M, Berenguer J, Rodriguez-Rosado R, Carbo J, Asensi V, et al. Hepatocellular carcinoma in HIV-infected patients with chronic hepatitis C. *Am J Gastroenterol*. 2001;96:179–83.
27. Brau N, Fox RK, Xiao P, Marks K, Naqvi Z, Taylor LE, et al. Presentation and outcome of hepatocellular carcinoma in HIV-infected patients: a US—Canadian multicenter study. *J Hepatol*. 2007;47:527–37.
28. Maeda S. NF-kappaB, JNK, and TLR signaling pathways in hepatocarcinogenesis. *Gastroenterol Res Pract*. 2010;2010:367694.
29. Soresi M, Bonfissuto G, Sesti R, Riili A, Di Giovanni G, Carroccio A, et al. Perihepatic lymph nodes and antiviral response in chronic HCV-associated hepatitis. *Ultrasound Med Biol*. 2004;30:711–7.
30. Palmer C, Hampartzoumian T, Lloyd A, Zekry A. A novel role for adiponectin in regulating the immune responses in chronic hepatitis C virus infection. *Hepatology*. 2008;48:374–84.
31. Ohki T, Tateishi R, Shiina S, Goto E, Sato T, Nakagawa H, et al. Visceral fat accumulation is an independent risk factor for hepatocellular carcinoma recurrence after curative treatment in patients with suspected NASH. *Gut*. 2009;58:839–44.
32. Park EJ, Lee JH, Yu GY, He G, Ali SR, Holzer RG, et al. Dietary and genetic obesity promote liver inflammation and tumorigenesis by enhancing IL-6 and TNF expression. *Cell*. 2010;140:197–208.
33. Arano T, Nakagawa H, Tateishi R, Ikeda H, Uchino K, Enooku K, et al. Serum level of adiponectin and the risk of liver cancer development in chronic hepatitis C patients. *Int J Cancer*. 2011;129:2226–35.

Frequency, Risk Factors and Survival Associated with an Intrasubsegmental Recurrence after Radiofrequency Ablation for Hepatocellular Carcinoma

Ryosuke Tateishi^{1*}, Shuichiro Shiina¹, Masaaki Akahane², Jiro Sato², Yuji Kondo¹, Ryota Masuzaki¹, Hayato Nakagawa¹, Yoshinari Asaoka¹, Tadashi Goto¹, Kuni Otomo², Masao Omata³, Haruhiko Yoshida¹, Kazuhiko Koike¹

¹ Department of Gastroenterology, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan, ² Department of Radiology, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan, ³ Yamanashi Prefectural Hospital Organization, Tokyo, Japan

Abstract

Background: In the treatment of hepatocellular carcinoma (HCC), hepatic resection has the advantage over radiofrequency ablation (RFA) in terms of systematic removal of a hepatic segment.

Methods: We enrolled 303 consecutive patients of a single naïve HCC that had been treated by RFA at The University of Tokyo Hospital from 1999 to 2004. Recurrence was categorized as either intra- or extra-subsegmental as according to the Couinaud's segment of the original nodule. To assess the relationship between the subsegments of the original and recurrent nodules, we calculated the kappa coefficient. We assessed the risk factors for intra- and extra-subsegmental recurrence independently using univariate and multivariate Cox proportional hazard regression. We also assessed the impact of the mode of recurrence on the survival outcome.

Results: During the follow-up period, 201 patients in our cohort showed tumor recurrence distributed in a total of 340 subsegments. Recurrence was categorized as exclusively intra-subsegmental, exclusively extra-subsegmental, and simultaneously intra- and extra-subsegmental in 40 (20%), 110 (55%), and 51 (25%) patients, respectively. The kappa coefficient was measured at 0.135 (95% CI, 0.079–0.190; $P < 0.001$). Multivariate analysis revealed that of the tumor size, AFP value and platelet count were all risk factors for both intra- and extra-subsegmental recurrence. Of the patients in whom recurrent HCC was found to be exclusively intra-subsegmental, extra-subsegmental, and simultaneously intra- and extra-subsegmental, 37 (92.5%), 99 (90.8%) and 42 (82.3%), respectively, were treated using RFA. The survival outcomes after recurrence were similar between patients with an exclusively intra- or extra-subsegmental recurrence.

Conclusions: The effectiveness of systematic subsegmentectomy may be limited in the patients with both HCC and chronic liver disease who frequently undergo multi-focal tumor recurrence.

Citation: Tateishi R, Shiina S, Akahane M, Sato J, Kondo Y, et al. (2013) Frequency, Risk Factors and Survival Associated with an Intrasubsegmental Recurrence after Radiofrequency Ablation for Hepatocellular Carcinoma. PLoS ONE 8(4): e59040. doi:10.1371/journal.pone.0059040

Editor: Yujin Hoshida, Icahn School of Medicine at Mount Sinai, United States of America

Received: November 18, 2011; **Accepted:** February 12, 2013; **Published:** April 12, 2013

Copyright: © 2013 Tateishi et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by Health Sciences Research Grants of The Ministry of Health, Labour and Welfare of Japan (Research on Hepatitis). No additional external funding was received for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: tateishi-ky@umin.ac.jp

Introduction

Hepatic resection is regarded as the most appropriate first-line treatment for patients with solitary hepatocellular carcinoma (HCC) who are non-cirrhotic or cirrhotic without portal hypertension [1]. Hepatic resection is also indicated for HCC patients with more advanced cirrhosis in countries like Japan where the option of performing a liver transplantation is limited by the scarcity of cadaveric donor organs [2]. As a surgical procedure, anatomical resection, which is the systematic removal of a hepatic segment containing tumor tissue, is considered to be preferable based on the concept that tumor cells disseminate through the portal vein [3–8].

Percutaneous tumor ablation methods, such as ethanol injection and microwave coagulation, have played an important role as nonsurgical treatments that can achieve high local cure rates without reducing background liver function [9–12]. Radiofrequency ablation (RFA) is currently considered to be the most effective first-line percutaneous ablation protocol because of its greater efficacy in terms of local cure compared with ethanol injection [13–16]. The survival outcomes for patients who achieved a complete response by RFA are comparable to that among patients treated by hepatic resection [17–20].

Hepatic resection is supposed to have the advantage over RFA as an effective intervention as it involves the systematic removal of a hepatic segment containing the tumor. Indeed, microscopic

satellite nodules, not detected by radiological examination prior to resection, are often observed in the resected specimen [5,6,21]. However, this does not necessarily mean that microscopic lesions will have been confined to the resected segment. Indeed, even after anatomical resection, the cumulative recurrence rate at 5 years is as high as 50–70% [6–8], and it is not known to what extent anatomical resection can reduce HCC recurrence as compared with RFA.

Whereas RFA can reliably eliminate target nodules together with some of the surrounding tissue, most of the liver parenchyma of the tumor-bearing segment is left unablated. In contrast to anatomical resection, it is possible to observe and analyze intra- and extra-subsegmental recurrence by following up patients after ablation. The aim of our present study was to assess the frequency, risk factors and survival outcomes associated with intra-subsegmental HCC recurrence after RFA in comparison with extra-subsegmental recurrence.

Patients and Methods

Patients

This retrospective study was conducted according to the ethical guidelines for epidemiological research designed by the Japanese Ministry of Education, Culture, Sports, Science and Technology and Ministry of Health, Labour, and Welfare. The study design was included in a comprehensive protocol of retrospective study at the Department of Gastroenterology, The University of Tokyo Hospital approved by The University of Tokyo Medical Research Center Ethics Committee (approval number 2058). The following statements were posted at a website (<http://gastro.m.u-tokyo.ac.jp/med/0602A.htm>) and participants who do not agree to the use of their clinical data can claim deletion of them.

Department of Gastroenterology at The University of Tokyo Hospital contains data from our daily practice for the assessment of short-term (treatment success, immediate adverse events etc.) and long-term (late complications, recurrence etc.) outcomes. Obtained data were stored in an encrypted hard disk separated from outside of the hospital. When reporting analyzed data, we protect the anonymity of participants for the sake of privacy protection. If you do not wish the utilization of your data for the clinical study or have any question on the research content, please do not hesitate to make contact with us.

From 1999 to 2004 a total of 569 patients with HCC underwent RFA as the initial treatment for naïve HCC. Of them, 304 patients had a single nodule. We enrolled 303 of these patients in our current study excluding one patient who could not achieve complete ablation. The inclusion criteria for RFA had been as follows: a total bilirubin level of less than 3 mg/dL, a platelet count of no less than $50 \times 10^3/\text{mm}^3$ and prothrombin activity levels of no less than 50%. Patients with a portal vein tumor thrombosis, refractory ascites, or extrahepatic metastasis were excluded. In general, we performed RFA on patients with three or fewer lesions of 3 cm or less in diameter. However, we also performed ablation on patients beyond these criteria if it was predicted to be clinically effective [22,23]. We enrolled patients who underwent transcatheter arterial chemoembolization (TACE) prior RFA when the treatments were sequentially performed.

Diagnosis of HCC

HCC was diagnosed using dynamic computed tomography (CT), with a consideration of hyperattenuation in the arterial phase with washout in the late phase as a definite sign of this disease [24]. Most nodules were also confirmed histopathologically via an ultrasound-guided biopsy.

Treatment and evaluation

All patients received dynamic CT with a slice thickness of 5 mm within one month prior to ablation for comparison. The interval between the initiation of contrast material infusion and CT image recording was 30 and 120 sec for single detector-row spiral CT (Highspeed Advantage; GE Medical Systems; Milwaukee, WI) and 25, 40 and 120 sec for multidetector-row CT (LightSpeed QX/i GE Medical Systems). The images were presented after axial reconstruction with a slice thickness of 5 mm. RFA was performed on an in-patient basis using a cooled-tip electrode (Covidien, Mansfield, MA) under real-time ultrasound guidance. After 1 to 2 sessions of RFA, dynamic CT was performed to evaluate the treatment efficacy. During the treatment evaluation, we compared the CT findings for early and late phase before ablation and late phase after ablation. A lesion was judged to be completely ablated when the non-enhanced area shown in the late phase of CT post-ablation covered the entire lesion shown in both early and late phase of CT pre-ablation with a safety margin in the surrounding liver parenchyma. We confirmed complete ablation in all slices on which the target nodule was visualized. Patients received additional sessions until complete ablation was confirmed in each nodule. Finally, 303 of the 304 patients enrolled in this study were judged to be completely ablated.

Assessment of tumor recurrence

The follow-up regimen consisted of blood tests and monitoring of tumor markers in an outpatient setting. Ultrasonography and

Table 1. Baseline Characteristics of the HCC Patients analyzed in this study (n = 303).

Variable	n(%)
Age (y)	
mean \pm SD	67.5 \pm 8.2
Range	44–91
Male sex	191 (63.0)
Viral infection	
HBsAg positive only	28 (9.2)
anti HCVAb positive only	225 (74.3)
Both positive	5 (1.7)
Both negative	35 (11.6)
Alcohol consumption >80 g/day	43 (14.2)
Child-Pugh classification	
Class A	213 (70.3)
Class B	75 (24.8)
Class C	6 (2.0)
Size of tumor (cm)	
mean \pm SD	2.5 \pm 1.1
≤ 2.0	106 (35.0)
2.1–3.0	121 (40.0)
>3.0	76 (25.1)
AFP >100 ng/mL	68 (22.4)
DCP >100 mAU/mL	39 (12.9)
AFP-L3 >15%	44 (14.5)

AFP, alpha-fetoprotein; AFP-L3, lens culinaris agglutinin-reactive fraction of AFP; Anti-HCVAb, anti-hepatitis C virus antibody; DCP, des-gamma-carboxy prothrombin; HBsAg, hepatitis B surface antigen.

doi:10.1371/journal.pone.0061367.t001

dynamic CT were also performed every four months. Tumor recurrence was defined as a newly developed lesion on a dynamic CT that showed hyperattenuation in the arterial phase with washout in the late phase. The nomenclature used for the hepatic segments conformed to *The General Rules for the Clinical and Pathological Study of Primary Liver Cancer, Second English Edition* [25]. According to these rules, subsegments 1 to 8 correspond to Couinaud's segment 1 to 8, respectively [26]. All images were independently reviewed by two experienced radiologists (M.A. and J.S.), and a consensus reading was subsequently performed. Recurrence was categorized as either intra- or extra-subsegmental based on the subsegment of the original nodule. When a tumor was located on two or more subsegments, the subsegment where the major part of the tumor was present was adopted. Local tumor progression and neoplastic seeding through a needle tract were considered to be an intrasubsegmental recurrence. Extrahepatic recurrence was defined as extrasubsegmental.

Treatment of recurrent HCC and Survival Outcomes

When HCC recurrence was identified, patients who met the same criteria used for primary HCC underwent RFA. Survival analysis was performed on a per patient basis. Patients without an indication for RFA due to a multiplicity of recurrent nodules underwent TACE if liver function was categorized as Child-Pugh class B or better. Patients with localized portal tumor invasion were treated by radiotherapy [27]. Patients with tumor invasion to the first branch or main tract of the portal vein were treated with intra-arterial 5-fluorouracil and systemic interferon- α combination therapy [28]. Those with extrahepatic tumor metastasis received systemic chemotherapy if they had well-preserved liver function and a good performance status. Survival time was defined as the interval between the diagnosis of recurrence and the last visit to the outpatient clinic or death up to December 31, 2010. We also

analyzed overall survival after the initial RFA. For the analysis start date was set at the day when we perform the first RFA for each patient.

Statistical analysis

Data were expressed as the mean \pm standard deviation (SD) unless otherwise indicated. To assess whether the location of recurrent nodules was independent of the subsegment of the original nodule, we calculated the kappa coefficient and its 95% confidence interval (CI) [29]. A coefficient of 1 indicates that the subsegments of the original and recurrent nodules are identical, whereas a kappa coefficient of 0 indicates that tumor recurrence occurs completely at random. P values were also calculated on the null hypothesis of kappa equal to zero.

To assess the exclusively intra-subsegmental recurrence rate separately from all kinds of recurrence, we used cumulative incidence estimation with competing risk methods [30]. On this analysis, all types of recurrence were categorized as exclusively intra-subsegmental recurrence, exclusively extrasubsegmental recurrence, or simultaneously intra- and extra-subsegmental recurrence. The hazard function of each type of recurrence was estimated using kernel-based methods described by Muller and Wang [31].

We assessed the risk factors for intra- and extra-subsegmental recurrence independently using univariate and multivariate Cox proportional hazard regression. In assessing the risk factor for intra-subsegmental recurrence, patients with exclusively extra-subsegmental recurrence were treated as censored data and vice versa. The following factors were used for these analyses: age, gender, hepatitis B surface antigen positivity, hepatitis C antibody positivity, platelet count, alanine aminotransferase (ALT), tumor size, alpha-fetoprotein (AFP), des-gamma-carboxyprothrombin (DCP) and lens culinaris agglutinin-reactive fraction of AFP




Figure 1. Patient enrollment flow.
doi:10.1371/journal.pone.0059040.g001




Figure 2. Figure 2A: Overall recurrence. Figure 2B: Recurrence rates of according to the mode of recurrence.
doi:10.1371/journal.pone.0059040.g002

(AFP-L3). Factors showing statistical significance as a predictor in univariate analysis were further analyzed using a multivariate Cox proportional hazard regression model with stepwise selection of variables based on the Akaike information criterion (AIC).

We plotted survival curves according to the mode of recurrence (i.e., intra-, extra-subsegmental or both) using the Kaplan-Meier method. Statistical significance among these three groups was assessed using the log-rank test. We also calculated adjusted hazard ratios for survival according to the mode of recurrence using multivariate Cox proportional hazard regression with factors that showed statistical significance in a univariate analysis of survival. Differences with a *P* value of less than 0.05 were considered statistically significant. All statistical analyses were

performed with S-Plus Ver. 7 (TIBCO Software Inc., Palo Alto, CA) and R 2.13.0 (<http://www.R-project.org>).

Results

Patient profiles

The enrolled HCC patient cohort in this study consisted of 191 males and 112 females with a mean age of 67.5 years (Table 1). The mean tumor size was 2.5±1.1 cm in diameter. The number of the nodules distributed in subsegments 1 to 8 was 7 (2.3%), 12 (4.0%), 30 (9.9%), 43 (14.2%), 37 (12.2%), 32 (10.6%), 46 (15.2%), and 96 (31.7%), respectively. One hundred one patients underwent TACE before RFA. The median (range) interval between TACE and RFA was 23 (6–71) days.




Figure 3. The estimated hazard function over time according to the mode of recurrence.
doi:10.1371/journal.pone.0059040.g003

HCC recurrence

During the follow-up period (mean, 2.3 years; range 0.2 to 7.3 years), tumor recurrence in the HCC patient cohort was identified in 201 cases. The recurrent nodules were distributed in a total of 340 subsegments. Recurrent nodules were exclusively intra-subsegmental in 40 patients (20%), and exclusively extra-subsegmental in 110 patients (55%, Fig. 1). Simultaneous intra- and extra-subsegmental recurrence was observed in the remaining 51 patients (25%). The diagnosis of recurrence revealed that 104, 39, 22, 17 and 19 patients had 1, 2, 3, 4–5, and >5 tumors, respectively. Local tumor progression was identified in 10 patients, among which two individuals had simultaneous extra-subsegmental recurrent nodules. Two patients with extrahepatic recurrence (one lymph node and one left adrenal gland) were categorized as extra-subsegmental. Neoplastic seeding, which was categorized as intra-subsegmental recurrence, was observed as the first recurrence in two patients. Details of the distribution of original and recurrent nodules based on subsegments are listed in Table 2. The observed proportion of recurrent nodules in the same subsegment as the original nodule was 0.268, whereas the expected probability that the subsegments of original and recurrent nodules were the

same, assuming a random distribution, was 0.154. The kappa coefficient was calculated as 0.135 (95% CI, 0.079–0.190; $P < 0.001$). When patients with a local tumor progression or neoplastic seeding were excluded from this calculation, the kappa statistic decreased to 0.101 (95% CI, 0.046–0.156; $P < 0.001$). The cumulative rates of overall recurrence at 1, 3, and 5 years were 19.6%, 61.8%, and 78.3%, respectively (Fig. 2A). Cumulative rates of exclusively intra-subsegmental, exclusively extra-subsegmental and simultaneously intra- and extra-subsegmental recurrence were 3.4%, 8.1%, and 7.1% at 1 year, 12.7%, 32.7%, and 16.4% at 3 years, and 15.3%, 43.6%, and 19.4% at 5 years, respectively (Fig. 2B). The estimated hazard function curves according to the three types of recurrence showed a similar pattern over the first 4 years. Then only the hazard rate of exclusively extra-subsegmental recurrence increased whereas the hazard rate of the other two types of recurrence decreased (Fig. 3).

Risk factors related to intra- and extra-subsegmental recurrence

Univariate Cox proportional regressions revealed that the following factors were significantly associated with intra-subsegmental recurrence: tumor size, AFP, DCP, AFP-L3, platelet count and anti-HCV antibody positivity. The final model for predicting intra-subsegmental recurrence with stepwise variable selection included tumor size, AFP, platelet count and anti-HCV antibody positivity (Table 3). Factors related to extra-subsegmental recurrence that were found to be significant by univariate Cox proportional hazard regression were age, platelet count, tumor size, AFP and AFP-L3. Multivariate analysis with step-wise variable selection showed that the risk factors for extra-subsegmental recurrence were age, platelet count, tumor size, and AFP (Table 4).

Treatment of recurrent HCC and associated survival outcomes

Among the 40, 110 and 51 patients in whom recurrent HCC was found to be exclusively intra-subsegmental, exclusively extra-subsegmental, and simultaneously intra- and extra-subsegmental, 37 (92.5%), 99 (90.8%) and 42 (82.3%), respectively, were treated using RFA. Of the three patients with an exclusively intra-subsegmental recurrence, one individual was treated by hepatic resection and one patient was treated by TACE. The remaining patient received best supportive care because of deterioration in liver function. During the follow up period up to December 31,

Table 2. Distribution of the Original and Recurrent Tumors Divided by Subsegment.

Subsegment of original tumor, n	Subsegment of recurrent tumor, n								sum
	S1	S2	S3	S4	S5	S6	S7	S8	
S1	2		2	1			1	2	8
S2	1		1		4	1	1		8
S3		7	10	8	3	3	4	12	47
S4		8	7	11	3	3	6	6	44
S5		2	2	5	10	2	4	7	32
S6	2	1		4	4	10	5	5	31
S7	3	8	11	9	9	9	15	7	71
S8	4	9	9	10	6	13	15	33	99
sum	12	35	42	48	39	41	51	72	340

doi:10.1371/journal.pone.0061367.t002

Table 3. Univariate and Multivariate Analysis of Intrasubsegmental Recurrences (n = 303).

Variable	Univariate		Multivariate	
	HR (95% CI)	P	HR (95% CI)	P
Age per year	1.00 (0.97–1.02)	0.77		
Male gender	1.05 (0.68–1.63)	0.82		
HBsAg, positive	0.64 (0.30–1.40)	0.27		
anti-HCVAb, positive	2.08 (1.13–3.84)	0.02	2.04 (1.09–3.81)	0.03
Platelet count, $\times 10^9/\mu\text{L}$	0.95 (0.91–0.99)	0.009	0.97 (0.93–1.01)	0.09
ALT >80 IU/L	0.99 (0.58–1.71)	0.98		
Size per 1 cm	1.29 (1.08–1.55)	0.006	1.28 (1.06–1.54)	0.009
log(AFP)	1.93 (1.53–2.45)	<0.001	1.29 (1.16–1.44)	<0.001
log(DCP)	1.66 (1.18–2.33)	0.003		
AFP-L3>15%	2.02 (1.20–3.41)	0.009		

HR, hazard ratio; CI, confidence interval; HBsAg, hepatitis B surface antigen; Anti-HCVAb, anti-hepatitis C virus antibody; ALT, alanine aminotransferase, AFP, alpha-fetoprotein; DCP, des-gamma-carboxyprothrombin; AFP-L3, lens culinaris agglutinin-reactive fraction of AFP. doi:10.1371/journal.pone.0061367.t003

2010, 130 patients died and 9 patients were lost to follow-up. The median survival time (95% CI) was 5.72 (3.51-NA) years in patients with exclusively intra-subsegmental recurrence, 4.95 (4.19–5.76) years in patients with exclusively extra-subsegmental recurrence, and 2.43 (1.90–4.26) years in patients with simultaneously intra- and extra-subsegmental recurrence, respectively ($P < 0.001$ by log-rank test, Fig. 4). Univariate Cox regression analysis revealed that patients with simultaneously intra- and extra-subsegmental recurrences had a significantly poorer survival than those with an exclusively intra-subsegmental recurrence (hazard ratio, 2.39; 95% CI, 1.32–4.02; $P = 0.001$), whereas this difference became non-significant (HR, 1.91; 95% CI, 0.96–3.80; $P = 0.07$) when adjusted using other significant factors in univariate analysis (Table 5). No differences in the survival outcomes between patients with exclusively intra- and extra-subsegmental recurrences were observed by univariate and multivariate analysis. Finally overall survival rates after the initial RFA at 1, 3, 5, 7 and 10 years were 96.7%, 81.4%, 62.4%, 49.0%, and 31.1%, respectively.

Discussion

Recurrences of HCC are more complicated than those of other solid tumors as they can arise in two distinct forms: de novo carcinogenesis and intrahepatic metastasis [32]. Systematic subsegmentectomy may be effective in treating such patients if the distribution of the hematogenous spread of cancer cells correlates with the physical distance from the original tumor or local portal venous flow. Indeed, in the present study we showed from our data that the location of recurrent nodules was weakly but significantly related to that of the original tumor, even after the exclusion of local tumor progression from the analysis. Given that exclusively intrasubsegmental recurrence in this study could be prevented by subsegmentectomy, through a simple calculation, one fifth of patients who received locally curative RFA might have benefitted if they had received systematic subsegmentectomy. However, it should be mentioned in this regard that those patients who had avoided an intra-subsegmental recurrence owing to a systematic subsegmentectomy would have subsequently encoun-

Table 4. Univariate and Multivariate Analysis of Extrasubsegmental Recurrences (n = 303).

Variable	Univariate		Multivariate	
	HR (95% CI)	P	HR (95% CI)	P
Age per year	1.02 (1.00–1.04)	0.03	1.03 (1.01–1.05)	0.001
Male gender	1.13 (0.83–1.56)	0.44		
HBsAg, positive	0.91 (0.30–1.40)	0.69		
anti-HCVAb, positive	1.49 (1.00–2.20)	0.049		
Platelet count, $\times 10^9/\mu\text{L}$	0.94 (0.91–0.97)	<0.001	0.94 (0.92–0.97)	<0.001
ALT >80 IU/L	1.05 (0.72–1.56)	0.78		
Size per 1 cm	1.32 (1.16–1.51)	<0.001	1.39 (1.21–1.60)	<0.001
log(AFP)	1.53 (1.27–1.85)	<0.001	1.37 (1.12–1.68)	0.03
log(DCP)	1.29 (0.96–1.73)	0.1		
AFP-L3>15%	1.66 (1.09–2.52)	0.018		

HR, hazard ratio; CI, confidence interval; HBsAg, hepatitis B surface antigen; Anti-HCVAb, anti-hepatitis C virus antibody; AST, aspartate aminotransferase; ALT, alanine aminotransferase, AFP, alpha-fetoprotein; DCP, des-gamma-carboxy prothrombin; AFP-L3, lens culinaris agglutinin-reactive fraction of AFP. doi:10.1371/journal.pone.0061367.t004

tered tumor recurrence in the remnant liver, and the actual risk reduction of recurrence would therefore be smaller. Actually recurrence-free survival at 10 years after systematic subsegmentectomy was reported to be only 9.4% in a previous nation-wide survey [33].

The risk factors related to de novo carcinogenesis and hematogenous intrahepatic metastasis would be expected to be different. The factors responsible for HCC development, such as fibrosis stage, age, gender, and presence of viral hepatitis, may also affect de novo carcinogenesis [34,35]. On the other hand, factors related to the primary tumor, such as the size and number of tumor nodules, pathological grade(s), the presence of vascular invasion, and positivity of tumor markers, may affect the possibility of intrahepatic occult metastasis at the time of initial treatment. We speculated that there would be differences between the risk factors for intra- and extrasubsegmental recurrence since the former would more strongly correlate with hematogenous intrahepatic metastasis. However the risk factors related to intra- and extrasubsegmental recurrence were found to be quite similar except that old age was a risk factor for only extrasubsegmental recurrence.

Previous reports suggested the hazard function of de novo carcinogenesis and hematogenous intrahepatic metastasis would be different [36,37]. The hazard function of the former is assumed to be gradually increasing over time whereas that of the latter has a peak within two years. And the actual hazard function represents the sum of the two curves. The estimated hazard function of exclusively extra-subsegmental recurrence in this study seemed compatible with the previous reports. However we should be careful to interpret the results because the number at risk at year 4 or 5 was limited.

A previous large scale cohort study of the prognosis of patients with HCC treated by liver transplantation has reported that microvascular invasion is the most important predictor of a poor outcome [38]. This suggests that even if the whole liver is removed, there may be remaining circulating tumor cells that have resulted from tumor nodule invasion of the microvessels. It has also been reported that microsatellite metastatic nodules sur-




Figure 4. Cumulative survival probability after the diagnosis of recurrence according to the mode of recurrence.
doi:10.1371/journal.pone.0059040.g004

rounding the main tumor are associated with microvascular invasion and indicate a higher risk of tumor recurrence after liver transplantation [39]. Hence, the impact of removing a tumor-bearing subsegment, including microvascular invasions or micro-satellite nodules, which is thought to be a major advantage of resection over RFA, might be more limited than previously considered.

In this study factors that were supposed to be related to de novo carcinogenesis (e.g., lower platelet count and HCV infection) were risk factors for intra-subsegmental recurrence as well as extra-subsegmental recurrence. The risk of recurrence due to de novo carcinogenesis might be reduced by a subsegmentectomy according to the resected liver volume. However, as most patients with HCC have chronic liver disease, removal of non-cancerous liver

Table 5. Univariate and Multivariate Analysis of Survival after Recurrence*.

Variable	Univariate		Multivariate	
	HR (95% CI)	P	HR (95% CI)	P
Exclusively extra-subsegmental recurrence vs. exclusively intra-subsegmental recurrence	1.18 (0.73–1.92)	0.50	1.28 (0.76–2.16)	0.35
Simultaneously intra- and extra-subsegmental recurrence vs. exclusively intra-subsegmental recurrence	2.39 (1.32–4.02)	0.001	1.91 (0.96–3.80)	0.07
Age, per 1year	1.02 (1.00–1.05)	0.04	1.04 (1.02–1.07)	0.001
Male gender	1.06 (0.75–1.52)	0.73		
HBsAg, positive	0.74 (0.40–1.38)	0.34		
anti-HCVAb, positive	1.24 (0.77–1.99)	0.39		
Child-Pugh Score, per 1 point	1.45 (1.28–1.63)	<0.001	1.44 (1.27–1.63)	<0.001
Platelet count, per 10 ⁴ /μL	0.98 (0.95–1.01)	0.15		
ALT >80 IU/L	0.81 (0.49–1.33)	0.40		
Size >2cm	1.57 (1.11–2.23)	0.01	1.54 (1.06–2.23)	0.02
Multinodular	1.66 (1.18–2.35)	0.004	1.02 (0.63–1.66)	0.92
log(AFP)	1.45 (1.20–1.85)	<0.001	1.13 (1.01–1.26)	0.04
log(DCP)	1.61 (1.21–2.15)	0.001	1.21 (1.06–1.38)	0.004
AFP-L3 >15%	1.93 (1.25–2.98)	0.003	1.17 (0.70–1.96)	0.56

*Clinical data at the diagnosis of recurrence were adopted. HR, hazard ratio; CI, confidence interval; HBsAg, hepatitis B surface antigen; Anti-HCVAb, anti-hepatitis C virus antibody; AST, aspartate aminotransferase; ALT, alanine aminotransferase; AFP, alpha-fetoprotein; DCP, des-gamma-carboxy prothrombin; AFP-L3, lens culinaris agglutinin-reactive fraction of AFP.
doi:10.1371/journal.pone.0061367.t005

parenchyma may have a negative impact on long-term survival, especially for those individuals with impaired liver regeneration. Therefore, the key issue is to what extent the liver parenchyma should be removed to sufficiently treat the patient on a case by case basis. It may be speculated that extensive resection could be tolerable and beneficial to those who have a well-preserved capacity for liver regeneration [40].

There is no doubt that tumor recurrence deteriorates the long-term prognosis for HCC patients. However it is also true that there are effective, sometimes potentially curative treatments for recurrent HCC. The re-resection after recurrence of HCC is indicated in 10–30% of patients [41–43] and percutaneous ablation can be repeatedly performed [20,44,45]. Indeed, 37 of 40 patients analyzed in this study who had recurrent nodules confined to the same subsegment as the original tumor were successfully re-treated with RFA. No differences in the survival outcomes were observed between patients with solely intra- or extra-subsegmental recurrences. Hence, the impact of the first recurrence on overall survival may be smaller for HCC compared with other gastrointestinal malignancies such as stomach cancer or colorectal cancer.

References

- Bruix J, Sherman M (2005) Management of hepatocellular carcinoma. *Hepatology* 42: 1208–1236.
- Ikai I, Arii S, Kojiro M, Ichida T, Makuuchi M, et al. (2004) Reevaluation of prognostic factors for survival after liver resection in patients with hepatocellular carcinoma in a Japanese nationwide survey. *Cancer* 101: 796–802.
- Makuuchi M, Hasegawa H, Yamazaki S (1985) Ultrasonically guided subsegmentectomy. *Surg Gynecol Obstet* 161: 316–350.
- Castaing D, Garden OJ, Bismuth H (1989) Segmental liver resection using ultrasound-guided selective portal venous occlusion. *Ann Surg* 210: 20–23.
- Fuster J, Garcia-Valdecasas JC, Grande L, Tabet J, Bruix J, et al. (1996) Hepatocellular carcinoma and cirrhosis. Results of surgical treatment in a European series. *Ann Surg* 223: 297–302.
- Imamura H, Matsuyama Y, Miyagawa Y, Ishida K, Shimada R, et al. (1999) Prognostic significance of anatomical resection and des-gamma-carboxy prothrombin in patients with hepatocellular carcinoma. *Br J Surg* 86: 1032–1038.
- Regimbeau JM, Kiamanesh R, Farges O, Dondero F, Sauvanet A, et al. (2002) Extent of liver resection influences the outcome in patients with cirrhosis and small hepatocellular carcinoma. *Surgery* 131: 311–317.
- Hasegawa K, Kokudo N, Imamura H, Matsuyama Y, Aoki T, et al. (2005) Prognostic impact of anatomic resection for hepatocellular carcinoma. *Ann Surg* 242: 252–259.
- Ebara M, Ohto M, Sugiura N, Kita K, Yoshikawa M, et al. (1990) Percutaneous ethanol injection for the treatment of small hepatocellular carcinoma. Study of 95 patients. *J Gastroenterol Hepatol* 5: 616–626.
- Livraghi T, Giorgio A, Marin G, Salmi A, de Sio I, et al. (1995) Hepatocellular carcinoma and cirrhosis in 716 patients: long-term results of percutaneous ethanol injection. *Radiology* 197: 101–108.
- Shiina S, Tagawa K, Niwa Y, Umuma T, Komatsu Y, et al. (1993) Percutaneous ethanol injection therapy for hepatocellular carcinoma: results in 146 patients. *AJR Am J Roentgenol* 160: 1023–1028.
- Seki T, Wakabayashi M, Nakagawa T, Imamura M, Tamai T, et al. (1999) Percutaneous microwave coagulation therapy for patients with small hepatocellular carcinoma: comparison with percutaneous ethanol injection therapy. *Cancer* 85: 1691–1702.
- Rossi S, Di Stasi M, Buscarini E, Quaretti P, Garbagnati F, et al. (1996) Percutaneous RF interstitial thermal ablation in the treatment of hepatic cancer. *AJR Am J Roentgenol* 167: 759–768.
- Allgaier HP, Deibert P, Zuber I, Olschewski M, Blum HE (1999) Percutaneous radiofrequency interstitial thermal ablation of small hepatocellular carcinoma. *Lancet* 353: 1676–1677.
- Livraghi T, Goldberg SN, Lazzaroni S, Meloni F, Solbiati L, et al. (1999) Small hepatocellular carcinoma: treatment with radio-frequency ablation versus ethanol injection. *Radiology* 210: 655–661.
- Curley SA, Izzo F, Ellis LM, Nicolas Vauthey J, Vallone P (2000) Radiofrequency ablation of hepatocellular cancer in 110 patients with cirrhosis. *Ann Surg* 232: 381–391.
- Huang GT, Lee PH, Tsang YM, Lai MY, Yang PM, et al. (2005) Percutaneous ethanol injection versus surgical resection for the treatment of small hepatocellular carcinoma: a prospective study. *Ann Surg* 242: 36–42.
- Livraghi T, Meloni F, Di Stasi M, Rolle E, Solbiati L, et al. (2008) Sustained complete response and complications rates after radiofrequency ablation of very early hepatocellular carcinoma in cirrhosis: Is resection still the treatment of choice? *Hepatology* 47: 82–89.
- Chen MS, Li JQ, Zheng Y, Guo RP, Liang HH, et al. (2006) A prospective randomized trial comparing percutaneous local ablative therapy and partial hepatectomy for small hepatocellular carcinoma. *Ann Surg* 243: 321–328.
- N'Kontchou G, Mahamoudi A, Aout M, Ganne-Carrie N, Grando V, et al. (2009) Radiofrequency ablation of hepatocellular carcinoma: long-term results and prognostic factors in 235 Western patients with cirrhosis. *Hepatology* 50: 1475–1483.
- Nakashima Y, Nakashima O, Tamaka M, Okuda K, Nakashima M, et al. (2003) Portal vein invasion and intrahepatic micrometastasis in small hepatocellular carcinoma by gross type. *Hepatol Res* 26: 142–147.
- Tateishi R, Shiina S, Teratani T, Ohi S, Sato S, et al. (2005) Percutaneous radiofrequency ablation for hepatocellular carcinoma. An analysis of 1000 cases. *Cancer* 103: 1201–1209.
- Teratani T, Yoshida H, Shiina S, Ohi S, Sato S, et al. (2006) Radiofrequency ablation for hepatocellular carcinoma in so-called high-risk locations. *Hepatology* 43: 1101–1108.
- Torzilli G, Minagawa M, Takayama T, Inoue K, Hui AM, et al. (1999) Accurate preoperative evaluation of liver mass lesions without fine-needle biopsy. *Hepatology* 30: 889–893.
- Liver Cancer Study Group of Japan (2003) General Rules for the Clinical and Pathological Study of Primary Liver Cancer, Second English Edition. Tokyo: Kasahara.
- Couinaud C (1951) Lobes et segments hépatiques. Note sur l'architecture anatomique et chirurgicale du foie. *Presse Med*, pp. 709–711.
- Nakagawa K, Yamashita H, Shiraishi K, Nakamura N, Tago M, et al. (2005) Radiation therapy for portal venous invasion by hepatocellular carcinoma. *World J Gastroenterol* 11: 7237–7241.
- Ohi S, Yoshida H, Toume R, Umuma T, Kanda M, et al. (2006) Combination therapy of intraarterial 5-fluorouracil and systemic interferon-alpha for advanced hepatocellular carcinoma with portal venous invasion. *Cancer* 106: 1990–1997.
- Cohen J (1960) A Coefficient of Agreement for Nominal Scales. *Educational and Psychological Measurement* 20: 37–46.
- Gray R (1988) A class of K-sample tests for comparing the cumulative incidence of a competing risk. *Annals of Statistics* 16: 1141–1154.
- Muller HG, Wang JL (1994) Hazard rate estimation under random censoring with varying kernels and bandwidths. *Biometrics* 50: 61–76.
- Sakon M, Umeshita K, Nagano H, Eguchi H, Kishimoto S, et al. (2000) Clinical significance of hepatic resection in hepatocellular carcinoma: analysis by disease-free survival curves. *Arch Surg* 135: 1456–1459.
- Eguchi S, Kanematsu T, Arii S, Okazaki M, Okita K, et al. (2008) Comparison of the outcomes between an anatomical subsegmentectomy and a non-anatomical minor hepatectomy for single hepatocellular carcinomas based on a Japanese nationwide survey. *Surgery* 143: 469–475.
- Koike Y, Shiratori Y, Sato S, Ohi S, Teratani T, et al. (2000) Risk factors for recurring hepatocellular carcinoma differ according to infected hepatitis virus-an analysis of 236 consecutive patients with a single lesion. *Hepatology* 32: 1216–1223.
- Ikeda K, Saitoh S, Tsubota A, Arase Y, Chayama K, et al. (1993) Risk factors for tumor recurrence and prognosis after curative resection of hepatocellular carcinoma. *Cancer* 71: 19–25.

36. Imamura H, Matsuyama Y, Tanaka E, Ohkubo T, Hasegawa K, et al. (2003) Risk factors contributing to early and late phase intrahepatic recurrence of hepatocellular carcinoma after hepatectomy. *J Hepatol* 38: 200–207.
37. Mazzaferro V, Romito R, Schiavo M, Mariani L, Camerini T, et al. (2006) Prevention of hepatocellular carcinoma recurrence with alpha-interferon after liver resection in HCV cirrhosis. *Hepatology* 44: 1543–1551.
38. Mazzaferro V, Llovet JM, Miceli R, Bhoori S, Schiavo M, et al. (2009) Predicting survival after liver transplantation in patients with hepatocellular carcinoma beyond the Milan criteria: a retrospective, exploratory analysis. *Lancet Oncol* 10: 35–43.
39. Plessier A, Codes L, Consigny Y, Sommacale D, Dondero F, et al. (2004) Underestimation of the influence of satellite nodules as a risk factor for post-transplantation recurrence in patients with small hepatocellular carcinoma. *Liver Transpl* 10: S86–90.
40. Shi M, Guo RP, Lin XJ, Zhang YQ, Chen MS, et al. (2007) Partial hepatectomy with wide versus narrow resection margin for solitary hepatocellular carcinoma: a prospective randomized trial. *Ann Surg* 245: 36–43.
41. Nagasue N, Kohno H, Hayashi T, Uchida M, Ono T, et al. (1996) Repeat hepatectomy for recurrent hepatocellular carcinoma. *Br J Surg* 83: 127–131.
42. Poon RT, Fan ST, Lo CM, Liu CL, Wong J (1999) Intrahepatic recurrence after curative resection of hepatocellular carcinoma: long-term results of treatment and prognostic factors. *Ann Surg* 229: 216–222.
43. Nakajima Y, Ko S, Kanamura T, Nagao M, Kanchiro H, et al. (2001) Repeat liver resection for hepatocellular carcinoma. *J Am Coll Surg* 192: 339–344.
44. Yamashiki N, Tateishi R, Yoshida H, Shiina S, Teratani T, et al. (2005) Ablation therapy in containing extension of hepatocellular carcinoma: a simulative analysis of dropout from the waiting list for liver transplantation. *Liver Transpl* 11: 508–514.
45. Liang HH, Chen MS, Peng ZW, Zhang YJ, Zhang YQ, et al. (2008) Percutaneous radiofrequency ablation versus repeat hepatectomy for recurrent hepatocellular carcinoma: a retrospective study. *Ann Surg Oncol* 15: 3481–3493.

Identification of a Functional Variant in the *MICA* Promoter Which Regulates *MICA* Expression and Increases HCV-Related Hepatocellular Carcinoma Risk

Paulisally Hau Yi Lo¹, Yuji Urabe^{1,2}, Vinod Kumar¹, Chizu Tanikawa¹, Kazuhiko Koike³, Naoya Kato⁴, Daiki Miki^{2,5}, Kazuaki Chayama^{2,5}, Michiaki Kubo⁵, Yusuke Nakamura^{1,6}, Koichi Matsuda^{1*}

1 Laboratory of Molecular Medicine, Human Genome Center, Institute of Medical Science, The University of Tokyo, Tokyo, Japan, **2** Departments of Medical and Molecular Science, Division of Frontier Medical Science, Programs for Biomedical Research, Graduate School of Biomedical Sciences, Hiroshima University, Hiroshima, Japan, **3** Department of Gastroenterology, Graduate School of Medicine, University of Tokyo, Tokyo, Japan, **4** Unit of Disease Control Genome Medicine, The Institute of Medical Science, The University of Tokyo, Tokyo, Japan, **5** Center for Genomic Medicine, The Institute of Physical and Chemical Research (RIKEN), Kanagawa, Japan, **6** Departments of Medicine and Surgery, and Center for Personalized Therapeutics, The University of Chicago, Chicago, Illinois, United States of America

Abstract

Hepatitis C virus (HCV) infection is the major cause of hepatocellular carcinoma (HCC) in Japan. We previously identified the association of SNP rs2596542 in the 5' flanking region of the *MHC class I polypeptide-related sequence A (MICA)* gene with the risk of HCV-induced HCC. In the current study, we performed detailed functional analysis of 12 candidate SNPs in the promoter region and found that a SNP rs2596538 located at 2.8 kb upstream of the *MICA* gene affected the binding of a nuclear protein(s) to the genomic segment including this SNP. By electrophoretic mobility shift assay (EMSA) and chromatin immunoprecipitation (ChIP) assay, we identified that transcription factor Specificity Protein 1 (SP1) can bind to the protective G allele, but not to the risk A allele. In addition, reporter construct containing the G allele was found to exhibit higher transcriptional activity than that containing the A allele. Moreover, SNP rs2596538 showed stronger association with HCV-induced HCC ($P = 1.82 \times 10^{-5}$ and $OR = 1.34$) than the previously identified SNP rs2596542. We also found significantly higher serum level of soluble MICA (sMICA) in HCV-induced HCC patients carrying the G allele than those carrying the A allele ($P = 0.00616$). In summary, we have identified a functional SNP that is associated with the expression of *MICA* and the risk for HCV-induced HCC.

Citation: Lo PHY, Urabe Y, Kumar V, Tanikawa C, Koike K, et al. (2013) Identification of a Functional Variant in the *MICA* Promoter Which Regulates *MICA* Expression and Increases HCV-Related Hepatocellular Carcinoma Risk. *PLoS ONE* 8(4): e61279. doi:10.1371/journal.pone.0061279

Editor: Erica Villa, University of Modena & Reggio Emilia, Italy

Received: September 24, 2012; **Accepted:** March 11, 2013; **Published:** April 11, 2013

Copyright: © 2013 Lo et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by the Ministry of Education, Culture, Sports, Science and Technology of the Japanese government, the Ministry of Health, Labour, and Welfare of the Japanese government. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: koichima@ims.u-tokyo.ac.jp

Introduction

Hepatocellular carcinoma (HCC) is one of the common cancers in the world. It is well-known to be associated with the chronic infection of Hepatitis B (HBV) and Hepatitis C (HCV) viruses. In Japan, nearly 70% of HCC patients are infected with HCV [1]. The annual rate of developing HCC among patients with HCV-related liver cirrhosis in Japan is estimated to be about 4–8 percent [2]. Recent analyses have identified various genetic factors that are related with viral induced liver diseases [3–5]. In our previous two-stage genome-wide association study (GWAS) using a total number of 1,394 cases and 5,486 controls, a SNP rs2596542 located on chromosome 6p21.33 was shown to be significantly associated with HCV-induced HCC ($P = 4.21 \times 10^{-13}$ and $OR = 1.39$) [6]. This SNP is located within the class I major histocompatibility complex (MHC) region and is at about 4.8 kb upstream of *MHC class I polypeptide-related sequence A (MICA)* gene. We also identified that the risk A allele of SNP rs2596542 was strongly associated with the low expression of soluble MICA (sMICA) in the serum of HCV-related HCC patients [6].

MICA is a membrane protein which is up-regulated in various tumor cells and also induced in response to various cellular stresses such as infection, hypoxia, and heat shock [7]. It is an important component of the innate immune response, as *MICA* can bind to the NKG2D receptor and subsequently activate natural killer (NK) cells, CD8+ cells, and $\gamma\delta$ T cells [8,9]. Moreover, membrane *MICA* can be shed by metalloproteinases, including MMP9, ADAM10, and ADAM17, and secreted into serum as a soluble form [10,11]. Since these metalloproteinases are often activated in HCC, the expressions of both membrane-bound *MICA* and sMICA are increased [12,13]. SNP rs2596542 was found to be associated with the progression from chronic hepatitis C (CHC) to HCC and also with serum sMICA level. Hence, both rs2596542 and sMICA would be possible prognostic biomarkers for CHC patients. However, their underlying molecular mechanisms were not fully elucidated so far.

We hypothesize that *MICA* variations could affect sMICA level by either one or both of the following two possible mechanisms: (1) the genetic variation(s) in the coding region affecting the protein stability and (2) the transcriptional regulation. Previously, variable

numbers of tandem repeats (VNTRs) in exon 5 of *MICA* were identified to affect *MICA* subcellular localization and serum *MICA* level [14]. The exon 5 of *MICA* encodes the transmembrane domain and the insertion of an extra G nucleotide in the domain would result in a premature stop codon that would generate *MICA* protein without a transmembrane domain and subsequently affect s*MICA* level [14]. However, our previous results indicated that *MICA* VNTR was not significantly associated with the s*MICA* level or HCC risk [6]. Therefore, in the current study, we have tried to investigate whether the *MICA* variations would affect the *MICA* transcription in the liver cancer cells. Through the functional analysis of genetic variations in the *MICA* promoter region, we here report a causative SNP rs2596538 that increases the binding affinity of the transcription factor Specificity Protein 1 (SP1) and the risk of progression of the disease.

Materials and Methods

Samples and genotyping

DNA samples for direct sequencing (50 HCV-related HCC cases), imputation analysis (721 HCV-related HCC cases and 5,486 HCV-negative controls), and serum samples for s*MICA* ELISA (246 HCV-related HCC) were obtained from BioBank Japan [15,16]. Genotyping of SNPs from 1,394 HCC patients and measurement of s*MICA* expression by ELISA were performed in the previous study [6]. Genotyping of SNP rs2596542 in 1,043 CHC was performed previously in RIKEN using Illumina HumanHap610-Quad BeadChip [17]. All CHC subjects had abnormal levels of serum alanine transaminase for more than 6 months and were positive for both HCV antibody and serum HCV RNA. The SNP rs2596542 in liver cirrhosis samples without hepatocellular carcinoma from BioBank Japan ($n = 420$) and the University of Tokyo ($n = 166$) were genotyped using Illumina HumanHap610-Quad BeadChip or invader assay [18]. All subjects were either subjected to liver biopsy or diagnosed by non-invasive methods including hepatic imaging, biochemical data, and the presence/absence of clinical manifestations of portal hypertension [18]. The samples used in the current project were listed in Table S1. Case samples with HBV co-infection were excluded from this study. The subjects with cancers, chronic hepatitis B, diabetes or tuberculosis were excluded from non-HCV controls. All subjects were Japanese origin and provided written informed consent. This research project was approved by the ethical committees of the University of Tokyo and RIKEN.

Imputation study

The imputation study was performed by using a hidden Markov model programmed in MACH [19] and haplotype information from 1000 genomes database [20]. The imputation results were confirmed by direct DNA sequencing in 50 randomly selected samples.

Cell culture

Human liver cancer cell lines HLE and HepG2 were purchased from JHSF (Osaka, Japan) and ATCC. These cells were grown in Dulbecco's modified Eagle's medium (Invitrogen) with 10% fetal bovine serum. Cells were cultured at 37°C with 5% CO₂.

EMSA

HLE cells were grown in 15 cm culture plate until they reached 95% confluency. The plate was then sealed with parafilm and immersed in a water bath at 42.5°C for 1.5 hours [21]. Nuclear extracts from these cells were prepared according to the standard

protocol [22]. EMSA was carried out using DIG Gel Shift Kit, 2nd Generation (Roche) according to the manufacturer's instructions. The sequences of the 12 probes were listed in the Table S2. In brief, 30 fmol of labeled probes were hybridized with 5 µg nuclear extract for 15 minutes at room temperature. The mixtures were then loaded into a 6% TBE gel, separated by electrophoresis at 4°C and transferred onto a nylon membrane. The membrane was then hybridized with anti-digoxigenin-AP antibody and developed by CSPD solution. For competition study, nuclear extracts were incubated with non-labeled oligonucleotides first before adding labeled probe. For supershift assay, SP1 antibody (SC-59X, Santa Cruz Biotechnology) was added into the nuclear extract and incubated on ice for 30 minutes first before adding labeled probe. The mixtures were then separated by electrophoresis using 4% TBE gel. All EMSAs were repeated twice for reconfirmation of the results.

ChIP

The HLE cells (G allele homozygote) and HepG2 cells (heterozygote) were used in the ChIP assay. The plasmid pCAGGS-SP1 was transfected into both cells by using FuGENE6 Transfection Reagent (Roche). The ChIP assays were carried out using Chromatin Immunoprecipitation Assay Kit (Millipore) according to the manufacturer's protocol. In brief, the cells were treated with formaldehyde to crosslink DNA-protein complexes at 48 hours post-transfection. DNA-protein complexes were then sheared by sonication and immunoprecipitated by rabbit polyclonal anti-SP1 antibody (SC-59X, Santa Cruz Biotechnology). The resulting DNAs were analyzed by PCR (Table S2). In order to determine the binding specificity of SP1 to the SNP rs2596538 allele, the PCR products from HepG2 cells were further subcloned into pCR 2.1 vector and sequenced to assess G to A ratio in both input DNA and immunoprecipitant.

Dual luciferase reporter assay

Three copies of 31 bp DNA fragments equivalent to the EMSA oligonucleotides of SNP rs2596538 were cloned into pGL3-promoter vector (Promega). The plasmids were co-transfected with pCAGGS-SP1 and pRL-TK plasmids (Promega) into HLE cells by FuGENE6 Transfection Reagent (Roche). The pCAGGS-SP1 plasmid provided the expression of transcription factor SP1, and pRL-TK plasmid served as internal control for transfection efficiency [23]. The cells were lysed at 48 hours post-transfection, and relative luciferase activities were measured by Dual Luciferase Assay System (Toyo B-Net).

Western blotting

Cancer cell lysates were prepared by using pre-chilled RIPA buffer, and 25 µg of each lysate was loaded into the gel and separated by SDS-PAGE. Western blotting was performed according to the standard protocol. Rabbit anti-*MICA* antibody (ab63709, abcam: 1/1000) and rabbit anti-SP1 antibody (17-601, Upstate Biotechnology: 1/500) were used in the experiment.

Statistical analysis

The case-control association was analyzed by Student's *t*-test and Fisher's exact test as appropriate. The association of allele dependent s*MICA* expression was studied by Kruskal-Wallis test using R statistical environment version 2.8.1. The LD and coefficients (D' and r^2) were calculated by Haploview version 4.2 [24].

Table 1. Association of rs2596542 with the progression from CHC to LC and HCC.

	Case MAF	Control MAF	P	OR	95% C.I.
LC vs CHC	0.3797	0.3442	0.04842	1.166	1.01–1.35
HCC vs LC	0.4012	0.3797	0.20296	1.094	0.95–1.26

MAF, minor allele frequency; OR, odds ratio for minor allele. C.I., confidence interval. SNP rs2596542 was analyzed in 1,043 chronic hepatitis C (CHC), 586 liver cirrhosis without hepatocellular carcinoma (LC) and 1,394 HCV-induced hepatocellular carcinoma patients (HCC). Calculated by Armitage trend test. doi:10.1371/journal.pone.0061279.t001

Results

Analyses of SNP rs2596542 in HCV-infected patients at different disease stages

Since the development of HCC consists of multiple steps, we investigated the role of SNP rs2596542 with disease progression. SNP rs2596542 was genotyped in patients at three different disease categories of CHC (chronic hepatitis C) without liver cirrhosis (LC) or HCC, LC without HCC, and HCC. The statistical analysis indicated that SNP rs2596542 was significantly associated with disease progression from CHC to LC with P-value of 0.048 and odds ratio of 1.17 (Table 1). The risk allele frequency among HCC patients (40.1%) was higher than that among LC patients (38.0%), but the association was not statistically significant (P-value of 0.203 and odds ratio of 1.09). These results suggested the involvement of *MICA* with both liver fibrosis and hepatocellular carcinogenesis.

HCV-HCC risk is not associated with *MICA* copy number variation

A previous report has indicated the deletion of the entire *MICA* locus in 3.2% of Japanese population [25] and this deletion was shown to be associated with the risk of nasopharyngeal carcinoma (NPC), especially in male [26]. To identify the functional SNP that may affect *MICA* mRNA expression, we analyzed the relation between the *MICA* copy number variation (CNV) and the HCC

susceptibility. We quantified this CNV by real-time PCR in 375 HCV-related HCC patients and 350 HCV-negative controls. As shown in Table S3, we found no difference in the copy numbers between HCC cases and controls, indicating that this CNV is unlikely to be causative genetic variation for the risk of HCC.

Direct sequencing of 5' flanking region of *MICA*

We then focused on the variations in the 5' flanking region of the *MICA* gene which may be associated with its promoter activity. We had conducted direct DNA sequencing of the 5-kb promoter region which included the marker SNP rs2596542 using genomic DNAs of 50 HCC subjects and identified 11 SNPs showing strong linkage disequilibrium with the marker SNP rs2596542 ($D' > 0.953$ and $r^2 > 0.832$) (Fig. S1, Table 2).

Allele specific binding of nuclear protein to genomic region including SNP rs2596538

To investigate whether these genetic variations would affect the binding affinity of some transcription factors, we had conducted the electrophoretic mobility shift assay (EMSA) using the nuclear extract of HLE human hepatocellular carcinoma cells. Since *MICA* is a stress-inducible protein [21], we first treated the cells with heat shock treatment at 42°C for 90 minutes and confirmed significant induction of *MICA* expression as shown in Fig. 1a. Then we performed EMSA using 24 labeled-oligonucleotides corresponding to each allele of the 12 candidates' SNPs. The results of EMSA demonstrated that an oligonucleotide corresponding to a G allele of SNP rs2596538 exhibited stronger binding affinity to a nuclear protein(s) than that to an A allele (Fig. 1b). We then confirmed the specific binding of nuclear proteins to the G allele by competitor assay using non-labeled oligonucleotides (Fig. 1c). The self (G allele) oligonucleotides inhibited the formation of DNA-protein complex in a dose-dependent manner, but the non-self (A allele) oligonucleotides showed no inhibition effect. Taken together, some nuclear protein(s) in hepatocellular carcinoma cells would interact with a DNA fragment including the G allele of SNP rs2596538.

Table 2. Linkage disequilibrium between 11 candidate SNPs and SNP rs2596542.

SNP ID	Relative position ^a	A1	A1 frequency	D'	r ²
rs2596542	-4815	A	0.36		
rs2428475	-4788	G	0.36	1	1
rs28366144	-4586	T	0.36	1	1
rs2428474	-4387	G	0.39	1	0.88
rs2251731	-4045	A	0.39	1	0.88
rs2844526	-3703	C	0.38	1	0.918
rs2596541	-3572	A	0.38	1	0.918
rs2523453	-3285	G	0.38	1	0.918
rs2544525	-3259	C	0.38	1	0.918
rs2523452	-2870	G	0.34	0.953	0.832
rs2596538	-2778	A	0.34	0.953	0.832
rs2844522	-2710	C	0.34	0.953	0.832

Note: Direct DNA sequence of 5-kb promoter region of *MICA* from 50 HCV-HCC subjects. D' and r² were calculated by comparing the genotypes of these SNPs to the marker SNP rs2596542 by Haploview. A1, minor allele; ^aRelative position to exon 1 of the *MICA* gene. doi:10.1371/journal.pone.0061279.t002