

Autoimmune Disorders Associated with Gain of Function of the Intracellular Sensor MDA5

Masahide Funabiki,^{1,2,11} Hiroki Kato,^{1,3,4,11} Yoshiaki Miyachi,² Hideaki Toki,⁵ Hiromi Motegi,⁵ Maki Inoue,⁵ Osamu Minowa,⁵ Aiko Yoshida,⁶ Katashi Deguchi,⁶ Hiroshi Sato,⁷ Sadayoshi Ito,⁸ Toshihiko Shiroishi,⁹ Kunio Takeyasu,⁶ Tetsuo Noda,^{5,10} and Takashi Fujita^{1,3,*}

¹Laboratory of Molecular Genetics, Institute for Virus Research

²Department of Dermatology, Graduate School of Medicine

³Laboratory of Molecular Cell Biology, Graduate School of Biostudies

Kyoto University, 53 Shogoin-Kawaharacho, Sakyo-ku, Kyoto City, Kyoto 606-8507, Japan

⁴PRESTO, Science and Technology Agency (JST), Kawaguchi, Saitama 332-0012, Japan

⁵Team for Advanced Development and Evaluation of Human Disease Models, Riken BioResource Center (BRC), 3-1-1 Koyadai, Tsukuba, Ibaraki 305-0074, Japan

⁶Laboratory of Plasma Membrane and Nuclear Signaling, Graduate School of Biostudies, Kyoto University, Yoshida-konoe-cho, Sakyo-ku, Kyoto 606-8501, Japan

⁷Department of Clinical Pharmacology and Therapeutics, Tohoku University Graduate School of Pharmaceutical Sciences and Faculty of Pharmaceutical Sciences, 6-3 Aramaki-aza-aoba, Aoba-ku, Sendai 980-8578, Japan

⁸Division of Nephrology, Endocrinology and Vascular Medicine, Tohoku University Graduate School of Medicine, 1-1 Seiryō-cho, Aoba-ku, Sendai 980-8574, Japan

⁹Division of Mammalian Genetics Laboratory, National Institute of Genetics, Yata 1111, Mishima, Shizuoka 411-8540, Japan

¹⁰Department of Cell Biology, Cancer Institute, The Japanese Foundation for Cancer Research, 3-8-31, Ariake, Koto, Tokyo 135-8550, Japan

¹¹These authors contributed equally to this work

*Correspondence: tfujita@virus.kyoto-u.ac.jp

<http://dx.doi.org/10.1016/j.immuni.2013.12.014>

SUMMARY

MDA5 is an essential intracellular sensor for several viruses, including picornaviruses, and elicits antiviral interferon (IFN) responses by recognizing viral dsRNAs. MDA5 has been implicated in autoimmunity. However, the mechanisms of how MDA5 contributes to autoimmunity remain unclear. Here we provide direct evidence that dysregulation of MDA5 caused autoimmune disorders. We established a mutant mouse line bearing MDA5 mutation by ENU mutagenesis, which spontaneously developed lupus-like autoimmune symptoms without viral infection. Inflammation was dependent on an adaptor molecule, MAVS indicating the importance of MDA5-signaling. In addition, intercrossing the mutant mice with type I IFN receptor-deficient mice ameliorated clinical manifestations. This MDA5 mutant could activate signaling in the absence of its ligand but was paradoxically defective for ligand- and virus-induced signaling, suggesting that the mutation induces a conformational change in MDA5. These findings provide insight into the association between disorders of the innate immune system and autoimmunity.


INTRODUCTION

Innate immune responses including induction of type I interferons (IFNs) are initiated by receptors, which sense the invasion

of microorganisms such as viruses and facilitate eradication of pathogens. Several transmembrane protein members, Toll-like receptors (TLRs), have been demonstrated to play an important role in antiviral responses by recognizing pathogen-specific molecular patterns (PAMPs) derived from viral species (Beutler, 2009; Imler and Hoffmann, 2001; Kawai and Akira, 2011) at the cell surface or in endosomes. In the cytoplasm, it has been shown that viral RNAs are sensed by RIG-I-like receptors (RLRs), which consist of three family member molecules, RIG-I, MDA5, and LGP2 (Yoneyama and Fujita, 2009). Cytosolic DNA also activates DNA sensors such as DDX41, IFI16, and cGAS to produce type I IFN (Sun et al., 2013; Unterholzner et al., 2010; Zhang et al., 2011).

RLRs are encoded by IFN-inducible genes and are expressed in various types of cells, including mouse embryonic fibroblasts (MEFs), dendritic cells (DCs), and macrophages. In conventional DCs (cDCs), RLRs play an essential role in antiviral defense, whereas TLRs play an essential role in plasmacytoid DCs (pDCs) for antiviral responses (Kato et al., 2005). These sensors discriminate non-self and self-nucleic acids by their molecular structure in a sequence-independent manner. For instance, RIG-I is activated by double-stranded RNA (dsRNA) and is also known to recognize RNAs with a triphosphate moiety at 5' end and/or polypyrimidine (Hornung et al., 2006; Pichlmair et al., 2006; Schlee et al., 2009; Takahashi et al., 2008). MDA5 also detects dsRNA, such as poly(I:C), especially longer than 2 kbp in length (Kato et al., 2008).

Type I IFNs produced by the innate immune response, so called IFN responses, are the first line and major source of host antiviral defense. However, there is also an implication that excessive production of type I IFNs could rather cause a



(legend on next page)

Immunity

MDA5 Mutant Mice Cause Spontaneous Nephritis

predisposition to autoimmunity (Trinchieri, 2010). There is also a murine model in which nucleic acid accumulates within cells, leading to excessive production of type I IFNs to cause autoimmunity (Yoshida et al., 2005). Some studies revealed the involvement of TLRs in autoimmunity. For example, TLR7 and TLR9 are important for autoantibody production in a murine model of lupus (Christensen et al., 2006), and spontaneous gene duplication of murine TLR7 causes a predisposition to autoimmunity (Pisitkun et al., 2006).

MDA5 (encoded by *IFIH1* in human) is associated with specific autoimmune diseases. A genome-wide association study (GWAS) revealed that SNPs of *IFIH1* (E627* and I923V), which are supposed to be loss-of-function mutations (Shigemoto et al., 2009), are significantly associated with resistance to type 1 diabetes (Nejentsev et al., 2009). In amyopathic dermatomyositis patients, anti-CADM140 antibodies, which could potentially target MDA5, were detected in patients' sera (Nakashima et al., 2010). These findings imply a link between MDA5 and autoimmune diseases; however, the link remains to be determined and little is known about the specific mechanisms that lead to disease.

Here, we investigated the association of MDA5 with autoimmunity in a mouse model. We found that mutant mice, harboring a single missense mutation G821S in MDA5 generated by N-ethyl-N-nitrosourea (ENU) mutagenesis, spontaneously developed lupus-like nephritis and systemic autoimmune symptoms without viral infection. Here we provide direct evidence that dysregulation of an innate immune sensor, MDA5, causes autoimmune disorders.

RESULTS

ENU Mutagenesis and Identification of Responsible Gene

ENU mutagenesis and phenotype screenings were carried out as previously reported (Toki et al., 2013). Briefly, male C57BL/6JJcl (B6) mice were treated with ENU and then mated with DBA/2JJcl (D2) females. In the process of the dominant phenotype screenings in the F1 progeny, we detected phenodeviants showing lupus-like nephritis. The inheritance of this phenotype was confirmed in N2 progeny backcrossed to D2, and the phenotype was mapped to 1.04 megabase (Mb) region in chromosome 5 (see Figure S1A available online). We conducted genomic DNA sequencing of all exons and splice sites of all genes in this region. In consequence, one single-base substitution in exon 13 of *Ifih1*, resulting in amino acid exchange G821S (gs), was identified (Figure S1B). Thus, we concluded that this

mutation was responsible for the phenotype. The F1 founder was backcrossed five generations to D2, and N6 progeny were subjected to a series of analyses.

Ifih1 gs Mutant Mice Develop Spontaneous Inflammation in Multiorgans


Major organs in mutant mice were analyzed histologically. Among several clinical manifestations, we found that nephritis was significant and reproducible in all heterozygous (*Ifih1*^{gs/+}) descendants, developing from 6 to 8 weeks after birth (Figures 1A and S1C). In addition to nephritis, most *Ifih1*^{gs/+} mice with a DBA/2 background developed severe multiorgan autoimmune disease. They showed growth retardation (Figure 1B), and 60% of the mutant died within 24 weeks after birth (Figure 1C). There was mild infiltration of polymorphonuclear leukocytes in the haired skin of the forehead (Figure S1D). We also found calcification in the liver, which indicates chronic inflammation (Figure S1D). Of note, homozygotes developed more severe inflammation and rarely survived for 4 weeks after birth (data not shown). In the affected kidney, deposition of immunoglobulin and complement was observed (Figure 1D). Anti-nuclear antibodies (ANA) and anti-dsDNA antibodies were detected in the sera of *Ifih1*^{gs/+} mice (Figure 1E). *Ifih1*^{gs/+} mice developed hyperimmunoglobulinemia of all tested immunoglobulin classes (Figure 1F). These pathological and serological features were consistent with lupus-like nephritis. In the kidney, production of inflammatory cytokines and chemokines including IFN- β , interleukin-6 (IL-6), CXCL10, Isg56, and tumor necrosis factor- α (TNF- α) was significantly upregulated (Figure 1G). Upregulation of IFN- β and IL-6 in systemic organs was observed (Figure S1E), suggesting that chronic inflammation occurred in multiple organs, including the kidney of *Ifih1*^{gs/+}, although nephritis was especially evident. Consistent with the upregulation of IFN in multiorgans, increased mRNA of RLRs, as well as protein was detected (Figures S1F and S1G). In summary, the mice exhibit continued upregulation of IFN and IFN-stimulated genes in multiple organs.

Intrinsically Activated DCs and Macrophages Are Possible Sources of Chronic Inflammation

MDA5 protein is ubiquitously expressed in multiorgans and various types of cells, including nonimmune cells such as fibroblasts and immune cells including DCs and macrophages. Abnormal function of innate immune cells is considered a predisposing factor for systemic lupus erythematosus (SLE) because they produce robust type I IFNs and activate adaptive immune system to trigger autoimmunity (Liu and Davidson, 2012). To

Figure 1. MDA5 G821S Mutant Mice Spontaneously Develop Lupus-like Autoimmune Disorder

- (A) Histology of kidney from *Ifih1*^{+/+} and *Ifih1*^{gs/+} mice (left panels, Haematoxylin-eosin staining; right panels, PAS).
(B) Body weight change of *Ifih1*^{+/+} and *Ifih1*^{gs/+} male and female mice following time course. *Ifih1*^{+/+} male, n = 14; *Ifih1*^{+/+} female, n = 15; *Ifih1*^{gs/+} male, n = 12; *Ifih1*^{gs/+} female, n = 13.
(C) Survival curve of *Ifih1*^{+/+} (n = 30) and *Ifih1*^{gs/+} (n = 28) mice in DBA/2JJcl background (upper) and of male (n = 12) and female (n = 16) *Ifih1*^{gs/+} mice (lower). Statistical significance was determined by Log rank test. *p < 0.03, NS; not significant.
(D) Immunoglobulin and complement 3(C3) depositions on *Ifih1*^{gs/+} kidney.
(E) Production of anti-nuclear antibodies (ANA) and anti-double-stranded DNA (anti-dsDNA) antibodies in *Ifih1*^{gs/+} sera.
(F) Hypergammaglobulinemia in *Ifih1*^{gs/+} mice. Serum immunoglobulin levels are shown. Statistical significance in (E) and (F) was determined by Student's t test. *p < 0.03.
(G) Amounts of mRNA in kidney for indicated genes were quantified by quantitative real-time PCR. Mean relative expression against littermate WT kidney is shown with standard as mean \pm SD of duplicate samples of a representative from three independent experiments. Open histogram, *Ifih1*^{+/+}; closed, *Ifih1*^{gs/+} kidney.



(legend on next page)

Immunity

MDA5 Mutant Mice Cause Spontaneous Nephritis

determine which cell types are responsible for systemic inflammation, we analyzed spleen cells by flow cytometry. *Ifih1*^{gs/+} mice, both cDCs (CD11c^{hi}B220^{lo}) and pDCs (CD11c^{int}B220^{hi}), and macrophages (CD11b^{hi}F4/80^{int}) were increased in numbers and their cell surface marker, CD86, was upregulated (Figures 2A and 2B), suggesting that these DCs and macrophages are activated. Next, we sorted these cells and examined the induced amount of type I IFN and Cxcl10 (Figure 2C), as well as IL-6 (Figure S2A). As we expected, marked upregulation of these genes was detected in DCs and macrophages, whereas no upregulation was observed in T and B cells. Of note, when cDCs and macrophages were differentiated from bone marrow cells in the presence of GM-CSF, IFN induction and slight upregulation of CD86 were clearly observed (Figure 2D; data not shown), suggesting that DCs and macrophages in *Ifih1*^{gs/+} mice are activated intrinsically, as well as through a bystander activation. Next we examined other cell types, including T cells and B cells. The ratio of T cells and B cells was normal in *Ifih1*^{gs/+} splenocytes (Figure S2B). Effector (CD44^{hi}CD62L^{lo}) and memory (CD44^{lo}CD62L^{hi}) T cells were increased, whereas naive T (CD44^{lo}CD62L^{hi}) cells were decreased in *Ifih1*^{gs/+} mice (Figure S2C). We detected upregulation of an activation marker CD69 in CD4⁺ and CD8⁺ T cells. Although no significant activation was observed in the B cell population with the cell surface markers, GL7 and CD69 (Figure S2D), plasma cell population was slightly increased and absolute cell number was higher in *Ifih1*^{gs/+} mice (Figure 2E), which could lead to augmentation of antibody production (Figures 1E and 1F). Consistent with systemic inflammation, granulocytes were significantly increased in *Ifih1*^{gs/+} mice (Figures S2E and S2F). Thus, we identified that DCs and macrophages were primary source of type I IFN and inflammatory cytokines, which could induce effector T cells and plasma cells, consequently leading to antibody production.

Hematopoietic Cells Are Involved in Nephritis

To access the contribution of hematopoietic cells to the chronic inflammation, we transferred bone marrow cells by reconstituting irradiated DBA/2 wild-type (WT) mice with *Ifih1*^{+/+} or *Ifih1*^{gs/+} bone marrow cells (Figure 3A). The mice with transferred bone marrow cells from *Ifih1*^{gs/+} displayed nephritis (Figure 3B) and immunoglobulin deposition in the kidney (Figure 3C) beginning 6 weeks after reconstitution. In the kidney, significant induction of inflammatory cytokines was detected (Figure 3D). In the spleen, upregulation of CD86 in both cDCs and pDCs was observed (Figure 3E). These results indicate that hematopoietic cells are sufficient for MDA5 mutation-dependent nephritis. Of note, those recipient mice exhibited a relatively milder phenotype than *Ifih1*^{gs/+} mice, implicating that not only hematopoietic

cells but also nonhematopoietic cells could possibly be involved in MDA5 mutation-dependent nephritis.

Critical Role of MAVS in the Development of Nephritis in *Ifih1*^{gs/+} Mice

MAVS is the key adaptor molecule for the RLR pathway to transduce signaling downstream (Kawai et al., 2005; Meylan et al., 2005; Seth et al., 2005; Xu et al., 2005). To clarify the involvement of G821S MDA5 mutation in chronic inflammation, we intercrossed *Ifih1*^{gs/+} mice with *Mavs*^{-/-} mice with a C57BL/6 background. *Ifih1*^{gs/+}*Mavs*^{+/-} mice developed nephritis with a less severe clinical score than *Ifih1*^{gs/+}*Mavs*^{+/+} mice (Figures 4A and 4B). In contrast, *Ifih1*^{gs/+}*Mavs*^{-/-} mice were devoid of nephritis and remained healthy. Moreover, production of type I IFNs and inflammatory cytokines in the kidney was revealed to be dependent on the gene dosage of *Mavs*; notably, virtually no production of these cytokines was detected in the absence of MAVS (Figure 4C). Consistent with our observation, virus-induced type I IFN production is reported to be dependent on *Mavs* gene dosage (Seth et al., 2005). Further, when CD11c⁺ cells in the spleen were examined, type I IFN gene expression observed in *Ifih1*^{gs/+}*Mavs*^{+/-} cells was abolished in *Ifih1*^{gs/+}*Mavs*^{-/-} cells (Figure 4D). Although the number of effector CD4⁺ T and effector CD8⁺ T cells was increased in the *Ifih1*^{gs/+}*Mavs*^{+/-} spleen compared with the *Ifih1*^{+/+}*Mavs*^{+/-} spleen, additional deletion of the *Mavs* gene (*Ifih1*^{gs/+}*Mavs*^{-/-}) abolished the increase of these effector cells (Figure S3A). In addition, activation of cDC and pDC as judged by the increase of CD86 expression, was similarly corrected by deletion of the *Mavs* gene (Figure S3B). These results strongly suggest that the observed kidney pathology, constitutive production of type I IFN and cytokines, and increase and activation of effector T cells and DCs are caused by a mutated *Ifih1* allele. Furthermore, the MDA5-MAVS axis is the key molecular basis for the pathogenesis in MDA5 gs mutant mice.

Type I IFN Signaling Contributes to But Is Not Essential for MDA5-Dependent Nephritis

Emerging evidence has implicated dysregulated type I IFN production as a key predisposing factor for autoimmune diseases including SLE (Banchereau and Pascual, 2006). MDA5 signaling causes type I IFN induction as well as other cytokines. To further investigate the contribution of type I IFNs in the current autoimmune model, we intercrossed *Ifih1*^{gs/+} mice with *Ifnar1*^{-/-} mice on a C57BL/6 background. *Ifih1*^{gs/+}*Ifnar1*^{-/-} mice had partially ameliorated nephritis; however, they still developed substantial pathology in the kidney (Figures 5A and 5B). In the affected kidney, slight induction of type I IFNs and inflammatory cytokines


Figure 2. Intrinsically Activated DCs and Macrophages Are Possible Sources of Chronic Inflammation

(A and B) Splenocytes were stained with CD11c and B220 (A) or CD11b and F4/80 (B) after MACS depletion of CD3⁺ and CD19⁺ cells and analyzed by flow cytometry. pDCs and cDCs (A) and macrophages (B) were additionally gated and their CD86 expression was accessed in histogram and quantification of mean fluorescent intensity (MFI) was quantified in bar graphs. Closed histogram, *Ifih1*^{+/+}; red open, *Ifih1*^{gs/+}. Statistical significance was determined by Student's t test. *p < 0.03.

(C) Indicated immune cells were sorted from spleen and indicated mRNA expression levels were determined by quantitative real-time PCR. Data are shown as mean ± SD of duplicate samples of a representative from two independent experiments.

(D) Bone-marrow-derived dendritic cells were sorted by MACS on day 6 and their *Ifnb* mRNA expression levels were determined by quantitative real-time PCR.

(E) Left shows splenocytes were stained with CD19/CD138 and analyzed by FACS. Right shows absolute cell counts of plasma cells were shown in bar graphs. Statistical significance was determined by Student's t test. **p < 0.05.



(legend on next page)

Immunity

MDA5 Mutant Mice Cause Spontaneous Nephritis

was observed (Figure 5C), suggesting that type I IFNs are partially responsible for the pathogenesis, whereas the other inflammatory cytokines are potentially responsible for the pathology of MDA5 gs mutant mice.

G821S Mutation of MDA5 Is Unresponsive to RNA Ligands but Constitutively Active

We examined *Irfnb* expression in MEFs expressing the G821S MDA5 mutation. *Irfh1^{gs/gs}* MEFs clearly expressed *Irfnb* gene in the absence of cellular stimuli and *Irfh1^{gs/+}* MEFs exhibited lower but significant upregulation compared to *Irfh1^{+/+}* MEFs (Figure 6A), suggesting that the G821S mutation confers gain of function. To further confirm this notion, expression vectors for human MDA5 and its mutants were transiently expressed in cultured cells. In addition to G821S, which was conserved at the same amino acid position in human *IFIH1*, another mutation, A946T, which has been implicated in the relation with SLE, was tested (Gateva et al., 2009). Expression of human G821S and human A946T mutants strongly activated IFN- β promoter in Huh-7 cells (Figure 6B), suggesting that these mutations conferred intrinsic activation function on MDA5. Next, to examine cell-type specificity in the mouse system, we introduced mutant MDA5 to *Irfh1^{-/-}* DCs with retroviral vectors (Figure 6C). WT MDA5 expression resulted in constitutive expression of *Irfnb*, which is consistent with a previous notion that MDA5 exhibits constitutive activity when overexpressed. However, MDA5 gs exhibited significantly higher activity than the WT. These data indicate the importance of a conserved G821 region in mouse and human MDA5 to suppress autoactivation in both hematopoietic and nonhematopoietic cells. We next investigated IFN- β production in WT and mutant MEFs upon viral infection (Figures 6D and 6E). EMCV infection is exclusively detected by MDA5, whereas infection by Newcastle disease virus (NDV) infection is exclusively detected by RIG-I (Kato et al., 2006). Upon EMCV infection, *Irfh1^{+/+}* and *Irfh1^{gs/+}* MEFs produced high levels of IFN- β ; however, *Irfh1^{gs/gs}* MEFs did not (Figure 6D). When these MEFs were infected with NDV, efficient production of IFN- β was observed due to the function of RIG-I in these cells (Figure 6E). Further, to clarify the response of A946T mutant, we introduced the mutant MDA5 to *Irfh1^{-/-}* MEFs with expression vector and concluded that A946T could not respond to EMCV (Figure 6F). These results demonstrate that although G821S and A946T mutations confer constitutive activity, they disrupt responsiveness to viral infection.

G821S and A946T Mutations Lead to Conformational Change and Abrogate ATPase Activity of MDA5

To characterize the physicochemical properties of MDA5 gs and A946T, we produced recombinant MDA5 proteins. The purity of

the recombinant MDA5 proteins were confirmed by SDS-PAGE (Figure 7A). First, the recombinant proteins were examined by an atomic force microscope (AFM) (Figure 7B). WT MDA5 appeared as three closely associated lumps, whereas MDA5 gs and A946T were observed as a single blunt lump. Quantification confirmed that MDA5 gs, A946T, and WT have distinct molecular shapes, suggesting open and closed conformations, respectively. We next examined ATPase activity because MDA5 is a dsRNA-dependent ATPase (Sumpter et al., 2005). In the absence of dsRNA, WT MDA5 exhibited very low ATPase activity, which was markedly enhanced in the presence of polyI:C. In contrast, ATPase activity of neither MDA5 gs nor A946T was augmented even in the presence of polyI:C (Figure 7C), consistent with the result that MDA5 gs and A946T failed to respond to viral infection (Figures 6D and 6F). Thus, the constitutive activity of MDA5 gs and A946T is likely a result of conformational alteration rather than hyperresponsiveness to endogenous RNA ligand.

DISCUSSION

Type I IFNs are associated with a number of human autoimmune diseases including SLE. Here we described the molecular mechanisms of a type I IFN-mediated autoimmune disorder in a mouse model. We identified lupus-like features, as well as upregulation of type I IFNs and cytokines, in multiorgans of MDA5 mutant mice. We showed that MAVS is essential for autoimmune disorders in MDA5 gs mice; thus, MDA5-dependent signaling through MAVS is critical for the pathogenesis of autoimmune disease associated with MDA5 mutation. Of note, MDA5 transgenic mice have been reported to upregulate the production of anti-nuclear antibodies in sera on an Fc γ R2b-deficient background, leading to glomerulonephritis (Crampton et al., 2012); however, MDA5 overexpression was not sufficient to develop autoimmune symptoms, suggesting that the magnitude of constitutive signaling by WT MDA5-overexpression is negligible compared to that by single copy of MDA5 gs. In accordance with this, no gene amplification but a point mutation of *Irfh1^{gs}* has been implicated in human autoimmune diseases (Gateva et al., 2009).

It was surprising that the amount of IFN- β produced by *Irfh1^{gs/+}* and *Irfh1^{gs/gs}* cells was far lower than that produced upon viral infection. However, it was sufficient to cause severe symptoms in mice. We can explain this by hypothesizing a cell-intrinsic mechanism: MDA5 gs possibly takes an autoactivated form even in the absence of RNA ligands, subsequently leading to constitutive signal transduction via MAVS. Consequently, it results in chronic activation of the type I IFN pathway through IRF3, IRF7, NF- κ B, and MAP kinases, resulting in upregulation

Figure 3. Hematopoietic Cells Are Involved in Renal Pathogenesis


(A) Schematic diagram of bone-marrow-transfer experiments.

(B) Histology of kidney from *Irfh1^{+/+}* and *Irfh1^{gs/+}* bone-marrow-transferred mice (left panels, Haematoxylin-eosin staining; right panels, PAS).

(C) Immunoglobulin and C3 depositions on kidney from *Irfh1^{+/+}* and *Irfh1^{gs/+}* bone-marrow-transferred mice.

(D) mRNA expression of kidney from *Irfh1^{+/+}* and *Irfh1^{gs/+}* bone-marrow transferred mice was determined quantitatively real-time PCR. Data are shown as mean \pm SD of duplicate samples of a representative from three independent experiments. Open histogram shows *Irfh1^{+/+}* bone-marrow transferred mice; closed shows *Irfh1^{gs/+}* bone-marrow transferred mice.

(E) Whole splenocytes were prepared and stained with CD11c and B220. pDCs and cDCs were gated (left) and their CD86 expression was accessed in histogram (middle) and quantification of mean fluorescent intensity (MFI) was quantified in bar graphs (right). Closed histogram shows *Irfh1^{+/+}*; red open shows *Irfh1^{gs/+}* splenocytes. Statistical significance was determined by Student's *t* test. **p* < 0.03.



(legend on next page)

Immunity

MDA5 Mutant Mice Cause Spontaneous Nephritis

of the genes of type I and III IFN, inflammatory cytokines, and IFN-inducible genes.

Such chronic activation should take place in all cell types and is therefore sufficient to trigger biological consequences. This hypothesis is consistent with the result that deletion of MAVS but not IFNAR1 completely abrogated the pathology. Because hematopoietic *Iflh1*^{gs/+} cell transfer is sufficient to reproduce the disorder and DCs and macrophages upregulate a cell activation marker, CD86, we propose DCs and macrophages as the primary cell types for initiating autoimmunity in *Iflh1*^{gs/+} mice. The partial effect of type I IFN receptor deletion on rescue from the symptoms suggests the potential involvement of other inflammatory cytokines, including IL-6 and TNF- α , for amplifying the activation cascade, leading to the effector T cells and antibody production.

These observations suggest possible therapies targeting the pathway to arrest disease progression. For example, reagents that modify the molecular structure of MDA5 or direct blocking of the interaction between MDA5 and MAVS or inhibition of downstream kinases will provide fundamental therapy. Temporal depletion of DCs and macrophages would potentially ameliorate inflammation and tissue damage. Conversely, blocking ATPase activity of MDA5 gs and A946T would not provide therapeutic effect because gs and A946T are negative for ATPase activity.

Crystal structures of RIG-I suggest that free RIG-I has an open structure in which CARD is masked by interactions with helicase domain 2, thus being inactive for signaling (Kowalinski et al., 2011). Upon binding with dsRNA, it markedly changes the conformation into a more packed structure, thereby releasing CARD. We propose a similar scenario for WT MDA5, which has an open structure with CARD masked; however, G821S mutation results in conformational change as revealed by AFM analysis.

Paradoxically, in the response to viral infection, gs is unable to further upregulate IFN. We propose that the gs mutation resulted in constitutive activity rather than being hypersensitive to endogenous RNA ligands. This hypothesis is supported by the observations; gs does not exhibit ATPase activity, even in the presence of dsRNA; AFM analysis revealed that purified gs protein takes distinct conformation from WT protein. It appears that gene induction by virus infection in WT cells is far more efficient than constitutive gene expression resulting from *Iflh1*^{gs/+} mutation. This is possibly because WT MDA5 is efficiently activated by binding with viral dsRNA to form a fiber-like polymer to facilitate a signaling relay (Wu et al., 2013); however, MDA5 gs is activated without such polymerization.

Within human *IFIH1* SNPs, A946T has been reported to be a risk variant for SLE (Gateva et al., 2009). This missense mutation is implied to be a gain-of-function mutation in SLE (Robinson

et al., 2011). We found that A946T, like G821S, constitutively activated IFN signaling. In addition, MDA5 A946T takes packed structure as observed by AFM and does not exhibit ATPase activity even in the presence of dsRNA. These features resemble those of G821S. Consistent with our observation, it was reported that mutations in MDA5 helicase motif I, III, and VI lead to constitutive IFN signaling with inactivation of ATPase activity (Bamming and Horvath, 2009). Furthermore, constitutive active mutants of RIG-I (mut1 and 2) exhibit no ATPase activity (Kageyama et al., 2011).

Increasing number of reports revealed that the molecules involved in innate immune signal pathways are associated with the development of inflammatory diseases. For example, NOD1 and NOD2 sense cytosolic peptidoglycan and muramyl dipeptide (MDP), respectively, and activate MAPK and NF- κ B-dependent pathway (Elinav et al., 2011). Variants of NOD2 are reported to increase the susceptibility to Crohn's disease (CD) (Hugot et al., 2001; Ogura et al., 2001). These mutations abolish the ability to sense MDP and fail to activate NF- κ B in peripheral blood mononuclear cells (van Heel et al., 2005), whereas overproduction of NF- κ B-regulated genes including TNF- α , IL-1 β , and IL-6 in epithelial cells and lamina propria macrophages are detected in CD patients. These paradoxical immune responses might resemble those mediated by MDA5 gs. In a basal state, MDA5 gs weakly but constitutively activates IFN signaling in the absence of ligand leading to persistent chronic inflammation. Conversely, in viral infection, MDA5 gs is defective in antiviral immune response for limiting inflammation, which could potentially underlie susceptibility to viral infection. For another inflammatory disease-associated molecule in NOD family, NALP-3 mutations are associated with three different autosomal diseases, Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome, and familial cold autoinflammatory syndrome (FACS) (Bihl et al., 2005; Hoffman et al., 2001). These NALP-3 mutations result in an increased processing of pro-IL-1 β and subsequent release of the active cytokine (Dinarello, 2004). These observations support gain-of-functional model for triggering chronic inflammation and provide a clue for understanding the mechanism of autoinflammatory disorders.

Some intracellular molecules that regulate degradation of nucleic acid have been linked to autoimmune disorders. For example, TREX1 is a major exonuclease that degrades endogenous DNA. In *Trex1*^{-/-} mice, cell-intrinsic ligands activate IFN signaling, consequently leading to autoimmunity (Gall et al., 2012; Stetson et al., 2008), linking the association that human genetic mutations in *Trex1* cause Aicardi-Goutieres syndrome (AGS) and chilblain lupus. In comparison, we defined MDA5 gs mutation as gain of function in basal state, but not hypersensitive to endogenous RNA ligands (above).

Figure 4. MAVS Is Critical for the Development of Nephritis in *Iflh1*^{gs/+} Mice

(A) Histology of kidney from WT, *Iflh1*^{gs/+}, *Mavs*^{+/-}, or *Mavs*^{-/-} mice (left panels, Haematoxylin-eosin staining; right panels, PAS).

(B) Histological scores and percentage of sclerosis in kidney from WT, *Iflh1*^{gs/+}, *Mavs*^{+/-}, *Mavs*^{+/-}, or *Mavs*^{-/-} mice were evaluated. Statistical significance was determined by Student's t test. *p < 0.03.

(C) Indicated mRNA gene expression levels in kidney from WT, *Iflh1*^{gs/+}, *Mavs*^{+/-}, *Mavs*^{+/-}, or *Mavs*^{-/-} mice were determined by quantitative real-time PCR. Littermate mice were subjected and data are shown as mean \pm SD of duplicate samples of a representative from three independent experiments.

(D) CD11c⁺ cells were sorted from splenocytes of indicated mice and their mRNA expression was determined by quantitative real-time PCR. Data are shown as mean \pm SD of duplicate samples of a representative from three independent experiments.