

研究成果の刊行に関する一覧表(平成 23 ~ 25 年度)

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Kubo S, Yamaoka K, Kondo M, Yamagata K, Zhao J, Iwata S, <u>Tanaka Y.</u>	The JAK inhibitor tofacitinib reduces the T cell stimulatory capacity of human monocyte-derived dendritic cells.	Ann Rheum Dis		in press	
<u>Tanaka Y</u> , Hirata S, Kubo S, Fukuyo S, Hanami K, Sawamukai N, Nakano K, Nakayamada S, Yamaoka K, Sawamura F, Saito K.	Discontinuation of adalimumab after achieving remission in patients with established rheumatoid arthritis: 1-year outcome of the HONOR study.	Ann Rheum Dis		in press	
<u>Tanaka Y.</u>	Next stage of RA treatment: TNF-inhibitor-free remission will be a possible treatment goal ?	Ann Rheum Dis	72	ii124-ii127	2013
<u>Tanaka Y</u> , Maeshima Y, Yamaoka K.	In vitro and in vivo analysis of a Jak inhibitor in rheumatoid arthritis.	Ann Rheum Dis	71	i70-i74	2012
Iwata S, Yamaoka K, Niuro H, Nakano K, Wang S-P, Akashi K, <u>Tanaka Y.</u>	Amplification of toll-like receptor-mediated signaling through Syk in human B cell activation.	J Allergy Clin Immunol	129	1594-1601	2012

Ikeda K, Seto Y, Narita A, kawakami A, Kawahito Y, Ito H, Matsushita I, Ohno S, Nishida K, Suzuki T, Kaneko A, Ogasawara M, Fukae J, Henmi M, Sumida T, Kamishima T, Koike T.	Ultrasound assessment of synovial pathologies in rheumatoid arthritis using comprehensive multi-plane images of the 2nd metacarpophalangeal joint-Identification of the components which are reliable and influential on the global assessment of the whole joint.	Arthritis Rheum	in press		
Koike T.	IgG4-related disease: why high IgG4 and fibrosis. Arthritis Res Ther.	Arthritis Res Ther	15(1)	103	2013
Takeuchi T, Yamanaka H, Ishiguro N, Miyasaka N, Mukai M, Matsubara T, Uchida S, Akama H, Hartmut K, Vipin A, and <u>Tanaka Y.</u>	Adalimumab, a Human anti-TNF monoclonal antibody, outcome study for the prevention of joint damage in Japanese patients with early RA; the HOPEFUL 1 study.	Ann Rheum Dis	in press		
Katayama M, Ohmura K, Yukawa N, Terao C, Hashimoto M, Yoshifuji H, Kawabata D, Fujii T, Iwakura Y, Mimori T	Neutrophils are essential as a source of IL-17 in the effector phase of arthritis.	PLos One	8(5)	e62231	2013
Terao C, Hashimoto M, Yamamoto K, Murakami K, Ohmura K, Nakashima R, Yamakawa N, Yoshifuji H, Yukawa N, Kawabata D, Usui T, Yoshitomi H, Furu M, Yamada R, Matsuda F, Ito H, Fujii T, Mimori T	Three Groups in the 28 Joints for Rheumatoid Arthritis Synovitis - Analysis Using More than 17,000 Assessments in the KURAMA Database.	PLos One	8(3)	e59341	2013

Sumitomo S, Fujio K, Okamura T, Morita K, Ishigaki K, Suzukawa K, Kanaya K, Kondo K, Yamasoba T, Furukawa A, Kitahara N, Shoda H, Shibuya M, Okamoto A, Yamamoto K.	Transcription factor early growth response 3 is associated with the TGF- 1 expression and the regulatory activity of CD4-positive T cells in vivo.	J Immunol	191(5)	2351-9	2013
Iwasaki Y, Fujio K, Okamura T, Yanai A, Sumitomo S, Shoda H, Tamura T, Yoshida H, Charnay P, Yamamoto K.	Egr-2 transcription factor is required for Blimp-1-mediated IL-10 production in IL-27-stimulated CD4+ T cells.	Eur J Immunol	43(4)	1063-73	2013
Sawamukai N, Satake A, Schmidt AM, Lamborn IT, <u>Tanaka Y</u> , Kambayashi T.	Cell autonomous role of TGF and IL-2 receptor in CD4+ and CD8+ inducible regulatory T cell generation during graft-versus-host disease.	Blood	119	5575-83	2012
Iwata S, Yamaoka K, Niuro H, Nakano K, Wang S-P, Akashi K, <u>Tanaka Y</u> .	Amplification of toll-like receptor-mediated signaling through Syk in human B cell activation.	J Allergy Clin Immunol	129	1594-1601	2012
Iwata S, Saito K, Tokunaga M, <u>Tanaka Y</u> .	Phenotypic changes of lymphocyte in a patients with IgG4-related disease after corticosteroid therapy.	Ann Rheum Dis	71	2058-2059	2012
Tak PP, Mease PJ, Genovese MC, Kremer J, Haraoui B, Tanaka Y, Bingham III CO, Ashrafzadeh A, Travers H, Safa-Leathers S, Kumar S, Dummer W.	Efficacy and safety of ocrelizumab in patients with active rheumatoid arthritis with an inadequate response to at least one TNF inhibitor: results from the phase III SCRIPT trial.	Arthritis Rheum	64	360-370	2012

Sonomoto K, Yamaoka K, Oshita K, Fukuyo S, Zhang X, Nakano K, Okada Y, <u>Tanaka Y.</u>	IL-1 induces differentiation of human mesenchymal stem cells into osteoblasts via the Wnt5a/Ror2 pathway.	Arthritis Rheum	64	3355-3363	2012
Maeshima K, Yamaoka K, Kubo S, Nakano K, Iwata S, Saito K, Ohishi M, Miyahara H, Tanaka S, Ishi K, Yoshimatsu H, <u>Tanaka Y.</u>	A JAK inhibitor tofacitinib regulates synovitis through inhibition of IFN-g and IL-17 production by human CD4+ T cells.	Arthritis Rheum	64	1790-1798	2012
Kawashiri SY, Arima K, Suzuki T, Nakashima Y, Horai Y, Okada A, Tamai M, Nakamura H, <u>Kawakami A.</u>	Presence of ultrasound subclinical synovitis and increment of serum vascular endothelial growth factor in a patient with rheumatoid arthritis achieved in sustained clinical remission by treatment with adalimumab and methotrexate.	Mod Rheumatol	In Press		
Kawashiri SY, Fujikawa K, Nishino A, Suzuki T, Okada A, Nakamura H, <u>Kawakami A.</u>	Usefulness of ultrasonography-proven tenosynovitis to monitor disease activity of a patient with very early rheumatoid arthritis treated by abatacept.	Mod Rheumatol	23 (3)	582-586	2013
Kita J, Tamai M, Arima K, Kawashiri SY, Horai Y, Iwamoto N, Okada A, Koga T, Nakashima Y, Suzuki T, Yamasaki S, Nakamura H, Origuchi T, Ida H, Aoyagi K, Uetani M, Eguchi K, <u>Kawakami A.</u>	Significant improvement in MRI-proven bone edema is associated with protection from structural damage in very early RA patients managed using the tight control approach.	Mod Rheumatol	23 (2)	254-259	2013
Kita J, Tamai M, Arima K, Nakashima Y, Suzuki T, Kawashiri SY, Iwamoto N, Okada A, Koga T, Yamasaki S, Nakamura H, Origuchi T, Ida H, Aoyagi K, Uetani M, Eguchi K, <u>Kawakami A.</u>	Treatment discontinuation in patients with very early rheumatoid arthritis in sustained simplified disease activity index remission after synthetic disease-modifying anti-rheumatic drug administration.	Mod Rheumatol	22 (3)	346-352	2012

<p>Tamai M, <u>Kawakami A</u>, Uetani M, Fukushima A, Arima K, Fujikawa K, Iwamoto N, Aramaki T, Kamachi M, Nakamura H, Ida H, Origuchi T, Aoyagi K, Eguchi K.</p>	<p>Magnetic resonance imaging (MRI) detection of synovitis and bone lesions of the wrists and finger joints in early-stage rheumatoid arthritis: comparison of the accuracy of plain MRI-based findings and gadolinium-diethylenetriamine pentaacetic acid-enhanced MRI-based findings.</p>	<p>Mod Rheumatol</p>	<p>22 (5)</p>	<p>654-658</p>	<p>2012</p>
<p>Fukae J, Isobe M, Kitano A, Henmi M, Sakamoto F, Narita A, Ito T, Mitsuzaki A, Shimizu M, Tanimura K, Matsuhashi M, Kamishima T, Atsumi T, <u>Koike T.</u></p>	<p>Radiographic prognosis of finger joint damage predicted by early alteration in synovial vascularity in patients with rheumatoid arthritis: Potential utility of power doppler sonography in clinical practice.</p>	<p>Arthritis Care Res</p>	<p>63(9)</p>	<p>1247-53</p>	<p>2011</p>
<p>Otomo K, Atsumi T, Amengual O, Fujieda Y, Kato M, Oku K, Horita T, Yasuda S, <u>Koike T.</u></p>	<p>The efficacy of antiphospholipid score for the diagnosis of antiphospholipid syndrome and its predictive value for thrombotic events.</p>	<p>Arthritis Rheum</p>	<p>64(2)</p>	<p>504-12</p>	<p>2012</p>
<p>Fukae J, Isobe M, Kitano A, Henmi M, Sakamoto F, Narita A, Ito T, Mitsuzaki A, Shimizu M, Tanimura K, Matsuhashi M, Kamishima T, Atsumi T, <u>Koike T.</u></p>	<p>Positive synovial vascularity in patients with low disease activity indicates smouldering inflammation leading to joint damage in rheumatoid arthritis: time-integrated joint inflammation estimated by synovial vascularity in each finger joint.</p>	<p>Rheumatology</p>	<p>52(3)</p>	<p>523-8</p>	<p>2013</p>
<p>Ikeda K, Seto Y, Narita A, kawakami A, Kawahito Y, Ito H, Matsushita I, Ohno S, Nishida K, Suzuki T, Kaneko A, Ogasawara M, Fukae J, Henmi M, Sumida T, Kamishima T, <u>Koike T.</u></p>	<p>Ultrasound assessment of synovial pathologic features in rheumatoid arthritis using comprehensive multi-plane images of the 2nd metacarpophalangeal joint - Identification of the components which are reliable and influential on the global assessment of the whole joint.</p>	<p>Arthritis Rheum</p>	<p>in press</p>		

Fukae J, Tanimura K, Atsumi T, <u>Koike T.</u>	Sonographic synovial vascularity of synovitis in rheumatoid arthritis.	Rheumatology	epub ahead of print		
Murakami Y, Mizoguchi F, Saito T, Miyasaka N, <u>Kohsaka H.</u>	p16(INK4a) exerts an anti-inflammatory effect through accelerated IRAK1 degradation in macrophages.	J Immunol.	189	5066-72	2012
細矢 匡, <u>上阪 等</u>	【低分子抗炎症薬の進歩】CDK 阻害薬の発見から臨床応用の可能性まで	炎症と免疫	20	278-283	2012
岩井 秀之, <u>上阪 等</u>	新規関節炎治療標的としての Triggering receptor expressed on myeloid cells-1 阻害	日本臨床免疫学会会誌	35	81-86	2012
溝口 史高, <u>上阪 等</u>	関節リウマチにおける microRNA の役割	日本臨床免疫学会会誌	35	69-74	2012
<u>上阪 等</u>	【関節リウマチの最新情報-寛解を目指した診断と治療の新展開-】新たな関節リウマチ治療のターゲット 滑膜線維芽細胞を標的とした治療法の開発	最新医学	67	281-286	2012
Umeda N, Matsumoto I, Ito I, Kawasaki A, Tanaka Y, Inoue A, Tanaka Y, Tsuboi H, Suzuki T, Hayashi T, Ito S, Tsuchiya N, <u>Sumida T.</u>	Anti-citrullinated glucose-6-phosphate isomerase peptide antibodies in patients with rheumatoid arthritis are associated with HLA-DRB1 shared epitope alleles and disease activity.	Clin Exp Immunol	172	44-53	2013
Horikoshi M, Goto D, Segawa S, Yoshiga Y, Iwanami K, Inoue A, Tanaka Y, Matsumoto I, <u>Sumida T.</u>	Activation of invariant NKT cells with glycolipid ligand -galactosylceramide ameliorates glucose-6-phosphate isomerase peptide-induced arthritis.	PLoS One	7	e51215	2012

Inoue A, Matsumoto I, Tanaka Y, Umeda N, Tanaka Y, Mihara M, Takahashi S, <u>Sumida T.</u>	Murine tumor necrosis factor -induced adipose-related protein (tumor necrosis factor -induced protein 9) deficiency leads to arthritis via interleukin-6 overproduction with enhanced NF- B, STAT-3 signaling, and dysregulated apoptosis of macrophages.	Arthritis Rheum	64	3877-85	2012
Kondo Y, Iizuka M, Wakamatsu E, Yao Z, Tahara M, Tsuboi H, Sugihara M, Hayashi T, Yoh K, Takahashi S, Matsumoto I, <u>Sumida T.</u>	Overexpression of T-bet gene regulates murine autoimmune arthritis.	Arthritis Rheum	64	162-72	2012
Yoshiga Y, Goto D, Segawa S, Horikoshi M, Hayashi T, Matsumoto I, Ito S, Taniguchi S, <u>Sumida T.</u>	Activation of natural killer T cells by -carba-GalCer (RCAI-56), a novel synthetic glycolipid ligand, suppresses murine collagen-induced arthritis.	Clin Exp Immunol	164	236-47	2011
Nishimoto N, Koichi Amano, <u>Tsutomu Takeuchi,</u> Nobuhiro Takagi, et al.	Retreatment efficacy and safety of tocilizumab in patients with rheumatoid arthritis in recurrence (RESTORE) study.	Mod Rheumatol	24	26-32	2014
Kaneko Y, Kondo H and <u>Takeuchi T.</u>	ACR/EULAR remission criteria maintains strict performance when evaluated in 44 joints.	J Rheumatology	40	1254-8	2013
Nishina N, Kaneko Y, Kameda H, Kuwana M, and <u>Takeuchi T.</u>	Reduction of plasma IL-6, but not TNF- by methotrexate in patients with early rheumatoid arthritis: a potential biomarker for radiographic progression.	Clinical Rheumatol	32	1661-7	2013
Nishina N, Kikuchi J, Hashizume M, Yoshimoto K, Kameda H, and <u>Takeuchi T.</u>	Baseline levels of soluble interleukin-6 receptor predict clinical remission in patients with rheumatoid arthritis treated with tocilizumab: implications for molecular targeted therapy.	Ann rheum Dis,		In press	

Takeuchi T, Miyasaka N, Tatsuki Y, Yano T, Yoshinari T, Abe T, and Koike T.	Baseline tumor necrosis factor alpha levels predict the necessity for dose escalation of infliximab therapy in patients with rheumatoid arthritis.	Ann Rheum Dis	70	1208-15	2011
Nakashima R, Imura Y, Hosono Y, Seto M, Murakami A, Watanabe K, Handa T, Mishima M, Hirakata M, Takeuchi T, Fujio K, Yamamoto K, Kohsaka H, Takasaki Y, Enomoto N, Suda T, Chida K, Hisata S, Nukiwa T, <u>Mimori T</u> .	The Multicenter Study of a New Assay for Simultaneous Detection of Multiple Anti-Aminoacyl-tRNA Synthetases in Myositis and Interstitial Pneumonia	PLoS One	9	e85062	2014
Tanizawa K, Handa T, Nakashima R, Kubo T, Hosono Y, Aihara K, Ikezoe K, Watanabe K, Taguchi Y, Hatta K, Oga T, Chin K, Nagai S, <u>Mimori T</u> , Mishima M	The prognostic value of HRCT in myositis-associated interstitial lung disease.	Respir Med.	107	745-752	2013
<u>Mimori T</u> , Nakashima R, Hosono Y	Interstitial lung disease in myositis: Clinical subsets, biomarkers, and treatment.	Curr Rheumatol Rep	14	264-274	2012
Watanabe K, Handa T, Tanizawa K, Hosono Y, Taguchi Y, Noma S, Kobashi Y, Kubo T, Aihara K, Chin K, Nagai S, <u>Mimori T</u> , Mishima M	Detection of antisynthetase syndrome in patients with idiopathic interstitial pneumonias.	Respir Med.	105	1238-1247	2011
Tanizawa K, Handa T, Nakashima R, Kubo T, Hosono Y, Watanabe K, Aihara K, Oga T, Chin K, Nagai S, <u>Mimori T</u> , Mishima M	HRCT features of interstitial lung disease in dermatomyositis with anti-CADM-140 antibody.	Resp Med	105	1380-1387	2011
Myouzen K, Kochi Y, Okada Y, Terao C, Suzuki A, Ikari K, Tsunoda T, Takahashi A, Kubo M, Taniguchi A, Matsuda F, Ohmura K, Momohara S, <u>Mimori T</u> , <u>Yamanaka H</u> , Kamatani N, Yamada R, Nakamura Y, Yamamoto K.	Functional variants in NFKBIE and RTKN2 involved in activation of the NF- B pathway are associated with rheumatoid arthritis in Japanese.	Plos Genet	8(9)	e1002949	2012

<p>Terao C, Ohmura K, Ikari K, Kochi Y, Maruya E, Katayama M, Yurugi K, Shimada K, Murasawa A, Honjo S, Takasugi K, Matsuo K, Tajima K, Suzuki A, Yamamoto K, Momohara S, <u>Yamanaka H</u>, Yamada R, Saji H, Matsuda F, Mimori T.</p>	<p>ACPA-negative RA consists of two genetically distinct subsets based on RF positivity in Japanese.</p>	<p>Plos One</p>	<p>7(7)</p>	<p>e40067</p>	<p>2012</p>
<p>Okada Y, Terao C, Ikari K, Kochi Y, Ohmura K, Suzuki A, Kawaguchi T, Stahl EA, Kurreeman FA, Nishida N, Ohmiya H, Myouzen K, Takahashi M, Sawada T, Nishioka Y, Yukioka M, Matsubara T, Wakitani S, Teshima R, Tohma S, Takasugi K, Shimada K, Murasawa A, Honjo S, Matsuo K, Tanaka H, Tajima K, Suzuki T, Iwamoto T, Kawamura Y, Tani H, Okazaki Y, Sasaki T, Gregersen PK, Padyukov L, Worthington J, Siminovitch KA, Lathrop M, Taniguchi A, Takahashi A, Tokunaga K, Kubo M, Nakamura Y, Kamatani N, Mimori T, Plenge RM, <u>Yamanaka H</u>, Momohara S, Yamada R, Matsuda F, Yamamoto K</p>	<p>Meta-analysis identifies nine new loci associated with rheumatoid arthritis in the Japanese population.</p>	<p>Nat Genet</p>	<p>25</p>	<p>511-6</p>	<p>2012</p>
<p>Yamanaka H, Tanaka Y, Inoue E, Hoshi D, Momohara S, Hanami K, Yunoue N, Saito K, Amano K, Kameda H, Takeuchi T.</p>	<p>Efficacy and tolerability of tocilizumab in rheumatoid arthritis patients seen in daily clinical practice in Japan: results from a retrospective study (REACTION study).</p>	<p>Mod Rheumatol</p>	<p>21</p>	<p>122-33</p>	<p>2011</p>

<p>Okada Y, Wu D, Trynka G, Raj T, Terao C, Ikari K, Kochi Y, Ohmura K, Suzuki A, Yoshida S, Graham RR, Manoharan A, Ortmann W, Bhangale T, Denny JC, Carroll RJ, Eyster AE, Greenberg JD, Kremer JM, Pappas DA, Jiang L, Yin J, Ye L, Su DF, Yang J, Xie G, Keystone E, Westra HJ, Esko T, Metspalu A, Zhou X, Gupta N, Mirel D, Stahl EA, Diogo D, Cui J, Liao K, Guo MH, Myouzen K, Kawaguchi T, Coenen MJ, van Riel PL, van de Laar MA, Guchelaar HJ, Huizinga TW, Dieudé P, Mariette X, Louis Bridges Jr S, Zhernakova A, Toes RE, Tak PP, Miceli-Richard C, Bang SY, Lee HS, Martin J, Gonzalez-Gay MA, Rodriguez-Rodriguez L, Rantapää-Dahlqvist S, Arlestig L, Choi HK, Kamatani Y, Galan P, Lathrop M; the RACI consortium; the GARNET consortium, Eyre S, Bowes J, Barton A, de Vries N, Moreland LW, Criswell LA, Karlson EW, Taniguchi A, Yamada R, Kubo M, Liu JS, Bae SC, Worthington J, Padyukov L, Klareskog L, Gregersen PK, Raychaudhuri S, Stranger BE, De Jager PL, Franke L, Visscher PM, Brown MA, <u>Yamanaka H</u>, Mimori T, Takahashi A, Xu H, Behrens TW, Siminovitch KA, Momohara S, Matsuda F, <u>Yamamoto K</u>, Plenge RM.</p>	<p>Genetics of rheumatoid arthritis contributes to biology and drug discovery.</p>	<p>Nature</p>	<p>506</p>	<p>376-381</p>	<p>2013</p>
<p>Okamoto A, Fujio K, Tsuno NH, Takahashi K, <u>Yamamoto K</u>.</p>	<p>Kidney-infiltrating CD4+ T-cell clones promote nephritis in lupus-prone mice.</p>	<p>Kidney Int</p>	<p>82</p>	<p>969-79</p>	<p>2012</p>

<p>Okada Y, Shimane K, Kochi Y, Tahira T, Suzuki A, Higasa K, Takahashi A, Horita T, Atsumi T, Ishii T, Okamoto A, Fujio K, Hirakata M, Amano H, Kondo Y, Ito S, Takada K, Mimori A, Saito K, Kamachi M, Kawaguchi Y, Ikari K, Mohammed O W, Matsuda K, Terao C, Ohmura K, Myouzen K, Hosono N, Tsunoda T, Nishimoto N, Mimori T, Matsuda F, Tanaka Y, Sumida T, Yamanaka H, Takasaki Y, Koike T, Horiuchi T, Hayashi K, Kubo M, Kamatani N. Yamada R, Nakamura Y, Yamamoto K.</p>	<p>Meta-analysis identifies nine new loci associated with rheumatoid arthritis in the Japanese population</p>	<p>Nat Genet.z</p>	<p>44</p>	<p>511-6</p>	<p>2012</p>
<p>Kochi Y, Thabet MM, Suzuki A, Okada Y, Daha NA, Toes REM, Huizinga TWJ, Myouzen K, Kubo M, Yamada R, Nakamura Y, Yamamoto K.</p>	<p>PADI4 polymorphism predisposes male smokers to rheumatoid arthritis.</p>	<p>Ann Rheum Dis.</p>	<p>70</p>	<p>515-5</p>	<p>2011</p>