

研究成果の刊行に関する一覧表

「みんなで考えましょう、アトピー性皮膚炎とおつきあい（市民公開講座）」
<http://www.kyudai-derm.org/atopy/openlec/index.html>

「エキスパートから学ぶ！子どものぜん息・アレルギーの予防と対策」
http://www.congre.co.jp/apapari2011_48jspaci/html/shimin/shimin.html

「アトピー性皮膚炎についていっしょに考えましょう」
<http://www.kyudai-derm.org/atopy/>

「アトピー性皮膚炎 よりよい治療のためのEvidence-based medicine (EBM)とデータ集」
2010年改定版 http://www.kyudai-derm.org/atopy_ebm/index.html

「アトピー性皮膚炎かゆみをやっつけよう！」
<http://www.kyudai-derm.org/kayumi/index.html>

「アトピー性皮膚炎の標準治療」
http://www.kyudai-derm.org/atopy_care/index.html

雑誌 発表者氏名、論文タイトル名、発表誌名、巻号、ページ、出版年

Morino-Koga S, Uchi H, Tsuji G, Takahara M, Kajiwara J, Hirata T, Furue M. Reduction of CC-chemokine ligand 5 by aryl hydrocarbon receptor ligands. *J Dermatol Sci*. 2013 Oct;72(1):9-15.

Furue M, Ebata T, Ikoma A, Takeuchi S, Kataoka Y, Takamori K, Satoh T, Saeki H, Augustin M, Reich A, Szepietowski J, Fleischer A, Blome C, Phan NQ, Weisshaar E, Yosipovitch G, Ständer S. Verbalizing Extremes of the Visual Analogue Scale for Pruritus: A Consensus Statement. *Acta Derm Venereol*. 2013 Mar 27;93(2):214-5.

Kashiwakura J, Okayama Y, Furue M, Kabashima K, Shimada S, Ra C, Siraganian RP, Kawakami Y, Kawakami T. Most Highly Cytokinergic IgEs Have Polyreactivity to Autoantigens. *Allergy Asthma Immunol Res*. 2012 Nov;4(6):332-40.

Kamo A, Tominaga M, Taneda K, Ogawa H, Takamori K. Neurotropin inhibits the increase in intraepidermal nerve density in the acetone-treated dry-skin mouse model. *Clin Exp Dermatol*. 38(6): 665-8, 2013.

Tominaga M, Takamori K. An update on peripheral mechanisms and treatments of itch. *Biol Pharm Bull*. , 36(8): 1241-7, 2013.

Tominaga M, Tenggara S, Kamo A, Ogawa H, Takamori K. Matrix metalloproteinase-8 is involved in dermal nerve growth: Implications for possible application to pruritus from *in vitro* models. *J Invest Dermatol* 131: 2105-12, 2011

Taneda K, Tominaga M, Negi O, Tenggara S, Kamo A, Ogawa H, Takamori K. Evaluation of epidermal nerve density and opioid receptor levels in psoriatic itch. *Br J Dermatol* 165: 277-284, 2011

Kamo A, Tominaga M, Negi O, Tenggara S, Taneda K, Ogawa H, Takamori K. Inhibitory effects of UV-based therapy on dry skin- inducible nerve growth in acetone-treated mice. *J Dermatol Sci* 62: 91-97, 2011

Taneda K, Tominaga M, Tenggara S, Ogawa H, Takamori K. Neurotropin inhibits both capsaicin-induced substance P release and nerve growth factor-induced neurite outgrowth in cultured rat dorsal root ganglion neurons. *Clin Exp Dermatol* 35: 73-77, 2010

Tengara S, Tominaga M, Kamo A, Taneda K, Negi O, Ogawa H, Takamori K. Keratinocyte-derived anosmin-1, an extracellular glycol-protein encoded by X-linked Kallmann syndrome gene, is involved in modulation of epidermal nerve density in atopic dermatitis. *J Dermatol Sci* 58: 64-71, 2010

Yano C, Saeki H, Ishiji T, Ishiujii Y, Sato J, Tofuku Y, Nakagawa H: Impact of disease severity on work productivity and activity impairment in Japanese patients with atopic dermatitis. *J Dermatol* 40: 736-9, 2013.

Saeki H, Hirota T, Nakagawa H, Tsunemi Y, Kato T, Shibata S, Sugaya M, Sato S, Doi S, Miyatake A, Ebe K, Noguchi E, Ebihara T, Amagai M, Esaki H, Takeuchi S, Furue M, Nakamura Y, Tamari M. Genetic polymorphisms in the IL22 gene are associated with psoriasis vulgaris in a Japanese population. *J Dermatol Sci*. 2013;71:148-150.

Yano C, Saeki H, Ishiji T, Ishiujii Y, Sato J, Tofuku Y, Nakagawa H: Impact of disease severity on sleep quality in Japanese patients with atopic dermatitis. *J Dermatol Sci* 72: 195-7, 2013.

Hiragun T, et al. Fungal protein MGL_1304 in sweat is an allergen for atopic dermatitis patients. *J Allergy Clin Immunol* 132: 608-615, 2013.

Hiragun M, et al. Elevated serum IgE against MGL_1304 in patients with atopic dermatitis and cholinergic urticaria. *Allergol Int*, in press.

Kagami S, Sugaya M, Suga H, Morimura S, Kai H, Ohmatsu H, Fujita H, Tsunemi Y, Sato S. Serum gastrin-releasing peptide levels correlate with pruritus in patients with atopic dermatitis. *J Invest Dermatol* 133: 1673-5, 2013.

Suga H, Sugaya M, Miyagaki T, Kawaguchi M, Morimura S, Kai H, Kagami S, Ohmatsu H, Fujita H, Asano Y, Tada Y, Kadono T, Sato S. Serum visfatin levels in patients with atopic dermatitis and cutaneous T-cell lymphoma. *Eur J Dermatol* 23: 629-35, 2013.

Kimura T, Sugaya M, Suga H, Morimura S, Miyamoto A, Kai H, Kagami S, Yanaba K, Fujita H, Asano Y, Tada Y, Kadono T, Sato S. Variations in serum TARC and I-TAC levels reflect minor changes in disease activity and pruritus in atopic dermatitis. *Acta Derm Venereol* in press.

Kawakami T, Soma Y. Questionnaire survey of the efficacy of emollients for adult patients with atopic dermatitis. *J Dermatol*. 2011;38(6): 531-535

Kawakami T, Kimura S, Haga T, Doi R, Kyoya M, Nakagawa K, Soma Y. Health-related quality of life assessed by the effect of bepotastine besilate in patients with pruritus: Importance of emotions score in atopic dermatitis. *J Dermatol*. 2012; 39(6): 527-530

Taniguchi K, Yamamoto S, Hitomi E, Inada Y, Suyama Y, Sugioka T, Hamasaki Y. Interleukin 33 is induced by tumor necrosis factor alpha and interferon gamma in keratinocytes and contributes to allergic contact dermatitis. *J Investig Allergol Clin Immunol* 2013;23(6):428-434

Taniguchi K, Yamamoto S, Hitomi E, Inada Y, Sugioka T, Hamasaki Y. Blockade of Interleukin-33 attenuates allergic contact dermatitis in model mice: possible mechanism via eosinophil infiltration. *Clin Exp Dermatol Res* 2013;4(3):1000183