

## REDUCTION OF HSPC AND MPC IN THE BM BY VEGF

9

- expands hematopoietic stem cells within the central but not endosteal bone marrow region. *Cytokine* 58:218–225.
32. Hoggatt J, KS Mohammad, P Singh, AF Hoggatt, BR Chitteti, JM Speth, P Hu, BA Poteat, KN Stilger, et al. (2013). Differential stem- and progenitor-cell trafficking by prostaglandin E2. *Nature* 495:365–369.
  33. Kook S, J Cho, SB Lee and BC Lee. (2013). The nucleotide sugar UDP-glucose mobilizes long-term repopulating primitive hematopoietic cells. *J Clin Invest* 123:3420–3435.
  34. Chamberlain G, J Fox, B Ashton and J Middleton. (2007). Concise review: mesenchymal stem cells: their phenotype, differentiation capacity, immunological features, and potential for homing. *Stem Cells* 25:2739–2749.
  35. Koizumi N, T Yamaguchi, K Kawabata, F Sakurai, T Sasaki, Y Watanabe, T Hayakawa and H Mizuguchi. (2007). Fiber-modified adenovirus vectors decrease liver toxicity through reduced IL-6 production. *J Immunol* 178:1767–1773.
  36. Liu L, Q Yu, J Lin, X Lai, W Cao, K Du, Y Wang, K Wu, Y Hu, et al. (2011). Hypoxia-inducible factor-1alpha is essential for hypoxia-induced mesenchymal stem cell mobilization into the peripheral blood. *Stem Cells Dev* 20:1961–1971.
  37. Pitchford SC, RC Furze, CP Jones, AM Wengner and SM Rankin. (2009). Differential mobilization of subsets of progenitor cells from the bone marrow. *Cell Stem Cell* 4:62–72.

Address correspondence to:

*Dr. Kenji Kawabata  
Laboratory of Stem Cell Regulation  
National Institute of Biomedical Innovation  
7-6-8, Saito-Asagi  
Ibaraki, Osaka 567-0085  
Japan*

*E-mail:* kawabata@nibio.go.jp

Received for publication September 27, 2013

Accepted after revision December 16, 2013

Prepublished on Liebert Instant Online XXXX XX, XXXX

