

参考文献

- 1) Nakamura M, Nishiwaki Y, Ushida T, Toyama Y. Prevalence and characteristics of chronic musculoskeletal pain in Japan. *J Orthop Sci* 2011 ; 16 : 424 - 32
- 2) O'Connor AB. Neuropathic pain : Quality - of - life, impact, costs and cost effectiveness of therapy. *Pharmacoeconomics* 2009 ; 27 : 95 - 112
- 3) Bouhassira D, et al. Prevalence of chronic pain with neuropathic characteristics in the general population. *Pain* 2008 ; 136 : 380 - 7
- 4) Treede RD, Jensen TS, Campbell JN, et al. Neuropathic pain : Redefinition and a grading system for clinical and research purposes. *Neurology*. 2008 ; 70 : 1630 - 5
- 5) Freynhagen R, Baron R, Gockel U, et al. PainDETECT : a new screening questionnaire to detect neuropathic components in patients with back pain. *Curr Med Res Opin*. 2006 ; 22 : 1911 - 20
- 6) Matsubayashi Y, Takeshita K, Sumitani M, et al. Validity and reliability of the Japanese version of the PainDETECT Questionnaire : A multicenter observational study. *Plos ONE* in press
- 7) Schmidt CO, et al. Modelling the prevalence and cost of back pain with neuropathic pain components in the general population. *Eur J Pain* 2009 ; 13 : 1030 - 5
- 8) Leeuw M, Goossens MEJB, Linton SJ, et al. The fear - avoidance model of musculoskeletal pain : Current state of scientific evidence. *J Behav Med* 2007 ; 30 : 77 - 94

