

- 43) 酒井規夫、ムコ多糖症と遺伝カウンセリング、第1回ムコ多糖症フォーラム、2013.7
- 44) Hossain MA¹, Higaki K², Nanba E², Suzuki Y³, Ozono K¹, Sakai N、CHAPERONE THERAPY FOR KRABBE DISEASE; JAPANESE LATE-ONSET MUTATIONS CAN BE TREATED EFFECTIVELY BY NOEV、ICIM2013.9
- 45) Hidehito Kondo¹, Michiko Shinpo¹, Mohammad Arif Hossain¹, Yusuke Hamada¹, Norio Sakai¹, Yoshihiro Asano², Takeshi Masaki², Tadayasu Togawa³, Keichi Ozono, A case report of Fabry disease with chronic heart failure treated with amiodarone, ACIM2013.11
- 46) Mohammad Arif Hossain¹, Katsumi Higaki², Eiji Nanba², Yoshiyuki Suzuki³, Keiichi Ozono¹, Norio Sakai¹, NOEV treatment option for Japanese Krabbe disease, ACIM2013.11
- 47) Mohammad Arif Hossain¹, Michiko Shinpo¹, Katsumi Higaki², Eiji Nanba², Yoshiyuki Suzuki³, Keiichi Ozono¹, Norio Sakai¹, NOEV can stabilize β -galactosidase in galactosialidosis patients' skin fibroblasts effectively and cause normalization of its activity, ACIM2013.11
- 48) Michiko Shinpo, Sayaka Nakano, Yusuke Hamada, Kouji Tominaga, Shin Nabatame, Takeshi Okinaga, Yoshiko Hashii, Norio Sakai, Keichi Ozono, Outcomes of hematopoietic stem cell transplantation for three patients with metachromatic leukodystrophy, ACIM2013.11
- 49) Yusuke Hamada¹, Hidehito Kondo¹, Michiko Shinpo¹, Yoshiro Wada², Norio Sakai¹, Yutaka Sumida², Kiichi Ozono¹, Different clinical course of propionic acidemia in two siblings, ACIM2013.11
- 50) Motohiro Akagi^{1, 2}, Mohammad Arif Hossain¹, Keiichi Ozono¹, Norio Sakai¹, Yoshinori Okumura³, Clinicogenetical features of a Japanese patient with static encephalopathy of childhood with neurodegeneration in adulthood (SENDA), ACIM2013.11
- 51) N. Sakai¹, X.F. Gu², H. Ida³ on behalf of the EDGE investigators, O. Kawaguchi⁴, Y. Xue⁵, A phase 3 study evaluating once versus twice daily dosing of eliglustat in patients with Gaucher disease type 1 (GD1): Interim results from the lead in period, ACIM2013.11
- 52) Akemi Tanaka¹, Takashi Hamazaki¹, Motomichi Kosuga², Torayuki Okuyama², Yasuyuki Suzuki³, Norio Sakai⁴, Tomo Sawada¹, Hiromasa Yabe⁵, Mika Ishige⁶, Hideo Mugishima⁶, Shunichi Kato⁵, Efficacy of hematopoietic stem cell transplantation versus enzyme replacement therapy on intelligence in the patients with Mucopolysaccharidosis type II, ACIM2013.11
- 53) Mohammad Arif Hossain, Chaperone therapy for Krabbe disease; Japanese late-onset mutations can be treated effectively by NOEV、第2回シヤペロン療法研究会、2013.12

G. 知的財産権の出願・登録状況

特許申請

特願 2011-101560 (発明の名称: リソソーム病治療用医薬組成物) として 2011 年 4 月 28 日に出願

研究成果一覽

原著論文

衛藤義勝、井田博幸、大橋十也、小林博司

- Takenori D, Takeo I, Minami M, Masahiro E, Toya O, Yoshikatu E. : A practical fluorometric assay method to measure lysosomal acid lipase activity in dried blood spots for the screening of cholesteryl ester storage disease and Wolman disease. *Mol Genet Metab*. Available online 16 November 2013
- Sato Y, Fujiwara M, Kobayashi H, Ida H. Massive accumulation of glycosaminoglycans in the aortic valve of a patient with hunter syndrome during enzyme replacement therapy. *Pediatr Cardiol*. 2013 Dec;34(8):2077-9.
- Akiyama K, Shimada Y, Higuchi T, Ohtsu M, Nakauchi H, Kobayashi H, Fukuda T, Ida H, Eto Y, Crawford BE, Brown JR, Ohashi T. Enzyme augmentation therapy enhances the therapeutic efficacy of bone marrow transplantation in mucopolysaccharidosis type II mice. *Mol Genet Metab*. 2013 Sep 21.
- Kawagoe S, Higuchi T, Otaka M, Shimada Y, Kobayashi H, Ida H, Ohashi T, Okano HJ, Nakanishi M, Eto Y. Morphological features of iPS cells generated from Fabry disease skin fibroblasts using Sendai virus vector (SeVdp). *Mol Genet Metab*. 2013 Aug;109(4):386-9.
- Sato T, Ikeda M, Yotsumoto S, Shimada Y, Higuchi T, Kobayashi H, Fukuda T, Ohashi T, Suda T, Ohteki T. Novel interferon-based pre-transplantation conditioning in the treatment of a congenital metabolic disorder. *Blood*. 2013 Apr 18;121(16):3267-73.
- J. Ito, T. Saito, C. Numakura, A. Iwaba, S. Sugahara, R. Ishii, C. Sato, H. Haga, K. Okumoto, Y. Nishise, H. Watanabe, H. Ida, K. Hayasaka, H. Togashi, S. Kawata, Y. Ueno: A Case of Adult Type1 Gaucher Disease Complicated by Temporal Intestinal Hemorrhage. *Case Rep Gastroenterol* 2013;7:340-346
- Higuchi T, Shimizu H, Fukuda T, Kawagoe S, Matsumoto J, Shimada Y, Kobayashi H, Ida H, Ohashi T, Morimoto H, Hirato T, Nishino K, Eto Y. : Enzyme replacement therapy (ERT) procedure for mucopolysaccharidosis type II (MPS II) by intraventricular administration (IVA) in murine MPS II. *Mol Genet Metab*. 2012 Sep;107(1-2):122-8. Epub 2012 May 18.
- Kobayashi M, Ohashi T, Fukuda T, Yanagisawa T, Inomata T, Nagaoka T, et al. No accumulation of globotriaosylceramide in the heart of a patient with the E66Q mutation in the α -Galactosidase A gene. *Mol Genet Metab*. 2012; 107:711-715.
- Sato Y, Fujiwara M, Kobayashi H, Ida H. Massive Accumulation of glycosaminoglycans in the aortic valve of a patient with Hunter syndrome during enzyme replacement therapy. *Pediatric Cardiology* (in press)
- Ohashi T, Iizuka S, Shimada Y, Higuchi T, Eto Y, Ida H, et al. Administration of anti-CD3 antibodies modulates the immune response to an infusion of alpha-glucosidase in mice. *Mol Ther* 2012;20:1924-1931.
- Nishiyama Y, Shimada Y, Yokoi T, Kobayashi H, Higuchi T, Eto Y, (Ohashi T) et al. Akt inactivation induces endoplasmic reticulum stress-independent autophagy in fibroblasts from patients with Pompe disease. *Mol Genet Metab* 2012;107:490-495.
- Kawagoe S, Higuchi T, Meng XL, Shimada Y, Shimizu H, Hirayama R, Fukuda T, Chang H, Nakahata T, Fukada S, Ida H, Kobayashi H, Ohashi T, Eto Y. : Generation of induced pluripotent stem (iPS) cells derived from a murine model of Pompe disease and differentiation of Pompe-iPS cells into skeletal muscle cells. *Mol Genet Metab*. 2011 Sep-Oct; 104(123-8). Epub 2011 Jun 2.
- Yoko Kato, Masaharu Akiyama, Fumiyuki Itoh, Hiroyuki Ida : A Study Investigating the Need and Impact

of Pediatric Palliative Care Education on Undergraduate Medical Students in Japan. *Journal of Palliative Medicine* 2011 ; 14(5) : 560-2

- Ohashi T, Iizuka S, Shimada Y, Eto Y, Ida H, Hachimura S, Kobayashi H. Oral administration of recombinant human acid α -glucosidase reduces specific antibody formation against enzyme in mouse. *Mol Genet Metab.* 2011;103(1):98-100. .
- Yokoi T, Kobayashi H, Shimada Y, Eto Y, Ishige N, Kitagawa T, Otsu M, Nakauchi H, Ida H, Ohashi T. Minimum requirement of donor cells to reduce the glycolipid storage following bone marrow transplantation in a murine model of Fabry disease. *J Gene Med.* 2011;13(5):262-8.
- Shimada Y, Kobayashi H, Kawagoe S, Aoki K, Kaneshiro E, Shimizu H, Eto Y, Ida H, Ohashi T. Endoplasmic reticulum stress induces autophagy through activation of p38 MAPK in fibroblasts from Pompe disease patients carrying c.546 G> T mutation. *Mol Genet Metab.* 2011;104(4):566-73.
- Shimada Y, Nishida H, Nishiyama Y, Kobayashi H, Higuchi T, Eto Y, Ida H, Ohashi T. Proteasome inhibitors improve the function of mutant lysosomal alpha-glucosidase in fibroblasts from Pompe disease patient carrying c.546G>T mutation. *Biochem. Biophys. Res. Commun.*, 2011;415(2):274-8.
- Kobayashi H, Shimada Y, Ikegami M, Kawai T, Sakurai K, Urashima T, Ijima M, Fujiwara M, Kaneshiro E, Ohashi T, Eto Y, Ishigaki K, Osawa M, Kyosen SO, Ida H. Pathology of the first autopsy case diagnosed as mucopolidosis type III α /beta suggesting autophagic dysfunction. *Mol Genet Metab.* 2011;102(2):170-5
- Eto Y. : Single gene disorder: recent advances of research. *Nippon Rinsho.* 2010 Aug; 68 Suppl 8:117-28. Japanese.
- Xing-Li Meng, Jin-Song Shen, Shiho Kawagoe, Toya Ohashi, Roscoe O Brady, Yoshikatu Eto : Induced pluripotent stem cells derived from mouse models of lysosomal storage disorders. *PNAS* 2010 vol.17(17):7886-7891.
- Tajima A, Ohashi T, Hamano S, Higurashi N, Ida H. Gaucher Disease Patient With Myoclonus Epilepsy and a Novel Mutation *Pediatr Neurol* 2010;42:65-68
- Kobayashi H, Shimada Y, Ikegami M, Kawai T, Sakurai K, Urashima T, Ijima M, Fujiwara M, Kaneshiro E, Ohashi T, Eto Y, Ishigaki K, Osawa M, Kyosen SO, Ida H. Prognostic factors for the late onset Pompe disease with enzyme replacement therapy: from our experience of 4 cases including an autopsy case. *Mol Genet Metab.* 2010; 100(1):14-9.
- Kobayashi H, Shimada Y, Ikegami M, Kawai T, Sakurai K, Urashima T, Ijima M, Fujiwara M, Kaneshiro E, Ohashi T, Eto Y, Ishigaki K, Osawa M, Kyosen SO, Ida H. Pathology of the first autopsy case diagnosed as mucopolidosis type III α /beta suggesting autophagic dysfunction. *Mol Genet Metab.* 2011;102(2):170-5

田中あけみ

- Hwu WL, Okuyama T, But WM, Estrada S, Gu X, Hui J, Kosuga M, Lin SP, Ngu LH, Shi H, Tanaka A, Thong MK, Wattanasirichaigoon D, Wasant P, McGill J, Current diagnosis and management of mucopolysaccharidosis VI in the Asia-Pacific region. *Mol. Genet. Metab.* 107: 136-44, 2012
- Tanaka A, Okuyama T, Suzuki Y, Sakai N, Takakura H, Sawada T, Tanaka T, Otomo T, Ohashi T,

Ishige-Wada M, Yabe H, Ohura T, Suzuki N, Kato K, Adachi S, Kobayashi R, Mugishima H, Kato S, Long-term efficacy of hematopoietic stem cell transplantation on brain involvement in patients with mucopolysaccharidosis type II: A nationwide survey in Japan. *Mol. Genet. Metab.* 107:513-20, 2012

島田 隆

- Sugano, H., Matsumoto, T., Miyake, K., Watanabe, A., Iijima, O., Migita, M., Narisawa, S., Millán, J.L., Fukunaga, Y., Shimada, T. (2012) Successful gene therapy *in utero* for lethal murine hypophosphatasia. *Hum. Gene Ther.* 23:399-406
- Tamai, H., Miyake, K., Yamaguchi, H., Takatori, M., Dan, K., Inokuchi, K., Shimada, T. (2012) AAV-8 vector expressing IL-24 efficiently suppresses tumor growth mediated by specific mechanisms in MLL/AF4-positive ALL model mice. *Blood* 119:64-71
- Takeichi, N., Midorikawa, S., Watanabe, A., Naing, B.T., Tamura, H., Kano, T., Sugihara, H., Nissato, S., Saito, Y., Aita, Y., Ishii, K., Igarashi, T., Kawakami, Y., Hara, H., Ikeda, T., Shimizu, K., Suzuki, S., Shimano, H., Kawamoto, M., Shimada, T., Watanabe, T., Oikawa, S., Takekoshi, K. (2012) Identical germline mutations in the *TMEM127* gene in 2 unrelated Japanese patients with bilateral pheochromocytoma. *Clin Endocrinol.* 77:707-714
- Masuno, M., Watanabe, A., Naing, B.T., Shimada, T., Fujimoto, W., Ninomiya, S., Ueda, Y., Kondo, E., Yamanouchi, Y., Ouchi, K., Kuroki, Y. (2012) Ehlers-Danlos Syndrome, Vascular Type: A Novel Missense Mutation in *COL3A1* Gene. *Congenit Anom.* 52:207-210
- Miyake, K., Miyake, N., Yamazaki, Y., Shimada, T., Hirai, Y. (2012) Serotype-independent method of recombinant adeno-associated virus (AAV) vector production and purification. *J Nippon Med Sch.* 79:394-402
- Yamamoto, S., Orimo, H., Matsumoto, T., Iijima, O., Narisawa, S., Maeda, T., Millán, J., Shimada, T. (2011) Prolonged survival and phenotypic correction of *Akp2*^{-/-} hypophosphatasia mice by lentiviral gene therapy. *J. Bone Miner. Res.* 26:135-142
- Kato, S., Kobayashi, K., Inoue, K., Kuramochi, M., Okada, T., Yaginuma, H., Morimoto, K., Shimada, T., Takada, M., Kobayashi, K. (2011) A lentiviral strategy for highly efficient retrograde gene transfer by pseudotyping with fusion envelope glycoprotein. *Hum. Gene Ther.* 22:197-206
- Kubodera, T., Yamada, H., Anzai, M., Ohira, S., Yokota, S., Hirai, Y., Mochizuki, H., Shimada, T., Mitani, T., Mizusawa, H., Yokota, T. (2011) *In Vivo* Application of an RNAi Strategy for the Selective Suppression of a Mutant Allele. *Hum. Gene Ther.* 22:27-34
- Watanabe A, Karasugi T, Sawai H, Banyar Tang Naing, Ikegawa S, Orimo H, Shimada S. (2011) Prevalence of c.1559delT in ALPL, a common mutation resulting in the perinatal (lethal) form of hypophosphatasia in Japanese and effects of the mutation on heterozygous carriers. *J Hum Genet* 56:166-168
- Mayra, A., Tomimitsu, H., Kubodera, T., Kobayashi, M., Piao, W., Sunaga, F., Hirai, Y., Shimada, T., Mizusawa, H., Yokota, T. (2011) Intraperitoneal AAV9-shRNA inhibits target expression in neonatal skeletal and cardiac muscles. *Biochem. Biophys. Res. Commun.* 405:204-209

- Naing, B.T., Watanabe, A., Shimada, T. (2011) A novel mutation screening system for Ehlers-Danlos Syndrome, vascular type by high-resolution melting curve analysis in combination with small amplicon genotyping using genomic DNA. *Biochem. Biophys. Res. Commun.* 405:368-372
- Yasuda, T., Hayakawa, H., Nihira, T., Ren, Y.R., Nakata, Y., Nagai, M., Hattori, N., Miyake, K. Takada, M. Shimada, T., Mizuno, Y., Mochizuki, H. (2011) Parkin-Mediated Protection of Dopaminergic Neurons in a Chronic MPTP-Minipump Mouse Model of Parkinson Disease. *J Neuropathol Exp Neurol.* 70:686-697
- Miyake, N., Miyake, K., Yamamoto, M., Hirai, Y., Shimada, T. (2011) Global gene transfer into the CNS across the BBB after neonatal systemic delivery of single-stranded AAV vectors. *Brain Res.* 1389:19-26
- Matsumoto, T., Miyake, K., Yamamoto, S., Orimo, H., Miyake, N., Odagaki, Y., Adachi, K., Iijima, O., Narisawa, S., Millán, J.L., Fukunaga, Y., Shimada, T. (2011) Rescue of Severe Infantile Hypophosphatasia Mice by AAV Mediated Sustained Expression of Soluble Alkaline Phosphatase. *Hum. Gene Ther.* 22:1355-1364
- Isotani, M., Miyake, K., Miyake, N., Hirai, Y., Shimada, T. (2011) Direct comparison of four adeno-associated virus serotypes in mediating the production of anti-angiogenic proteins in mouse muscle. *Cancer Invest.* 29:353-359
- Kato, S., Kuramochi, M., Takasumi, K., Kobayashi, K., Inoue, K., Takahara, D., Hitoshi, S., Takenaka, K., Shimada, T., Takada, M., Kobayashi, K. (2011) Neuron-Specific Gene Transfer through Retrograde Transport of Lentiviral Vector Pseudotyped with a Novel Type of Fusion Envelope Glycoprotein. *Hum. Gene Ther.* 22:1511-1523
- Nihira, T., Yasuda, T., Hirai, Y., Shimada, T., Mizuno, Y., Mochizuki, H. (2011) Adeno-associated viral vector-mediated gene transduction in mesencephalic slice culture. *J Neurosci Methods.* 201:55-60
- Tamai, H., Miyake, K., Takatori, M., Miyake, N., Yamaguchi, H., Dan, K., Shimada, T., Inokuchi, K. (2011) Activated K-Ras protein accelerates human MLL/AF4-induced leukemo-lymphomogenicity in a transgenic mouse model. *Leukemia* 25:888-891
- Tamai, H., Miyake, K., Yamaguchi, H., Inokuchi, K., Dan, K., Shimada, T. (2011) Resistance of MLL/AF4-positive acute lymphoblastic leukemia to tumor necrosis factor-alpha is mediated by S100A6 up-regulation. *Blood Cancer J.* 1:e38
- Takasu, K., Sakai, A., Hanawa, H., Shimada, T., Suzuki, H. (2011) Overexpression of GDNF in the Uninjured DRG Exerts Analgesic Effects on Neuropathic Pain Following Segmental Spinal Nerve Ligation in Mice. *J. Pain* 12:1130-1139
- Fujita, A., Migita, M., Ueda, T., Ogawa, R., Fukunaga, Y., Shimada, T. (2010) Hematopoiesis in regenerated bone marrow within hydroxyapatite scaffold. *Pediatr. Res.* 68:35-40
- Igarashi, T., Miyake, K., Masuda, I., Takahashi, H., Shimada, T. (2010) Adeno-associated vector (type 8)-mediated expression of soluble Flt-1 efficiently inhibits neovascularization in a murine choroidal neovascularization model. *Hum. Gene Ther.* 21:631-637
- Miyake, N., Miyake, K., Karlsson, S., Shimada, T. (2010) Successful treatment of metachromatic leukodystrophy using bone marrow transplantation of HoxB4 overexpressing cells. *Mol. Ther.* 18:1373-1378

- Yasuniwa, Y., Izumi, H., Wang, K.-Y., Shimajiri, S., Sasaguri, Y., Kawai, K., Kasai, H., Shimada, T., Miyake, K., Kashiwagi, E., Hirano, G., Kidani, A., Akiyama, M., Han, B., Wu, Y., Ieiri, I., Higuchi, S., Kohno, K. (2010) Circadian disruption accelerates tumor growth and angio/stromagenesis through a Wnt signaling pathway. *PloS ONE* 5:e15330

酒井規夫

- Hara M, Inokuchi T, Taniwaki T, Otomo T, Sakai N, Matsuishi T, Yoshino M., An adult patient with mucopolipidosis III alpha/beta presenting with parkinsonism., *Brain Dev.* 35(5):462-5, 2013
- Tokushige SI, Sonoo T, Maekawa R, Shiota Y, Hanajima R, Terao Y, Matsumoto H, Hossain MA, Sakai N, Shiio Y., Isolated pyramidal tract impairment in the central nervous system of adult-onset Krabbe disease with novel mutations in the GALC gene., *Brain Dev.* 35(6):579-81, 2013
- Chang B, Gorbea C, Lezin G, Li L, Shan L, Sakai N, Kogaki S, Otomo T, Okinaga T, Hamaoka A, Yu X, Hata Y, Nishida N, Yost HJ, Bowles NE, Brunelli L, Ichida F., 14-3-3ε Gene variants in a Japanese patient with left ventricular noncompaction and hypoplasia of the corpus callosum., *Gene.* 515(1):173-80, 2013
- Kardas F, Uzak AS, Hossain A, Sakai N, Canpolat M, Yikılmaz A., A novel homozygous GALC mutation: Very early onset and rapidly progressive Krabbe disease., *Gene.* 517(1):125-7, 2013
- Eto K, Sakai N, Shimada S, Shioda M, Ishigaki K, Hamada Y, Shinpo M, Azuma J, Tominaga K, Shimojima K, Ozono K, Osawa M, Yamamoto T., Microdeletions of 3p21.31 characterized by developmental delay, distinctive features, elevated serum creatine kinase levels, and white matter involvement., *Am J Med Genet A.* 2013 Dec;161(12):3049-56.
- Hossain MA, Otomo T, Saito S, Ohno K, Sakuraba H, Hamada Y, Ozono K, Sakai N., Late-onset Krabbe disease is predominant in Japan and its mutant precursor protein undergoes more effective processing than the infantile-onset form., *Gene.* 2013 Nov 16. doi:pii: S0378-1119(13)01515-1. 10.1016/j.gene.2013.11.003.
- Akagi M, Mohri I, Iwatani Y, Kagitani-Shimono K, Okinaga T, Sakai N, Ozono K, Taniike M., Clinicogenetical features of a Japanese patient with giant axonal neuropathy., *Brain Dev.* 34(2): 156-62, 2012
- Otomo T, Hossain MA, Ozono K, Sakai N., Genistein reduces heparan sulfate accumulation in human mucopolipidosis II skin fibroblasts., *Mol Genet Metab.* 105(2): 266-9, 2012
- Lee T, Takeshima Y, Okizuka Y, Hamahira K, Kusunoki N, Awano H, Yagi M, Sakai N, Matsuo M, Iijima K., A Japanese child with geleophysic dysplasia caused by a novel mutation of FBN1. *Gene.* 2012 Nov 2. pii: S0378-1119(12)01341-8. [Epub ahead of print]
- Otomo T, Higaki K, Nanba E, Ozono K, Sakai N., Lysosomal storage causes cellular dysfunction In mucopolipidosis II skin fibroblasts., *J Biol Chem.* 286(40): 35283-90, 2011
- Hashimoto N, Kagitani-Shimono K, Sakai N, Otomo T, Tominaga K, Nabatame S, Mogami Y, Takahashi Y, Imai K, Yanagihara K, Okinaga T, Nagai T, Taniike M and Ozono K, SLC2A1 gene analysis of Japanese patients with glucose transporter 1 deficiency syndrome, *J Hum Genet* 56(12): 846-51, 2011
- Miyoshi Y, Sakai N, Hamada Y, Tachibana M, Hasegawa Y, Kiyohara Y, Yamada H, Murakami M, Kondou H, Kimura-Ohba S, Mine J, Sato T, Kamio N, Ueda H, Suzuki Y, Shiomi M, Ohta H, Shimozawa

N, Ozono K., Clinical aspects and adrenal functions in eleven Japanese children with X-linked adrenoleukodystrophy., *Endocr J.* 57(11): 965-72, 2010

- Terao M, Sakai N, Higashiyama S, Kotobuki Y, Tanemura A, Wataya-Kaneda M, Yutsudo M, Ozono K, Katayama I., Cutaneous symptoms in a patient with cardiofaciocutaneous syndrome and increased ERK phosphorylation in skin fibroblasts., *Br J Dermatol.* 2010 Oct;163(4):881-4.

高橋 勉

- Takamura A, Sakai N, Shinpo M, Noguchi A, Takahashi T, Matsuda S, Yamamoto M, Narita A, Ohno K, Ohashi T, Ida H, Eto Y: The useful preliminary diagnosis of Niemann-Pick disease type C by filipin test in blood smear. *Mol Genet metab* 111: 401-404, 2013
- Sato, Y., Ishida-Nakajima, W., Kawamura, M., Miura, S., Oguma, R., Arai, H., Takahashi, T. Hypoxia-ischemia induces hypo-phosphorylation of collapsing response mediator protein 2 in a neonatal rat model of periventricular leukomalacia(2011) *Brain Res*, 1386, 165-174.

高柳正樹

- Wakiya T, Sanada Y, Urahashi T, Ihara Y, Yamada N, Okada N, Ushijima K, Otomo S, Sakamoto K, Murayama K, Takayanagi M, Hakamada K, Yasuda Y, Mizuta K. Impact of enzyme activity assay on indication in liver transplantation for ornithine transcarbamylase deficiency. *Mol Genet Metab.* 2012 Mar;105(3):404-7. Epub 2012 Jan 4
- Purevsuren J, Kobayashi H, Hasegawa Y, Yamada K, Takahashi T, Takayanagi M, Fukao T, Fukuda S, Yamaguchi S. Intracellular in vitro probe acylcarnitine assay for identifying deficiencies of carnitine transporter and carnitine palmitoyltransferase-1. *Anal Bioanal Chem.* 2012
- Kido J, Nakamura K, Mitsubuchi H, Ohura T, Takayanagi M, Matsuo M, Yoshino M, Shigematsu Y, Yorifuji T, Kasahara M, Horikawa R, Endo F. Long-term outcome and intervention of urea cycle disorders in Japan. *J Inherit Metab Dis.* 2011
- Sanayama Y, Nagasaka H, Takayanagi M, Ohura T, Sakamoto O, Ito T, Ishige-Wada M, Usui H, Yoshino M, Ohtake A, Yorifuji T, Tsukahara H, Hirayama S, Miida T, Fukui M, Okano Y. Experimental evidence that phenylalanine is strongly associated to oxidative stress in adolescents and adults with phenylketonuria. *Mol Genet Metab.* 103:220-5, 2011 Epub 2011 Mar 29.
- Hori T, Fukao T, Kobayashi H, Teramoto T, et al. Carnitine palmitoyltransferase 2 deficiency: the time-course of blood and urinary acylcarnitine levels during initial L-carnitine supplementation. *Tohoku J Exp Med.* 2010;221(3):191-5.
- Fukao T, Horikawa R, Naiki Y, et al. A novel mutation in an exonic splicing enhancer results in exon 10 skipping in the human mitochondrial acetoacetyl-CoA thiolase gene. *Mol Genet Metab.* 2010 Aug;100(4):339-44.
- Kimura A, Kage M, Nagata I, et al. Histological findings in the livers of patients with neonatal intrahepatic cholestasis caused by citrin deficiency. *Hepatol Res.* 2010 Mar 1;40(3):295-303.
- Fukao T, Ishii T, Amano N, Kursula P, et al. A neonatal-onset succinyl-CoA:3-ketoacid CoA transferase (SCOT)-deficient patient with T435N and c.658-666dupAACGTGATT p.N220_I222dup mutations in the OXCT1 gene. *J Inherit Metab Dis.* 2010

大野耕策、成田 綾

- ○Suzuki Y, Ichinomiya S, Kurosawa M, Matsuda J, Ogawa S, Iida M, Kubo T, Tabe M, Itoh M, Higaki K, Nanba E, Ohno K. Therapeutic chaperone effect of N-octyl 4-epi-β-valienamine on murine G(M1)-gangliosidosis. *Mol Genet Metab.* 106(1):92-8, 2012
- ○Aguilar-Moncayo M, Takai T, Higaki K, Mena-Barragán T, Hirano Y, Yura K, Li L, Yu Y, Ninomiya H, García-Moreno MI, Ishii S, Sakakibara Y, Ohno K, Nanba E, Ortiz Mellet C, García Fernández JM, Suzuki Y. Tuning glycosidase inhibition through aglycone interactions: pharmacological chaperones for Fabry disease and GM1 gangliosidosis. *Chem Commun (Camb).* 48(52):6514-6,2012
- ○Luan Z, Li L, Higaki K, Nanba E, Suzuki Y, Ohno K. The chaperone activity and toxicity of ambroxol on Gaucher cells and normal mice. *Brain Dev*35: 317-322, 2013
- ○Castilla J, Rísquez R, Cruz D, Higaki K, Nanba E, Ohno K, Suzuki Y, Díaz Y, Ortiz Mellet C, García Fernández JM, Castellón S. Conformationally-Locked N-Glycosides with Selective β-Glucosidase Inhibitory Activity: Identification of a New Non-Iminosugar-Type Pharmacological Chaperone for Gaucher Disease. *J Med Chem.* 55(15):6857-65,2012
- ○Takamura A, Higaki K, Ninomiya H, Takai T, Matsuda J, Iida M, Ohno K, Suzuki Y, Nanba E. Lysosomal accumulation of Trk protein in brain of G(M1) -gangliosidosis mouse and its restoration by chemical chaperone. *J Neurochem.* 118(3):399-406, 2011
- ○Higaki K, Li L, Udin Bahrudin, Okuzawa S, Takamura A, Yamamoto K, Adachi K, Rubigilda C. Paraguison, Takai T, Ikehata H, Tominaga L, Hisatome I, Iida M, Ogawa S, Matsuda J, Ninomiya H, Sakakibara Y, Ohno K, Suzuki Y, Nanba E. Chemical chaperone therapy: chaperone effect on mutant enzyme and cellular pathophysiology in β-galactosidase deficiency. *Human Mutation.* 32(7):843-52,2011
- ○Ohno K. Kunihiro Suzuki and sphingolipidoses. *J Biochem* 150(6):597-605,2011
- ○Luan Z, Ninomiya H, Ohno K, Ogawa S, Kubo T, Iida M, Suzuki Y. The effect of N-octyl-β-valienamine on β-glucosidase activity in tissues of normal mice. *Brain Dev* 32(10):805-9,2010
- ○Luan Z, Higaki K, Aguilar-Moncayo M, Li L, Ninomiya H, Nanba E, Ohno K, Garcia-Moreno M, Mellet CO, Fernandez JMG, Suzuki Y. A fluorescent SP2-iminosugar with pharmacological chaperone activity for Gaucher disease: Cellular uptake and intracellular distribution studies. *ChemBioChem* 11(17):2453-64,2010
- ○Li L, Higaki K, Ninomiya H, Luan Z, Iida M, Ogawa S, Suzuki Y, Nanba E, Ohno K. Chemical chaperone therapy: Luciferase assay for screening of β-galactosidase mutations. *Mol Genet Metab* 101(4) : 364-369,2010

辻 省二

- Matsukawa T, Asheuer M, Takahashi Y, Goto J, Suzuki Y, Shimozawa N, Takano H, Onodera O, Nishizawa M, Aubourg P and Tsuji S. Identification of novel SNPs of ABCD1, ABCD2, ABCD3 and ABCD4 genes in patients with X-linked adrenoleukodystrophy (ALD) based on comprehensive resequencing and association studies with ALD phenotypes. *Neurogenetics* 12: 41-50, 2011

難波 栄二

- X Luan Z, Li L, Higaki K, Nanba E, Suzuki Y, Ohno K, The chaperone activity and toxicity of ambroxol on GD cells and normal mice. *Brain Dev*, 2013 35: 317-322.
- ○Takai T, Higaki K, Aguilar-Moncayo M, Mena-Barragán T, Hirano Y, Yura K, Yu L, Ninomiya H,

García-Moreno I, Sakakibara Y, Ohno K, Nanba E, Ortiz Mellet C, García Fernández JM, Suzuki Y, A bicyclic 1-deoxygalactonojirimycin derivative as a novel pharmacological chaperone for GM₁ gangliosidosis. *Mol Ther*, 2013 21: 526-532.

- Xiong H, Higaki K, Wei C, Bao CW, Zhanf YH, Fu N, Qui J, Adachi K, Kumura Y, Ninomiya H, Nanba E, Wu XR. Genotype/phenotype of 6 Chinese cases with Niemann-Pick disease type C. *Gene*, 498, 332-335, 2012
- ○Suzuki Y, Ichinomiya S, Kurosawa M, Matsuda J, Ogawa S, Iida M, Kubo T, Tabe M, Itoh M, Higaki K, Nanba E, Ohno K: Therapeutic chaperone effect of *N*-octyl 4-epi-β-valienamine on murine GM₁-gangliosidosis. *Mol Genet Metab* 106: 92-8, 2012.
- ○Aguilar-Moncayo M, Takai T, Higaki K, Mena-Barragán T, Hirano Y, Yura K, Li L, Yu Y, Ninomiya H, García-Moreno I, Ishii S, Sakakibara Y, Ohno K, Nanba E, Ortiz Mellet C, García Fernández JM, Suzuki Y: Tuning glycosidase inhibition through aglycone interactions: Pharmacological chaperones for Fabry disease and GM₁ gangliosidosis, *Chem Commun* 48: 6514-6, 2012.
- Castilla J, Riquez R, Cruz D, Higaki K, Nanba E, Ohno K, Suzuki Y, Diaz Y, Ortiz Mellet C, Garcia Fernandez JM, Castillon S: Conformationally-locked N-glycosides with selective β-glucosidase inhibitory activity: Identification of a new non-iminosugar-type pharmacological chaperone for Gaucher disease. *J Med Chem* 55:6857-65, 2012.
- ○Higaki K, Li L, Bahrudin U, Okuzawa S, Takamura A, Yamamoto K, Adachi K, Paraguison RC, Takai T, Ikehata H, Tominaga L, Hisatome I, Iida M, Ogawa S, Matsuda J, Ninomiya H, Sakakibara Y, Ohno K, Suzuki Y, Nanba E. Chemical chaperone therapy: chaperone effect on mutant enzyme and cellular pathophysiology in β-galactosidase deficiency. *Hum Mutat*, 32, 843-852, 2011
- ○Takamura A, Higaki K, Ninomiya H, Takai T, Matsuda J, Iida M, Ohno K, Suzuki Y, Nanba E. Lysosomal accumulation of Trk protein in brain of GM1-gangliosidosis mouse and its restoration by chemical chaperone. *J Neurochem*, 118, 399-406, 2011
- ○Li L, Higaki K, Ninomiya H, Luan Z, Iida M, Ogawa S, Suzuki Y, Ohno K, Nanba E. Chemical chaperone therapy: Luciferase assay for screening of β-galactosidase mutations. *Mol Genet Metab*, 101, 354-369, 2010
- Luan Z, Higaki K, Aquilar-Moncayo M, Li L, Ninomiya H, Nanba E, Ohno K, García-Moreno MI, Ortiz Mellet C, García Fernández JM, Suzuki Y, A fluorescent sp²-iminosugar with pharmacological chaperone activity for Gaucher disease: Synthesis and intracellular distribution studies. *ChemBioChem*, 11, 2453-2463, 2010

鈴木康之

- Yasuda E, Suzuki Y, Tomatsu S et al. Pathogenesis of Morquio A syndrome: an autopsied case reveals systemic storage disorder. *Mol Genet Metab*. 2013 ;109:301-11
- Pravin Patel, Yasuyuki Suzuki², Miho Maeda, Eriko Yasuda, Tsutomu Shimada, Kenji E. Orii , Tadao Orii, Shunji Tomatsu. Growth charts for patients with Hunter Syndrome. *Molecular Genetics and Metabolism* (in press)
- Tomatsu S, Montaña AM, Oikawa H, Giugliani R, Harmatz P, Smith M, Suzuki Y, and Orii T. Impairment

of Body Growth in Mucopolysaccharidoses. Preedy VR ed. Handbook of Growth and Growth Monitoring in Health and Disease. Springer, 2012

- Tomatsu S, Mackenzie WG, Theroux MC, Mason RW, Thacker MM, Shaffer TH, Montaña AM, Rowan D, Sly W, Alméjiga-Díaz CJ, Barrera LA, Chinen Y, Yasuda E, Ruhnke K, Suzuki Y, Orii T. Current and emerging treatments and surgical interventions for Morquio A syndrome: a review. *Research and Reports in Endocrine Disorders* 2012; 265-77
- Hintze JP, Tomatsu S, Fujii T, Montaña AM, Yamaguchi S, Suzuki Y, Fukushi M, Ishimaru T, Orii T. Comparison of liquid chromatography-tandem mass spectrometry and sandwich ELISA for determination of keratan sulfate in plasma and urine. *Biomark Insights*. 2011;6:69-78.
- Muenzer J, Beck M, Giugliani R, Suzuki Y, Tytki-Szymanska A, Valayannopoulos V, Vellodi A, Wraith JE. Idursulfase treatment of Hunter syndrome in children younger than 6 years: results from the Hunter Outcome Survey. *Genet Med*. 2011 Feb;13(2):102-9.
- Tomatsu S, Montaña AM, Oikawa H, Smith M, Barrera L, Chinen Y, Thacker MM, Mackenzie WG, Suzuki Y, Orii T. Mucopolysaccharidosis type IVA (Morquio A disease): clinical review and current treatment. *Curr Pharm Biotechnol*. 2011 Jun;12(6):931-45.
- Morita M, Shimozawa N, Kashiwayama Y, Suzuki Y, Imanaka T. ABC subfamily D proteins and very long chain fatty acid metabolism as novel targets in adrenoleukodystrophy. *Curr Drug Targets*. 2011 May;12(5):694-706.
- Tomatsu S, Montaña AM, Oguma T, Dung VC, Oikawa H, Gutiérrez ML, Yamaguchi S, Suzuki Y, Fukushi M, Barrera L, Orii T. Validation of disaccharide composition derived from dermatan sulfate and heparan sulfate in mucopolysaccharidoses and mucopolipidoses II and III by tandem mass spectrometry. *Molecular Genetics and Metabolism* 2010;99:124-131
- Tomatsu S, Montaña AM, Oguma T, Dung VC, Oikawa H, Carvalho TG, Gutiérrez MG, Yamaguchi S, Suzuki Y, Fukushi M, Kida K, Kubota M, Kida K, Kubota M, Orii T. Validation of keratan sulfate level in Mucopolysaccharidosis IVA by liquid tandem mass spectrometry method. *J Inherit Metab Dis* (Epub 2010).
- Tomatsu S, Montaña AM, Oguma T, Dung VC, Oikawa H, de Carvalho TG, Gutiérrez ML, Yamaguchi S, Suzuki Y, Fukushi M, Sakura N, Barrera L, Kida K, Kubota M, Orii T. Dermatan sulfate and heparan sulfate as a biomarker for mucopolysaccharidosis I. *J Inherit Metab Dis*. 2010; 33:141-150

櫻庭 均

- Nakano S, Morizane Y, Makisaka N, et al : Development of a highly sensitive immuno-PCR assay for the measurement of α -galactosidase A protein levels in serum and plasma. *PLoS ONE* 8: e78588, 2013
- Sakuraba H : High risk screening for Fabry disease. Asian Congress for Lysosomal Storage Disease Screening, Kumamoto, Japan, May. 2013
- Sakuraba H : E66Q: Biochemical, pathological and structural studies. 3rd Update on Fabry Nephropathy: Biomarkers, Progression and Treatment Opportunities, Hong Kong, China, Jun. 2013
- Sakuraba H : Genotype/Phenotype correlation in Fabry disease. The 15th Annual Asia LSD

Symposium, Chiba, Japan, Nov. 2013

- Kawashima I, Mitobe S, Kodama T, Tsukimura T, Togawa T, Sakuraba H: Development of enzyme replacement therapy with a modified enzyme and an activator for Fabry disease. The 3rd Asian Congress for Inherited Metabolic Diseases / The 55th Annual Meeting of The Japanese Society for Inherited Metabolic Diseases. Chiba, Japan, Nov. 2013
- Shibasaki F, Nakano S, Togawa T, Tsukimura T, Kawashima I, Sakuraba H.: Development of a highly sensitive immuno-PCR measurement of α -galactosidase A protein levels in serum and plasma. The 3rd Asian Congress for Inherited Metabolic Diseases / The 55th Annual Meeting of The Japanese Society for Inherited Metabolic Diseases. Chiba, Japan, Nov. 2013
- Nakano S, Togawa T, Tsukimura T, Kawashima I, Sakuraba H, Futoshi S.: Rapid Immunochromatographic measurement of anti- α -galactosidase A antibodies in Fabry patients Treated with enzyme replacement therapy. The 3rd Asian Congress for Inherited Metabolic Diseases / The 55th Annual Meeting of The Japanese Society for Inherited Metabolic Diseases. Chiba, Japan, Nov. 2013
- Togawa T, Tsukimura T, Katayama M, Mitobe S, Sakuraba H.: Fabry patients exhibiting no elevation in plasma globotriaosylsphingosine level. The 3rd Asian Congress for Inherited Metabolic Diseases / The 55th Annual Meeting of The Japanese Society for Inherited Metabolic Diseases. Chiba, Japan, Nov. 2013
- Tsukimura T, Takada M, Aizawa Y, Suzuki T, Katayama M, Sakuraba H, Togawa T.: Comparative study on the content of mannose 6-phosphate residues of recombinant lysosomal enzymes. The 3rd Asian Congress for Inherited Metabolic Diseases / The 55th Annual Meeting of The Japanese Society for Inherited Metabolic Diseases. Chiba, Japan, Nov. 2013
- Itoh K, Tsuji D, Namba K, Yamaguchi S, Imataki I, Ishimaru N, Sakuraba H.: Establishment of human neural cell culture systems induced from ips cells derived from Tay-Sachs disease patient for drug discovery. The 3rd Asian Congress for Inherited Metabolic Diseases / The 55th Annual Meeting of The Japanese Society for Inherited Metabolic Diseases. Chiba, Japan, Nov. 2013
- Kitakaze K, Kawano K, Tsuji D, Asanuma D, Kamiya M, Urano Y, Sakuraba H, Itoh K.: Imaging of enzyme replacement with a novel fluorescent probe and purified lysosomal β -hexosaminidase carrying M6P-type glycans. The 3rd Asian Congress for Inherited Metabolic Diseases / The 55th Annual Meeting of The Japanese Society for Inherited Metabolic Diseases. Chiba, Japan, Nov. 2013
- Tsukimura T, Kawashima I, Togawa T, et al: Efficient uptake of recombinant α -galactosidase A produced with a gene-manipulated yeast by Fabry mice kidneys. *Mol Med* 18: 76-82, 2012
- Togawa T, Tsukimura T, Kodama T, et al: Fabry disease: Biochemical, pathological and structural studies of the α -galactosidase A with E66Q amino acid substitution. *Mol Genet Metab* 105: 615-620, 2012
- Doi K, Noiri E, Ishizu T, et al: High-throughput screening identified disease-causing mutants and functional variants of α -galactosidase A gene in Japanese male hemodialysis patients. *J Hum Genet* 57:

575-579, 2012

- Mitobe S, Togawa T, Tsukimura T, et al: Mutant α -galactosidase A with M296I does not cause elevation of the plasma globotriaosylsphingosine level. *Mol Genet Metab* 107: 623-626, 2012
- Maita N, Taniguchi H, Sakuraba H: Crystallization, X-ray diffraction analysis and SIRAS phasing of human α -L-iduronidase. *Acta Cryst F*68: 1363-1366, 2012
- Mawatari K, Yasuoka H, Oba T, et al: Screening for Fabry disease in patients with left ventricular hypertrophy. *Int J Cardiol*, in press
- Ogawa Y, Tanaka M, Tanabe M, et al: Impaired neural differentiation of induced pluripotent stem cells generated from a mouse model of Sandhoff disease. *PLoS ONE*, in press

芳野 信、渡邊順子

- Ihara K, Yoshino M, Watanabe Y, et al. : Coagulopathy in patients with late-onset ornithine transcarbamylase deficiency in remission state: a previously unrecognized complication. *Pediatrics*. 2013 Jan;131(1):e327-30.
- Miyake N, Yano S, Watanabe Y, et al. : Mitochondrial complex III deficiency caused by a homozygous UQCRC2 mutation presenting with neonatal-onset recurrent metabolic decompensation. *Hum Mutat*. 2013 Mar;34(3):446-52.
- Okano Y, Yoshino M, Watanabe Y, et al. : Fatigue and quality of life in citrin deficiency during adaptation and compensation stage. *Mol Genet Metab*. 2013 May;109(1):9-13.
- Hara M, Matsuishi T, Yoshino M, et al. : An adult patient with mucopolidosis III alpha/beta presenting with parkinsonism. *Brain Dev*. 2013 May;35(5):462-5.
- Miyake N, Watanabe Y, Yoshino M, et al. : MLL2 and KDM6A mutations in patients with Kabuki syndrome. *Am J Med Genet A*. 2013 Sep;161(9):2234-43.
- Higashimoto K, Watanabe Y, Soejima H, et al. : Homozygous deletion of DIS3L2 exon 9 due to non-allelic homologous recombination between LINE-1s in a Japanese patient with Perlman syndrome. *Eur J Hum Genet*. 2013 Nov;21(11):1316-9.
- Nakashima S, Watanabe Y, Ogata T, et al. : Critical role of Yp inversion in PRKX/PRKY-mediated Xp;Yp translocation in a patient with 45,X testicular disorder of sex development. *Endocr J*. 2013 Oct 3. [Epub ahead of print]
- Yatsuki H, Higashimoto K, Jozaki K, Koide K, Okada J, Watanabe Y, Okamoto N, Tsuno Y, Yoshida Y, Ueda K, Shimizu K, Ohashi H, Mukai T, Soejima H. Novel mutations of CDKN1C in Japanese patients with Beckwith-Wiedemann syndrome. *Genes Genom* (2013) 35:141-147
- Higashimoto K, Watanabe Y, Soejima H, et al. : Aberrant methylation of H19-DMR acquired after implantation was dissimilar in soma versus placenta of patients with Beckwith-Wiedemann syndrome. *Am J Med Genet A*. 2012 May 10. *Am J Med Genet A*. 2012 Part A 9999:1-6
- Mitobe S, Yoshino M, Sakuraba H, et al. : Mutant α -galactosidase A with M296I does not cause elevation of the plasma globotriaosylsphingosine level. *Molecular Genetics and Metabolism*, 2012 Jul 14. [Epub ahead of print]
- Okada J, Matsuishi T, Iwata O, et al. : Levothyroxine replacement therapy and refractory hypotension out of transitional period in preterm infants. *Clin Endocrinol* 2011; 74: 354-364. (doi:

10.1111/j.1365-2265.2010.03927.x.)

- Sanayama Y, Yoshino M, Okano Y, et al. : Experimental evidence that phenylalanine is strongly associated to oxidative stress in adolescents and adults with phenylketonuria. *Mol Genet Metab* 2011; 103: 220-225.
- Harada H, Yoshino M, Ikeda H, et al. : Laronidase replacement therapy improves myocardial function in mucopolysaccharidosis I. *Mol Genet Metab* 2011; 103: 215-219.
- Watanabe Y, Yano S, Yoshino M, et al. : A familial case of LEOPARD syndrome associated with a high-functioning autism spectrum disorder. *Brain Dev* 2011; 33: 576-57
- Segawa Y, Watanabe Y, Yoshino M, et al. : A long-term survival case of arginase deficiency with severe multicystic white matter and compound mutations. *Brain Dev* 2011; 33: 45-48.
- Yano S, Bagheri A, Watanabe Y, et al. : Familial Simpson-Golabi-Behmel syndrome: Studies of X-chromosome inactivation and clinical phenotypes in two female individuals with GPC3 mutations. *Clin Genet*. 2011 Nov;80(5):466-71.
- Shimizu N, Yoshino M, Aoki T, et al : Effects of long-term zinc treatment in Japanese patients with Wilson disease: efficacy, stability, and copper metabolism. *Transl Res* 2010; 156: 350-357.
- Fujii C, Umehashi H, Yoshino M, et al. : Attitude to extended use and long-term storage of newborn screening blood spots in Japan. *Pediatr Int* 2010; 52: 393-397.
- Pinner JR, Kirk EP, Yoshino M, et al. : Female heterozygotes for the hypomorphic R40H mutation can have ornithine transcarbamylase deficiency and present in early adolescence: a case report and review of the literature. *J Med Case Reports* 2010; 4: 361.
- Narumi S, Watanabe Y, Hasegawa T, et al. : Various types of LRP5 mutations in four patients with osteoporosis-pseudoglioma syndrome: identification of a 7.2-kb microdeletion by oligonucleotide tiling microarray. *Am J Med Genet A*. 2010 Jan;152A(1):133-40.
- Komatsuzaki S, Watanabe Y, Matsubara Y, et al. : Mutation analysis of the SHOC2 gene in Noonan-like syndrome and in hematologic malignancies. *J Hum Genet*. 2010 Dec;55(12):801-9.

北川照男

- T. Kitagawa. Newborn Screening for Inborn Errors of Metabolism in Japan -A History of the Department of Newborn Screening-. *Pediatric Endocrinology Reviews*, 10; Suppl 1; 8-25, 2012.
- Teruo Kitagawa. Hepatorenal tyrosinemia. *Proc. Jpn. Acad., Ser. B*, Vol.88, 192-200, 2012. Togawa T, Kodama T, Kitagawa T, et al. Plasma globotriaosylsphingosine as a biomarker of Fabry disease. *Mol Genet Metab* 100; 257-261, 2010.

奥山虎之

- T Tajima G, Sakura N, Kosuga M et al. Effects of idursulfase enzyme replacement therapy for Mucopolysaccharidosis type II when started in early infancy: comparison in two siblings. *Mol Genet Metab*. 2013;108:172-7.
- Sasaki T, Niizeki H, Shimizu A, Shiohama A, Hirakiyama A, Okuyama T, Seki A, Kabashima K, Otsuka A, Ishiko A, Tanese K, Miyakawa SI, Sakabe JI, Kuwahara M, Amagai M, Okano H, Suematsu M, Kudoh J. Identification of mutations in the prostaglandin transporter gene SLCO2A1 and its phenotype-genotype

correlation in Japanese patients with pachydermoperiostosis. *J Dermatol Sci.* 2012;68:36-44.

- Hwu WL, Okuyama T, But WM, Estrada S, Gu X, Hui J, Kosuga M, Lin SP, Ngu LH, Shi H, Tanaka A, Thong MK, Wattanasirichaigoon D, Wasant P, McGill J. Current diagnosis and management of mucopolysaccharidosis VI in the Asia-Pacific region. *Mol Genet Metab.*2012;107:136-144.
- D'Aco K, Underhill L, Rangachari L, Arn P, Cox GF, Giugliani R, Okuyama T, Wijburg F, Kaplan P. Diagnosis and treatment trends in mucopolysaccharidosis I: findings from the MPS I Registry. *Eur J Pediatr.*2012;171:911-919
- Furujo M, Kubo T, Kosuga M, Okuyama T. Enzyme replacement therapy attenuates disease progression in two Japanese siblings with mucopolysaccharidosis type VI. *Mol Genet Metab* 2011;104:597-602.
- Oda E, Tanaka T, Migita O, Kosuga M, Fukushi M, Okumiya T, Osawa M, Okuyama T. Newborn Screening for Pompe disease in Japan. *Mol Genet Metab.*2011;104:560-565.
- Furukawa Y, Hamaguchi A, Nozaki I, Iizuka T, Sasagawa T, Shima Y, Demura S, Murakami H, Kawahara N, Okuyama T, Iwasa K, Yamada M. Cervical pachymeningeal hypertrophy as the initial and cardinal manifestation of mucopolysaccharidosis type I in monozygotic twins with a novel mutation in the alpha-l-iduronidase gene. *J Neurol Sci.* 2011;15:121-125.
- Hayashi S, Imoto I, Aizu Y, Okamoto N, Mizuno S, Kurosawa K, Okamoto N, Honda S, Araki S, Mizutani S, Numabe H, Saitoh S, Kosho T, Fukushima Y, Mitsubuchi H, Endo F, Chinen Y, Kosaki R, Okuyama T, Ohki H, Yoshihashi H, Ono M, Takada F, Ono H, Yagi M, Matsumoto H, Makita Y, Hata A, Inazawa J. Clinical application of array-based comparative genomic hybridization by two-stage screening for 536 patients with mental retardation and multiple congenital anomalies. *J Hum Genet.* 2011;56:110-124.
- Kondoh T, Kanno A, Itoh H, Nakashima M, Honda R, Kojima M, Noguchi M, Nakane H, Nozaki H, Sasaki H, Nagai T, Kosaki R, Kakee N, Okuyama T, Fukuda M, Ikeda M, Shibata Y, Moriuchi H. Donepezil significantly improves abilities in daily lives of female Down syndrome patients with severe cognitive impairment: a 24-week randomized, double-blind, placebo-controlled trial. *J Psychiatry Med.* 2011;41:71-89.
- Okuyama T, Tanaka A, Suzuki Y, Ida H, Tanaka T, Cox GF, Eto Y, Orii T. Japan Elaprase((R)) Treatment (JET) study: Idursulfase enzyme replacement therapy in adult patients with attenuated Hunter syndrome (Mucopolysaccharidosis II, MPS II). *Mol Genet Metab.* 2010;99:18-25.

坪井一哉

- Tsuboi, K., S. Suzuki, et al. "Descriptive Epidemiology of Fabry Disease Among Beneficiaries of the Specified Disease Treatment Research Program in Japan." *Journal of Epidemiology* 22(4): 370-374.2012
- Tsuboi, K. and H. Yamamoto (2012). "Clinical observation of patients with Fabry disease after switching from agalsidase beta (Fabrazyme) to agalsidase alfa (Replagal)." *Genetics in Medicine advance online publication* (in press).
- Yamamoto, H. and K. Tsuboi (2012). "Age-influence upon Auditory Dysfunction in Patients with Fabry Disease." *Journal of Transportation Medicine* 64(3,4): 25-29.

松田純子

- Murakami, I, Mitsutake, S, Kobayashi, N, Matsuda, J, Suzuki, A, Shigyo, T, Igarashi, Y.: Improved

high-fat diet-induced glucose intolerance by an oral administration of phytosphingosine. *Biosci. Biotechnol. Biochem.* 77, 194-197. 2013.

- ○ Hisaki H, Matsuda J, Tadano-Aritomi K, Uchida S, Okinaga H, Miyagawa M, Tamamori- Adachi M, Iizuka M, Okazaki T. Primary polydipsia, but not accumulated ceramide, causes lethal renal damage in saposin D-deficient mice. *Am. J. Physiol. Renal Physiol.* (2012) 303:F1049-1059.
- ○ Hojo H, Tanaka H, Hagiwara M, Asahina Y, Ueki A, Katayama H, Nakahara Y, Yoneshige A, Matsuda J, Ito Y, Nakahara Y. Chemoenzymatic synthesis of hydrophobic glycoprotein: synthesis of saposin C carrying complex-type carbohydrate. *J. Org. Chem.* (2012) 77:9437-9446.
- ○ Toyofuku T, Nojima S, Ishikawa T, Takamatsu H, Tsujimura T, Uemura A, Matsuda J, Seki T, Kumanogoh A. Endosomal sorting by Semaphorin 4A in retinal pigment epithelium supports photoreceptor survival. *Genes Dev.* (2012) 26:816-829.
- Suzuki A, Miyazaki M, Matsuda J, Yoneshige A. High-performance thin-layer chromatography/mass spectrometry for the analysis of neutral glycosphingolipids. *Biochimica et Biophysica Acta. (Molecular and Cell Biology of Lipids)* (2011) 1811:861-874.
- ○ Hojo H, Katayama H, Tano C, Nakahara Y, Yoneshige A, Matsuda J, Sohma Y, Kiso Y, Nakahara Y. Synthesis of the sphingolipid activator protein, saposin C, using an azido-protected O-acyl isopeptide as an aggregation-disrupting element. *Tetrahedron Lett.* (2011) 52:635-639.
- ○ Yoneshige A, Suzuki K, Suzuki K, Matsuda J. A Mutation in the Saposin C Domain of the Sphingolipid Activator Protein (Prosaposin) Gene Causes Neurodegenerative Disease in Mice. *J. Neurosci. Res.* (2010) 88: 2118-2134.
- Yoneshige A, Sasaki A, Miyazaki M, Kojima N, Suzuki A, Matsuda J. Developmental changes in glycolipids and synchronized expression of nutrient transporters in the mouse small intestine. *J. Nutr. Biochem.*

遠藤 文夫

- Kido J, Nakamura K, Matsumoto S, Mitsubuchi H, Ohura T, Shigematsu Y, Yorifuji T, Kasahara M, Horikawa R and Endo F. Current status of hepatic glycogen storage disease in Japan: clinical manifestations, treatments and long-term outcomes. *J. Hum. Genet.* 58, 285-292 (2013)
- Lee D, Oka T, Hunter B, Robinson A, Papp S, Nakamura K, Srisakuldee W, Nickel BE, Light PE, Dyck JRB, Lopaschuk GD, Kardami E, Opas M, and Michalak M. Calreticulin induces dilated cardiomyopathy. *Plos One* 8, e56387 (2013)
- Yamamoto A, Nakamura K, Matsumoto S, Iwai M, Shigematsu Y, Tajima G, Tsumura M, Okada S, Mitsubuchi H, Endo F. VLCAD deficiency in a patient who recovered from VF, but died suddenly of an RSV infection. *Pediatr Int.* 55, 775-778 (2013)
- Nakamura K, Sekijima Y, Nakamura K, Hattori K, Nagamatsu K, Shimizu Y, Yazaki M, Sakurai A, Endo F, Fukushima Y, Ikeda S. p.E66Q Mutation in the GLA Gene is Associated with a High Risk of Cerebral Small-Vessel Occlusion in Elderly Japanese Males. *Eur J Neurol* (2013 in press)
- Inoue T, Hattori K, Ihara K, Ishii A, Nakamura K, Hirose S. Newborn screening for

Fabry disease in Japan: Prevalence and genotypes of Fabry disease in a pilot study. *J. Hum. Genet.* (2013 in press)

- Tanaka T, Mochida T, Maki Y, Shiraki Y, Mori H, Matsumoto S, Shimbo K, Ando T, Nakamura K, Endo F, Okamoto M. Interactive network analysis of the plasma amino acids profile in a mouse model of hyperglycemia. *Springerplus.* (2013 in press)
- Fujisawa D, Nakamura K, Mitsubuchi H, Ohura T, Shigematsu Y, Yorifuji T, Kasahara M, Horikawa R and Endo F. Clinical features and management of organic acidemias in Japan. *J. Hum. Genet.* (2013 in press)
- Kido J, Nakamura K, Mitsubuchi H, Ohura T, Takayanagi M, Matsuo M, Yoshino M, Shigematsu Y, Yorifuji T, Kasahara M, Horikawa R, Endo F.; Long-term outcome and intervention of urea cycle disorders in Japan. *J Inherit Metab Dis.* (2012)35:777-785.
- Mochida T, Tanaka T, Shiraki Y, Tajiri H, Matsumoto S, Shimbo K, Ando T, Nakamura K, Okamoto M, Endo F. Time-dependent changes in the plasma amino acid concentration in diabetes mellitus. *Mol. Genet. Metab.* 103(2011)406-409
- Nakano S, Ozasa S, Yoshioka K, Fujii I, Mitsui K, Nomura K, Kosuge H, Endo F, Matsukura M, Kimura S. Checking exon-skipping events in candidates for clinical trials of morpholino. *Pediatr Int.* 2011 Feb 22.
- ○Shigeto S, Katafuchi T, Okada Y, Nakamura K, Endo F, Okuyama T, Takeuchi H, Kroos MA, Verheijen FW, Reuser AJJ, Okumiya T.; Improved assay for differential diagnosis between Pompe disease and acid alpha-glucosidase pseudodeficiency on dried blood spots. *Mol. Genet. Metab.* 103(2011) 12-17
- Nakamura K, Hattori K, Endo F.: Newborn Screening for lysosomal disorders. *Am J Med Genet.* 157(2011) 63-71
- Kato I, Umeda K, Awaya T, Yui Y, Niwa A, Fujino H, Matsubara H, Watanabe K, Heike T, Adachi N, Endo F, Mizukami T, Nunoi H, Nakahata T, Adachi S. Successful treatment of refractory donor lymphocyte infusion-induced immune-mediated pancytopenia with rituximab. *Pediatr Blood Cancer.* 2010 Feb;54(2):329-31
- Hayashi S, Imoto I, Aizu Y, Okamoto N, Mizuno S, Kurosawa K, Okamoto N, Honda S, Araki S, Mizutani S, Numabe H, Saitoh S, Kosho T, Fukushima Y, Mitsubuchi H, Endo F, Chinen Y, Kosaki R, Okuyama T, Ohki H, Yoshihashi H, Ono M, Takada F, Ono H, Yagi M, Matsumoto H, Makita Y, Hata A, Inazawa J. Clinical application of array-based comparative genomic hybridization by two-stage screening for 536 patients with mental retardation and multiple congenital anomalies. *J Hum Genet.* 2011 Feb;56(2):110-24.
- ○Nakamura K, Sekijima Y, Nakamura K, Hattori K, Nagamatsu K, Shimizu Y, Yasude T, Ushiyama M, Endo F, Fukushima Y, Ikeda S.: Cerebral hemorrhage in Fabry's disease. *J Hum Genet.* 55(2010) 259-261
- Numata S, Koda Y, Ihara K, Sawada T, Okano Y, Matsuura T, Endo F, Yoo HW, Arranz JA, Rubio V, Wermath B, Mew NA, Tuchman M, Pinner JR, Kirk EP Yoshino M.: Mutant alleles associated with late-onset ornithine transcarbamylase deficiency in male patients have recurrently arisen and have been retained in some populations. *J Hum Genet.* 55(2010) 18-22

下澤伸行

- Vu Chi Dung, Nobuyuki Shimozawa, Nguyen Ngoc Khanh, et al. Mutations of ABCD1 gene

and phenotype of Vietnamese patients with X-linked adrenoleukodystrophy (X-ALD). *International Journal of Pediatric Endocrinology Suppl* 1: 127, 2013.

- Ohba C, Osaka H, Shimozawa N, et al. Diagnostic utility of whole exome sequencing in patients showing cerebellar and/or vermis atrophy in childhood. *Neurogenetics* 14: 225-32, 2013.
- Hama K, Nagai T, Shimozawa N, et al. Molecular Species of Phospholipids with Very Long Chain Fatty Acids in Skin Fibroblasts of Zellweger Syndrome. *Lipids* 48: 1253-1267, 2013.
- Shuji Matsui, Masuko Funahashia, Nobuyuki Shimozawa, et al. Newly identified milder phenotype of peroxisome biogenesis disorder caused by mutated PEX3 gene. *Brain Dev*; 35: 842-8, 2013.
- Yumi Mizuno, Yuichi Ninomiya, Nobuyuki Shimozawa, et al. Tysnd1 deficiency in mice interferes with the peroxisomal localization of PTS2 enzymes, causing lipid metabolic abnormalities and male infertility. *PLOS Genetics* 9 :e1003286, 2013.
- Masashi Morita, Junpei Kobayashi, Nobuyuki Shimozawa, et al. A novel double mutation in the ABCD1 gene in a patient with X-linked adrenoleukodystrophy: Analysis of the stability and function of the mutant ABCD1 protein. *J Inher Metab Dis, Rep* 10: 95-102, 2013.
- Iwasa M, Yamagata T, Shimozawa N et al. Contiguous ABCD1 DXS1357E deletion syndrome: Report of an autopsy case. *Neuropathology* 33: 292-8, 2013.
- Noriyuki Kanzawa, Nobuyuki Shimozawa, Ronald J.A. Wanders, Kazutaka Ikeda, Yoshiko Murakami, Hans R. Waterham, Satoru Mukai, Morihisa Fujita, Yusuke Maeda, Ryo Taguchi, Yukio Fujiki, and Taroh Kinoshita. Defective lipid remodeling of GPI anchors in peroxisomal disorders, Zellweger syndrome and rhizomelic chondrodysplasia punctata. *J Lipid Res* 53: 653-63, 2012
- Mizumoto H, Akashi R, Hikita N, Kumakura A, Yoshida Y, Honda A, Shimozawa N, Hata D. Mild case of D-bifunctional protein deficiency associated with novel gene mutations. *Pediatr Int* 54(2): 303-4, 2012.
- Nobuyuki Shimozawa, Ayako Honda, Naomi Kajiwara, Sachi Kozawa, Tomoko Nagase, Yasuhiko Takemoto, Yasuyuki Suzuki X-linked adrenoleukodystrophy: Diagnostic and follow-up system in Japan *J Hum Genet* 56 (2): 106-109, 2011
- Nobuyuki Shimozawa. Molecular and clinical findings and diagnostic flowchart of peroxisomal diseases *Brain Dev* 33: 770-776, 2011
- Kozawa S, Honda A, Kajiwara N, Takemoto Y, Nagase T, Nikami H, Okano Y, Nakashima S, Shimozawa N. Induction of peroxisomal lipid metabolism in mice fed a high-fat diet. *Mol Med Report* 4: 1157-1162, 2011
- Shaheen R, Al-Dirbashi OY, Al-Hassnan ZN, Al-Owain M, Makhshed N, Basheeri F, Seidahmed MZ, Salih MAM, Faqih E, Zaidan H, Al-Sayed M, Rahbeeni Z, Al-Sheddi T, Hashem M, Kurdi W, Shimozawa N, Alkuraya FS. Clinical, biochemical and molecular characterization of peroxisomal diseases in Arabs. *Clin Genet* 79 (1): 60-70, 2011
- Matsukawa T, Asheuer M, Takahashi Y, Goto J, Suzuki Y, Shimozawa N, Takano H, Onodera O, Nishizawa M, Aubourg P, Tsuji S. Identification of novel SNPs of ABCD1, ABCD2, ABCD3, and ABCD4

genes in patients with X-linked adrenoleukodystrophy (ALD) based on comprehensive resequencing and association studies with ALD phenotypes. *Neurogenetics*. 12 (1) :41-50, 2011.

- Masashi Morita, Nobuyuki Shimozawa, Yoshinori Kashiwayama, Yasuyuki Suzuki, Tsune Imanaka ABC subfamily D proteins and very long chain fatty acid metabolism as novel targets in adrenoleukodystrophy *Current Drug Targets* 12 (5): 694-706, 2011.
- Miyoshi Y, Sakai N, Hamada Y, Tachibana M, Hasegawa Y, Kiyohara Y, Yamada H, urakami M, Kondou H, Kimura-Ohba S, Mine J, Sato T, Kamio N, Ueda H, Suzuki Y, Shiomi M, Ohta H, Shimozawa N, Ozono K. Clinical aspects and adrenal functions in eleven Japanese children with X-linked adrenoleukodystrophy. *Endocr J*. 57 (11): 965-972, 2010

今中常雄

- Morita M, Kobayashi J, Yamazaki K, et al.: A novel double mutation in the ABCD1 gene in a patient with X-linked adrenoleukodystrophy: Analysis of the stability and function of the mutant ABCD1 protein. *J Inherit Metab Dis Rep* 10: 95-102, 2013
- Hama K, Nagai T, Nishizawa C, *et al.*: Molecular species of phospholipids with very long chain fatty acids in skin fibroblasts of Zellweger syndrome. *Lipids* 48: 1253-1267, 2013
- Morita M., Imanaka T.: Peroxisomal ABC transporters: Structure, function and role in disease. *Biochem. Biophys. Acta* 1822, 1387-1396, 2012. (review)
- Morita M., Shinbo S., Asahi A., Imanaka T.: Very long chain fatty acid β -oxidation in astrocytes: Contribution of the ABCD1-dependent and -independent pathways. *Biol. Pharm. Bull.* 35, 1972-1979, 2012.

加我牧子

- Yasumira A, Kokubo N, Kaga M, et al : Neurobehavioral and hemodynamic evaluation of Stroop and reverse Stroop interference in children with attention-deficit/hyperactivity disorder. *Brain & Development*. (in press).
- Tsujimoto S, Yasumura A, Kaga M et al. Increased prefrontal oxygenation related to distractor-resistant working memory. *Child Psychiatry Hum Development* 44:678-688, 2013.
- Inoue Y, Ito K, Kaga M, et al, Psychometric properties of Japanese version of the Swanson, Nolan, and Pelham, version-IV Scale-Teacher Form: A study of school children in community samples. *Brain & Development*. (in press)
- Kokubo N, Inagaki M, Gunji A, Kobayashi T, Ohta H, Kajimoto O, Kaga M: Developmental change of visuo-spatial working memory in children: Quantitative evaluation through and Advanced Trail Making Test. *Brain Dev* 34: 799-805, 2012.
- Inoue Y, Sakihara K, Gunji A, Ozawa H, Kimiya S, Shinoda S, Kaga M, Inagaki M: Reduced prefrontal hemodynamic response in children with AD/HD during the Go/NoGo task: A NIRS study. *Neuroreport* 23: 55-60, 2012.
- Yatabe K, Goto T, Watanabe K, Kaga M, Inagaki M: Reading and Writing Achievement Tests for Assessing Orthographical and Phonological Impairments of Japanese Children with Developmental

Disorders. In W. Sittiprapaporn (Ed.), Learning Disabilities. Rijeka, Croatia: InTech Publishing. ISBN 978-953-51-0269-4. pp. 69-86, 2012.

- Kita Y, Gunji A, Inoue Y, Goto T, Sakihara K, Kaga M, Inagaki M, Hosokawa T: Self-face recognition in children with autism spectrum disorders: A near-infrared spectroscopy study. *Brain Dev.* 33: 494-503, 2011.

横山和明

- Hama K, Nagai T, Nishizawa C, et al. Molecular Species of Phospholipids with Very Long Chain Fatty Acids in Skin Fibroblasts of Zellweger Syndrome. *Lipids* 48, 1253–1267 (2013).

大澤真木子、石垣景子

- Ishigaki K, Yoshikawa Y, Kuwatsuru R, Oda E, Murakami T, Sato T, Saito T, Umezu R, Osawa M. High-density CT of muscle and liver may allow early diagnosis of childhood-onset Pompe disease., *Brain Dev.* 2012 Feb;34(2):103-6.
- Ishigaki K, Murakami T, Nakanishi T, Oda E, Sato T, Osawa M., Close monitoring of initial enzyme replacement therapy in a patient with childhood-onset Pompe disease., *Brain Dev.* 2012 Feb;34(2):98-102.
- Eri Oda , Toju Tanaka , Ohsuke Migita , Motomichi Kosuga , Masaru Fukushi , Toshika Okumiya , Makiko Osawa , Torayuki Okuyama, Newborn screening for Pompe disease in Japan, *Molecular Genetics and Metabolism* 104 (2011) 560–565,
- Hiroi A, Yamamoto T, Shibata N, Osawa M, Kobayashi M., Roles of Fukutin, the Gene responsible for Fukuyama-type congenital muscular dystrophy, in neurons: possible involvement in Synaptic function and neuronal migration., *Acta Histochem Cytochem* 2011;44(2):91-101
- Ito Y, Shibata N, Saito K, Kobayashi M, Osawa M., New insights into the pathogenesis of spinal muscular atrophy., *Brain Dev* 2011;33:321-31.
- Ito Y, Oguni H, Ito S, Oguni M, Osawa M., A modified Atkins diet is promising as a treatment for glucose transporter type 1 deficiency syndrom., *Developmental Medicine and Child Neurology* 2011;(53)7:658-63
- Tsuruta T, Aihara Y, Kanno H, Kiyotani C, Maebayashi K, Sakauchi M, Osawa M, Fujii H, Kubo O, Okada Y., High-dose chemotherapy followed by autologous and allogeneic peripheral blood stem cell transplantation for recurrent disseminated trilateral retinoblastoma., *Chil Nerv Syst* 2011;27:1019-24
- Tsuruta T, Aihara Y, Kanno H, Funase M, Murayama T, Osawa M, Fujii H, Kubo O, Okada Y., Shared molecular targets in pediatric gliomas and Ependymomas., *Pediatr Blood Cancer* 2011;57:1117-23.
- ○Shioda M , Hayashi M, Takanashi J, Osawa M, Lesions in the central tegmental tract in autopsy cases of developmental brain disorders, *Brain & Development* (2010)
- ○Ishigaki K, Mitsuhashi S, Kuwatsuru R, Murakami T, Shishikura K, Suzuki H, Hirayama Y, Nonaka I, Osawa M., High-density areas on muscle CT in childhood-onset Pompe disease are caused by excess calcium accumulation., *Acta Neuropathol.* 2010 Oct;120(4):537-43.
- Kobayashi H, Shimada Y, Ikegami M, Kawai T, Sakurai K, Urashima T, Ijima M, Fujiwara M, Kneshiro E, Ohashi T, Eto Y, Ishigaki K, Osawa M, Kyosen SO, Ida H., Prognostic factors for the late onset Pompe disease with enzyme replacement therapy: From our experience of 4 cases including an autopsy case. *Mol Genet Metab* 2010;100:14-9.