

TABLE 1. Clinical Severity Grading Criteria of the Ocular Surface Findings

	Grade 0	Grade 1	Grade 2	Grade 3	Comments
Assessment of Corneal Complications					
SPK	A1D1	A1D2, A2D1	A1D3, A2D2, A3D1	A2D3, A3D2, A3D3	Using fluorescein staining based on the area and density of the lesions as described by Miyata and associates ²⁹
Corneal epithelial defect	Absent	Less than 1/4 of the corneal surface	1/4 to 1/2 of the corneal surface	More than 1/2 of the corneal surface	
Conjunctivalization	Absent	Less than 1/4 of the corneal surface	1/4 to 1/2 of the corneal surface	More than 1/2 of the corneal surface	
Neovascularization	Absent	Confined to the corneal periphery	Extending beyond the pupil margin	Extending beyond the pupil margin into the central cornea	In eyes where significant opacification or extensive symblepharon formation made it difficult to evaluate corneal neovascularization, a grade of 3 was assigned
Corneal opacification	Clear cornea with easily visible iris details	Partial obscuration of the iris details	Iris details poorly seen with barely visible pupil margins	Complete obscuration of iris and pupil details	
Keratinization	Absent	Less than 1/4 of the corneal surface	1/4 to 1/2 of the corneal surface	More than 1/2 of the corneal surface	
Assessment of Conjunctival Complications					
Conjunctival hyperemia	Absent	Mild or sectoral engorgement of the conjunctival vessels	Moderate or diffuse engorgement of the conjunctival vessels	Severe or significant engorgement of the conjunctival vessels	
Symblepharon	Absent	Involving only the conjunctival surface	Less than 1/2 of the corneal surface	More than 1/2 of the corneal surface	
Assessment of Eyelid Complications					
Trichiasis	Absent	Less than 1/4 of the lid margin	1/4 to 1/2 of the lid margin	More than 1/2 of the lid margin	

Continued on next page

TABLE 1. Clinical Severity Grading Criteria of the Ocular Surface Findings (Continued)

	Normal MJ	Mild irregularity of MJ	Moderate irregularity of MJ	Severe irregularity of MJ	Bron grading was employed for the classification of MJ changes. ³⁰
MJ involvement	Normal MJ	Mild irregularity of MJ	Moderate irregularity of MJ	Severe irregularity of MJ	Bron grading was employed for the classification of MJ changes. ³⁰
MG involvement	Oily expressible secretion	Expressible yellowish-white oily secretion	Expressible thick cheesy material	Inability to express any secretion	Fluorescein staining of the conjunctiva was performed for evaluating the MJ involvement. In eyes where significant keratinization of the lid margin or extensive symblepharon formation made it difficult to evaluate mucocutaneous junction involvement, Grade 3 was assigned.
Punctal involvement	Normal	Iatrogenic punctal occlusion	Either superior or inferior punctal occlusion	Both superior and inferior punctal occlusion	The severity was determined clinically by the nature of the meibomian gland secretion expressed manually at the center of the upper lid.

A = area; D = density; MG = mucocutaneous gland; MJ = mucocutaneous junction; SPK = superficial punctuate keratopathy.

in the study received training to standardize the conduct of examinations performed at each center. The conduct of examinations was checked by trained coordinators at each center for consistency of the examination procedures. The diagnosis of SJS or TEN was based on the history of the presence of cryptogenic fever and acute inflammation of mucosal membranes most commonly after taking cold remedies, antibiotics, or anti-inflammatory drugs, and on the presence of chronic ocular surface complications such as symblepharon, entropion, trichiasis, xerophthalmia, and/or corneal vascularization.^{1,3-5} Two hundred eight eyes of 104 healthy normal subjects (30 male, 74 female; mean age: 36.2 ± 12.0 years; range: 20–72 years) without dry eye disease and 132 eyes of 66 SS patients (66 female; mean age: 62.8 ± 11.1 years; range: 28–82 years) who were diagnosed according to Fox criteria were also investigated in this multicenter study.²⁵ Patients or control subjects with other systemic or ocular diseases, history of ocular surgery within 6 months, history of ocular cicatricial pemphigoid, or chemical, thermal, or radiation injury that would have adverse ocular surface effects were excluded according to the study exclusion criteria. SJS patients with a baseline best-corrected Landolt conventional visual acuity of less than 20/2000 attributable to cataract in both eyes, ocular surface keratinization, glaucoma, or posterior segment disease were excluded from this study, since the functional visual acuity measurement system cannot assess functional visual acuity at such low visual acuity levels.

- **SLIT-LAMP EXAMINATIONS:** All study subjects underwent slit-lamp examinations observing 12 components of 3 categories of ocular complications, such as corneal complications consisting of superficial punctuate keratopathy (SPK), epithelial defect, conjunctivalization, neovascularization, opacification, and keratinization; conjunctival complications consisting of hyperemia and symblepharon formation; and eyelid complications consisting of trichiasis, mucocutaneous junction involvement, meibomian gland involvement, and punctal damage. Each component was graded on a scale from 0 to 3, depending on the severity of involvement.²⁶

The severity gradings and ocular surface tests were performed under the same single protocol by the researchers of all contributing study centers. Table 1 shows the clinical severity grading criteria of the ocular surface findings.

- **TEAR FUNCTION AND OCULAR SURFACE EXAMINATIONS:** The standard Schirmer test without topical anesthesia was performed as previously reported.⁷ A vital staining severity grading was also assigned. A 2-μL volume of 1% fluorescein dye was instilled in the conjunctival sac by a micropipette. The minimum score for corneal fluorescein staining was 0 points and the maximum score was 9 points.²⁷

- **STANDARD VISUAL ACUITY MEASUREMENTS:** Standard visual acuity testing using Landolt charts placed 5 m away from subjects was performed. Landolt visual acuity

TABLE 2. Standard Visual Acuity and Visual Parameters Assessed by Functional Visual Acuity Measurement System in Eyes of Patients With Sjögren Syndrome, Stevens-Johnson Syndrome Patients, and Healthy Normal Subjects

	SJS	SS	Normal
Conventional visual acuity			
logMAR	0.76 ± 0.76	-0.004 ± 0.13	-0.10 ± 0.10
Decimal	0.17	1.01	1.26
Functional visual acuity			
logMAR	0.98 ± 0.62 ^a	0.28 ± 0.27 ^a	-0.008 ± 0.13
Decimal	0.10	0.52	1.02
Maximal visual acuity			
logMAR	0.83 ± 0.65	0.10 ± 0.26	-0.15 ± 0.12
Decimal	0.15	0.79	1.41
Minimal visual acuity			
logMAR	1.19 ± 0.60	0.53 ± 0.36	0.17 ± 0.19
Decimal	0.06	0.30	0.68
Visual maintenance ratio	0.86 ± 0.12	0.91 ± 0.07 ^b	0.98 ± 0.05 ^{c,d}
Reaction time	1.0 ± 0.2	1.1 ± 0.2	1.0 ± 0.2
Blink number	11.2 ± 9.3	17.2 ± 9.6 ^b	16.4 ± 8.7 ^c

logMAR = logarithm of minimal angle of resolution; logMAR = logarithm of minimal angle of resolution; SJS = Stevens-Johnson syndrome; SS = Sjögren syndrome; VA = visual acuity.

^a*P* < .05 between conventional VA and functional VA.

^b*P* < .05 between groups of SJS and SS.

^c*P* < .05 between groups of SJS and Normal.

^d*P* < .05 between groups of SS and Normal.




FIGURE 1. Visual function and Visual Function Questionnaire-25 in Stevens-Johnson syndrome (SJS) with minimal corneal complications and Sjögren syndrome (SS) patients. (Left) Conventional and functional visual acuity and visual maintenance ratio in Stevens-Johnson syndrome with minimal corneal complications and Sjögren syndrome patients. (Right) Total composite NEI VFQ-25 scores in Stevens-Johnson syndrome with minimal corneal complications and Sjögren syndrome patients. logMAR = logarithm of minimal angle of resolution.

was employed instead of Snellen chart since it is a standard test in Japan, and because the optotypes in Landolt and functional visual acuity testing are similar.

• **FUNCTIONAL VISUAL ACUITY MEASUREMENTS:** Continuous visual acuity testing during a 60-second period under natural blinking was performed as previously reported.²⁴

• **FUNCTIONAL VISUAL ACUITY INDICES:** Briefly, the outcome parameters of the functional visual acuity measurement system were functional visual acuity (defined as the average of visual acuities measured during a

60-second testing), visual maintenance ratio (defined as the ratio of logMAR values of the functional visual acuities over the time frame for testing divided by the logarithm of minimal angle of resolution (logMAR) baseline visual acuity),⁷ maximal corrected visual acuity, minimal corrected visual acuity, standard deviation of functional visual acuity, mean reaction time (defined as the mean of the response time taken by a subject to respond to an optotype), and blink numbers during a 60-second functional visual acuity test.

• **VISUAL FUNCTION QUESTIONNAIRE-25:** We used the Japanese version of the NEI VFQ-25 (National Eye Insti-

tute Visual Function Questionnaire 25) to evaluate the vision-related quality of life.²⁸ NEI VFQ-25 measures the following 12 vision-targeted subscales: general health, general vision, ocular pain, near activities, distant activities, social functioning, mental health, role difficulties, dependency, driving, color vision, and peripheral vision. A scale of 0 to 100 points is used for subscale scores. A score of 100 indicates the best possible score, while 0 indicates the worst possible score.

• **STATISTICAL ANALYSIS:** A 1-way ANOVA was performed for the comparison of conventional visual acuities, functional visual acuities, visual maintenance ratios, ocular surface grading scores, and VFQ-25 scores among SJS patients, SS patients, and normal control subjects. The Bonferroni test was used for further multiple comparisons. A paired t test was performed for the comparison between conventional and functional visual acuities in SJS patients, SS patients, and normal control subjects alone. To investigate whether the visual disturbance or quality of life are similarly affected in SJS patients compared to SS patients, conventional visual acuities, functional visual acuities, visual maintenance ratios, and VFQ-25 scores were compared among SJS patients with minimal corneal complications and SS patients by paired t test. Minimal corneal complication was defined as a grading score ≤ 4 points, in relation to keratinization, conjunctivalization, opacification, corneal epithelial defect, neovascularization, and SPK. Severe corneal complication was defined as a grading score >4 points. To investigate the effect of tear functions on the ocular surface complications, visual disturbance, or quality of life in SJS and SS patients, ocular surface grading scores, conventional visual acuities, functional visual acuities, visual maintenance ratios, and VFQ-25 scores were compared in SJS patients with and without aqueous tear deficiency by 1-way ANOVA. Aqueous tear deficiency was defined as a Schirmer test score ≤ 5 mm. The relation between ocular surface grading scores, conventional visual acuities, and functional visual acuities was analyzed by Pearson correlation analysis. The relation between ocular surface complications and conventional visual acuities, functional visual acuities, visual maintenance ratios, or VFQ-25 scores was also analyzed by Pearson correlation analysis in SJS patients with and without aqueous tear deficiency and SS patients. In the correlation analysis between ocular surface grading scores and conventional visual acuities or functional visual acuities in SJS patients, eyes were divided into 3 visual groups: good conventional visual acuity group ($\log\text{MAR}$ conventional visual acuity score ≤ 0), intermediate conventional visual acuity group ($0 < \log\text{MAR}$ conventional visual acuity ≤ 0.3), and poor conventional visual acuity group ($0.3 < \log\text{MAR}$ conventional visual acuity score ≤ 2.0). The relation between VFQ-25 score, conventional visual acuity, and functional visual acuity was analyzed by the same methodology, using the eye with better conventional




FIGURE 2. Visual function, ocular surface grading score, and Visual Function Questionnaire-25 in Stevens-Johnson syndrome (SJS) patients with and without aqueous tear deficiency and Sjögren syndrome (SS) patients. (Top) Conventional and functional visual acuity and visual maintenance ratio in Stevens-Johnson syndrome patients with and without aqueous tear deficiency and Sjögren syndrome patients. (Middle) Total ocular surface grading scores in Stevens-Johnson syndrome patients with and without aqueous tear deficiency and Sjögren syndrome patients. (Bottom) Total composite NEI VFQ-25 scores in Stevens-Johnson syndrome patients with minimal corneal complications and Sjögren syndrome patients. $\log\text{MAR}$ = logarithm of minimal angle of resolution.

visual acuity. The correlation between clinical findings, conventional visual acuities, and functional visual acuities was also investigated by multiple linear regression analysis. A probability level of $P < .05$ was considered statistically significant. SPSS (SPSS Inc, Chicago, Illinois, USA) was used as the statistical analysis software.

RESULTS

• **TEAR FUNCTION TESTS:** The mean Schirmer test values were 9.1 ± 9.3 mm in SJS patients, 4.6 ± 4.5 mm in SS patients, and 18.6 ± 9.5 mm in healthy control subjects, respectively. The Schirmer test values were significantly higher in SJS patients compared to SS patients

TABLE 3. Percentages of Ocular Surface Grading Score in Sjögren Syndrome Patients, Stevens-Johnson Syndrome Patients, and Healthy Normal Subjects

	SJS				SS				Normal			
	Grade 0	Grade 1	Grade 2	Grade 3	Grade 0	Grade 1	Grade 2	Grade 3	Grade 0	Grade 1	Grade 2	Grade 3
Assessment of corneal complications												
SPK	15.1%	25.9%	25.2%	33.8%	31.1%	25.6%	20.2%	23.3%	99.0%	1.0%	0	0
Corneal epithelial defect	92.2%	2.8%	0	5.0%	100%	0	0	0	100%	0	0	0
Conjunctivalization	32.1%	14.3%	12.9%	40.7%	95.4%	4.6%	0	0	100%	0	0	0
Neovascularization	25.7%	27.1%	24.3%	22.9%	97.7%	2.3%	0	0	98.1%	1.9%	0	0
Opacification	28.4%	50.4%	13.5%	7.8%	99.2%	0.8%	0	0		0	0	0
Keratinization	88.7%	4.3%	3.4%	3.4%	100%	0	0	0	100%	0	0	0
Assessment of conjunctival complications												
Conjunctival hyperemia	24.1%	58.9%	14.9%	2.1%	82.3%	16.9%	0.8%	0	100%	0	0	0
Symblepharon	38.8%	48.2%	7.9%	5.0%	99.2%	0.8%	0	0	100%	0	0	0
Assessment of eyelid complications												
Trichiasis	40.3%	25.2%	14.4%	20.1%	100%	0	0	0	99.0%	1.0%	0	0
MJ involvement	15.6%	42.6%	27.0%	14.9%	86.2%	12.3%	1.5%	0	100%	0	0	0
MG involvement	17.0%	22.0%	17.7%	43.3%	81.5%	14.6%	0	3.8%	100%	0	0	0
Punctal involvement	27.7%	19.9%	9.9%	42.6%	75.4%	23.1%	0.8%	0.8%	100%	0	0	0

MG = meibomian gland; MJ = mucocutaneous junction; SJS = Stevens-Johnson syndrome; SPK = superficial punctate keratopathy; SS = Sjögren syndrome.

($P < .05$). A total of 49.6 % of the patients with SJS had Schirmer test values greater than 5 mm.

• **STANDARD CONVENTIONAL VISUAL ACUITY:** Table 2 shows the mean logMAR conventional visual acuity in SJS and SS patients and the normal subjects. The mean logMAR conventional visual acuity in SJS patients was significantly lower compared to the mean logMAR conventional visual acuity in SS patients and normal controls ($P < .05$).

The mean logMAR conventional visual acuity in SJS patients with severe corneal complications was 0.74 ± 0.57 . The mean logMAR conventional visual acuity in SJS patients with minimal corneal complications and SS patients was 0.21 ± 0.42 and -0.001 ± 0.12 , respectively. The logMAR conventional visual acuities in SJS patients were significantly higher compared to SS patients (Figure 1, Left).

The mean logMAR conventional visual acuity in SJS patients with and without aqueous tear deficiency and SS patients was 0.47 ± 0.53 , 0.65 ± 0.63 , and -0.004 ± 0.13 , respectively. The logMAR conventional visual acuities in SJS patients were significantly higher compared to SS patients (Figure 2, Top).

• **FUNCTIONAL VISUAL ACUITY INDICES:** Table 2 shows the results of all indices measured by the functional visual acuity measurement system. The mean logMAR functional visual acuity was significantly lower compared to the mean logMAR conventional visual acuity in pa-

tients with SJS and SS ($P < .05$). The mean logMAR standard deviation of functional visual acuity was significantly greater in patients with SJS and SS compared to normal subjects ($P < .05$). The mean visual maintenance ratio in the SJS patients was significantly lower than in SS patients, and the mean visual maintenance ratio in SS patients was significantly lower than in normal subjects ($P < .05$). There were no significant differences in reaction times among SJS patients, SS patients, and normal subjects. The mean blink number in the SJS patients was significantly lower compared to SS patients and normal subjects ($P < .05$).

The mean logMAR functional visual acuity in SJS patients with severe corneal complications was 1.16 ± 0.45 . The mean logMAR functional visual acuity in SJS and SS patients with minimal corneal complications was 0.50 ± 0.55 and 0.28 ± 0.27 , respectively. The functional visual acuities in SJS patients were significantly higher compared to in SS patients (Figure 1, Left). The mean visual maintenance ratio in SJS and SS patients with minimal corneal complications was 0.88 ± 0.10 and 0.91 ± 0.07 , respectively. Visual maintenance ratios in SJS patients were significantly lower compared to SS patients (Figure 1, Left).

The mean logMAR functional visual acuity in SJS patients with and without aqueous tear deficiency and SS patients was 0.83 ± 0.54 , 0.99 ± 0.63 , and 0.28 ± 0.27 , respectively. The functional visual acuities in SJS patients with and without aqueous tear deficiency were significantly




FIGURE 3. Correlation between visual function and ocular surface grading score. (Left) Correlation between logMAR conventional visual acuity scores and total ocular surface grading scores. (Right) Correlation between logarithm of minimal angle of resolution (logMAR) functional visual acuity scores and total ocular surface grading scores. SJS: Stevens-Johnson syndrome; SS: Sjögren syndrome.

TABLE 4. Multivariable Regression Analyses Between Ocular Surface Grading Score, logMAR Conventional Visual Acuity, and logMAR Functional Visual Acuity

Complication	logMAR Conventional Visual Acuity ^a		logMAR Functional Visual Acuity ^b	
	Standard Partial Regression	P value	Standard Partial Regression	P value
Neovascularization	0.509	<.001	0.229	<.001
Opacification	0.385	<.001	0.308	<.001
Keratinization	-0.088	.002	-0.131	<.001
SPK	0.054	.061	0.193	<.001
Symblepharon	0.059	.103	0.162	<.001
Conjunctivalization	-0.058	.262	0.168	<.001
Corneal epithelial defect	0.034	.198	0.027	.319

logMAR = logarithm of minimal angle of resolution; SPK = superficial punctate keratopathy.

^aConditioned multiple correlation coefficient for logMAR conventional visual acuity = 0.81.

^bConditioned multiple correlation coefficient for logMAR functional visual acuity = 0.84.

higher compared to SS patients (Figure 2, Top). The mean visual maintenance ratios in SJS patients with and without aqueous tear deficiency and SS patients were 0.86 ± 0.10 , 0.83 ± 0.14 , and 0.91 ± 0.07 , respectively. The visual maintenance ratios in SJS patients both with and without

aqueous tear deficiency were significantly lower compared to SS patients (Figure 2, Top).

• **CLINICAL FINDINGS:** Table 3 shows the mean ocular surface grading scores in SS and SJS patients and normal subjects. The mean ocular surface grading scores in all 12 components of clinical findings was significantly higher in SJS patients compared to SS patients and normal subjects ($P < .05$).

The mean ocular surface grading scores in SJS patients with and without aqueous tear deficiency and SS patients were 12.0 ± 8.1 , 12.8 ± 5.7 , and 2.3 ± 1.8 , respectively. The total ocular surface grading scores in SJS both with and without aqueous tear deficiency were significantly higher compared to SS patients (Figure 2, Middle).

• **CORRELATION BETWEEN VISUAL FUNCTION AND CLINICAL FINDINGS:** Figure 3 shows the correlation between visual function and ocular surface grading score in SJS patients, SS patients, and normal subjects overall. A strong significant correlation was observed between total ocular surface grading scores and best-corrected logMAR Landolt conventional visual acuities ($r = 0.78$, $P < .001$), as well as best-corrected logMAR Landolt functional visual acuities ($r = 0.82$, $P < .001$).

Table 4 shows the correlation of visual function and ocular surface grading scores. The results of multiple linear regression analysis between the clinical findings and log-

TABLE 5. Correlations Between Ocular Complications and Visual Function or the Composite National Eye Institute Visual Function Questionnaire Scores in Stevens-Johnson Syndrome Patients With Aqueous Tear Deficiency and Sjögren Syndrome Patients

	SJS With Aqueous Tear Deficiency				SS			
	Pearson CC				Pearson CC			
	Log Conventional Visual Acuity	Log Functional Visual Acuity	Visual Maintenance Ratio	NEI VFQ-25	Log Conventional Visual Acuity	Log Functional Visual Acuity	Visual Maintenance Ratio	NEI VFQ-25
Trichiasis	0.09	0.08	0.16	-0.02	—	—	—	—
Symblepharon	0.43 ^b	0.53 ^b	-0.30 ^a	-0.46 ^a	0.08	0.15	-0.07	-0.12
Punctal involvement	0.55 ^b	0.57 ^b	-0.28	-0.49 ^b	0.09	0.13	-0.13	0.01
MG involvement	0.48 ^b	0.44 ^b	-0.23	-0.55 ^b	0.06	0.07	-0.04	-0.42 ^b
MJ involvement	0.25	0.33 ^b	-0.26	-0.39 ^a	-0.06	0.20 ^a	-0.32 ^a	-0.05
Conjunctival hyperemia	0.28 ^a	0.31 ^a	-0.23	-0.48 ^b	0.22 ^b	0.11	0.01	-0.24
Keratinization	0.06	0.09	-0.03	-0.03	—	—	—	—
Conjunctivalization	0.53 ^b	0.52 ^b	-0.19	-0.53 ^b	0.29 ^b	0.41 ^b	-0.28 ^b	-0.22
Opacification	0.59 ^b	0.67 ^b	-0.47 ^b	-0.69 ^b	0.17	0.20	-0.19	—
Corneal epithelial defect	0.06	0.16	-0.15	-0.11	—	—	—	—
Neovascularization	0.64 ^b	0.63 ^b	-0.20	-0.63 ^b	0.21 ^a	0.23 ^b	-0.14	-0.2 ^a
SPK	0.35 ^a	0.35 ^a	-0.08	-0.41 ^a	0.04	0.212	-0.11	-0.22
Total ocular complications	0.55 ^b	0.58 ^b	-0.26	-0.61 ^b	0.12	0.25 ^b	-0.15	-0.40 ^b

CC = correlation coefficient; MG = meibomian gland; MJ = mucocutaneous junction; NEI VFQ-25 = National Eye Institute visual function questionnaire; SJS = Stevens-Johnson syndrome; SPK = superficial punctuate keratopathy; SS = Sjögren syndrome.

^a*P* < .05.

^b*P* < .01.

TABLE 6. Correlations Between Visual Function and Ocular Surface Grading Scores or Composite National Eye Institute Visual Function Questionnaire Scores in Good, Intermediate, or Poor Conventional Visual Acuity Group of Stevens-Johnson Syndrome Patients

	All Groups		Good Conventional Visual Acuity Group		Intermediate Conventional Visual Acuity Group		Poor Conventional Visual Acuity Group	
	Pearson CC	<i>P</i> Value	Pearson CC	<i>P</i> Value	Pearson CC	<i>P</i> Value	Pearson CC	<i>P</i> Value
Conventional visual acuity vs clinical finding scores	0.59 ^b	.001	0.37	.08	0.24	.15	0.40 ^b	.001
Functional visual acuity vs clinical finding scores	0.63 ^b	.001	0.56 ^b	.005	0.49 ^b	.002	0.34 ^b	.007
Conventional visual acuity vs composite NEI VFQ-25 scores	-0.74 ^b	.001	-0.44	.06	-0.25	.25	-0.56 ^a	.03
Functional visual acuity vs composite NEI VFQ-25 scores	-0.74 ^b	.001	-0.55 ^b	.02	-0.20	.37	-0.57 ^a	.03

CC = correlation coefficient; NEI VFQ-25 = National Eye Institute Visual Function Questionnaire.

^a*P* < .05.

^b*P* < .01.

MAR conventional visual acuity showed a significant and strong correlation with neovascularization, opacification, and keratinization grades. Clinical findings such as SPK, symblepharon, and conjunctivalization also had a significant and strong correlation with the functional visual acuities. The

multiple regression equation of logMAR conventional visual acuity was expressed as follows: logMAR conventional visual acuity = -0.084 + neovascularization × 0.509 + opacification × 0.385 + keratinization × -0.088. Likewise, the multiple regression equation of logMAR functional visual




FIGURE 4. Correlations between visual function and ocular surface grading score in the good and intermediate conventional visual acuity group of Stevens-Johnson syndrome patients. (Left) Correlation in the good conventional visual acuity group of Stevens-Johnson syndrome patients. (Right) Correlation in the intermediate conventional visual acuity group of Stevens-Johnson syndrome patients. logMAR = logarithm of minimal angle of resolution.




FIGURE 5. Visual Function Questionnaire-25 results in patients with Stevens-Johnson syndrome (SJS), patients with Sjögren syndrome (SS), and healthy normal subjects.

acuity was expressed as follows: $\text{logMAR functional visual acuity} = -0.061 + \text{neovascularization} \times 0.229 + \text{opacification} \times 0.308 + \text{keratinization} \times -0.131 + \text{SPK} \times 0.193 + \text{symblepharon} \times 0.162 + \text{conjunctivalization} \times 0.168$.

Table 5 shows the correlation between ocular complications and visual function in SJS patients with aqueous

tear deficiency and SS patients. Strong significant correlations were observed between total ocular surface grading score and logMAR conventional visual acuities or logMAR functional visual acuities in SJS patients with aqueous tear deficiency, and similar strong significant correlations in SJS patients without aqueous tear defi-




FIGURE 6. Relation between visual function and composite NEI VFQ-25 scores. (Left) Correlation between logMAR conventional visual acuity scores and total composite NEI VFQ-25 scores. (Right) Correlation between logarithm of minimal angle of resolution (logMAR) functional visual acuity scores and total composite NEI VFQ-25 scores. SJS: Stevens-Johnson syndrome; SS: Sjögren syndrome.

ciency ($r = 0.66, P < .001$) (data not shown), while significant correlations were observed only between total ocular surface grading score and logMAR functional visual acuities in SS patients (Table 5).

Table 6 shows the correlations between visual function and ocular surface grading scores in the good, intermediate, and poor conventional visual acuity groups of SJS patients. A strong positive significant correlation was observed between total ocular surface grading scores and logMAR Landolt functional visual acuities in the good conventional visual acuity group ($r = 0.56, P = .005$) and intermediate conventional visual acuity group ($r = 0.49, P = .002$), while no correlation was observed between total ocular surface grading scores and logMAR Landolt conventional visual acuities in these groups (Figure 4).

• **VISUAL FUNCTION QUESTIONNAIRE-25:** Mean subscale and composite NEI VFQ scores for SJS and SS patients and normal subjects are presented in Figure 5. All 12 subscale NEI VFQ scores were significantly lower in the SJS patients compared to the normal subjects ($P < .05$). Likewise, all subscale scores were significantly lower in the SS patients compared to the normal subjects ($P < .05$). The subscale of “ocular pain” was remarkably low in SS patients, while all subscale scores were remarkably lower in SJS patients. The mean composite NEI VFQ scores of the 12 subscales were 49.1 ± 21.6 in SJS patients, 72.8 ± 12.8 in SS patients, and 89.4 ± 8.1 in the normal subjects.

The mean total composite NEI VFQ score in SJS patients with severe corneal complications was 45.2 ± 20.9 . The mean total composite NEI VFQ scores in SJS patients with minimal corneal complications and SS patients were 62.2 ± 19.8 and 73.0 ± 12.8 , respectively. The total composite NEI VFQ scores in SJS patients were significantly lower compared to SS patients (Figure 1, Right).

The mean total composite NEI VFQ scores in SJS patients with and without aqueous tear deficiency were 51.6 ± 20.2 , 49.4 ± 23.5 , and 72.8 ± 12.8 , respectively. The total composite NEI VFQ scores in SJS both with and without aqueous tear deficiency were significantly lower compared to SS patients (Figure 2, Right).

• **CORRELATION OF VISUAL FUNCTION AND NEI VFQ-25 SCORES:** Figure 6 shows the correlation between visual function and the composite NEI VFQ-25 scores in SJS patients, SS patients, and normal subjects overall. A strong negative correlation was detected between the composite NEI VFQ-25 scores and best-corrected logMAR Landolt conventional visual acuities ($r = -0.76, P < .01$), and best-corrected logMAR Landolt functional visual acuities ($r = -0.81, P < .01$).

Table 6 shows the correlations between visual function and the composite NEI VFQ-25 scores in the good, intermediate, and poor conventional visual acuity groups in SJS patients. A positive significant correlation was




FIGURE 7. Correlations between visual function and the total composite NEI VFQ-25 scores in the good conventional visual acuity group of Stevens-Johnson syndrome patients. logMAR = logarithm of minimal angle of resolution.

observed between the composite NEI VFQ-25 scores and logMAR Landolt functional visual acuities in the good conventional visual acuity group ($r = 0.55$, $P = .02$), while no correlation was observed between the composite NEI VFQ-25 scores and logMAR Landolt conventional visual acuities in this group ($r = 0.44$, $P = .06$) (Figure 7).

• **CORRELATION BETWEEN NEI VFQ-25 SCORES AND CLINICAL FINDINGS:** Table 5 shows the correlation between ocular complications and the total composite NEI VFQ-25 scores in SJS patients with aqueous tear deficiency and SS patients. Strong significant correlations were observed between total ocular surface grading score and the composite NEI VFQ-25 scores in SJS patients with aqueous tear deficiency and SS patients (Table 5), and similarly in SJS patients without aqueous tear deficiency ($r = 0.51$, $P = .002$) (data not shown).

DISCUSSION

SEVERE OCULAR SURFACE DISEASE ASSOCIATED WITH SJS has been reported to cause visual deterioration. However, quantifying visual acuity in SJS patients has not been assessed, although an interest in the quantitative interpre-

tation of visual function has been rising over the last few years, especially in the fields of refractive surgery, cataract, and dry eyes, through analyses by contrast sensitivity, contrast visual acuity, and wavefront analysis.⁹⁻¹⁹ In this report, we measured the functional visual acuity in addition to conventional visual acuity testing and evaluated the relations between visual functions, ocular surface clinical findings, and the vision-related quality of life in SJS patients. We chose functional visual acuity testing for the assessment of the visual function, which has been shown to be efficient in the detection of "masked impairment of visual function" in dry eye patients, since SJS is known to be associated with severe dry eyes.²⁰⁻²⁴

Visual function testing revealed several interesting findings. First, visual acuities measured by conventional Landolt visual acuity testing were low in the SJS patients as compared with SS patients and normal subjects. When we focused on the visual function of the SS patients and normal subjects, the functional visual acuity scores in the SS patients were significantly lower than in the normal subjects, although there were no differences in the conventional visual acuities. In addition, the mean visual maintenance ratios in the SJS patients were significantly lower than in the SS patients, indicating that ability to

maintain the best visual acuity in SJS patients had deteriorated more than in the SS patients.

The functional visual acuity examination has been shown to be useful for the assessment of visual function related to dry eyes in our previous reports.²⁰⁻²² A previous report has also suggested the possibility that the functional visual acuity examination might reflect the effect of ocular surface findings and dry eye states on visual functions.⁸ In this report, we analyzed the relation of ocular surface findings with visual function and quality of life in detail by grading the severity of ocular surface findings. The clinical severity scores of the examined ocular surface findings were much higher in the SJS patients. We analyzed whether visual disturbance and quality of life were similarly affected in SJS and SS patients without corneal complications or only with minimal corneal complications. Interestingly, we observed that visual function and quality of life were deteriorated in SJS patients with minimal corneal complications compared with SS patients. Moreover, we noted more visual dysfunction and declined quality of life in SJS patients with similar aqueous tear deficiency compared to SS patients. According to the multiple linear regression analysis, neovascularization, opacification, and keratinization involving the optical axis appeared to have a significant effect on the logMAR conventional visual acuities. SPK, symblepharon, and conjunctivalization also had a significant effect on the logMAR functional visual acuities.

Our findings that a stronger correlation existed between ocular surface grading score and logMAR functional visual acuity compared to logMAR conventional visual acuity suggest that functional visual acuity testing can indeed reflect the effect of clinical complications of ocular surface disease on visual function in SJS. In the correlations between visual function and ocular surface grading scores in the good, intermediate, and poor conventional visual acuity groups in SJS patients, a strong positive significant correlation was observed between total ocular surface grading scores and logMAR Landolt functional visual acuities in the good conventional visual acuity group and intermediate conventional visual acuity group, while no correlation was observed between total ocular surface grading scores and logMAR Landolt conventional visual acuities in these groups. These results suggest that functional visual acuity reflects the effect of ocular surface complications on visual function more sensitively in SJS patients with good and intermediate conventional visual acuities. The strong correlation of logMAR functional visual acuity with ocular surface grading score also suggests that functional visual acuity may be detecting the effect of ocular surface disease severity on other visual functions such as contrast, glare, or higher-order aberrations (compared to conventional visual acuity testing), which needs to be investigated in future studies employing the above-

mentioned methodologies in conjunction with functional visual acuity testing.

The mean of all VFQ-25 subscale scores was remarkably worse in the SJS patients compared to normal subjects. Likewise, the mean of all subscale scores in SS patients was significantly lower than in the normal subjects. When analyzed in detail, only the subscale scores of "general health" and "ocular pain" were worse, without marked changes in other subscale scores. In SJS patients, as compared with normal subjects, all VFQ-25 subscale scores, especially "ocular pain," "near activities," "distance activities," "mental health," "role difficulties," and "driving," were very low. These findings suggest that SJS patients suffer from an actual limitation of vision-related daily activity rather than a sense of decreased visual performance and health decline.

A strong negative correlation was observed in the relation between logMAR conventional visual acuities and the VFQ-25 composite scores in this study, with a strong correlation detectable for the relation between the VFQ-25 composite scores and the logMAR functional visual acuities.

We had noteworthy observations that patients with SJS had significantly worse dry eye and visual symptom scores compared to SS patients. We believe these observations owe to the presence of a higher incidence of ocular surface complications in SJS such as symblepharon, corneal opacification, and SPK.

One of the weak points of the current study is that SJS patients, SS patients, and normal subjects were not age-matched. However, it was actually difficult to recruit subjects with age matching in the current study. In fact, the age of onset of SS is usually beyond middle age, while individuals have a risk to be involved with SJS at any age. Moreover, recruitment of elderly individuals with normal tear functions as normal control subjects is another challenging task. It should be noted that the VFQ-25 subscale scores might have been affected by sex and age differences. Another weakness was the lack of definitive diagnosis of SJS/TEN by skin biopsy. We diagnosed SJS or TEN on the history of the presence of cryptogenic fever and acute inflammation of mucosal membranes, most commonly after taking cold remedies, antibiotics, or anti-inflammatory drugs, and on the presence of the chronic ocular surface complications.

Overall, although standard visual acuity testing is a good measurement of one aspect of visual function, the functional visual acuity examination provided other important and detailed information on visual functions related with clinical findings and vision-related quality of life. In conclusion, SJS patients with good or intermediate visual acuity scores measured by conventional visual acuity testing were found to suffer from lower vision-related quality of life, as assessed by functional visual acuity testing and VFQ scores.

ALL AUTHORS HAVE COMPLETED AND SUBMITTED THE ICMJE FORM FOR DISCLOSURE OF POTENTIAL CONFLICTS OF Interest. None of the authors received lecture fees or equity payments from Nidek. Drs Kazuo Tsubota and Minako Kaido both hold patent rights for the method and the apparatus for the measurement of functional visual acuity (US patent no: 7470026). All study centers received and shared an official grant from the Japanese Ministry of Health and Welfare, Tokyo, Japan during the conduct of the study. Involved in conception and design (M.K., C.S., S.K., K.T.); analysis and interpretation (M.K., C.S., K.T.); writing the article (M.K.); critical revision of the article (M.Y., C.S., S.K., J.S., Y.T., Y.H., T.C., K.T.); final approval of the article (M.K., M.Y., C.S., S.K., J.S., Y.T., Y.H., T.C., K.T.); data collection (M.K., M.Y., C.S., J.S., Y.T., Y.H., T.C.); provision of materials, patients, or resources (M.K., M.Y., C.S., J.S., Y.T., Y.H., T.C.); statistical expertise (M.K.); obtaining funding (M.K., M.Y., C.S.); literature search (M.K., C.S.); and administrative, technical, or logistical support (C.S., S.K., K.T.). Ethics committee approvals for the examination procedures and study protocol were obtained at each center for this prospective study (IRB approval number: 17-129, Keio University School of Medicine, 20. 6, 2006). Written informed consent was obtained from each patient to participate in this study.

REFERENCES

- Arsujo OE, Floweres FP. Steven-Johnson syndrome. *J Emerg Med* 1984;2(2):129-135.
- Tsubota K, Toda I, Saito H, Shinozaki N, Shimazaki J. Reconstruction of the corneal epithelium by limbal allograft transplantation for severe ocular surface disorders. *Ophthalmology* 1995;102(10):1486-1496.
- Power WJ, Ghoraishi M, Merayo-Llovers J, Neves RA, Foster CS. Analysis of the acute ophthalmic manifestations of the erythema multiforme/Stevens-Johnson syndrome/toxic epidermal necrolysis disease spectrum. *Ophthalmology* 1995; 102(11):1669-1676.
- Puangrichareon V, Tseng SCG. Cytologic evidence of corneal disease with limbal stem cell deficiency. *Ophthalmology* 1995;102(19):1476-1485.
- Tsubota K, Satake Y, Kaido M, et al. Treatment of severe ocular-surface disorders with corneal epithelial stem-cell transplantation. *N Engl J Med* 1999;340(22):1697-1703.
- Kaido M, Goto E, Dogru M, Tsubota K. Punctal occlusion in the management of chronic Stevens-Johnson syndrome. *Ophthalmology* 2004;111(5):895-900.
- Kaido M, Dogru M, Yamada M, et al. Functional visual acuity in Stevens-Johnson syndrome. *Am J Ophthalmol* 2006; 142(6):917-922.
- Sotozono C, Ueta M, Koizumi N, et al. Diagnosis and treatment of Stevens-Johnson syndrome and toxic epidermal necrolysis with ocular complications. *Ophthalmology* 2009; 116(4):685-690.
- Rieger G. The importance of the precorneal tear film for the quality of topical imaging. *Br J Ophthalmol* 1992;76(3):157-158.
- Niesen U, Businger U, Hartmann P, Senn P, Schipper I. Glare sensitivity and visual acuity after excimer laser photorefractive for myopia. *Br J Ophthalmol* 1997;81(2):136-140.
- Ghaith AA, Daniel J, Stulting RD, Thompson KP, Lynn M. Contrast sensitivity and glare disability after radial keratotomy and photorefractive keratectomy. *Arch Ophthalmol* 1998; 116(1):12-18.
- Roland M, Lester M, Maci A, Calabria G. Low spatial-contrast sensitivity in dry eyes. *Cornea* 1998;17(4):376-379.
- Liu Z, Pflugfelder SC. Corneal surface regularity and the effect of artificial tears in aqueous tear deficiency. *Ophthalmology* 1999;106(5):939-943.
- Nakamura K, Bissen-Miyajima H, Toda I, Hori Y, Tsubota K. Effect of laser in situ keratomileusis correction on contrast visual acuity. *J Cataract Refract Surg* 2001;27(3):357-361.
- Huang FC, Tseng SH, Shih MH, Chen FK. Effect of artificial tears on corneal surface regularity, contrast sensitivity, and glare disability in dry eyes. *Ophthalmology* 2002;109(10): 1934-1940.
- Bron AJ, Tiffany JM, Gouveia SM, Yokoi N, Voon LW. Functional aspects of the tear film lipid layer. *Exp Eye Res* 2004;78(3):347-360.
- Kojima T, Ishida R, Dogru M, et al. A new noninvasive tear stability analysis system for the assessment of dry eyes. *Invest Ophthalmol Vis Sci* 2004;45(5):1369-1374.
- Puell MC, Benitez-del-Castillo JM, Martinez-de-la-Casa J, et al. Contrast sensitivity and disability glare in patients with dry eye. *Acta Ophthalmol Scand* 2006;84(4):527-531.
- Koh S, Maeda N, Hirohara Y, et al. Serial measurements of high-order aberrations after blinking in patients with dry eye. *Invest Ophthalmol Vis Sci* 2008;49(1):133-138.
- Goto E, Yagi Y, Matsumoto Y, Tsubota K. Impaired functional visual acuity of dry eye patients. *Am J Ophthalmol* 2002;133(2):181-186.
- Goto E, Yagi Y, Kaido M, Matsumoto Y, Konomi K, Tsubota K. Improved functional visual acuity after punctal occlusion in dry eye patients. *Am J Ophthalmol* 2003;135(5):704-705.
- Ishida R, Kojima T, Dogru M, et al. The application of a new continuous functional visual acuity measurement system in dry eye syndrome. *Am J Ophthalmol* 2005;139(2):253-258.
- Goto E, Ishida R, Kido M, et al. Optical aberrations and visual disturbances associated with dry eye. *Ocul Surf* 2006; 4(4):207-213.
- Kaido M, Ishida R, Dogru M, Tamaoki T, Ysubota K. Efficacy of punctum plug treatment in short break-up time dry eye. *Optom Vis Sci* 2008;85(8):758-763.
- Fox RI, Robinson CA, Curd JG, Kozin F, Howell FV. Sjogren's syndrome: proposed criteria for classification. *Arthritis Rheum* 1986;29(5):577-585.
- Sotozono C, Ang LP, Koizumi N, et al. New grading system for the evaluation of chronic ocular manifestations in patients with Stevens-Johnson syndrome. *Ophthalmology* 2007; 114(7):1294-1302.
- Van Bijsterveld OP. Diagnostic tests in the Sicca syndrome. *Arch Ophthalmol* 1969;82(1):10-14.
- Suzukamo Y, Oshika T, Yuzawa M, et al. Psychometric properties of the 25-item National Eye Institute Visual Function Questionnaire (NEI VFQ-25), Japanese version. *Health Qual Life Outcome* 2005;3:65.
- Miyata K, Amano S, Sawa M, Nishida T. A novel grading method for superficial punctate keratopathy magnitude and its correlation with corneal epithelial permeability. *Arch Ophthalmol* 2003;121(11):1537-1539.
- Bron AJ, Benjamin L, Snibson GR. Meibomian gland diseases. Classification and grading of lid changes. *Eye* 1991; 5(Pt 4):395-411.

