

研究成果の刊行に関する一覧表

書籍

著者氏名	論文タイトル名	書籍全体の編集者名	書籍名	出版社名	出版地	出版年	ページ
該当なし							

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Itakura H, Yokoyama M, Matsuzaki M, Saito Y, Origasa H, Ishikawa Y, Oikawa S, Sasaki J, Hishida H, Kita T, Kitabatake A, Nakaya N, Sakata T, Shimada K, Shirato K,	Relationships between Plasma Fatty Acid Composition and Coronary Artery Disease.	J Atheroscler Thromb.	18(2)	99-107	2011
Sone H, Tanaka S, Tanaka S, Iimuro S, Oida K, Yamasaki Y, Oikawa S, Ishibashi S, Katayama S, Ohashi Y, Akanuma Y, Yamada N; for the Japan Diabetes Complications Study Group	Serum Level of Triglycerides Is a Potent Risk Factor Comparable to LDL Cholesterol for Coronary Heart Disease in Japanese Patients with Type 2 Diabetes:	J Clin Endocrinol Metab.	96	3448-3456	2011 Aug 24. [Epub ahead of print]
Itakura H, Yokoyama M, Matsuzaki M	The Change in Low-Density Lipoprotein Cholesterol Concentration is Positively Related to Plasma Docosahexaenoic Acid but not Eicosapentaenoic Acid.	J Atheroscler Thromb.	19(7)	673-9	2012
Sasaki J, Yokoyama M, Matsuzaki M, et al	Relationship between coronary artery disease and non-HDL-C, and effect of highly purified EPA on the risk of coronary artery disease in	J Atheroscler Thromb.	19(2)	194-204	2012
Dohi T, Miyauchi K, Ohkawa R, Nakamura K, Kishimoto T, Miyazaki T, Nishino A, Nakajima N, Yaginuma K, Tamura H, Kojima T, Yokoyama K, Kurata T, Shimada K, Yatomi Y, Daida H.	Increased circulating plasma lysophosphatidic acid in patients with acute coronary syndrome. Clin Chim Acta.	Chim Acta	413	207-212	2011

Inoue K, Kodama T, Daida H.	Pentraxin 3: a novel biomarker for inflammatory cardiovascular disease.	Int J Vasc Med	657025, doi:10.1155/2012/657025	2012
Fukushima Y, Ohmura H, Mokuno H, et al.	Non-high-density lipoprotein cholesterol is a practical predictor of long-term cardiac death after coronary artery bypass grafting.	Atherosclerosis	221 206-211	2012
Kasai T, Miyauchi K, Kubota N, et al.	Probucol therapy improves long-term (>10-year) survival after complete revascularization: a propensity analysis.	Atherosclerosis	220 463-469	2012
Teramoto T, Sasaki J, Ishibashi S, Birou S, Daida H, Dohi S, Egusa G, Hiro T, Hirobe K, Iida M, Kihara S, Kinoshita M, Maruyama C, Ohta T, Okamura T, Yamashita S, Yokode M, Yokote K; Japan Atherosclerosis Society.	Executive summary of the Japan Atherosclerosis Society (JAS) guidelines for the diagnosis and prevention of atherosclerotic cardiovascular diseases in Japan -2012 version.	J Atheroscler Thromb	20 517-523	2013
Sai E, Shimada K, Yokoyama T, Sato S, Miyazaki T, Hiki M, Tamura Y, Aoki S, Watada H, Kawamori R, Daida H.	Association between myocardial triglyceride content and cardiac function in healthy subjects and endurance athletes.	PLoS One	8(4) 1-6	2013
Matsumori R, Miyazaki T, Shimada K, Kume A, Kitamura Y, Oshida K, Yanagisawa N, Kiyonagi T, Hiki M, Fukao K, Hirose K, Ohsaka H, Mokuno H, Daida H.	High levels of very long-chain saturated fatty acid in erythrocytes correlates with atherogenic lipoprotein profiles in subjects with metabolic syndrome.	Diabetes Res Clin Pract	99 12-18	2013

Kasai T, Miyauchi K, Yanagisawa N, Kajimoto K, Kubota N, Ogita M, Tsuboi S, Amano A, Daida H.	Mortality risk of triglyceride levels in patients with coronary artery disease.	Heart	99	22-29	2013
Nakatani K, Sugimoto T, Masuda D, Okano R, Oya T, Monden Y, Yamashita T, Kawase R, Nakaoka H, Inagaki M, Yuasa-Kawase M, Tsubakio-Yamamoto K, Ohama T, Nishida M, Ishigami M, Komuro I, Yamashita S.	Serum apolipoprotein B-48 levels are correlated with carotid intima-media thickness in subjects with normal serum triglyceride levels.	Atherosclerosis	218(1)	226-32	2011
Masuda D, Sugimoto T, Tsujii K, Inagaki M, Nakatani K, Yuasa-Kawase M, Tsubakio-Yamamoto K, Ohama T, Nishida M, Ishigami M, Kawamoto T, Matsuyama A, Sakai N, Komuro I, Yamashita S.	Correlation of fasting serum apolipoprotein B-48 with coronary artery disease prevalence.	Eur J Clin Invest	42(9)	992-9	2012
Mugii S, Hanada H, Okubo M, Masuda D, Takeoka K, Hiedaka Y, Ohama T, Matsuyama A, Nakagawa-Toyama Y, Nishida M, Ishigami M, Komuro I, Yamashita S.	Thyroid function influences serum apolipoprotein B-48 levels in patients with thyroid disease.	<i>J Atheroscler Thromb.</i>	19(10)	890-6	2012
Yuasa-Kawase M, Masuda D, Kitazume-Taneike R, Yamashita T, Kawase R, Nakaoka H, Inagaki M, Nakatani K, Tsubakio-Yamamoto K, Ohama T, Toyama-Nakagawa Y, Nishida M, Ishigami M, Saito M, Eto M, Matsuyama A, Komuro I, Yamashita S.	Apolipoprotein B-48 to triglyceride ratio is a novel and useful marker for detection of type III hyperlipidemia after anti-hyperlipidemic intervention.	<i>J Atheroscler Thromb.</i>	19(9)	862-71	2012

Katsuren K, Nakamura K, Ohta T.	Effect of body mass index Z-score on adverse levels of cardiovascular disease risk factors.	Pediatrics International	54	200-204	2012
Asada S, Kuroda M, Aoyagi Y, Fukaya Y, Tanaka S, Konno S, Tanio M, Aiso M, Satoh K, Okamoto Y, Nakayama T, Saito Y, Bujo H.	Ceiling culture-derived proliferative adipocytes retain high adipogenic potential suitable for use as a vehicle for gene transduction therapy.	<i>Am J Physiol Cell Physiol.</i>	301	C181-C185	2011
Naito S, Kamata M, Furuya M, Hayashi M, Kuroda M, Bujo H, Kamata K.	Amelioration of circulating lipoprotein profile and proteinuria in a patient with LCAT deficiency due to a novel mutation (Cys74Tyr) in the lid region of LCAT under a fat-restricted diet and ARB treatment.	Atherosclerosis	228	193-197	2013
Arai H, Kokubo Y, Watanabe M, Sawamura T, Ito Y, Minagawa A, Okamura T, Miyamoto Y,	Small Dense Low-Density Lipoproteins Cholesterol can Predict Incident Cardiovascular Disease in an Urban Japanese Cohort	The Suita Study, <i>J Aththeroscler Thromb,</i>	20	195-203	2013
Arai H, Ishibashi S, Bujo H, Hayashi T, Yokoyama S, Oikawa S, Kobayashi J, Shirai K, Ota T, Yamashita S, Gotoda T, Harada-Shiba M, Sone H, Eto M, Suzuki H, Yamada N	Management of type IIb dyslipidemia,	<i>J Atheroscler Thromb,</i>	19	105-114	2012

Harada-Shiba M, Arai H, Okamura T, Yokote K, Oikawa S, Nohara A, Okada T, Ohta T, Bujo H, Watanabe M, Wakatsuki A, and Yamashita S,	Multicenter Study to Determine the Diagnostic Criteria of Heterozygous Familial Hypercholesterolemia in Japan,	J Atheroscler Thromb,	19	1019-26	2012
Harada-Shiba M, Arai H, Oikawa S, Ohta T, Okada T, Okamura T, Nohara A, Bujo H, Yokote K, Wakatsuki A, Ishibashi S, and Yamashita S,	Guidelines for the Management of Familial Hypercholesterolemia,	J Atheroscler Thromb,	19	1043-60	2012
Yokoyama S, Yamashita S, Ishibashi S, Sone H, Oikawa S, Shirai K, Ohta T, Bujo H, Kobayashi J, Arai H, Harada-Shiba M, Eto M, Hayashi T, Gotoda T, Suzuki H, Yamada N,	Background to Discuss Guidelines for Control of Plasma HDL-Cholesterol in Japan,	J Atheroscler Thromb,	19	207-212	2012
Mabuchi H, Nohara A, Noguchi T, Kobayashi J, Kawashiri MA, Tada H, Nakanishi C, Mori M, Yamagishi M, Inazu A, Koizumi J; Hokuriku FH Study Group	Molecular genetic epidemiology of homozygous familial hypercholesterolemia in the okuriku district of Japan.	Atherosclerosis	214	404-407	2011
Tada H, Kawashiri MA, Ohtani R, Noguchi T, Nakanishi C, Konno T, Hayashi K, Nohara A, Inazu A, Kobayashi J, Mabuchi H, Yamagishi M	A novel type of familial hypercholesterolemia: Double heterozygous mutations in LDL receptor and LDL receptor adaptor protein 1 gene	Atherosclerosis	219	663-666	2011
Tada H, Kawashiri MA, Ikegami K, Terao Y, Noguchi T, Nakanishi C, Tsuchida M, Takata M, Miwa K, Konno T, Hayashi K, Nohara A, Inazu A, Kobayashi J, Mabuchi H, Yamagishi M	Altered metabolism of low-density lipoprotein and very-low-density lipoprotein remnant in autosomal recessive hypercholesterolemia	results from stable isotope kinetic study in vivo. Circ Cardiovasc Genet	5(1)	35-41	2012 Feb 1

Harada-Shiba M, Arai H, Okamura T, Yokote K, Oikawa S, Nohara A, Okada T, Ohta T, Bujo H, Watanabe M, Wakatsuki A, Yamashita S	Multicenter study to determine the diagnosis criteria of heterozygous familial hypercholesterolemia in Japan	J Atheroscler Thromb	19(11)	1019-26	2012 Nov 26
Harada-Shiba M, Arai H, Okamura T, Yokote K, Oikawa S, Nohara A, Okada T, Ohta T, Bujo H, Watanabe M, Wakatsuki A, Yamashita S	Multicenter study to determine the diagnosis criteria of heterozygous familial hypercholesterolemia in Japan	J Atheroscler Thromb	19(11)	1019-26	2012 Nov 26
Sone H, Tanaka S, Tanaka S, Iimuro S, Ishibashi S, Oikawa S, Shimano H, Katayama S, Ohashi Y, Akanuma Y, Yamada N; Japan Diabetes Complications Study Group.	Comparison of various lipid variables as predictors of coronary heart disease in Japanese men and women with type 2 diabetes: subanalysis	Diabetes Care.	35(5)	1150-7	2012
Fujihara K, Suzuki H, Sato A, Kodama S, Heianza Y, Saito K, Iwasaki H, Kobayashi K, Yatoh S, Takahashi A, Yamada N, Sone H, Shimano H	Carotid Artery Plaque and LDL-to-HDL Cholesterol Ratio Predict Atherosclerotic Status in Coronary Arteries in Asymptomatic	J Atheroscler Thromb.	20(5)	452-464	2013
Yokoyama S, Yamashita S, Ishibashi S, Sone H, Oikawa S, Shirai K, Ohta T, Bujo H, Kobayashi J, Arai H, Harada-Shiba M, Eto M, Hayashi T, Goto da T, Suzuki H, Yamada	Background to Discussion Guidelines for Control of Plasma HDL-Cholesterol in Japan. J	Atheroscler Thromb	19	207-212	2012
Ina K, Hayashi T, Araki A, Kawashima S, Sone H, Watanabe H, Ohru T, Yokote K, Takemoto M, Kubota K, Noda M, Notomi H, Ding QF, Zhang J, Yu ZY, Yoon BK, Nomura H, Kuzuya M; Japan CD M Group.	Importance of high-density lipoprotein cholesterol levels in elderly diabetic individuals with type IIb dyslipidemia: A 2-year survey of cardiovascular events.	Geriatr Gerontol Int		1-5	2013
Sun L, Ishida T, Miyashita K, Kinoshita N, Mori K, Yasuda T, Toh R, Nakajima K, Imamura S, Hirata K.	Plasma activity of endothelial lipase impacts high-density lipoprotein metabolism and coronary risk factors in humans.	J Atheroscler Thromb. In press.	20	1-9	2013

Ishida T, Miyashita K, Shimizu M, Kinoshita N, Mori K, Sun L, Yasuda T, Imamura S, Nakajima K, Stanhope KL, Havel PJ, Hirata K.	ELISA system for human Endothelial lipase.	Clin Chem	58	1656-64	2012
Gotoda T, Shirai K, Ohta T, Kobayashi J, Yokoyama S, Oikawa S, Bujo H, Ishibashi S, Arai H, Yamashita S, Harada-Shiba M, Eto M, Hayashi T, Sone H, Suzuki H, Yamada N.	Diagnosis and management of type I and type V hyperlipoproteinemia	J. Atheroscler. Thromb.	19(1)	1-12	2012
Takase S, Osuga JI, Fujita H, Hara K, Sekiya M, Igarashi M, Takanashi M, Takeuchi Y, Izumida Y, Ohta K, Kumagai M, Nishi M, Kubota M, Masuda Y, Taira Y, Okazaki S, Iizuka Y, Yahagi N, Ohashi K, Yoshida H, Yanai H, Tada N, Gotoda T, Ishibashi S, Kadowaki T, Okazaki H	Apolipoprotein C-II deficiency with no rare variant in the APOC2 gene.	J. Atheroscler. Thromb.	20(5)	481-493	2013
Yamamoto T, Obika S, Nakatani M, Yasuhara H, Wada F, Shibata E, Shibata MA, Harada-Shiba M	Locked nucleic acid antisense inhibitor targeting apolipoprotein C-III efficiently and preferentially removed triglyceride from large VLDL particles from murine plasma	European Journal of Pharmacology	723	353-359	2014
Teramoto T, Sasaki J, Harada-Shiba M, Arai H, Ishibashi S, Birou S, Bujo H, Daida H, Dohi S, Egusa G, Hiro T, Hirobe K, Iida M, Kihara S, Kinoshita M, Maruyama C, Nomura A, Ohta T, Oikawa S, Okada T, Okamura T, Watatsuki A, Yamashita S, Yokode M, Yokote	Familial hypercholesterolemia, Executive summary of the Japanese Atherosclerosis Society(JAS) Guidelines for diagnosis and prevention of atherosclerotic cardiovascular diseases in Japan-2012 version.	J Atheroscler Thromb	21(1)	6-10	2013

Terasaki F, Morita H, Harada-Shiba M, Ohta N, Otsuka K, Nogi S, Miyamura M, Suzuki S, Ito T, Shimomura H, Katsumata T, Miyamoto Y, Ishizaka N.	Familial Hypercholesterolemia with Multiple Large Tendinous Xanthomas and Advanced coronary Artery Atherosclerosis	Intern Med	52	577-581	2013
Sugisawa T, Okamura T, Makino H, Watanabe M, Kishimoto I, Miyamoto Y, Iwamoto N, Yamamoto A, Yokoyama S, Harada-Shiba M	Defining patients at extremely high risk for coronary artery disease in heterozygous familial hypercholesterolemia	J Atheroscler Thromb	19(4)	369-375	2012