

considered that age-correlated increases in p16 expression might be implicated in the decreased replicative potential of self-renewing compartments with ageing [18]. Importantly, age-related expression of p16 might be closely involved in carcinogenesis in older patients. Singer et al. [24] reported that aged lymphoid progenitors exhibit severe growth defects due to the preferential expression of p16 and p19ARF, while their downregulation restored susceptibility to transformation. Baker et al. [25] demonstrated that late-life clearance of p16-positive senescent cells attenuated progression of age-related disorders in mice, suggesting that some sort of factors secreted by p16-positive cells might be implicated in the cellular and tissue dysfunction. In this regard, Coppe et al. [26] revealed that p16 could promote cell senescence without inducing a senescence-associated secretory phenotype (SASP). To clarify the age-related effects of p16 in vivo, further studies using animal models are required.

Regulatory mechanism of p16 involved in carcinogenesis

Until recently, DNA methylation and genetic deletion have been widely regarded as the main factors for p16 inactivation in cancer cells. Extensive studies, however, have provided new insights into the regulatory mechanism of p16. For example, the RAS/B-RAF/ERK-mediated MAPK pathway is the most well-known signaling cascade for inducing p16 expression [18]. RAS stimulates the ERK-mediated activation of ETS1/2 transcriptional factors, leading to the induction of p16 [27]. Therefore, together with the fact that various types of cancer cells exhibit N-RAS or B-RAF mutations with homozygous INK4a/ARF deletion, inactivation of p16 might potentiate the early steps of carcinogenesis in cells with activated RAS [18].

Several molecules have been newly identified as direct inhibitors of p16. The first player is inhibitor of DNA binding 1 (ID-1), a family member of helix–loop–helix (HLH) transcriptional regulatory proteins. ID-1 inhibits transcriptional activation at E-boxes within the p16/INK4a promoter, and thereby can inactivate p16 without epigenetic changes at the INK4/ARF locus [27, 28]. We recently reported that ID-1 was highly expressed in 42 of 112 patients (38 %) with liver cirrhosis. We found that the cumulative incidence of HCC development in patients with high ID-1 expression was significantly high ($P = 0.0008$), suggesting that increased expression of ID-1 may play an important role in the early step of hepatocarcinogenesis. Because the level of ID-1 expression was not correlated with the status of p16, its involvement in the induction of HCC development might not be mediated by p16 [29].

Furthermore, as ID-1 has been reported to cooperate with oncogenic RAS to induce metastatic mammary carcinoma in animal models [30], it might also act as an independent oncogene. A second player is BMI-1 (B lymphoma Mo-MLV insertion region 1 homolog), a transcriptional repressor belonging to the mouse Polycomb group [31]. Through its activity, BMI-1 plays an essential role in the maintenance of adult self-renewing hematopoietic cells [32] and is now considered to be a useful target for treating cancer [33]. A third player is cell division cycle 6 (CDC6), a member of the pre-replicative complex together with the origin recognition complex (ORC), chromatin licensing and DNA replication factor 1 (CDT1), and mini chromosome maintenance (MCM). It has been reported that CDC6 induces RD(INK4/ARF)-dependent transcriptional repression, recruits histone deacetylases, and promotes heterochromatinization of the INK4/ARF locus, resulting in the decreased expression of p16 [34]. Because there have been no reports of the relationship between CDC6 and p16 in clinical samples, it would be intriguing to investigate the status of CDC6 along with p16 in HCC.

More recently, chromatin remodeling at the INK4a/ARF/INK4b locus has been denoted as a potential mechanism for p16 inactivation. Kotake et al. [35] reported that lysine 4 of histone 3 (H3K4) is methylated at the p16-encoding locus in cells with high expression of p16, and H3K4 methyltransferase MLL1 is required for the induction of p16 in RAS-activated cells. Similarly, Dhawan et al. [36] reported that age-related repression of the INK4a/ARF locus was associated with decreased BMI-1 binding, loss of histone H2A ubiquitination, increased MLL1 recruitment, and a concomitant increase in the trimethylation of H3K4. Further, Agger et al. [37] reported that expression of the histone H3 Lys27 (H3K27) demethylase jumonji domain containing 3 (JMJD3) was induced upon activation of the RAS–RAF signaling pathway, and contributed to the transcriptional activation of p16/INK4A. Moreover, Yao et al. [38] revealed that H3K27 trimethylation was an early event in the silencing of p16 during tumor development. Currently, only a few reports have examined H3K4 and H3K27 methylation in solid cancers, and there have been no reports in HCC to date. Follow-up studies of these p16 regulators in subjects with liver cirrhosis would be of value for developing useful biomarkers to enable prediction of patients with a high risk of HCC.

p16 and hepatocellular carcinoma

To date, many studies have led to the suggestion that p16 is mainly inactivated in HCC by epigenetic changes at the gene promoter region [10, 39]. Non-epigenetic changes of p16, such as allelic loss or point mutation, have only been

reported in specific geographic areas and in some inherited cases. For example, homozygous deletion of the *INK4a/ARF* locus has been detected in 25 of 41 HCCs (61 %) in Korean subjects with HCC [40], and it has been reported that four of 26 patients with familial HCC (15 %) carried hemizygous germ-line point mutations of the *p16* gene [41]. In contrast in Taiwan, homozygous deletion of the *p16* gene was only identified in one of 30 HCCs (3 %) and no aberrant DNA methylation and mutation were detected [42], suggesting that the genetic alteration pattern of *p16* is diverse among different geographic areas. Thorough examinations of their etiological backgrounds are important for further understanding.

With respect to the observations in Japanese cases, we previously examined 60 HCC patients and found no allelic loss of the *p16* gene [20]. Point mutation of the *p16* gene was only detected in four of 60 HCCs (6.6 %), suggesting that genetic changes of the *p16* gene are rare in Japanese subjects. Immunohistochemical analysis showed that *p16* expression was negative in 29 of 60 tumors (48 %), and the degree of *p16* expression was inversely correlated with the histopathological grading of the tumors: negative immunostaining was observed in one (20 %) of five well-differentiated HCCs, 13 (50 %) of 26 moderately differentiated HCCs, 11 (48 %) of 23 poorly differentiated HCCs, and four (67 %) of six undifferentiated HCCs. The results of methylation-specific PCR (MSP) revealed that 24 (83 %) of the 29 HCCs with negative *p16* immunostaining exhibited DNA methylation in the promoter region, indicating that DNA methylation is the main reason for *p16* inactivation. We addressed the degree of DNA methylation in the *p16* gene using a methylation-sensitive single nucleotide primer extension (Ms-SNuPE) method, which is a previously reported protocol based on the measurement of the band intensities of the methylated *p16* gene incorporating [³²P]dCTP and the un-methylated *p16* gene incorporating [³²P]TTP [43]. Our results showed that the degrees of CpG methylation in HCC with *p16*-negative expression ranged from 60 to 85 %, indicating that the *p16* gene was extensively methylated at multiple sites in HCC.

Intriguingly, epigenetic changes of the *p16* gene can be also observed in non-tumorous liver tissues. Kaneto et al. [44] reported that DNA methylation of the *p16* gene was detected in five of 17 liver cirrhosis (29.4 %) and four of 17 chronic hepatitis (23.5 %). Li et al. [45] observed that the *p16* gene was methylated in six of 38 (16 %) cases with chronic hepatitis or cirrhosis. We examined the genetic and epigenetic changes of the *p16* gene in chronic liver injury and found that it was not mutated in any of the cases (unreported data), while methylated *p16* promoter was detected in four of 112 (4 %) cases of liver cirrhosis [29]. Although the reported frequencies of *p16* gene methylation in non-tumorous liver tissues differ among the studies, it

should be kept in mind that techniques with higher sensitivity and specificity are required for the precise detection of methylated DNA. Methylation-specific PCR is a commonly used standard method for detecting methylated DNA, and is based on the examination of bisulfite-converted genomic DNA using methylated DNA-specific primers. This technique has been used in many studies to examine the methylation status of the *p16* gene; however, false-negative results due to contamination of the tissue samples with PCR inhibitors cannot be ruled out. Recently, bisulfite sequencing has been applied for the screening of the methylation status of different DNA regions, and the MethyLight method (a sensitive and quantitative fluorescent real-time PCR technique) has been reported as the best method for the practical screening of methylated PCR. Using this new technique, Csepreg et al. [46] identified that methylation of CpG islands at the first exon of *p16* gene could be detected by MethyLight but not by MSP. Therefore, to understand the clinical significance of *p16* in non-tumorous liver tissues, surveillance using highly sensitive methods such as MethyLight technique would be useful.

p27 as a tumor suppressor

p27 is a member of the Cip/Kip family of CDKs involved in G1 arrest and is encoded by the cyclin-dependent kinase inhibitor 1B (*CDKN1B*) gene [47, 48]. *p27* binds and inhibits cyclinE-CDK2 complexes, leading to the hypophosphorylation of pRb in arrested cells. During the cell cycle, the level of *p27* expression is stable and post-translational mechanisms mainly regulate its activity including protein expression levels, sequestration between the nucleus and cytoplasm, protein phosphorylation, and proteasome-mediated protein degradation. Of note, it has been unveiled that the role of *p27* is not restricted to a CDK inhibitor, and different phosphorylation sites independently determine its activity. For example, mitogenic stimuli result in phosphorylation of *p27* at serine 10 (S10) and sequestration of *p27* from the nucleus to the cytoplasm, which leads to repressed inhibition of the cyclin E/CDK2 complex. When *p27* is phosphorylated at threonine 157 and 198 (Thr157 and Thr198) through AKT signaling, *p27* is located in the cytoplasm and sequestered from cyclin E/CDK2 to cyclin D1/CDK4 complexes, which leads *p27* to act as an activator of CDK4. In turn, *p27* is phosphorylated at threonine 187 (T187) by cyclin E/CDK2 complexes in late G1–S, which causes protein degradation by the recruited S phase kinase-associated protein 2 (SKP2) ubiquitin ligase-bound SCF complexes (consisting of Cullin-1, Skp1 and RBX1) [49, 50]. Recent studies have reported that decreased expression of *p27* is correlated with poor prognosis in cancer patients, suggesting that it might




Fig. 2 **a** Key regulatory mechanisms controlling p27 activity. p27 inhibits cyclin E/CDK2 complexes at the G1–S phase. p27 is inactivated by translational modifications, including protein degradation, protein phosphorylation, cytoplasmic localization and sequestration into cyclin D/CDK4 complexes. **b** p27 is an adverse prognostic biomarker of HCC. Cases with high p27 expression can be categorized into two groups according to the labeling index of p16. In cases with high p27 and low p16 expression, CDK2 activities are strong and p27 is sequestered into cyclin D/CDK4 complexes, leading to poor prognoses

be a useful prognostic biomarker. In many types of cancer cells, gene mutations of p27 are rare but p27 expression is decreased through proteasome-mediated protein degradation. In fact, an inverse relationship between the levels of SKP2 and p27 has been observed in many types of human malignancies with poorer prognosis [51, 52] (Fig. 2).

p27 and hepatocellular carcinoma

To date, accumulating evidence has suggested that decreased p27 expression can be considered as an adverse prognostic biomarker in HCC [53–57]. Consistent with previous studies, we found that the 5-year survival rates were 62 and 93 % in low p27-expressing patients [labeling index (LI) of <50 by immunohistochemical analysis] and

high p27-expressing patients (LI of >50 %), respectively. Low-p27 expressers represented a relative risk of dying from HCC as compared to high-p27 expressers ($P = 0.002$) [10]. Intriguingly, we also found that the cases of high-p27 expressers represented diverse levels of Ki-67 LI (a cell proliferation marker) and could be categorized into two groups with low and high Ki-67 expression. Of 40 cases with high-p27 expressers, 26 (65 %) showed a Ki-67 LI above 20 % (22–42 %), while 14 (35 %) showed a Ki-67 LI less than 20 % (2–13 %). To address the diverse cell proliferation status among the cases of high-p27 expressers, we set out to examine the activities of CDK2 in these cases. As expected, a relatively close relationship between the level of CDK2 activities and Ki-67 LI was observed, and the CDK2 kinase activities were significantly strong in all HCC cases in a subgroup with a Ki-67 LI of >20 %. In cases of p27 expressers representing strong CDK2 activities, heating the lysates of the protein samples effectively recovered the inhibitory function of p27, indicating that p27 alone was functional. In these tumors, p27 was found to be exclusively bound to cyclin D1/CDK4 complexes, supporting the idea that p27 can be inactivated by inappropriate protein–protein interaction in HCC. More importantly, we found that p16 was undetectable in all HCCs ($n = 8$), representing exclusive binding of p27 with cyclin D1/CDK4 complexes. Taken together, our data may indicate that the level of p27 expression does not accurately indicate the degree of functional activity, and in tumors with positive p27 expression p16 might act as potent determinant of the conformational change of p27. Moreover, when the cases of high p27 expression ($n = 40$) were categorized according to p16-negative and p16-positive status, the 5-year overall survival rates were 70 and 100 %, respectively ($P = 0.0004$). Multivariate analysis showed that p16 was an independent adverse prognostic factor in HCC ($P = 0.003$). Taken together, we surmise that combined examination of p16 and p27 might be useful for the precise prognostic diagnosis of HCC.

Cytoplasmic p27: another aspect of cell cycle inhibition

Although p27 has been widely considered a favorable prognostic factor in patients with different types of cancer, it has been shown that p27 exerts cell cycle-independent oncogenic properties [58, 59]. Post-translational modifications through phosphatidylinositol-3 kinase (PI3K)/AKT and RAS/mitogen-activated protein kinase (MAPK) signaling have been reported to be the main determinants of the function of p27 [60–65]. In the case of HCC, several studies have reported that the level of p27 in the cytoplasmic fraction of tumor cells was associated with cancer progression. For example, Nan et al. [66] reported an association with

clinical stage and the degree of invasion of HCC. Zhou et al. [67] examined 528 cases of HCC and reported that phosphorylated AKT and cytoplasmic p27 expression showed the strongest associations with pathological parameters of HCC. We previously reported that seven of 40 (17.5 %) HCCs exhibited cytoplasmic p27 expression and all the cases with cytoplasmic p27 had a Ki-67 LI of >20 % (22–42 %) [68]. Therefore, it is highly likely that nuclear p27 is a favorable prognostic marker, while cytoplasmic p27 is an adverse prognostic marker. For precise evaluation of the clinical significance of p27 in HCC, a comprehensive analysis of p27 including its expression level, localization, and the site of phosphorylation is required.

Future perspective

Recent progress has enabled improved diagnosis and management of HCC, but its prognosis still remains dismal. Identification of biomarkers is an important step for developing an efficient treatment. Dysfunction of either p16 or p27 causes abnormal cell proliferation, eventually leading to cancer development. The modality and mechanism for inactivating p16 and p27 are unique: epigenetic changes are the main cause of p16 inactivation, and either protein degradation or inappropriate protein–protein interaction results in the inactivation of p27. We have extensively examined the statuses of both CDK inhibitors in HCC, and our results show that their role in tumor aggressiveness is complex. Our hypothesis that p16 rather than p27 might be a useful prognostic factor in HCC should be investigated in the near future.



Dr. Yasunobu Matsuda, of the Department of Medical Technology, Niigata University Graduate School of Health Sciences, Niigata, is the winner of the Japanese Society for Clinical Molecular Morphology Award for Promoting Young Researchers in 2011. Dr. Matsuda is recognized for his great contribution in molecular morphological analysis of human hepatocellular carcinoma and advancement, and also for his contribution in the prediction of the prognosis in hepatocellular carcinoma patients, using biomarkers such as p16 and p17.

References

- Llovet JM, Burroughs A, Bruix J (2003) Hepatocellular carcinoma. *Lancet* 362:1907–1917
- El-Serag HB (2011) Hepatocellular carcinoma. *New Engl J Med* 365:1118–1127
- Matsuda Y, Ichida T, Fukumoto M (2011) Hepatocellular carcinoma and liver transplantation: clinical perspective on molecular targeted strategies. *Med Mol Morphol* 44:117–124
- Shariff MIF, Cox IJ, Gomaa AI, Khan S, Gedroyc W, Taylor-Robinson SD (2009) Hepatocellular carcinoma: current trends in worldwide epidemiology, risk factors, diagnosis and therapeutics. *Expert Rev Gastroenterol Hepatol* 3:353–367
- Gao J, Xie L, Yang W-S, Zhang W, Gao S, Wang J, Xiang Y-B (2012) Risk factors of hepatocellular carcinoma—current status and perspectives. *Asian Pac J Cancer Prev* 13:743–752
- Feitelson M, Sun B, Satiroglu T, Liu J, Pan J, Lian Z (2002) Genetic mechanisms of hepatocarcinogenesis. *Oncogene* 21:2593–2604
- Effendi K, Sakamoto M (2010) Molecular pathology in early hepatocarcinogenesis. *Oncology* 78:157–160
- Gramantieri L, Fornari F, Callegari E, Sabbioni S, Lanza G, Croce CM, Bolondi L, Negrini M (2008) MicroRNA involvement in hepatocellular carcinoma. *J Cell Mol Med* 12:2189–2204
- Greenbaum LE (2004) Cell cycle regulation and hepatocarcinogenesis. *Cancer Biol Ther* 3:1200–1207
- Matsuda Y, Ichida T (2006) P16 and P27 are functionally correlated during the progress of hepatocarcinogenesis. *Med Mol Morphol* 39:169–175
- Matsuda Y (2008) Molecular mechanism underlying the functional loss of cyclin-dependent kinase inhibitors p16 and p27 in hepatocellular carcinoma. *World J Gastroenterol* 14:1734–1740
- Borzio M, Trerè D, Borzio F, Ferrara R, Bruno S, Roncalli M, Colloredo G, Leandro G, Oliveri F, Derenzini M (1998) Hepatocyte proliferation rate is a powerful parameter for predicting hepatocellular carcinoma development in liver cirrhosis. *Mol Pathol* 51:96–101
- Shibata M, Morizane T, Uchida T, Yamagami T, Onozuka Y, Nakano M (1998) Early reports irregular regeneration of hepatocytes and risk of hepatocellular carcinoma in chronic hepatitis and cirrhosis with hepatitis-C-virus infection. *Lancet* 351:1773–1777
- Murray AW (2004) Recycling the cell cycle: cyclins revisited. *Cell* 116:221–234
- Zhou Q, Hopp T, Fuqua SA, Steeg PS (2001) Cyclin D1 in breast premalignancy and early breast cancer: implications for prevention and treatment. *Cancer Lett* 162:3–17
- Poomsawat S, Buajeeb W, Khovichunkit S, Punyasingh J (2010) Alteration in the expression of cdk4 and cdk6 proteins in oral cancer and premalignant lesions. *J Oral Pathol Med* 39:793–799
- Sherr CJ, Roberts JM (1999) CDK inhibitors: positive and negative regulators of G1-phase progression. *Genes Dev* 13:1501–1512
- Kim WY, Sharpless NE (2006) The regulation of INK4/ARF in cancer and aging. *Cell* 127:265–275
- Pagano M, Tam SW, Theodoras AM, Beer-Romero P, Del Sal G, Chau V, Yew PR, Draetta GF, Rolfe M (1995) Role of the ubiquitin–proteasome pathway in regulating abundance of the cyclin-dependent kinase inhibitor p27. *Science* 269:682–685
- Matsuda Y, Ichida T, Matsuzawa J, Sugimura KAH (1999) p16(INK4) is inactivated by extensive CpG methylation in human hepatocellular carcinoma. *Gastroenterology* 116:394–400
- Li H, Collado M, Villasante A, Strati K, Ortega S, Cañamero M, Blasco MA, Serrano M (2009) The Ink4/Arf locus is a barrier for iPS cell reprogramming. *Nature* 460:1136–1139
- Zindy F, Quelle DE, Roussel MF, Sherr CJ (1997) Expression of the p16INK4a tumor suppressor versus other INK4 family members during mouse development and aging. *Oncogene* 15:203–211
- Krishnamurthy J, Torrice C, Ramsey MR, Kovalev GI, Al-Regaiey K, Su L, Sharpless NE (2004) Ink4a/Arf expression is a biomarker of aging. *J Clin Invest* 114:1299–1307

24. Signer RAJ, Montecino-Rodriguez E, Witte ON, Dorshkind K (2008) Aging and cancer resistance in lymphoid progenitors are linked processes conferred by p16Ink4a and Arf. *Genes Dev* 22:3115–3120
25. Baker DJ, Wijshake T, Tchkonja T, LeBrasseur NK, Childs BG, Van de Sluis B, Kirkland JL, Van Deursen JM (2011) Clearance of p16Ink4a-positive senescent cells delays ageing-associated disorders. *Nature* 479:232–236
26. Coppé J-P, Rodier F, Patil CK, Freund A, Desprez P-Y, Campisi J (2011) Tumor suppressor and aging biomarker p16(INK4a) induces cellular senescence without the associated inflammatory secretory phenotype. *J Biol Chem* 286:36396–36403
27. Ohtani N, Zebedee Z, Huot T, Stinson J, Sugimoto M, Ohashi Y, Sharrocks A, Peters G, Hara E (2001) Opposing effects of Ets and Id proteins on p16 INK4a expression during cellular senescence. *Nature* 409:1067–1070
28. Alani RM, Young AZ, Shifflett CB (2001) Id1 regulation of cellular senescence through transcriptional repression of p16/Ink4a. *Proc Natl Acad Sci USA* 98:7812–7816
29. Matsuda Y, Yamagiwa S, Takamura M, Honda Y, Ishimoto Y, Ichida T, Aoyagi Y (2005) Overexpressed Id-1 is associated with a high risk of hepatocellular carcinoma development in patients with cirrhosis without transcriptional repression of p16. *Cancer* 104:1037–1044
30. Swarbrick A, Roy E, Allen T, Bishop JM (2008) Id1 cooperates with oncogenic Ras to induce metastatic mammary carcinoma by subversion of the cellular senescence response. *Proc Natl Acad Sci USA* 105:5402–5407
31. Jacobs JJ, Kieboom K, Marino S, DePinho RA, Van Lohuizen M (1999) The oncogene and Polycomb-group gene bmi-1 regulates cell proliferation and senescence through the ink4a locus. *Nature* 397:164–168
32. Park I, Qian D, Kiel M, Becker MW, Pihalja M, Weissman IL, Morrison SJ, Clarke MF (2003) Bmi-1 is required for maintenance of adult self-renewing haematopoietic stem cells. *Nature* 423:302–305
33. Cao L, Bombard J, Cintron K, Sheedy J, Weetall ML, Davis TW (2011) BMI1 as a novel target for drug discovery in cancer. *J Cell Biochem* 112:2729–2741
34. Gonzalez S, Klatt P, Delgado S, Conde E, Lopez-Rios F, Sanchez-Cespedes M, Mendez J, Antequera F, Serrano M (2006) Oncogenic activity of Cdc6 through repression of the INK4/ARF locus. *Nature* 440:702–706
35. Kotake Y, Zeng Y, Xiong Y (2009) DDB1-CUL4 and MLL1 mediate oncogene-induced p16INK4a activation. *Cancer Res* 69:1809–1814
36. Dhawan S, Tschen S-I, Bhushan A (2009) Bmi-1 regulates the Ink4a/Arf locus to control pancreatic beta-cell proliferation. *Genes Dev* 23:906–911
37. Agger K, Cloos PAC, Rudkjaer L, Williams K, Andersen G, Christensen J, Helin K (2009) The H3K27me3 demethylase JMJD3 contributes to the activation of the INK4A-ARF locus in response to oncogene- and stress-induced senescence. *Genes Dev* 23:1171–1176
38. Yao J-Y, Zhang L, Zhang X, He Z-Y, Ma Y, Hui L-J, Wang X, Hu Y-P (2010) H3K27 trimethylation is an early epigenetic event of p16INK4a silencing for regaining tumorigenesis in fusion reprogrammed hepatoma cells. *J Biol Chem* 285:18828–18837
39. Zang J-J, Xie F, Xu J-F, Qin Y-Y, Shen R-X, Yang J-M, He J (2011) P16 gene hypermethylation and hepatocellular carcinoma: a systematic review and meta-analysis. *World J Gastroenterol* 17:3043–3048
40. Piao Z, Park C, Lee JS, Yang CH, Choi KY, Kim H (1998) Homozygous deletions of the CDKN2 gene and loss of heterozygosity of 9p in primary hepatocellular carcinoma. *Cancer Lett* 122:201–207
41. Chaubert P, Gayer R, Zimmermann A, Fontollet C, Stamm B, Bosman F, Shaw P (1997) Germ-line mutations of the p16INK4(MTS1) gene occur in a subset of patients with hepatocellular carcinoma. *Hepatology* 25:1376–1381
42. Lin Y, Chen C, Huang G, Lee P, Wang J, Chen D, Lu F, Sheu J (1998) Infrequent mutations and no methylation of CDKN2A (P16/MTS1) and CDKN2B (p15/MTS2) in hepatocellular carcinoma in Taiwan. *Eur J Cancer* 34:1789–1795
43. Gonzalgo ML, Jones PA (1997) Rapid quantitation of methylation differences at specific sites using methylation-sensitive single nucleotide primer extension (Ms-SNuPE). *Nucl Acids Res* 25:2529–2531
44. Kaneto H, Sasaki S, Yamamoto H et al (2001) Detection of hypermethylation of the p16(INK4A) gene promoter in chronic hepatitis and cirrhosis associated with hepatitis B or C virus. *Gut* 48:372–377
45. Li X, Hui A-M, Sun L, Hasegawa K, Torzilli G, Minagawa M, Takayama T, Makuuchi M (2004) p16INK4A hypermethylation is associated with hepatitis virus infection, age, and gender in hepatocellular carcinoma. *Clin Cancer Res* 10:7484–7489
46. Csepregi A, Ebert MP, Röcken C, Schneider-Stock R, Hoffmann J, Schulz H-U, Roessner A, Malfertheiner P (2010) Promoter methylation of CDKN2A and lack of p16 expression characterize patients with hepatocellular carcinoma. *BMC cancer* 10:317
47. Slingerland J, Pagano M (2000) Regulation of the cdk inhibitor p27 and its deregulation in cancer. *J Cell Physiol* 183:10–17
48. Sgambato A, Cittadini A, Faraglia B, Weinstein B (2000) Multiple functions of p27(Kip1) and its alterations in tumor cells: a review. *J Cell Physiol* 183:18–27
49. Nakayama KI, Hatakeyama S, Nakayama K (2001) Regulation of the cell cycle at the G1-S transition by proteolysis of cyclin E and p27Kip1. *Biochem Biophys Res Commun* 282:853–860
50. Pagano M (2004) Control of DNA synthesis and mitosis by the Skp2-p27-Cdk1/2 axis. *Mol Cell* 14:414–416
51. Carrano A, Eytan E, Hershko A, Pagano M (1999) SKP2 is required for ubiquitin-mediated degradation of the CDK inhibitor p27. *Nat Cell Biol* 1:193–199
52. Lloyd RV, Erickson LA, Jin L, Kulig E, Qian X, Chevillat JC, Scheithauer BW (1999) p27Kip1: a multifunctional cyclin-dependent kinase inhibitor with prognostic significance in human cancers. *Am J Pathol* 154:313–323
53. Fiorentino M, Altamari A, Errico AD (2000) Acquired expression of p27 is a favorable prognostic indicator in patients with hepatocellular carcinoma. *Clin Cancer Res* 6:3966–3972
54. Tannapfel A, Grund D, Katalinic A, Uhlmann D, Köckerling F, Haugwitz U, Wasner M, Hauss J, Engeland K, Wittekind C (2000) Decreased expression of p27 protein is associated with advanced tumor stage in hepatocellular carcinoma. *Int J Cancer* 355:350–355
55. Qin L, Ng I (2001) Expression of p27(KIP1) and p21(WAF1/CIP1) in primary hepatocellular carcinoma: clinicopathologic correlation and survival analysis. *Hum Pathol* 32:778–784
56. Armengol C, Boix L, Bachs O, Sole M, Fuster J, Sala M, Llovet JM, Rodes J, Bruix J (2003) p27 Kip1 is an independent predictor of recurrence after surgical resection in patients with small hepatocellular carcinoma. *J Hepatol* 38:591–597
57. Zhou Q, Wang G, Liang L, Zheng W, Pang Z (2003) Expression of p27, cyclin A in hepatocellular carcinoma and its clinical significance. *World J Gastroenterol* 9:2450–2454
58. Sgambato A, Zhang YJ, Arber N, Hibshoosh H, Doki Y, Ciamporrone M, Santella RM, Cittadini A, Weinstein IB (1997) Deregulated expression of p27Kip1 in human breast cancers. *Clin Cancer Res* 3:1879–1887
59. Fredersdorf S, Burns J, Milne AM, Packham G, Fallis L, Gillett CE, Royds JA, Peston D, Hall PA, Hanby AM, Barnes DM, Shousha S, O'Hare MJ, Lu X (1997) High level expression of

- p27Kip1 and cyclin D1 in some human breast cancer cells: inverse correlation between the expression of p27Kip1 and degree of malignancy in human breast and colorectal cancers. *Proc Natl Acad Sci USA* 94:6380–6385
60. Liu X, Sun Y, Ehrlich M, Lu T, Kloog Y, Weinberg RA, Lodish HF, Henis YI (2000) Disruption of TGF-beta growth inhibition by oncogenic ras is linked to p27Kip1 mislocalization. *Oncogene* 19:5926–5935
 61. Donovan JC, Milic A, Slingerland JM (2001) Constitutive MEK/MAPK activation leads to p27(Kip1) deregulation and antiestrogen resistance in human breast cancer cells. *J Biol Chem* 276:40888–40895
 62. Liang J, Zubovitz J, Petrocelli T, Kotchetkov R, Connor MK, Han K, Lee JH, Ciarallo S, Catzavelos C, Beniston R, Franssen E, Slingerland JM (2002) PKB/Akt phosphorylates p27, impairs nuclear import of p27 and opposes p27-mediated G1 arrest. *Nat Med* 8:1153–1160
 63. Viglietto G, Motti ML, Bruni P, Melillo RM, D'Alessio A, Califano D, Vinci F, Chiappetta G, Tschlis P, Bellacosa A, Fusco A, Santoro M (2002) Cytoplasmic relocalization and inhibition of the cyclin-dependent kinase inhibitor p27(Kip1) by PKB/Akt-mediated phosphorylation in breast cancer. *Nat Med* 8:1136–1144
 64. Liang J, Slingerland JM (2003) Multiple roles of the PI3K/PKB (Akt) pathway in cell cycle progression. *Cell Cycle* 2:339–345
 65. Kim J, Jonasch E, Alexander A, Short JD, Cai S, Wen S, Tsavachidou D, Tamboli P, Czerniak BA, Do KA, Wu KJ, Marlow LA, Wood CG, Copland JA, Walker CL (2009) Cytoplasmic sequestration of p27 via AKT phosphorylation in renal cell carcinoma. *Clin Cancer Res* 15:81–90
 66. Nan KJ, Jing Z, Gong L (2004) Expression and altered subcellular localization of the cyclin-dependent kinase inhibitor p27Kip1 in hepatocellular carcinoma. *World J Gastroenterol* 10:1425–1430
 67. Zhou L, Huang Y, Li J, Wang Z (2010) The mTOR pathway is associated with the poor prognosis of human hepatocellular carcinoma. *Med Oncol* 27:255–261
 68. Matsuda Y, Ichida T, Genda T, Yamagiwa S, Aoyagi Y, Asakura H (2003) Loss of p16 contributes to p27 sequestration by cyclin D(1)-cyclin-dependent kinase 4 complexes and poor prognosis in hepatocellular carcinoma. *Clin Cancer Res* 9:3389–3396

Value of Highly Sensitive Fucosylated Fraction of Alpha-Fetoprotein for Prediction of Hepatocellular Carcinoma Recurrence After Curative Treatment

Yasushi Tamura · Takeshi Suda · Shigeki Aarii · Michio Sata · Fuminori Moriyasu · Hiroshi Imamura · Seiji Kawasaki · Namiki Izumi · Tadatoshi Takayama · Norihiro Kokudo · Masakazu Yamamoto · Hiroko Iijima · Yutaka Aoyagi

Received: 13 February 2013 / Accepted: 20 March 2013 / Published online: 18 April 2013
© Springer Science+Business Media New York 2013

Abstract

Background The fucosylated fraction of alpha-fetoprotein (AFP-L3) has been used as a diagnostic marker for hepatocellular carcinoma (HCC). Recently, a highly sensitive immunoassay using an on-chip electrokinetic reaction and separation by affinity electrophoresis (micro-total analysis system; μ TAS) has been developed.

Aim The aim of this study was to investigate the relationship between changes in the serum AFP-L3 level measured by μ TAS assay and recurrence of HCC after curative treatment.

Methods A total of 414 HCC patients who met the Milan criteria and underwent hepatectomy or radiofrequency ablation were investigated prospectively for the relationship between HCC recurrence and values of tumor markers.

Results There were significant differences in recurrence-free survival between groups with and without AFP-L3

elevation measured before and after treatment ($p = 0.024$ and $p = 0.001$ for before and after treatment, respectively). Multivariate analysis revealed that AFP-L3 status ($p = 0.002$) measured 1 month after treatment was a significant independent predictor of HCC recurrence after curative treatment.

Conclusions Elevation of the serum AFP-L3 level before treatment is a predictor of HCC recurrence, and sustained elevation of the AFP-L3 level after treatment is an indicator of HCC recurrence. Repeated measurement of μ TAS AFP-L3 should be performed for surveillance of HCC recurrence after curative treatment.

Keywords Hepatocellular carcinoma · Alpha-fetoprotein · Fucosylated fraction of alpha-fetoprotein · Des-gamma-carboxy prothrombin

Y. Tamura (✉) · T. Suda · Y. Aoyagi
Division of Gastroenterology and Hepatology, Niigata University Graduate School of Medical and Dental Sciences, 757 Asahimachi Dori-1-Bancho, Chuo-ku, Niigata 951-8520, Japan
e-mail: ytamura@med.niigata-u.ac.jp

S. Aarii
Department of Hepato-Biliary-Pancreatic Surgery, Graduate School of Medicine, Tokyo Medical and Dental University, Tokyo, Japan

M. Sata
Division of Gastroenterology, Department of Medicine, Kurume University School of Medicine, Kurume, Japan

F. Moriyasu
Department of Gastroenterology and Hepatology, Tokyo Medical University, Tokyo, Japan

H. Imamura · S. Kawasaki
Department of Hepatobiliary-Pancreatic Surgery, Juntendo University School of Medicine, Tokyo, Japan

N. Izumi
Department of Gastroenterology and Hepatology, Musashino Red Cross Hospital, Tokyo, Japan

T. Takayama
Department of Digestive Surgery, Nihon University School of Medicine, Tokyo, Japan

N. Kokudo
Hepato-Biliary-Pancreatic Surgery Division, Artificial Organ and Transplantation Division, Department of Surgery, University of Tokyo, Tokyo, Japan

M. Yamamoto
Department of Surgery, Institute of Gastroenterology, Tokyo Women's Medical University, Tokyo, Japan

H. Iijima
Ultrasound Imaging Center, Division of Hepatobiliary and Pancreatic Medicine, Department of Internal Medicine, Hyogo College of Medicine, Hyogo, Japan

Introduction

Hepatocellular carcinoma (HCC) is the sixth most common malignancy and the third most common cause of cancer-related death [1]. Three serum markers, alpha-fetoprotein (AFP), fucosylated fraction of AFP (AFP-L3), and des-gamma-carboxy prothrombin (DCP), have been widely used for HCC surveillance and diagnosis. Among these markers, AFP is the most frequently used as a diagnostic marker for HCC in Japan [2, 3]. However, AFP levels are sometimes elevated in patients with chronic hepatitis and cirrhosis who have no evidence of HCC [4–6]. In contrast, AFP-L3 has been accepted as a specific marker for HCC by several investigators [7–9]. Moreover, its level predicts the malignant potential of HCC, and thus the expected outcome after treatment [10–15]. However, measurement of AFP-L3 has not always been reliable using serum samples with a low total AFP concentration determined by conventional lectin affinity electrophoresis, or using a liquid-phase binding assay system (LiBASys assay) [16]. Recently, a highly sensitive immunoassay for AFP-L3 using the on-chip electrokinetic reaction and separation by affinity electrophoresis (micro-total analysis system; μ TAS) has been developed [17, 18]. We have already reported that the μ TAS AFP-L3 is more sensitive for discriminating HCC from benign liver disease than the conventional LiBASys AFP-L3, particularly in subgroups with lower AFP concentrations and early stage HCC [19]. However, there has been little information about the μ TAS AFP, AFP-L3 and DCP measured after curative treatment of HCC.

In the present study, we investigated the relationship between the tumor markers μ TAS AFP, AFP-L3, and DCP, measured before and after curative treatment, and the likelihood of HCC recurrence. Additionally, we examined the changes in these markers in relation to treatment outcome.

Methods

Patients

A total of 420 HCC patients who met the Milan criteria (single tumor ≤ 5 cm in size or ≤ 3 tumors each ≤ 3 cm in size, and no macrovascular invasion) and who underwent hepatectomy or radiofrequency ablation at ten participating hospitals (Niigata University Medical and Dental Hospital, Medical Hospital of Tokyo Medical and Dental University, Kurume University Hospital, Tokyo Medical University Hospital, Juntendo University Hospital, Musashino Red Cross Hospital, Nihon University Itabashi Hospital, The University of Tokyo Hospital, Tokyo Women's Medical

University Hospital, Hyogo Medical College Hospital), between May 2008 and November 2009, were investigated prospectively. HCC recurrence was assessed using several imaging modalities every 3 months after treatment, and all recurrences were evaluated up to the end of January 2012. The mean observation time after treatment was 594 ± 319 days. Six patients were excluded from the study because their follow-up periods were less than 1 month, giving a final total of 414 patients who were enrolled. Among these patients, HCC recurrence was observed in 236 patients, and the remaining 178 showed no HCC recurrence within the study period. Informed consent was obtained from each patient, and the study protocol conformed with the ethical guidelines of the 1975 Declaration of Helsinki, as reflected in the a priori approval granted by each institution's human research committee.

Diagnosis of HCC and Laboratory Examination

Hepatocellular carcinoma patients were diagnosed using several imaging modalities, including computed tomography (CT), magnetic resonance imaging (MRI), and CT, during hepatic arteriography, giving due consideration to hyperattenuation in the arterial phase with washout in the late phase. Vascular invasion was evaluated on the basis of imaging modalities. In some cases that showed atypical features on imaging, ultrasound-guided biopsies were performed.

Simultaneous quantitative measurement of AFP-L1 (ng/ml) and AFP-L3 (ng/ml) was performed using the μ TAS assay (Wako Pure Chemical Industries Ltd., Osaka, Japan) [17, 18]. Total AEP concentration (ng/ml) in the serum sample was determined by summation of AFP-L3 and AFP-L1, and then the percentage of the AFP-L3 level was calculated. The serum level of des-gamma-carboxy prothrombin (DCP) was also measured using a μ TAS assay, except for patients who had taken warfarin, a DCP-inducing agent. The lower limits of quantitation for AFP (L1 and L3) and DCP were 0.3 ng/ml and 5 mAU/ml, respectively [17, 18]. When AFP-L3 was not detectable, the percentage of AFP-L3 was defined as 0.5 %. AFP, AFP-L3, and DCP were measured in the same serum sample before and 1 month after treatment. Cut-off positivity values for AFP, AFP-L3, and DCP were set at 20 ng/ml, 10 %, and 40 mAU/ml, respectively. Hepatic functional reserve was ranked using the criteria of the Child-Pugh scoring system.

Treatment of HCC

Therapeutic modalities for HCC patients were chosen on the basis of hepatic functional reserve, tumor multiplicity, and tumor size. Among the 414 HCC patients, 228 underwent hepatic resection and 186 underwent

radiofrequency ablation (RFA). The efficacy of RFA treatment was evaluated using dynamic CT or dynamic MRI within a few days after treatment. Complete ablation of HCC was defined as non-enhancement of the lesion relative to the surrounding liver parenchyma. Patients received additional sessions of an ablative therapy until the treatment was judged as complete.

Statistical Analysis

Differences in the proportions of patients with elevated levels of AFP, AFP-L3, and DCP before and after treatment were determined by Fisher's exact test. Recurrence-free survival of patients with HCC was determined by the Kaplan–Meier method. Logrank test was used to test for equality of recurrence-free survival between the groups. Multivariate analysis of prognostic factors among the clinical features was performed using the Cox stepwise proportional hazards model. The factors included in multivariate analysis were patient age (years), gender (female/male), HBsAg (negative/positive), anti-HCV (negative/positive), alcohol abuse (negative/positive), Child-Pugh class (A/B,C), platelet count ($\times 10^4/\mu\text{l}$), tumor size (cm), number of tumors (single/multiple), treatment method (hepatic resection/RFA), pretreatment AFP (ng/ml) ($<20/\geq 20$), pretreatment AFP-L3 (%) ($<10/\geq 10$), pretreatment DCP (mAU/ml) ($<40/\geq 40$), post-treatment AFP (ng/ml) ($<20/\geq 20$), post-treatment AFP-L3 (%) ($<10/\geq 10$), and post-treatment DCP (mAU/ml) ($<40/\geq 40$). Statistical analyses were performed with the SPSS version 17.0 software package (SPSS Japan Inc., Tokyo, Japan). Differences at $p < 0.05$ were considered to be statistically significant.

Results

Clinical Features of Patients

Table 1 summarizes the demographics, etiology of liver disease, hepatic functional reserve ranked by the Child-Pugh classification, platelet count, serum concentration of AFP (ng/ml), AFP-L3 (%), DCP (mAU/ml) tumor size, number of tumors, and treatment modality for the study patients.

This population comprised 136 females and 278 males with a median age of 71 (range 33–89) years. The majority of patients had a viral etiology for their liver disease, anti-HCV being positive in 287 (69.3 %) and HBsAg being positive in 53 (12.8 %). Most patients (86.2 %) were Child-Pugh class A. The median tumor size was 1.9 (range, 0.6–5.0) cm and multiple tumors were present in 17.4 % of the patients. When the cut-off values were set at 20 ng/ml

Table 1 Clinical features of patients ($n = 414$)

Age (years) [median (range)]	71 (33–89)
Gender (female/male)	136/278
Etiology	
HBsAg negative/positive	361/53
Anti-HCV negative/positive	127/287
Alcohol abuse negative/positive	396/18
Child-Pugh classification	
A/B/C	357/37/4
Platelet ($10^4/\mu\text{l}$) [median (range)]	12.2 (3.6–59.2)
AFP (ng/ml) [median (range)]	11.1 (0.3–14,597)
<20 ng/ml/ ≥ 20 ng/ml	248/166
AFP-L3 (%) <10 %/ ≥ 10 %	309/105
DCP (mAU/ml) [median (range)]	23 (5–33,283)
<40 mAU/ml/ ≥ 40 mAU/ml	273/136
Tumor size (cm) [median (range)]	1.9 (0.6–5.0)
Tumor number (single/multiple)	342/72
Therapeutic modalities	
Hepatic resection/RFA	228/186

Range or percent are shown in *parentheses*

HCC hepatocellular carcinoma, HBsAg hepatitis B surface antigen, HCV hepatitis C virus, AFP alpha-fetoprotein, DCP des-gamma-carboxy prothrombin, RFA Radiofrequency ablation

for AFP, 10 % for AFP-L3 and 40 mAU/ml for DCP, the positivity rates for AFP, AFP-L3 and DCP were 40.1, 25.4, and 33.3 %, respectively.

Impact of AFP, AFP-L3, and DCP on Recurrence-Free Survival of Patients with HCC

Among the 414 patients enrolled in this study, HCC recurrence was observed in 236, and the remaining 178 showed no HCC recurrence within the study period. Recurrence-free survival rates of patients with HCC were evaluated in terms of serum AFP, AFP-L3, and DCP levels before and after treatment. Comparison of the three tumor markers measured before treatment showed that patients with an elevated AFP-L3 level had a significantly lower recurrence-free survival rate than patients without such an elevation of the AFP-L3 level ($p = 0.024$) (Fig. 1). In contrast, there was no significant difference in recurrence-free survival between the groups with and without elevation of the AFP and DCP levels before treatment (AFP, $p = 0.171$; DCP, $p = 0.208$) (Fig. 1). The recurrence-free survival rate of patients with an elevation of AFP, AFP-L3, and DCP measured after treatment were significantly lower than that of patients without such elevation (AFP, $p = 0.009$; AFP-L3 and DCP, $p = 0.001$) (Fig. 2).

In addition, we evaluated the recurrence-free survival rates of patients according to AFP-L3 status measured 9 months after treatment. In patients who had no




Fig. 1 Comparison of recurrence-free survival rates between patients with and without elevation of the AFP, AFP-L3, and DCP levels before treatment. Recurrence-free survival rates according to AFP (a), AFP-L3 (b), and DCP (c)




Fig. 2 Comparison of recurrence-free survival rates between patients with and without elevation of the AFP, AFP-L3, and DCP levels at one month after treatment. Recurrence-free survival rates according to AFP (a), AFP-L3 (b), and DCP (c)

recurrence for at least 9 months, a significant difference was also observed between those with and without elevation of the AFP-L3 level ($p = 0.003$) (Fig. 3).

Multivariate analysis using the Cox stepwise proportional hazard model revealed that post-treatment AFP-L3 status ($p = 0.002$) and post-treatment DCP status ($p = 0.004$) were significant independent factors predictive of recurrence-free survival in patients with HCC (Table 2).

Relationship Between HCC Recurrence and Change in Positivity Rates for Serum AFP, AFP-L3, and DCP Before and After Treatment

We evaluated the changes in positivity rates for serum tumor markers before and after treatment in 308 of the 414 patients who were enrolled in this study. These 308 patients comprised 193 who suffered recurrence within 2 years and 115 in whom no recurrence was observed for more than 2 years. Positivity rates for serum AFP, AFP-L3, and DCP were investigated in

relation to HCC recurrence, and the results are shown in Table 3. Regardless of HCC recurrence, the proportions of patients showing elevation of the AFP and DCP levels (AFP, ≥ 20 ng/ml; DCP, ≥ 40 mAU/ml) for 1 month after curative treatment were significantly lower in comparison to the situation before treatment (AFP, $p = 0.004$ for patients with HCC recurrence and $p = 0.001$ for patients without HCC recurrence; DCP, $p < 0.001$ for both patients with and without HCC recurrence). On the other hand, there was no significant decrease in the proportion of patients showing an elevated AFP-L3 level 1 month after treatment in comparison to the situation before treatment in both of the groups with and without HCC recurrence. Among the 193 patients in the HCC recurrence group, 15 (26.8 %) of 56 patients with an elevated AFP-L3 level (≥ 10 %) before treatment became negative for AFP-L3 (< 10 %) after treatment. In the recurrence-free group, eight (38.1 %) of 21 patients with an elevated AFP-L3 level before treatment became negative for AFP-L3 after treatment. In addition, we investigated the long-term changes




Fig. 3 Comparison of recurrence-free survival rates between patients with and without elevation of the AFP-L3 level at 9 months after treatment

Table 2 Multivariate analysis of factors associated with HCC recurrence after curative treatment

Variables	Hazard ratio (95 % CI)	<i>p</i> value
Post-treatment AFP-L3		
<10 %	1	0.002
≥10 %	1.592 (1.183–2.144)	
Post-treatment DCP		
<40 mAU/ml	1	0.004
≥40 mAU/ml	1.687 (1.187–2.398)	

Hazard ratio and *p* value were calculated using Cox's stepwise proportional hazard model

HCC hepatocellular carcinoma, AFP alpha-fetoprotein, DCP des-gamma-carboxy prothrombin, CI confidence interval

in the level of AFP-L3 after treatment. Among patients who suffered recurrence of HCC after 9 months or later of treatments, the proportion of those who were still AFP-L3-positive 6 and 9 months after treatment was 28.6 % (4/14) and 24.1 % (7/29), respectively, being similar to the proportion of such patients before treatment (29.2 %; 69/236). On the other hand, in the group who remained recurrence-free for more than 2 years, no patient was AFP-L3-positive at 6 and 9 months after treatment (6 months; 0 %, 0/5; 9 months; 0 %, 0/18).

Discussion

Many studies have shown that AFP-L3 status is a specific marker for HCC, and that the level of AFP-L3 predicts the

malignant potential of HCC, and thus the expected outcome after treatment [10–15]. Recently, μ TAS AFP-L3 has been newly developed as an automated immunoassay for AFP-L3. A number of investigators, including our group, have shown that the μ TAS AFP-L3 is more sensitive for discriminating HCC from benign liver diseases than the conventional LiBASys AFP-L3, particularly in subgroups with lower AFP concentrations and early stage HCC [19–21]. In addition, we have previously reported that μ TAS AFP-L3 status is a statistically significant independent prognostic indicator of long-term survival in patients with HCC [19]. On the other hand, there has been little information about the levels of μ TAS AFP, AFP-L3, and DCP measured after curative treatment of HCC.

In the present study, we demonstrated that there was a significant difference in recurrence-free survival between groups with and without elevation of the AFP-L3 level before and after treatment. Multivariate analysis revealed that AFP-L3 status measured 1 month after treatment was a significant independent predictor of HCC recurrence after curative treatment. On the other hand, we also found that, irrespective of HCC recurrence, there was no significant decrease in the proportion of patients showing an elevated AFP-L3 level 1 month after treatment in comparison to the situation before treatment. In our present examination of long-term changes in the level of AFP-L3 after treatment, the proportion of patients with an elevated level of AFP-L3 at 6 and 9 months after treatment was decreased relative to that before treatment in the recurrence-free group. By contrast, the proportion of patients who were still AFP-L3-positive after treatment was similar to that before treatment in the HCC recurrence group. Taken together, these results suggest that an elevated AFP-L3 level before treatment is a predictor of HCC recurrence, and that sustained elevation of AFP-L3 after treatment should be considered to indicate HCC recurrence.

The decrease in the total concentration of AFP within 1 month after treatment, in terms of half-life, is reported to be 4–6 days [22, 23]. However, the percentage decrease in the level of AFP-L3 basically does not change, and the lower limit for quantitation of μ TAS AFP-L3 is quite low (0.3 ng/ml). We suggest that the minimal change in AFP-L3 positivity in some patients without HCC recurrences within the first month after treatment was attributable to serum AFP that had been produced by HCC just before treatment. Therefore, repeated combination measurement of μ TAS AFP, AFP-L3, and DCP should be performed for surveillance of HCC recurrence after curative treatment.

Recently, Yamamoto et al. [24] have investigated the relationship between HCC recurrence and changes in the levels of three tumor markers—AFP, AFP-L3, and DCP—measured using the LiBASys assay. They reported that, among these markers, AFP-L3 positivity after treatment

Table 3 Serum AFP, AFP-L3, and DCP before and 1 month after treatment

Patient group and tumor markers	Before treatment	1 month after treatment	<i>p</i> value
Patients with HCC recurrence within 2 years (<i>n</i> = 193)			
AFP			
Patient number (≥ 20 ng/ml/ < 20 ng/ml)	83/110	55/138	0.004
Percent of patients with AFP ≥ 20 ng/ml	43.0 %	28.5 %	
AFP-L3			
Patient number (≥ 10 %/ < 10 %)	56/137	52/141	0.734
Percent of patients with AFP-L3 ≥ 10 %	29.0 %	26.9 %	
DCP			
Patient number (≥ 40 mAU/ml/ < 40 mAU/ml)	69/123	35/155	< 0.001
Percent of patients with DCP ≥ 40 mAU/ml	34.3 %	18.4 %	
Patients without HCC recurrence for more than 2 years (<i>n</i> = 115)			
AFP			
Patient number (≥ 20 ng/ml/ < 20 ng/ml)	43/72	20/95	0.001
Percent of patients with AFP ≥ 20 ng/ml	37.4 %	17.4 %	
AFP-L3			
Patient number (≥ 10 %/ < 10 %)	21/94	17/98	0.595
Percent of patients with AFP-L3 ≥ 10 %	18.3 %	14.8 %	
DCP			
Patient number (≥ 40 mAU/ml/ < 40 mAU/ml)	31/83	6/109	< 0.001
Percent of patients with DCP ≥ 40 mAU/ml	27.0 %	5.2 %	

p values were calculated by Fisher's exact test

HCC hepatocellular carcinoma, AFP alpha-fetoprotein, DCP des-gamma-carboxy prothrombin

had the highest risk ratio of 5.0 for HCC recurrence after curative treatment. Kobayashi et al. [25] have investigated the relationship between changes in the serum AFP-L3 level measured by μ TAS assay 30–120 days after curative treatment and HCC recurrence. They reported that 29 of 37 patients (78.4 %) with preoperative AFP elevation (> 20 ng/ml) showed a decrease in the AFP level to < 20 ng/ml, although 16 of 42 patients (38.1 %) with preoperative AFP-L3 elevation (> 5 %) showed a decrease in the level to < 5 %. On this basis, they concluded that it was rare for AFP-L3 to become negative after treatment. Toyoda et al. [26] also investigated the value of AFP, AFP-L3, and DCP, measured before and 1–2 months after treatment, for prediction of survival and recurrence in patients who had undergone hepatectomy for HCC. They concluded that the combination of tumor markers measured by μ TAS assay after hepatectomy had excellent ability to predict postoperative survival and recurrence. Our present results support their findings, although the AFP-L3 cut-off value used in our study was 10 %, unlike their value of 5 %. In our previous study, we suggested that a cut-off value of 7 % was most appropriate for discriminating HCC from benign liver disease using μ TAS AFP-L3 [19]. In our present study, however, we chose a cut-off value of 10 % for μ TAS AFP-L3 in view of a recently published report by Kanke et al. [27] in which intra-individual biological variation of the μ TAS AFP-L3 level was relatively high, at 29.0 %.

Some factors other than AFP-L3 and DCP could potentially have been associated with HCC recurrence. Several studies have implicated obesity and diabetes as risk factors for HCC [28–30]. However, we did not investigate these factors in the present study.

The difficulty in the treatment of HCC is related to the underlying impairment of hepatic functional reserve and high rate of recurrence, even after curative treatment. Therefore, early detection of HCC recurrence after treatment is an important issue for improving the survival of patients with HCC. From this viewpoint, the μ TAS assay is an extremely powerful tool for detection of HCC at an early stage. Additionally, clinicians should interpret the values of tumor markers measured by the μ TAS assay giving due consideration to their property. In conclusion, the present study has demonstrated that patients with an elevated AFP-L3 level determined before treatment have a high risk of HCC recurrence, and that sustained elevation of AFP-L3 after treatment is a strong predictor of HCC recurrence. In addition, short-term AFP-L3 status is not a reliable indicator of incomplete treatment for HCC. Accordingly, repeated and combined measurement of μ TAS AFP, AFP-L3, and DCP should be performed for surveillance of HCC recurrence after curative treatment.

Acknowledgments This study was supported in part by a Grant-in-Aid (20390205), Japan Society for the Promotion Science and a Grant-in-Aid from Health, Labor and Welfare, Japan.

Conflict of interest None.

References

1. El-Serag HB, Rudolph KL. Hepatocellular carcinoma: epidemiology and molecular carcinogenesis. *Gastroenterology*. 2007;132:2557–2576.
2. Abelev GI. Production of embryonal serum alpha-globulin by hepatoma: review of experimental and clinical data. *Cancer Res*. 1968;28:1344–1350.
3. Conor GI, Tatarinov YS, Abelev GI, Uriel J. A collaborative study for the evaluation of a serologic test for primary liver cancer. *Cancer*. 1970;25:1091–1098.
4. Bayati N, Silverman AL, Gordon SC. Serum alpha-fetoprotein levels and liver histology in patients with chronic hepatitis C. *Am J Gastroenterol*. 1998;93:2452–2456.
5. Goldstein NS, Blue DE, Hankin R, et al. Serum alpha-fetoprotein levels in patients with chronic hepatitis C. Relationships with serum alanine aminotransferase values, histologic activity index, and hepatocyte MIB-1 scores. *Am J Clin Pathol*. 1999;111:811–816.
6. Chu CW, Hwang SJ, Luo JC, et al. Clinical, virologic, and pathologic significance of elevated serum alpha-fetoprotein levels in patients with chronic hepatitis C. *J Clin Gastroenterol*. 2001;32:240–244.
7. Aoyagi Y, Suzuki Y, Isemura M, et al. The fucosylation index of alpha-fetoprotein and its usefulness in the early diagnosis of hepatocellular carcinoma. *Cancer*. 1988;61:769–774.
8. Aoyagi Y, Saitoh A, Suzuki Y, et al. Fucosylation index of alpha-fetoprotein, a possible aid in the early recognition of hepatocellular carcinoma in patients with cirrhosis. *Hepatology*. 1993;17:50–52.
9. Taketa K, Endo Y, Sekiya C, et al. A collaborative study for the evaluation of lectin-reactive alpha-fetoproteins in early detection of hepatocellular carcinoma. *Cancer Res*. 1993;53:5419–5423.
10. Aoyagi Y, Isokawa O, Suda T, et al. The fucosylation index of alpha-fetoprotein as a possible prognostic indicator for patients with hepatocellular carcinoma. *Cancer*. 1998;83:2076–2082.
11. Hayashi K, Kumada T, Nakano S, et al. Usefulness of measurement of *Lens culinaris* agglutinin-reactive fraction of alpha-fetoprotein as a marker of prognosis and recurrence of small hepatocellular carcinoma. *Am J Gastroenterol*. 1999;94:3028–3033.
12. Oka H, Saito A, Ito K, et al. Multicenter prospective analysis of newly diagnosed hepatocellular carcinoma with respect to the percentage of *Lens culinaris* agglutinin-reactive alpha-fetoprotein. *J Gastroenterol Hepatol*. 2001;16:1378–1383.
13. Tateishi R, Shiina S, Yoshida H, et al. Prediction of recurrence of hepatocellular carcinoma after curative ablation using three tumor markers. *Hepatology*. 2006;44:1518–1527.
14. Miyaaki H, Nakashima O, Kurogi M, Eguchi K, Kojiro M. *Lens culinaris* agglutinin-reactive alpha-fetoprotein and protein induced by vitamin K absence II are potential indicators of a poor prognosis: a histopathological study of surgically resected hepatocellular carcinoma. *J Gastroenterol*. 2007;42:962–968.
15. Tamura Y, Igarashi M, Suda T, et al. Fucosylated fraction of alpha-fetoprotein as a predictor of prognosis in patients with hepatocellular carcinoma after curative treatment. *Dig Dis Sci*. 2010;55:2095–2101.
16. Nakamura K, Imajo N, Yamagata Y, et al. Liquid-phase binding assay of alpha-fetoprotein using a sulfated antibody for bound/free separation. *Anal Chem*. 1998;70:954–957.
17. Kawabata T, Wada HG, Watanabe M, Satomura S. Electrokinetic analyte transport assay for alpha-fetoprotein immunoassay integrates mixing, reaction and separation on-chip. *Electrophoresis*. 2008;29:1399–1406.
18. Kagebayashi C, Yamaguchi I, Akinaga A, et al. Automated immunoassay system for AFP-L3% using on-chip electrokinetic reaction and separation by affinity electrophoresis. *Anal Biochem*. 2009;388:306–311.
19. Tamura Y, Igarashi M, Kawai H, Suda T, Satomura S, Aoyagi Y. Clinical advantage of highly sensitive on-chip immunoassay for fucosylated fraction of alpha-fetoprotein in patients with hepatocellular carcinoma. *Dig Dis Sci*. 2010;55:3576–3583.
20. Toyoda H, Kumada T, Tada T, et al. Clinical utility of highly sensitive *Lens culinaris* agglutinin-reactive alpha-fetoprotein in hepatocellular carcinoma patients with alpha-fetoprotein <20 ng/ml. *Cancer Sci*. 2011;102:1025–1031.
21. Morimoto M, Numata K, Nozaki A, et al. Novel *Lens culinaris* agglutinin-reactive fraction of α -fetoprotein: a biomarker of hepatocellular carcinoma recurrence in patients with low α -fetoprotein concentrations. *Int J Clin Oncol*. 2012;17:373–379.
22. Han SJ, Yoo S, Choi SH, Hwang EH. Actual half-life of alpha-fetoprotein as a prognostic tool in pediatric malignant tumors. *Pediatr Surg Int*. 1997;12:599–602.
23. Murray MJ, Nicholson JC. α -fetoprotein. *Arch Dis Child Educ Pract Ed*. 2011;96:141–147.
24. Yamamoto K, Imamura H, Matsuyama Y, et al. AFP, AFP-L3, DCP, and GP73 as markers for monitoring treatment response and recurrence and as surrogate markers of clinicopathological variables of HCC. *J Gastroenterol*. 2010;45:1272–1282.
25. Kobayashi M, Hosaka T, Ikeda K, et al. Highly sensitive AFP-L3% assay is useful for predicting recurrence of hepatocellular carcinoma after curative treatment pre- and postoperatively. *Hepatol Res*. 2011;41:1036–1045.
26. Toyoda H, Kumada T, Tada T, et al. Prognostic significance of a combination of pre- and post-treatment tumor markers for hepatocellular carcinoma curatively treated with hepatectomy. *J Hepatol*. 2012;57:1251–1257.
27. Kanke F, Kumada T, Toyoda H, Satomura S. Reference change values for *Lens culinaris* agglutinin-reactive α -fetoprotein and des- γ -carboxy prothrombin in patients with chronic hepatitis C. *Clin Chem Lab Med*. 2012;50:957–960.
28. Chen CL, Yang HI, Yang WS, et al. Metabolic factors and risk of hepatocellular carcinoma by chronic hepatitis B/C infection: a follow-up study in Taiwan. *Gastroenterology*. 2008;135:111–121.
29. Hassan MM, Hwang LY, Hatten CJ, et al. Risk factors for hepatocellular carcinoma: synergism of alcohol with viral hepatitis and diabetes mellitus. *Hepatology*. 2002;36:1206–1213.
30. Yuan JM, Govindarajan S, Arakawa K, Yu MC. Synergism of alcohol, diabetes, and viral hepatitis on the risk of hepatocellular carcinoma in blacks and whites in the U.S. *Cancer*. 2004;101:1009–1017.

VIRAL HEPATITIS

Hepatitis B virus X induces cell proliferation in the hepatocarcinogenesis via up-regulation of cytoplasmic p21 expression

Masahiko Yano^{1,4,*}, Shogo Ohkoshi^{1,4,*}, Yo-hei Aoki¹, Hiromichi Takahashi¹, Sou Kurita¹, Kazuhide Yamazaki¹, Kenta Suzuki¹, Satoshi Yamagiwa¹, Ayumi Sanpei², Shun Fujimaki³, Toshifumi Wakai², Shin-ei Kudo⁴, Yasunobu Matsuda^{1,3} and Yutaka Aoyagi¹

1 Division of Gastroenterology and Hepatology, Graduate School of Medical and Dental Sciences Niigata University, Niigata, Japan

2 Division of Digestive and General Surgery, Graduate School of Medical and Dental Sciences of Niigata University, Niigata, Japan

3 Department of Medical Technology, Graduate School of Health Science of Niigata University, Niigata, Japan

4 Digestive Disease Center, Showa University, Northern Yokohama Hospital, Kanagawa, Japan

Keywords

HBV X protein – hepatocarcinogenesis – IFN- β – p21 (Cip1/WAF1) – PKC α

Abbreviations

CDK, cyclin-dependent kinase; DMSO, dimethyl sulfoxide; ERK, extracellular signal-regulated kinase; FBS, foetal bovine serum; HBV, hepatitis B virus; HBx, HBV X protein; HCC, hepatocellular carcinoma; IFN, interferon; MEK, mitogen-activated protein kinase/ERK kinase; PI-3K/Akt, phosphatidylinositol 3-kinase/protein kinase B; PKC, protein kinase C; pRb, retinoblastoma protein; qRT-PCR, quantified real-time PCR; WM, wild mice; Xg, HBx transgenic mice.

Correspondence

Shogo Ohkoshi, Division of Gastroenterology and Hepatology, Graduate School of Medical and Dental Sciences of Niigata University, 1-754 Asahimachi – Dori, Chuo-ku, Niigata 951-8122, Japan
Tel: +81 25 227 2204
Fax: +81 25 227 0776
e-mail: okoshi@med.showa-u.ac.jp

Received 10 July 2012

Accepted 10 March 2013

DOI:10.1111/liv.12176

Liver Int. 2013; 33: 1218–1229

Persistent infection with hepatitis B virus (HBV) is a prevalent aetiology for chronic liver disease, and may lead to a premalignant condition preceding the development of hepatocellular carcinoma (HCC) (1, 2). Elimination of this virulent pathogen is a solid clinical strategy for prevention of HCC formation. Despite

Abstract

Background: Hepatitis B virus X protein (HBx) has been shown to induce hepatocarcinogenesis by disrupting the functions of intracellular molecules. Cyclin-dependent kinase inhibitor p21 (Cip1/WAF1), known as a tumour-suppressor gene, has been reported to have paradoxical function, that is, acting as an oncogene, particularly when expressed in the cytoplasm. The effects of HBx on the expression and function of p21 also remain controversial. **Aims:** We attempted to investigate the role of HBx in the hepatocarcinogenic process, focusing on the association with this paradoxical function of p21. The results obtained were further verified with experiments using the antihepatocarcinogenic action of interferon (IFN)- β . **Methods:** HBx transgenic mice (Xg) and HBx-transfected hepatoma cell lines were used. Intracellular localization of p21 was determined by Western blot analysis and immunofluorescence. **Results:** Xg and HBx-transfected cells exhibited increased expression of p21. Up-regulation of p21 was positively correlated with the expression of cyclin D1 and inactive phosphorylation of retinoblastoma protein (pRb). These HBx-induced cell proliferative responses were cancelled by knockdown of p21, which resulted in growth reduction in HBx-expressing cells, suggesting the oncogenic properties of HBx-induced p21. HBx induced accumulation of p21 in the cytoplasm, and activation of PKC α was involved. Finally, IFN- β -treated Xg liver, as well as hepatoma cells, showed a shift of cytoplasmic p21 to the nucleus, accompanied by the abrogation of HBx-induced oncogenic modulation. **Conclusions:** Our results suggest that HBx induces hepatocarcinogenesis via PKC α -mediated overexpression of cytoplasmic p21 and IFN- β suppressed these molecular events by shifting p21 to the nucleus.

recent advances in anti-HBV therapy, such as interferon (IFN)-based therapy and use of nucleic acid analogue drugs, there remain a number of problems in the establishment of preventive strategies for HBV-related HCC (3). These problems result from unresolved questions regarding the molecular mechanisms of HBV-induced hepatocarcinogenesis (4, 5). Among the four proteins (S, C, P and X) encoded by the four open reading frames in HBV genome, HBV X protein (HBx), consisting of 154 amino acids, has been found to disrupt cell proliferative and apoptotic processes through the modulation

*Present address: Digestive Disease Center, Showa University, Northern Yokohama Hospital, 35-1 Chigasaki-chuo, Tsuzuki-ku, Kanagawa 224-8503, Japan.

of intracellular signal transduction pathways, including the protein kinase C (PKC), phosphatidylinositol 3-kinase/protein kinase B (PI-3K/Akt) and extracellular signal-regulated kinase (ERK)1/2 pathways (6–8). These aberrant molecular effects of HBx during HBV infection are likely to render normal hepatocytes susceptible to neoplastic and malignant transformation.

Mammalian cell proliferation is controlled by various protein kinase complexes, including cyclin/cyclin-dependent kinase (CDK) complexes that regulate distinct phases of the cell cycle by repressing anticell proliferative functions of retinoblastoma protein (pRb) (9). Although induction of p21 (also known as Cip1/WAF1) as a universal inhibitor of cyclin/CDK activity downstream of active p53 generally results in inhibition of cell proliferation (10), more recent studies have revealed the opposite effect. Such studies have suggested that p21 renders normal cells susceptible to transformation by leading to an anti-apoptotic or a cell proliferative status, which is unrelated to p53 (11–13) and controlled by protein kinase activity (14–16). This oncogenic potential of p21 has also been confirmed *in vivo* and in clinical studies (17, 18). It has been suggested that the intracellular distribution pattern of p21 in particular may determine whether it has anti-oncogenic or oncogenic property. Zhou *et al.* reported that cytoplasmic accumulation of p21 resulted in a loss of the cell cycle arrest induced by p21 located in nuclei (19). Furthermore, forced cytoplasmic localization of p21, using a nuclear localization signal-deficient p21 construct, triggered cell cycle progression as well as protection from apoptosis (20).

Previous reports have revealed that HBx triggers hepatocarcinogenesis by modulation of p21 expression and function (21–24). However, they showed different results and interpretations. Kwun and Jang demonstrated that, using several HBx clones from the sera of patients with chronic HBV infection, natural variants of HBx determined the expression level of p21 in a p53-dependent or -independent pathway, suggesting that these variants show different effects on the cell cycle progression (25). However, these studies were not confirmed by comparative assays using experimental models, which showed the suppressive state against HBx-induced hepatocarcinogenesis after anticancerous treatment. Thus, the role of p21 in HBx-triggered HCC development remains uncertain.

PKC plays a central role in signal transduction and has been implicated in a wide range of physiological functions, including cell growth, transformation and differentiation. HBx protein was found to increase the levels of endogenous PKCs and subsequently activate AP-1 and NF- κ B transcription factors (6, 7). As oncogenic p21 was reported to be regulated by PKC α activation, resulting in cell cycle progression by incorporation of p21 into cyclin/CDK complexes (14), we hypothesize that HBx-induced p21 up-regulation may be mediated through active PKC α . We here report that HBx

promotes cell proliferation via up-regulation of PKC α -mediated overexpression of cytoplasmic p21 using both HBx transgenic mice (Xg) (26, 27) and HBx-transfected cells. Because our previous study found that treatment with IFN- β prevented HBx-induced HCC development in Xg (28), we verified our results using the anticell proliferative action of IFN- β .

Materials and methods

Mouse liver samples

Frozen mouse livers for quantified real-time PCR (qRT-PCR) and Western blot analyses were obtained as described previously (28). mRNA and proteins were extracted from the livers of 6-month-old wild mice (WM) and Xg treated with either saline (control) or IFN- β . RNA (1 μ g) purified with RNeasy Mini Kit (Qiagen, Valencia, CA, USA) was subjected to reverse transcription and qRT-PCR using a Light Cycler System (Roche, Mannheim, Germany). Specific primers and probes for the genes were designed based on the Universal Probe Library (Roche). All the experimental protocols were approved by the Institutional Animal Care and Use Committee of Niigata University and performed in accordance with the National Institute of Health Guidelines.

Cell culture and reagents

HepG2 and Hep3B were maintained in Dulbecco's modified Eagle's medium supplemented with 10% foetal bovine serum (FBS) (Invitrogen, Carlsbad, CA, USA) and antibiotic-antimycotic solution (Sigma Aldrich, St. Louis, MO, USA). IFN- β was kindly provided by Toray Medical Co., Ltd. (Chiba, Japan). Specific kinase inhibitors, GF109203X, U0126 and LY294002, were purchased from Calbiochem (San Diego, CA, USA), and dimethyl sulfoxide (DMSO) was from Wako Pure Chemical Ind., Ltd. (Osaka, Japan). Rabbit-polyclonal antibodies specific for p21, used for mouse liver samples, and p53 and anti-GAPDH mouse-monoclonal antibody were purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Mouse-monoclonal antibodies specific for FLAG M2 and β -actin were purchased from Sigma, and other specific antibodies were from Cell Signaling Technology (Beverly, MA, USA).

Plasmid construction and transfection

For HBx overexpression in cell lines, we generated pCAG-FLAG-HBx expressing FLAG-tagged HBx under the CAG promoter. The blunted *NotI*-*BglII* fragment containing the FLAG-HBx coding sequence from pNK-FLAG-HBx (kind gift from Dr Murakami, Kanazawa University, Japan) was inserted into the blunted *EcoRI* site downstream of the CAG promoter in pCAG-IRES-Puro. After sequence verification, transfection with

pCAG-IRES-Puro, used for mock transfection, or pCAG-FLAG-HBx was performed using FuGENE6 (Roche) and MA lipofection enhancer (IBA, St. Louis, MO, USA) according to the manufacturer's protocols. To establish stable expressing cell lines, pCAG-IRES-Puro and pCAG-FLAG-HBx were linearized with *PvuII* and transfected into HepG2, followed by selection with 2 µg/ml puromycin (Cayla, Toulouse, France).

Western blot analysis

Frozen mouse livers or culture cells were lysed using Tissue Ruptor or Cell Scraper with RIPA buffer (Thermo Scientific, Rockford, IL, USA) containing protease inhibitor cocktail and phospho-stop tablets (Roche). Lysates were sonicated and collected by centrifugation to remove debris. Nuclear and cytoplasmic fractions of liver tissues and culture cells were prepared by NE-PER Nuclear and Cytoplasmic Extraction Reagents (Thermo Scientific) according to the manufacturer's protocols. Next, 20 µg of mouse samples and 10 µg of cellular lysates, respectively, were subjected to Western blotting.

Knockdown of p21 with siRNA

For knockdown assay of p21, the FlexiTube GeneSolution for CDKN1A (Entrez gene ID: 1026) containing four different siRNA duplexes of p21 or negative control siRNA was transfected into HepG2 cells using HiPerFect Transfection Reagent (Qiagen) according to the manufacturer's protocols. At 24 h of siRNA delivery, mock or HBx expression plasmids were transiently expressed. To examine the effects of p21 knockdown on cell growth or cell cycle, 5×10^5 HepG2 cells were plated onto six-well plates in triplicate or a 10 cm dish and transfected with siRNA. For cell growth analysis, at 24 or 48 h after plasmid expression, cells were harvested and then the number of viable cells was counted after trypan blue dye treatment as described previously (29). For cell cycle analysis, 48-h HBx-expressed cells were stained with BrdU using a FITC BrdU Flow Kit (BD Pharmingen, San Jose, CA, USA) according to the manufacturer's protocols. The FITC intensity and total DNA content were measured by FACScan (BD Pharmingen) and analysed using FlowJo software (Ver. 7; Tree Star Inc., Ashland, OR, USA).

Immunofluorescence

Cultured cells were fixed in 4% paraformaldehyde and permeabilized with 0.1% Triton X-100 in PBS. After washing, cells were blocked in Image-iT FX Signal Enhancer (Invitrogen). For double immunofluorescence staining, fixed cells were reacted with anti-FLAG and anti-p21 antibody at a dilution of 1/50, followed by incubation with fluorescence-conjugated AlexaFluor secondary antibody (Cell Signaling) at a dilution of 1/

1000. Nuclei were counterstained with DAPI (Invitrogen) and slides were examined by fluorescence microscopy.

Immunoprecipitation

Cell extracts were prepared by incubation in 0.5% modified NP40 Cell Lysis Buffer (Invitrogen) for 30 min on ice. After centrifugation, 500 µg of cell extracts was immunoprecipitated with anti-FLAG or negative control mouse IgG (Thermo Scientific) using Catch and Release v2.0 system (Millipore, Billerica, MA, USA) according to the manufacturer's protocols. Eluted samples were separated by SDS-PAGE and subjected to immunoblotting.

Statistical analysis

Statistical analysis was performed using commercial SPSS software (Version 15.0; SPSS, Chicago, IL, USA). One-way analysis of variance (ANOVA) was used to compare the differences among mRNA levels on qRT-PCR analysis and cell growth data. *P* values <0.05 were considered to be statistically significant.

Results

Effects of HBx on p21 *in vivo* and *in vitro*

Using qRT-PCR, we first compared the intrahepatic expression of p21 mRNA between 6-month-old WM and Xg. p21 mRNA was highly up-regulated in Xg liver, showing about 10-fold more expression than in WM (Fig. 1A). As shown in Fig. 1B, we also verified this difference by assessing p21 protein levels in 6- and 12-month-old mouse liver lysates. Notably, expression of p53, a general upstream regulator of p21, was not enhanced in Xg liver, suggesting that p21 was p53-independently overexpressed in Xg.

Next, using *in vitro* assays, we determined whether HBx-induced p21 overexpression could be demonstrated in HBx-expressing hepatoma-derived cells as shown in Xg liver. Transient transfection with HBx constructs, but not mock constructs, resulted in dose-dependent enhancement of p21 expression without changes in p53 expression, in HepG2 and in p53-null Hep3B cells (Fig. 1C). We further confirmed that HBx caused p21 up-regulation in stable cell lines, similar to that in transiently expressing cells (Fig. 1D). These results indicate that culture cells transfected with HBx constructs also showed enhancement of p21 expression in a p53-independent manner.

Effects of HBx on p21-associated molecules

p21 has been well known as a tumour suppressor directly inhibiting activation of cyclin D/CDK complexes, which promote the cell cycle following




Fig. 1. p53-independent p21 expression is up-regulated by HBx in Xg liver and HBx-expressing hepatoma cells. (A) Comparative analysis of p21 mRNA in livers from WM and Xg at age 6 months by qRT-PCR; GAPDH, internal control ($*P < 0.01$). Liver mRNAs from seven mice in each group were used. (B) Western blot analysis showing p53-independent up-regulation of p21 in Xg liver. Two liver samples from each mouse group at age 6 months (6 M; left panel) or 12 months (12 M; right) were immunoblotted with antibodies specific for p21 and p53. (C, D) Western Blot analysis showing p53-independent up-regulation of p21 in HBx-expressing hepatoma cells. HepG2 (left panel) or Hep3B (right) cells were transiently transfected with mock or increasing amounts of HBx plasmid (pHBx) for 40 h (C), and mock- or HBx-expressing stable HepG2 cells (sHBx) were established as described in Materials and methods (D). After incubation and harvest, levels of p21, p53 and FLAG (HBx) were analysed by Western blot. Membranes were stripped and reprobed with anti- β -actin antibody to control for protein-loading variations.

phosphorylation and inactivation of pRb (10). However, recent studies focusing on intracellular functions of p21 found that it could also facilitate the assembly of cyclin D/CDK complexes via a p53-independent pathway (12, 13). Hence, we examined the expression and phosphorylation of these proteins in Xg liver and HBx-transfected cells. As shown in Fig. 2A, Xg liver exhibited increased the expression of cyclin D1, while cyclin D3 and CDKs expression were not enhanced. Consistent with this, HBx-transfected hepatoma cells showed increases in cyclin D1 and phosphorylation, inactivated form, of pRb, which may be mediated by active cyclin D1, in both transient and stable cell lines (Fig. 2B,C). These observations implied a positive link among p21, cyclin D1 expression and pRb inactivation in HBx-induced hepatocarcinogenesis.

Knockdown and localization analysis of p21 in HBx-transfected cells

Both *in vivo* and *in vitro* experiments indicated the involvement of p53-independent up-regulation of p21 expression in the hepatocarcinogenic action of HBx, but the mechanisms of HBx-induced p21 remain uncertain. To evaluate whether HBx-induced p21 promotes cell growth and cell cycle, we investigated the effects of p21 knockdown with siRNA on cell proliferation and oncogenic events in this setting. Figure 3A shows attenuations of pRb phosphorylation and cyclin D1 expression by p21 knockdown in HBx-transfected cells, and reciprocally increased induction of those in mock-transfected cells. These results suggest that the opposite function of p21, acceleration of the cell cycle, was induced by HBx.




Fig. 2. Up-regulations of cyclin D1 expression and pRb phosphorylation are induced by HBx. (A) Western blot analysis showing cyclin D1 up-regulation in Xg liver. The same liver samples as shown in Fig. 1B were immunoblotted with antibodies specific for cyclin D/CDKs. (B, C) Western blot analysis showing increases in cyclin D1 and phosphorylated-pRb (P-pRb) in HBx-expressing cells. The same HBx-expressing transient (B) or stable (C) cell samples as shown in Fig. 1C or 1D were immunoblotted with antibodies specific for cyclin D/CDKs, pRb (phosphorylated form and total) and FLAG (HBx). Anti- β -actin antibody was used as protein-loading control.

Reflecting these observations, in cell growth and cell cycle assays, significant reductions in cell proliferation and S phase cell population were observed in HBx-transfected cells with the knockdown of p21 in contrast to control siRNA (Fig. 3B,C).

Previous reports indicated that the regulatory role of p21 in the cell cycle may be dependent upon its intracellular localization (19, 20). We therefore performed Western blot analysis using nuclear and cytoplasmic fractions to verify the intracellular localization of p21 in liver lysates from Xg and in HBx-transfected cells. We found that intracellular localization of p21 became predominantly cytoplasmic in both Xg liver and in HBx-expressing cells (Fig. 4A,B). These results suggest that the aberrant overexpression of cytoplasmic p21 by HBx results in pRb inactivation and cell cycle progression, which may lead to an oncogenic state.

Involvement of PKC α in HBx-induced cytoplasmic p21 overexpression

As there are previous reports that this oncogenic localization of p21 in cytoplasm requires the activation of PKC (14), mitogen-activated protein kinase/ERK kinase (MEK)-ERK1/2 (16) or Akt (15), which have been

known to be aberrantly stimulated by HBx and lead to hepatocarcinogenesis (6–8), we tested the effects of an inhibitor specific for each kinase on molecular results caused by HBx in HBx-expressing stable cells. Treatment with GF109203X (PKC inhibitor) or U0126 (MEK1/2 inhibitor), but not LY294002 (PI-3K inhibitor), attenuated HBx-enhanced p21 expression, and only GF109203X treatment resulted in the attenuations of both cyclin D1 and phosphorylated-pRb (Fig. 5A). As shown in Fig. 5B, GF109203X also inhibited HBx-induced overexpression of cytoplasmic p21, whereas U0126, which decreased total expression of p21 in HBx cells (Fig. 5A), did not alter the intracellular distribution of p21, indicating that HBx enhanced this oncogenic cytoplasmic p21 via PKC activation, not through MEK-ERK1/2 or PI-3K/Akt. We confirmed the up-regulation of PKC α phosphorylation in Xg liver as well as in HBx-expressing cells (Fig. 5C,D), but other PKC isoforms were not phosphorylated in any experimental models (data not shown).

HBx reportedly exerts hepatocarcinogenic functions via direct association with various molecules, including p53 (7). We hypothesized that up-regulation of cell proliferative p21 by HBx is mediated through their direct interaction in cytoplasm (see Fig. 4C). However, HBx




Fig. 3. HBx-induced p21 overexpression positively regulates cell proliferation and cell cycle. (A) Effects of p21 knockdown with siRNA on HBx-caused cell proliferative up-regulations of cyclin D1 and phosphorylated-pRb (P-pRb). HepG2 cells were transfected overnight with negative control (iNC) or p21 siRNA (ip21) and then mock or HBx plasmid (pHBx) was transiently expressed in iNC or ip21-transfected cells respectively. After incubation for 40 h, cells were analysed by Western blot for levels of p21, cyclin D1, pRb (phosphorylated form and total) and FLAG (HBx). Anti- β -actin antibody controlled for protein loading. (B, C) Effects of p21 knockdown on cell growth and cell cycle. Transfection with siRNA and expression plasmid into HepG2 cells was performed as described in (A); at 24 and 48 h post-transfection, cells were harvested and number of viable cells was determined. *P* values were compared for pHBx-transfected cells with iNC (black bars) and those with ip21 (white bars) (B). At 48 h post-pHBx transfection, cells with iNC (left) or ip21 (right) were labelled with 10 μ M BrdU for 30 min and harvested at 3 h after changing the medium to remove BrdU. The samples were subjected to incubation with a FITC-conjugated anti-BrdU antibody and DNA staining with 7-AAD. FACS analysis was performed as described in Materials and methods (C).




Fig. 4. HBx induces overexpression of cytoplasmic p21 in Xg liver and HBx-expressing cells. (A) Intracellular localization of p21 in livers from WM and Xg. Cytoplasmic (C) and nuclear (N) fractions of two mouse groups (WM and Xg at the age of 6 months) were immunoblotted with antibodies specific for p21, GAPDH (cytoplasmic protein marker) and p53 (nuclear marker). (B) Intracellular localization of p21 in transiently mock- or HBx plasmid (pHBx)-transfected HepG2 cells. After incubation for 40 h, C/N fractions were immunoblotted with antibodies specific for p21, Hsp90 (C), p53 (N) and FLAG (HBx). (C) Immunofluorescence of p21 localization in transiently pHBx-transfected HepG2 cells. Intracellular localization after 40 h of expression; HBx (FLAG; red) and p21 (green). Nuclei were stained with DAPI (blue).

did not associate with p21, whereas direct interaction between HBx and PKC α was found by immunoprecipitation assay (Fig. 5E). Immunofluorescence staining further revealed colocalization of HBx with PKC α in the cytoplasm of HBx-transfected cells (Fig. 5F). These results suggest that HBx indirectly promotes cell-proliferative accumulation of p21 in the cytoplasm via direct stimulation of PKC α .

Analysis using antihepatocarcinogenic models provided by IFN- β

We previously reported that treatment with IFN- β prevented the occurrence of HCC in Xg (28). To confirm the involvement of PKC α -mediated cytoplasmic p21 overexpression in hepatocarcinogenesis by HBx, we used this antihepatocarcinogenic state model induced by IFN- β . Xg, as well as HBx-expressing cells, treated with IFN- β exhibited suppressions of HBx-induced oncogenic modulations (i.e. reductions in p21, phosphorylated-PKC α , cyclin D1 and phosphorylated-pRb)

(Fig. S1A,B). Finally, we found that Xg liver treated with IFN- β showed higher levels of nuclear p21 than non-treated liver (Fig. 6A). Shifts of cytoplasmic p21 to the nucleus by IFN- β were also observed in HBx-expressing cells (Fig. 6B,C), a finding consistent with our *in vivo* experimental results. Interestingly, this regulation of intracellular localization of p21 by IFN- β was also observed in mock-transfected cells.

Discussion

This study indicates that p21 plays an important role in hepatocarcinogenicity by HBx. p21 mediates the anti-proliferative response through arrest of the G1/S transition in the cell cycle under the control of the p53 tumour suppressor (10). Although many previous studies have explored the relevance of HBx-modified p21 expression to HBV-triggered hepatocarcinogenesis, the results and interpretations are contradictory. Some studies revealed that HBx promotes the hepatocarcinogenic process by suppression of anti-oncogenic p21