

- validation of a histological scoring system for nonalcoholic fatty liver disease, *Hepatology* 41 (2005) 1313–1321.
- [28] N. Ando, M. Shimizu, M. Okuno, R. Matsushima-Nishiwaki, H. Tsurumi, T. Tanaka, H. Moriwaki, Expression of retinoid X receptor alpha is decreased in 3'-methyl-4-dimethylaminoazobenzene-induced hepatocellular carcinoma in rats, *Oncol. Rep.* 18 (2007) 879–884.
- [29] D. Terakura, M. Shimizu, J. Iwasa, A. Baba, T. Kochi, T. Ohno, M. Kubota, Y. Shirakami, M. Shiraki, K. Takai, H. Tsurumi, T. Tanaka, H. Moriwaki, Preventive effects of branched-chain amino acid supplementation on the spontaneous development of hepatic preneoplastic lesions in C57BL/KsJ-db/db obese mice, *Carcinogenesis* 33 (2012) 2499–2506.
- [30] F. Kassie, M. Uhl, S. Rabot, B. Grasl-Kraupp, R. Verkerk, M. Kundi, M. Chabicovsky, R. Schulte-Hermann, S. Knasmüller, Chemoprevention of 2-amino-3-methylimidazo[4,5-f]quinoline (IQ)-induced colonic and hepatic preneoplastic lesions in the F344 rat by cruciferous vegetables administered simultaneously with the carcinogen, *Carcinogenesis* 24 (2003) 255–261.
- [31] J. Xiao, Y.P. Ching, E.C. Liong, A.A. Nanji, M.L. Fung, G.L. Tipoe, Garlic-derived S-allylmercaptocysteine is a hepato-protective agent in non-alcoholic fatty liver disease in vivo animal model, *Eur. J. Nutr.* 52 (2013) (2012) 179–191.
- [32] J. Iwasa, M. Shimizu, M. Shiraki, Y. Shirakami, H. Sakai, Y. Terakura, K. Takai, H. Tsurumi, T. Tanaka, H. Moriwaki, Dietary supplementation with branched-chain amino acids suppresses diethylnitrosamine-induced liver tumorigenesis in obese and diabetic C57BL/KsJ-db/db mice, *Cancer Sci.* 101 (2010) 460–467.
- [33] H. Yoshiji, R. Noguchi, Y. Ikenaka, K. Kaji, Y. Aihara, H. Fukui, Impact of renin-angiotensin system in hepatocellular carcinoma, *Curr. Cancer Drug Targets* 11 (2011) 431–441.
- [34] M.A. Abdelmegeed, A. Banerjee, S.H. Yoo, S. Jang, F.J. Gonzalez, B.J. Song, Critical role of cytochrome P450 2E1 (CYP2E1) in the development of high fat-induced non-alcoholic steatohepatitis, *J. Hepatol.* 57 (2012) 860–866.
- [35] Y. Wang, L.M. Ausman, R.M. Russell, A.S. Greenberg, X.D. Wang, Increased apoptosis in high-fat diet-induced nonalcoholic steatohepatitis in rats is associated with c-Jun NH2-terminal kinase activation and elevated proapoptotic Bax, *J. Nutr.* 138 (2008) 1866–1871.
- [36] H. Chen, G. Sullivan, L.Q. Yue, A. Katz, M.J. Quon, QUICKI is a useful index of insulin sensitivity in subjects with hypertension, *Am. J. Physiol. Endocrinol. Metab.* 284 (2003) E804–812.
- [37] H. Tsuda, S. Fukushima, H. Wanibuchi, K. Morimura, D. Nakae, K. Imaida, M. Tatematsu, M. Hirose, K. Wakabayashi, M.A. Moore, Value of GST-P positive preneoplastic hepatic foci in dose-response studies of hepatocarcinogenesis: evidence for practical thresholds with both genotoxic and nongenotoxic carcinogens. A review of recent work, *Toxicol. Pathol.* 31 (2003) 80–86.
- [38] Y. Nabeshima, S. Tazuma, K. Kanno, H. Hyogo, M. Iwai, M. Horiuchi, K. Chayama, Anti-fibrogenic function of angiotensin II type 2 receptor in CCl4-induced liver fibrosis, *Biochem. Biophys. Res. Commun.* 346 (2006) 658–664.
- [39] A. Hirose, M. Ono, T. Saibara, Y. Nozaki, K. Masuda, A. Yoshioka, M. Takahashi, N. Akisawa, S. Iwasaki, J.A. Oben, S. Onishi, Angiotensin II type 1 receptor blocker inhibits fibrosis in rat nonalcoholic steatohepatitis, *Hepatology* 45 (2007) 1375–1381.
- [40] S. Yokohama, M. Yoneda, M. Haneda, S. Okamoto, M. Okada, K. Aso, T. Hasegawa, Y. Tokusashi, N. Miyokawa, K. Nakamura, Therapeutic efficacy of an angiotensin II receptor antagonist in patients with nonalcoholic steatohepatitis, *Hepatology* 40 (2004) 1222–1225.
- [41] I. Sakaida, K. Hironaka, K. Uchida, C. Suzuki, K. Kayano, K. Okita, Fibrosis accelerates the development of enzyme-altered lesions in the rat liver, *Hepatology* 28 (1998) 1247–1252.
- [42] R. Bataller, E. Gabele, R. Schoonhoven, T. Morris, M. Lehnert, L. Yang, D.A. Brenner, R.A. Rippe, Prolonged infusion of angiotensin II into normal rats induces stellate cell activation and proinflammatory events in liver, *Am. J. Physiol. Gastrointest. Liver Physiol.* 285 (2003) G642–651.
- [43] R. Bataller, E. Gabele, C.J. Parsons, T. Morris, L. Yang, R. Schoonhoven, D.A. Brenner, R.A. Rippe, Systemic infusion of angiotensin II exacerbates liver fibrosis in bile duct-ligated rats, *Hepatology* 41 (2005) 1046–1055.
- [44] R.M. Pereira, R.A. dos Santos, F.L. da Costa Dias, M.M. Teixeira, A.C. Simoes e Silva, Renin-angiotensin system in the pathogenesis of liver fibrosis, *World J. Gastroenterol.* 15 (2009) 2579–2586.
- [45] M. Moreno, T. Gonzalo, R.J. Kok, P. Sancho-Bru, M. van Beuge, J. Swart, J. Prakash, K. Temming, C. Fondevila, L. Beljaars, M. Lacombe, P. van der Hoeven, V. Arroyo, K. Poelstra, D.A. Brenner, P. Gines, R. Bataller, Reduction of advanced liver fibrosis by short-term targeted delivery of an angiotensin receptor blocker to hepatic stellate cells in rats, *Hepatology* 51 (2010) 942–952.
- [46] M.O. Rakowski, A.G. Singal, M.A. Rogers, H. Conjeevaram, Meta-analysis: insulin sensitizers for the treatment of non-alcoholic steatohepatitis, *Aliment. Pharmacol. Ther.* 32 (2010) 1211–1221.
- [47] A.J. Sanyal, N. Chalasani, K.V. Kowdley, A. McCullough, A.M. Diehl, N.M. Bass, B.A. Neuschwander-Tetri, J.E. Lavine, J. Tonascia, A. Unalp, M. Van Natta, J. Clark, E.M. Brunt, D.E. Kleiner, J.H. Hoofnagle, P.R. Robuck, NASH CRN, Pioglitazone, vitamin E, or placebo for nonalcoholic steatohepatitis, *N. Engl. J. Med.* 362 (2010) 1675–1685.
- [48] M. Shimizu, Y. Fukutomi, M. Ninomiya, K. Nagura, T. Kato, H. Araki, M. Suganuma, H. Fujiki, H. Moriwaki, Green tea extracts for the prevention of metachronous colorectal adenomas: a pilot study, *Cancer Epidemiol. Biomarkers Prev.* 17 (2008) 3020–3025.
- [49] S. Bettuzzi, M. Brausi, F. Rizzi, G. Castagnetti, G. Peracchia, A. Corti, Chemoprevention of human prostate cancer by oral administration of green tea catechins in volunteers with high-grade prostate intraepithelial neoplasia: a preliminary report from a one-year proof-of-principle study, *Cancer Res.* 66 (2006) 1234–1240.
- [50] M. Antonello, D. Montemurro, M. Bolognesi, M. Di Pascoli, A. Piva, F. Grego, D. Sticchi, L. Giuliani, S. Garbisa, G.P. Rossi, Prevention of hypertension, cardiovascular damage and endothelial dysfunction with green tea extracts, *Am. J. Hypertens.* 20 (2007) 1321–1328.
- [51] I. Papparella, G. Ceolotto, D. Montemurro, M. Antonello, S. Garbisa, G. Rossi, A. Semplicini, Green tea attenuates angiotensin II-induced cardiac hypertrophy in rats by modulating reactive oxygen species production and the Src/epidermal growth factor receptor/Akt signaling pathway, *J. Nutr.* 138 (2008) 1596–1601.
- [52] I. Kurita, M. Maeda-Yamamoto, H. Tachibana, M. Kamei, Antihypertensive effect of Benifuuki tea containing O-methylated EGCG, *J. Agric. Food Chem.* 58 (2010) 1903–1908.

Obesity and hepatocellular carcinoma: targeting obesity-related inflammation for chemoprevention of liver carcinogenesis

Masahito Shimizu · Takuji Tanaka · Hisataka Moriwaki

Received: 25 June 2012 / Accepted: 16 August 2012 / Published online: 4 September 2012
© Springer-Verlag 2012

Abstract Obesity and related metabolic abnormalities, including a state of chronic inflammation, increase the risk of hepatocellular carcinoma (HCC). Adipose tissue constitutively expresses the proinflammatory cytokine tumor necrosis factor- α (TNF- α) and interleukin-6 (IL-6), which are important tumor promoters in inflammation-related carcinogenesis. Dysregulation of TNF- α and IL-6 is associated with the development of steatosis and inflammation within the liver. These cytokines also lie at the core of the association between obesity and insulin resistance, which is a key factor in the development of obesity-related HCC. Here we present a detailed review of the relationship between metabolic abnormalities and the development of HCC, focusing on the role played by inflammation. Drawing from our basic and clinical research, the present report also reviews evidence that targeting metabolic abnormalities, such as attenuation of chronic inflammation and improvement of insulin resistance by either pharmaceutical or nutritional intervention, may be an effective strategy in preventing the development of HCC in obese individuals.

Keywords Obesity · Inflammation · Hepatocellular carcinoma · Chemoprevention

This article is a contribution to the special issue on Inflammation and Cancer - Guest Editor: Takuji Tanaka

This article is published as part of the Special Issue on *Inflammation and Cancer* [35:2].

M. Shimizu (✉) · H. Moriwaki
Department of Gastroenterology, Gifu University Graduate School of Medicine,
1-1 Yanagido,
Gifu 501-1194, Japan
e-mail: shimim-gif@umin.ac.jp

T. Tanaka
The Tohukai Cytopathology Institute: Cancer Research and Prevention (TCI-CaRP),
Gifu 500-8285, Japan

Introduction

Obesity, a condition resulting from an excess of adipose tissue, is currently a serious health problem throughout the world, with approximately 1.6 billion overweight and 500 million obese adults [1]. Numerous health disorders complicate obesity, including cardiovascular disease, hypertension, insulin resistance, diabetes mellitus, and hyperlipidemia, which are collectively known as “metabolic syndrome.” Non-alcoholic fatty liver disease (NAFLD), which is known to be a hepatic manifestation of metabolic syndrome, is also the most common form of chronic liver disease in developed countries [2, 3]. In addition, recently, obesity and its related metabolic abnormalities, especially diabetes mellitus, have been recognized as major risk factors for the development of certain types of human malignancies, including hepatocellular carcinoma (HCC) [4–16]. A prospective study of a population of more than 900,000 American adults showed that a higher body mass index (BMI) is significantly associated with higher rates of death from cancer, including HCC [17].

Mounting evidence obtained from experimental and epidemiological studies indicates that several pathophysiological mechanisms link obesity and liver carcinogenesis, including the emergence of insulin resistance, alterations in the insulin-like growth factor-1 (IGF-1)/IGF-1 receptor (IGF-1R) axis, a state of chronic inflammation, induction of oxidative stress, and the occurrence of adipokine imbalance [4–8]. Insulin resistance leads to an increased expression of proinflammatory cytokine tumor necrosis factor- α (TNF- α) and interleukin-6 (IL-6), central mediators of chronic inflammatory diseases, and their dysregulation is associated with the development of steatosis and inflammation within the liver [4–8]. Therefore, among obesity-related pathophysiological conditions that cooperatively enhance the development of HCC, insulin resistance and the subsequent inflammatory cascade are thought to play a critical role in the development of HCC [4–8]. On the other hand, studies

of these conditions also suggest that such pathophysiological disorders might be critical targets for inhibiting obesity-related carcinogenesis [18]. For instance, experimental studies have revealed that improvement of chronic inflammation by inhibiting the expression of TNF- α and IL-6 plays a significant role in the prevention of obesity-related colorectal tumorigenesis [19–21].

The present review aims to summarize multiple pathogenic mechanisms by which obesity and related metabolic disorders influence the development of HCC, focusing on the emergence of insulin resistance and the subsequent inflammatory cascade. This article also aims to review the possibility that nutritional or pharmaceutical approaches targeting pathophysiological conditions caused by obesity might be effective in preventing obesity-related liver carcinogenesis.

Obesity, diabetes mellitus, and HCC

HCC, which is the dominant form of primary liver carcinoma worldwide, is one of the most frequently occurring cancers in the world, accounting for 750,000 annual cases; approximately the same number of individuals (700,000) die from this malignancy each year [22]. Although HCC development is frequently associated with chronic inflammation and subsequent cirrhosis of the liver induced by a persistent infection with hepatitis B virus (HBV) or hepatitis C virus (HCV), recent epidemiological and clinical studies have revealed that obesity and diabetes mellitus are major risk factors for the development of HCC [6–9, 12–16, 23]. In particular, a recent meta-analysis concluded that the summary relative risk of HCC was 117 % for overweight subjects (BMI 25–30 kg/m²) and 189 % for the obese individuals (BMI \geq 30 kg/m²) [14]. Obesity represents an independent HCC risk factor in patients with alcoholic and cryptogenic cirrhosis [15]. The association between HCC development and diabetes, which is characterized by hyperglycemia, insulin resistance, and hyperinsulinemia, has also been ascertained by repeated meta-analyses [10, 11]. In one population-based study, diabetes increased the risk of HCC by threefold [23]. Insulin resistance has also been shown to raise the risk for recurrence of HCC after curative radiofrequency ablation in HCV-positive patients [13].

The relationship between HCV infection and metabolic syndrome is clinically relevant because insulin resistance and subsequent diabetes and severe steatosis frequently occur in HCV-infected patients [24, 25]. Furthermore, there are synergistic effects between metabolic disorders (obesity and diabetes) and other HCC risk factors such as hepatitis virus infection and alcohol consumption [23, 26–29]. A long-term (14 years) follow-up study in Taiwan has shown that the combined presence of HCV and diabetes is

associated with a 37-fold increase in the rate of HCC development [23]. Moreover, HCC risk is increased by more than 100-fold in HBV or HCV carriers with both obesity and diabetes [23]. A recent prospective study showed that insulin resistance itself is associated with HCC in HCV-positive cirrhosis and is a strong predictor of liver-related death or transplantation [30]. Therefore, viral hepatitis patients with metabolic disorders would seem to be at high risk for the development of HCC and thus should be closely monitored for this malignancy.

NAFLD, nonalcoholic steatohepatitis, and HCC

NAFLD is the major hepatic manifestation of obesity and its related metabolic disorders, particularly diabetes mellitus and dyslipidemia, and has become one of the most common liver disorders in developed countries [2, 3, 31, 32]. The accumulation of fat caused by excess energy intake can result in liver dysfunction as the liver synthesizes more triglycerides but fails to export them. Triglyceride deposition in hepatocytes leads to hepatic steatosis. The overlap between the prevalence of NAFLD and diabetes is equally substantial [32]. On the other hand, NAFLD is commonly associated with insulin resistance and hyperinsulinemia even in the non-obese [33], indicating that insulin resistance might be a key factor in the development of NAFLD. In addition, NAFLD that has not yet progressed to nonalcoholic steatohepatitis (NASH) can induce hepatocyte proliferation and hepatic hyperplasia, both of which initiate the hepatic neoplastic process in obesity [34].

While most patients with NAFLD remain asymptomatic, 20 % progress to develop chronic hepatic inflammation or NASH, which in turn can lead to liver fibrosis, portal hypertension, cirrhosis, HCC development, and increased mortality [2, 3, 31, 32, 35]. A subsequent study of natural history in NAFLD indicates that steatohepatitis is a risk for the development of cirrhosis and HCC [36]. The exact prevalence of HCC in NASH remains unknown; however, some prospective studies found at least 2 to 3 % yearly cumulative incidence of HCC in patients with NASH [37, 38]. In 1998, Day and James proposed a “two-hit theory” to explain NAFLD/NASH pathogenesis [39]. The first hit, the flux of free fatty acids into the liver and subsequent hepatic steatosis, plays a role in lipotoxicity-induced mitochondrial abnormalities that sensitize the liver to additional proinflammatory insults, the second hit. These hits include enhanced lipid peroxidation and increased generation of reactive oxygen species. Insulin resistance is also regarded as a critical factor in the etiology of NASH [39, 40].

Potential pathophysiological mechanisms linking obesity and HCC development

Figure 1 shows several pathophysiological mechanisms linking obesity and its related metabolic abnormalities to liver carcinogenesis. Substantial evidence has shown that insulin resistance, among various obesity-related metabolic disorders, significantly contributes to the development of HCC. Insulin, which is a key regulator of glucose metabolism itself, and the signal transduction network it regulates play important roles in oncogenesis [41, 42]. Insulin induces HCC cells to proliferate and resist apoptosis [43, 44], suggesting that hyperinsulinemia directly contributes to the growth of HCC cells. In addition, insulin resistance increases the biological activity of IGF-1, an important endocrine and paracrine regulator of tissue growth and metabolism. Numerous pieces of evidence indicate that the IGF-1/IGF-1R axis plays an important role in the carcinogenesis of many cancer types, including HCC [41, 42]. Insulin receptor and IGF-1R are receptor tyrosine kinases, and the binding of insulin and IGF-1 to their respective receptors on tumors and precancerous cells activates the phosphatidylinositol 3-kinase (PI3K)/Akt pathway, which is responsible for cellular processes like growth, proliferation, and survival [41, 42]. IGF-1R activity is also required for oncogenic transformation by a number of oncogenes,

including RAS, and can promote tumor formation in vivo [41, 45]. Activation of the IGF/IGF-1R axis is critically involved in the growth of HCC cells and in liver carcinogenesis [46–48]. For HCC, IGF-1R activation is observed in a subgroup of tumor cells but not in adjacent cirrhotic tissue [48]. We have recently reported that insulin resistance and the activation of IGF/IGF-1R axis are involved in liver carcinogen *N*-diethylnitrosamine (DEN)-induced liver tumorigenesis in obese and diabetic *C57BL/KsJ-db/db (db/db)* mice [49, 50].

An adipokine imbalance caused by excess production of storage lipids may also be related to obesity-associated liver carcinogenesis. For instance, higher levels of serum leptin, which regulates energy homeostasis and is elevated in obese individuals [51], increase the risk of HCC recurrence after curative treatment [52]. Leptin stimulates the growth of HCC cells by upregulating cyclin D1 expression [53]. Treatment with leptin also increases the proliferation of HCC-derived cells by activating several signaling pathways: signal transducer and activator of transcription-3 (Stat3), AKT, and extracellular signal-regulated kinase (ERK) [54]. In animal models, leptin has been shown to promote angiogenesis and thus could facilitate the progression of NASH to HCC [55]. In addition, lack of adiponectin, the other member of the adipokine group that is significantly reduced in obese individuals [56], enhances the progression of hepatic




Fig. 1 Proposed mechanisms linking obesity and its related metabolic abnormalities to the development of HCC

steatosis and tumor formation in a mouse model of NASH [57]. However, this adipokine alleviates hepatic steatosis [58]. In vitro and in vivo studies show that adiponectin exerts antitumor effects in HCC cells [59]. Moreover, the induction of adiponectin plays a role in the suppression of chemically induced liver tumorigenesis in obese mice [60]. These findings suggest that obesity and its related metabolic abnormalities, such as sustained insulin resistance, activation of the IGF-1/IGF-1R axis, and adipokine imbalance, play an important role in the development of HCC and thus might be promising targets in the prevention of obesity-related liver tumorigenesis.

Obesity-induced insulin resistance and chronic inflammation

There is substantial evidence that obesity is associated with chronic low-grade systemic inflammation, which contributes to metabolic disorders and the progression from hepatic steatosis to NASH and subsequent HCC development [4–8]. Hypertrophic adipocytes, which are associated with the deposition and accumulation of excess lipids, secrete free fatty acids (FFAs); in addition, together with various immune cells, they release various proinflammatory cytokines, including TNF- α and IL-6 [4–8]. In particular, macrophage infiltration into white adipose tissue, which is accompanied by TNF- α and IL-6 production, is an early contributing event for the development of chronic low-grade systemic inflammation [61, 62]. In 1993, Hotamisligil et al. demonstrated that adipocytes constitutively express TNF- α and neutralization of TNF- α by soluble TNF- α receptor decreases insulin resistance in obese mice [63]. This suggests that TNF- α lies at the core of the association between obesity and insulin resistance. TNF- α enhances obesity-related systemic insulin resistance by inhibiting the tyrosine phosphorylation of insulin receptor [64]. On the other hand, the loss of TNF- α and its receptor improves insulin sensitivity in obese mice [65]. TNF- α contributes to obesity-induced IL-6 production, which causes hepatic inflammation and activates ERK and Stat3 [66]. TNF- α and IL-6 expressions in the liver are strongly induced in response to a high-fat diet, but inhibition of TNF- α signaling or ablation of IL-6 prevents hepatosteatosis [66]. Type 2 diabetes is an inflammatory condition, as evidenced by the elevated concentrations of IL-6, which induces cellular insulin resistance in hepatocytes, observed in these patients [67–69]. The concentration of IL-6 together with IL-1 β , which is another inflammatory cytokine that induces insulin resistance in liver-derived cells, is a more predictive risk factor for type 2 diabetes in humans than either cytokine alone [70, 71]. TNF- α and IL-6 increase the levels of leptin, whereas leptin influences inflammatory responses, possibly by triggering

the release of TNF- α and IL-6 [72, 73]. Hepatic steatosis has negative effects on liver function, which might be mediated by inflammation because the expression of TNF- α , IL-6, and IL-1 β mRNA increases in the liver with increasing adiposity [74].

Cytokine signaling pathway associated with obesity-induced inflammation and HCC development

Several specific intracellular signaling pathways, including c-Jun N-terminal kinase (JNK) and nuclear factor (NF)- κ B, have emerged as potential targets for many inflammatory cytokines and chemokines that promote obesity-related metabolic disorders such as insulin resistance [75]. For instance, activation of JNK inhibits normal tyrosine phosphorylation of insulin receptor substrate-1 and downstream insulin signal transduction [76]. The effects of obesity-induced activation of NF- κ B are mediated through the synthesis of NF- κ B target gene expression, including TNF- α , IL-6, and IL-1 β [77]. Therefore, activation of JNK and NF- κ B is associated with the induction of insulin resistance, whereas their inhibition provides glucose tolerance and protection from obesity in rodents [75]. Reactive oxygen species that are increased by adiposity have also been shown to activate JNK and NF- κ B [78]. In addition, saturated FFAs lead to JNK activation, which can, in turn, increase the production of inflammatory cytokines capable of causing insulin resistance [79]. Saturated FFAs have also been found to enhance NF- κ B activation in macrophages [80], suggesting that there is a potential link between elevated circulating or tissue lipid concentrations and the part of the immune system that mediates inflammation. In hepatocytes, saturated FFAs can induce time- and dose-dependent lipoapoptosis, which is the combination of lipid accumulation and induction of apoptosis in hepatocytes [81]. Experimental data have also shown that FFAs cause TNF- α production and subsequent NF- κ B activation by promoting hepatic lipotoxicity [82]. These findings appear significant because lipotoxicity and lipoapoptosis play a pivotal role in the progression of NAFLD to NASH [83]. JNK1 activation also promotes the development of NASH in mice fed with methionine- and choline-deficient diets [84], which indicates that JNK and NF- κ B are critical factors in the occurrence of NAFLD and its progression to NASH.

The role of obesity-induced inflammation in liver tumorigenesis has recently been demonstrated in several experimental models [50, 66, 85, 86]. For instance, administration of DEN was found to enhance the development of preneoplastic lesions in the livers of rats fed with high-fat diets and this was associated with elevated TNF- α /NF- κ B signaling and ERK-related hepatocyte proliferation [85]. Phosphorylation of ERK, Akt, Stat3, and JNK proteins and upregulation of

TNF- α , IL-6, and IL-1 β in the liver are involved in DEN-induced liver tumorigenesis in *db/db* obese mice [50]. Enhanced production of adipose-derived TNF- α and IL-6 and activation of Stat3 are critical in the development of obesity-related liver tumorigenesis [66]. This study [66], together with another recent study [87], clearly indicates that Stat3 activation, which is associated with TNF- α and IL-6 production in hepatocytes, is essential for liver carcinogenesis.

Targeting obesity-related metabolic abnormalities for cancer prevention

As mentioned earlier, obesity and its related metabolic abnormalities, such as a state of chronic inflammation, play a critical role in the development of HCC. On the other hand, these findings may suggest the possibility that the metabolic disorders caused by obesity might be effective targets in the prevention of liver carcinogenesis [18]. For instance, ablation of IL-6 or inhibition of TNF- α signaling can inhibit obesity-promoted hepatocarcinogenesis by reducing hepatosteatosis and steatohepatitis [66]. Treatment with adiponectin, an anti-inflammatory adipokine, also reduces liver tumorigenesis in nude mice [59].

To verify our hypothesis that targeting metabolic abnormalities caused by obesity might be an effective strategy for preventing cancer development in obese individuals, we have conducted several experimental studies. We initially performed chemopreventive studies using a mouse model of obesity-related colorectal carcinogenesis because increased body fat levels and BMI are associated with an increased risk of colorectal cancer [17, 88, 89]. The model used obese and diabetic *db/db* mice, which are susceptible to the colonic carcinogen azoxymethane (AOM) and thus easily develop colonic precancerous lesions [90]. We have found that pitavastatin and renin-angiotensin system inhibitors, which are drugs for hyperlipidemia and hypertension, respectively, suppress AOM-induced colonic preneoplastic lesions in *db/db* mice by inhibiting the levels of TNF- α and IL-6 in the serum and colonic mucosa [20, 21]. Curcumin, a component of turmeric, also exerts chemopreventive effects in the development of obesity-related colonic preneoplastic lesions in *db/db* mice, and this is associated with inhibition of NF- κ B activity and TNF- α and IL-6 expression in the colonic mucosa [19]. Furthermore, branched-chain amino acids (BCAA) and (-)-epigallocatechin gallate (EGCG) prevent obesity-related colorectal carcinogenesis by improving insulin resistance and inhibiting IGF/IGF-1R axis in these mice [91, 92].

Among these agents, BCAA is considered as one of the most promising candidates to prevent obesity-related liver tumorigenesis. This is because it is widely used for the treatment of protein energy malnutrition (PEM) that frequently occurs in

patients with liver cirrhosis [93–96]. EGCG, a major biologically active component of green tea, also seems to have a considerable effect given that green tea catechins (GTCs) improve metabolic abnormalities and possess anticancer and cancer chemopreventive properties [97–100]. In the following sections, we will discuss in detail the effects of BCAA and EGCG in the prevention of obesity-related liver tumorigenesis based on our recent experimental studies. In addition, we also discuss the effects of acyclic retinoid (ACR), which is a promising agent for the chemoprevention of HCC [101–104], on the prevention of liver tumorigenesis in obese mice.

Preventive effects of BCAA on obesity-related liver tumorigenesis

Because the liver is a critical organ for regulating metabolism, a variety of nutritional and metabolic disorders, such as PEM and insulin resistance, are frequently seen in patients with chronic liver diseases [93–96, 105, 106]. Decreased serum levels of BCAA (valine, leucine, and isoleucine) and albumin appear with a high incidence in liver cirrhosis, whereas supplementation with BCAA has been shown to improve PEM and increase the serum albumin concentration in cirrhotic patients. This subsequently improves the quality of life and prognosis in patients with liver cirrhosis by preventing complications associated with the disease [93–96]. In addition, recent clinical and experimental studies have revealed that BCAA improves insulin resistance and glucose tolerance [107–110]. In 2005, Muto et al. reported the results of a large-scale ($n=622$) multicenter randomized controlled trial, the Long-Term Survival Study, which investigated the effects of supplemental BCAA therapy on event-free survival in patients with decompensated cirrhosis. In the trial, oral supplementation with a BCAA preparation significantly prevented progressive hepatic failure and improved event-free survival [95], strongly suggesting that supplementation with BCAA can serve as a first-line therapy for patients with decompensated cirrhosis.

Moreover, it should be emphasized that the results of the subset analysis from this trial demonstrated that long-term oral supplementation with BCAA was associated with a reduced frequency of HCC in obese cirrhotic patients ($P=0.008$) [12]. To clarify the precise mechanisms of BCAA in the prevention of the development of HCC in obese cirrhotic patients, we performed an experimental study using the obesity-related liver carcinogenesis model in *db/db* mice [49]. In the study, BCAA supplementation significantly suppressed the development of DEN-induced hepatic preneoplastic lesions in *db/db* mice by inhibiting the expression of IGF-1, IGF-2, and IGF-1R in the liver. The development of liver neoplasms, including hepatic adenoma and HCC,

was also reduced by BCAA supplementation, and this was associated with improvement of insulin resistance, reduction of serum leptin levels, and attenuation of hepatic steatosis and fibrosis [49]. Obese cirrhotic patients generally have a particularly high incidence of hyperinsulinemia and insulin resistance [105, 106]. Therefore, our findings [49], together with the results of an *in vitro* study showing that BCAA suppresses insulin-induced proliferation of HCC cells by inhibiting the insulin-induced activation of the PI3K/Akt pathway [111], suggest that BCAA supplementation reduced the risk of developing HCC in obese cirrhotic patients. This was accomplished, at least in part, by targeting insulin resistance and its related signaling pathways (Fig. 2; Table 1). These findings are consistent with the results of an experimental study reported by Yoshiji et al. showing the chemopreventive effects of BCAA supplementation against liver tumorigenesis in obese and diabetic rats, which are also complicated with insulin resistance [112].

In addition, in our unpublished study, BCAA supplementation was shown to suppress the spontaneous development of hepatic preneoplastic lesions in *db/db* mice by inhibiting the expression of TNF- α , IL-6, and IL-1 β mRNA in the liver. BCAA supplementation also inhibited increased macrophage infiltration and the expression of TNF- α , IL-6, and monocyte chemoattractant protein-1 mRNA in the white adipose tissue, suggesting that chronic inflammation induced by obesity in the liver and adipose tissue could also serve as a critical target of BCAA in the inhibition of the

early phase of obesity-related liver tumorigenesis (unpublished data).

Preventive effects of GTCs on obesity-related liver tumorigenesis

Green tea is a beverage commonly consumed worldwide. Its component polyphenols, which are known as GTCs, have received great attention for their beneficial effects, particularly their involvement in the improvement of metabolic abnormalities and prevention of certain types of malignancies [97–100]. A recent meta-analysis of clinical trials reported that GTCs help reduce body weight [98]. Supplementation with GTCs was found to decrease plasma levels of insulin, TNF- α , and IL-6 and improve hepatic steatosis and liver dysfunction in a rodent model of obesity and diabetes. This indicated that treatment with GTCs is effective in the prevention of the progression of obesity-related metabolic disorders such as chronic inflammation [113–115]. The anti-inflammatory properties of GTCs are also responsible for the anticancer and cancer-preventive effects of the molecules [99]. EGCG, a type of GTC, suppresses inflammation-related colon carcinogenesis in mice by decreasing the mRNA expression of TNF- α and IL-6 in the colonic mucosa [116]. EGCG also inhibits proliferation and induces apoptosis in HCC- and colorectal cancer-derived cells by inhibiting the activation of IGF-1R and its




Fig. 2 Mechanisms of action of BCAA, EGCG, and ACR in the inhibition of obesity-related liver carcinogenesis

Table 1 Suppressive effects of BCAA, EGCG, ACR, and pitavastatin on obesity-related liver tumorigenesis in *db/db* mice

| Agent | Inhibition rate (%) | | Inhibition mechanisms | Reference number | |
|--------------|---------------------|--------------------|--|--|------|
| | Adenoma | FCA ^a | | | |
| BCAA | 75 ^{b, c} | 50 ^{b, c} | Hepatic IGF-1, IGF-2, and IGF-1R mRNAs ↓ Hepatic steatosis ↓ Hepatic fibrosis ↓ | Serum leptin and ALT levels ↓ Insulin sensitivity ↑ Hepatocyte proliferation ↓ | [49] |
| EGCG | 86 ^c | 48 ^c | Hepatic pIGF-1R, pERK, and pAkt proteins ↓ Hepatic steatosis ↓ Hepatic and systemic inflammation ↓ | Serum insulin, IGF-1, IGF-2, and FFA levels ↓ Hepatic pAMPK protein ↑ | [50] |
| ACR | 86 ^c | 81 ^c | Hepatic Ras activity ↓ Hepatic RARβ and p21 ^{CIP1} mRNAs ↑ Insulin sensitivity ↑ | Hepatic pRXRα, pERK, and pStat3 proteins ↓ Hepatic steatosis ↓ Hepatic and systemic inflammation ↓ | [86] |
| Pitavastatin | NE ^d | 29 ^e | Pro-apoptotic effect ↑ Hepatic steatosis ↓ Serum adiponectin level ↑ | Hepatocyte proliferation ↓ Hepatic pAMPK protein ↑ Hepatic and systemic inflammation ↓ | [60] |

^aFoci of cellular alteration

^bCompared to the casein supplementation mice (a nitrogen content-matched control for BCAA)

^cMice were treated with agent for 34 weeks

^dNot examined

^eMice were treated with agent for 14 weeks

downstream signaling pathways, including Ras/MAPK and PI3K/Akt [46, 117]. In addition, this agent prevents carbon tetrachloride-induced hepatic fibrosis in rats by inhibiting IGF-1R expression [118], indicating that the IGF/IGF-1R axis, which is critically involved in cancer development and obesity-related metabolic disorder, might be a critical target of GTCs. Several interventional studies also provide clear evidence for the chemopreventive effects and safety of tea preparations [119–121].

Because GTCs are expected to improve metabolic disorders and exert chemopreventive properties by targeting chronic inflammation and the IGF/IGF-1R axis, we examined whether EGCG treatment inhibits obesity-associated liver tumorigenesis [50]. We found that drinking water containing EGCG significantly inhibited the development of hepatic preneoplastic lesions and adenoma [50]. EGCG consumption also improved hepatic steatosis; decreased the serum levels of insulin, IGF-1, and IGF-2; and inhibited the phosphorylation of the IGF-1R, ERK, Akt, Stat3, and JNK proteins in the liver of obese mice [50]. The serum levels of FFA and TNF-α were also decreased by drinking EGCG, which additionally lowered the expression of TNF-α, IL-6, and IL-1β mRNAs in the liver [50]. These findings suggest that EGCG prevents obesity-related liver tumorigenesis by inhibiting the IGF/IGF-1R axis, improving hyperinsulinemia, and attenuating chronic inflammation (Fig. 2; Table 1). Thus, in addition to BCAA, GTCs may also be useful in the chemoprevention of liver tumorigenesis in obese individuals.

Preventive effects of ACR on obesity-related liver tumorigenesis

Retinoids, a group of structural and functional derivatives of vitamin A, play fundamental roles in cellular activities, including growth, differentiation, and apoptosis, as well as in morphology [122, 123]. Because of this, loss of retinoid activity or responsiveness is linked to the development of several types of human malignancies, including HCC; therefore, they might be critical targets for cancer chemoprevention and chemotherapy [103, 104, 124, 125]. Retinoids exert their biological functions primarily by regulating gene expression through two distinct nuclear receptors, the retinoic acid receptors (RARs) and retinoid X receptors (RXRs), both of which are composed of three subtypes (α, β, and γ) [122, 123]. Among the retinoid receptors, RXRα is thought to be one of the most important with respect to exerting fundamental effects on cellular activities. This is because it forms a heterodimer with other nuclear receptors and thereby acts as the master regulator of nuclear receptors [122, 123]. We have reported that abnormalities in the expression and function of RXRα are prominently involved in the development of HCC. The repression of RXRα was found to occur in the early stages of liver carcinogenesis in a rat model of chemically induced liver carcinogenesis [126]. Moreover, a malfunction of the RXRα due to phosphorylation by the Ras/MAPK signaling pathway is significantly

associated with liver carcinogenesis. That is, accumulation of phosphorylated RXR α protein, which is regarded as the nonfunctional form of RXR α , interferes with the function of normal (unphosphorylated) RXR α in a dominant-negative manner, thus playing a critical role in HCC development [103, 104, 127–130]. These findings therefore suggest that targeting RXR α phosphorylation may be an effective and important strategy for the prevention and treatment of HCC.

ACR, a synthetic retinoid that was initially developed as an agonist for RXR, is a possible candidate for this purpose because it can impede the development of HCC and it inhibits cancer cell growth by repressing the Ras/MAPK signaling pathway and subsequent RXR α phosphorylation [103, 104, 128, 131]. One early-phase randomized controlled clinical trial tested the chemopreventive effect of ACR on secondary HCC in patients who underwent potentially curative treatment for initial HCC. In this study, oral administration of ACR significantly reduced the incidence of recurrent or new HCC ($P=0.04$) and improved the recurrence-free survival ($P=0.002$) and overall survival rates ($P=0.04$) [101, 102]. Moreover, a large-scale ($n=401$) randomized placebo-controlled trial (phase II/III trial) also showed that ACR had a strong effect on the prevention of second primary HCC in HCV-positive patients. It showed a hazard ratio of 0.27 (95 % CI, 0.07–0.96) 2 years after the treatment, indicating that ACR reduced the recurrence of HCC, particularly after 2 years of treatment [132].

Because numerous preclinical experiments and clinical trials indicate that ACR is a promising agent for the chemoprevention of HCC, we investigated whether ACR could prevent obesity-related liver tumorigenesis [86]. In the study, treatment with ACR effectively prevented the development of obesity-related liver tumorigenesis by inhibiting the activation of Ras and the phosphorylation of ERK and RXR α , thus restoring RXR α function in the liver of DEN-treated *db/db* mice [86]. ACR administration also inhibits this tumorigenesis through attenuation of the chronic inflammation induced by excessive fatty deposits, as demonstrated by the improved liver steatosis and decreased serum TNF- α levels and expression levels of TNF- α , IL-6, and IL-1 β mRNA in the liver [86]. In addition, ACR administration improved insulin sensitivity, which was also associated with the prevention of obesity-related liver tumorigenesis [86] (Fig. 2; Table 1). Therefore, the results obtained from both clinical trials [101, 102, 132] and this preclinical experiment [86] encourage the clinical use of ACR for cirrhotic patients with obesity and diabetes who are at a notably higher risk of developing HCC.

Conclusion

Obesity and its related metabolic abnormalities, including increased cancer risk, are a serious public health problem worldwide. Among all cancers, HCCs are the malignancies most frequently affected by obesity. The liver disease influenced most by obesity is NAFLD, and this disease, by itself and in synergy with other risk factors such as hepatitis virus infection, is becoming one of the most common causes of HCC in developed countries. Therefore, there is an urgent need to develop more effective therapeutic strategies to prevent the development of obesity-related HCC or halt its progression. Obesity and diabetes enhance HCC development through insulin resistance, activation of the IGF/IGF-1R axis, and lipid accumulation within hepatocytes, thereby leading to a chronic low-grade systemic inflammation. This involves abnormalities of various types of cytokines and adipokines. Among them, TNF- α and IL-6 play a critical role in the onset of NASH and the initiation and promotion of HCC.

In this review, we indicate the possibility that pharmaceutical and nutraceutical approaches for targeting and restoring metabolic disorders, especially chronic low-grade inflammation involving increased levels of TNF- α and IL-6, may be an effective strategy for preventing the development of obesity-related HCC. We further indicate that BCAA, GTCs, and ACR are considered as some of the most promising agents for achieving this purpose. Therefore, further advanced translational research, such as pilot trials, to clarify whether active intervention using these agents can prevent the development and recurrence of HCC in patients with chronic liver disease and obesity is required. In addition, further experimental studies to determine whether specific drugs, such as antidiabetic drugs, anti-hypertensive drugs, and lipid-lowering drugs, can inhibit obesity-related liver carcinogenesis should be performed. Considering that these drugs are widely used for patients with metabolic syndrome, it would be beneficial if they could exert chemopreventive effects on obesity-associated carcinogenesis. Our recent findings that pitavastatin, a recently developed lipophilic statin, suppresses the development of chemically induced colonic and hepatic preneoplastic lesions in *db/db* mice by attenuating chronic inflammation may provide a basis for this attempt [21, 60] (Fig. 2; Table 1).

Acknowledgments This review was based on studies supported in part by Grants-in-Aid from the Ministry of Education, Science, Sports, and Culture of Japan (No. 22790638 to M. S. and No. 21590838 to H. M.) and by Grant-in-Aid for the 3rd Term Comprehensive 10-Year Strategy for Cancer Control from the Ministry of Health, Labour, and Welfare of Japan.

Conflict of interest The authors declare no conflict of interest.

References

- WHO: World Health Organization fact sheet for world wide prevalence of obesity. <http://www.who.int/mediacentre/factsheets/fs311/en/index.html>. Accessed on 27 December 2011
- Angulo P (2002) Nonalcoholic fatty liver disease. *N Engl J Med* 346(16):1221–1231
- Siegel AB, Zhu AX (2009) Metabolic syndrome and hepatocellular carcinoma: two growing epidemics with a potential link. *Cancer* 115(24):5651–5661
- Calle EE, Kaaks R (2004) Overweight, obesity and cancer: epidemiological evidence and proposed mechanisms. *Nat Rev Cancer* 4(8):579–591
- Anderson AS, Caswell S (2009) Obesity management—an opportunity for cancer prevention. *Surgeon* 7(5):282–285
- Sun B, Karin M (2012) Obesity, inflammation, and liver cancer. *J Hepatol* 56(3):704–713
- El-Serag HB, Rudolph KL (2007) Hepatocellular carcinoma: epidemiology and molecular carcinogenesis. *Gastroenterology* 132(7):2557–2576
- El-Serag HB, Tran T, Everhart JE (2004) Diabetes increases the risk of chronic liver disease and hepatocellular carcinoma. *Gastroenterology* 126(2):460–468
- Davila JA, Morgan RO, Shaib Y, McGlynn KA, El-Serag HB (2005) Diabetes increases the risk of hepatocellular carcinoma in the United States: a population based case control study. *Gut* 54(4):533–539
- El-Serag HB, Hampel H, Javadi F (2006) The association between diabetes and hepatocellular carcinoma: a systematic review of epidemiologic evidence. *Clin Gastroenterol Hepatol* 4(3):369–380
- Wang P, Kang D, Cao W, Wang Y, Liu Z (2012) Diabetes mellitus and risk of hepatocellular carcinoma: a systematic review and meta-analysis. *Diabetes Metab Res Rev* 28(2):109–122
- Muto Y, Sato S, Watanabe A, Moriwaki H, Suzuki K, Kato A, Kato M, Nakamura T, Higuchi K, Nishiguchi S, Kumada H, Ohashi Y, Long-Term Survival Study Group (2006) Overweight and obesity increase the risk for liver cancer in patients with liver cirrhosis and long-term oral supplementation with branched-chain amino acid granules inhibits liver carcinogenesis in heavier patients with liver cirrhosis. *Hepatol Res* 35(3):204–214
- Imai K, Takai K, Nishigaki Y, Shimizu S, Naiki T, Hayashi H, Uematsu T, Sugihara J, Tomita E, Shimizu M, Nagaki M, Moriwaki H (2010) Insulin resistance raises the risk for recurrence of stage I hepatocellular carcinoma after curative radiofrequency ablation in hepatitis C virus-positive patients: a prospective, case series study. *Hepatol Res* 40(4):376–382
- Larsson SC, Wolk A (2007) Overweight, obesity and risk of liver cancer: a meta-analysis of cohort studies. *Br J Cancer* 97(7):1005–1008
- Nair S, Mason A, Eason J, Loss G, Perrillo RP (2002) Is obesity an independent risk factor for hepatocellular carcinoma in cirrhosis? *Hepatology* 36(1):150–155
- Regimbeau JM, Colombat M, Mognol P, Durand F, Abdalla E, Degott C, Degos F, Farges O, Belghiti J (2004) Obesity and diabetes as a risk factor for hepatocellular carcinoma. *Liver Transpl* 10(2 Suppl 1):S69–S73
- Calle EE, Rodriguez C, Walker-Thurmond K, Thun MJ (2003) Overweight, obesity, and mortality from cancer in a prospectively studied cohort of U.S. adults. *N Engl J Med* 348(17):1625–1638
- Shimizu M, Kubota M, Tanaka T, Moriwaki H (2012) Nutritional approach for preventing obesity-related colorectal and liver carcinogenesis. *Int J Mol Sci* 13(1):579–595
- Kubota M, Shimizu M, Sakai H, Yasuda Y, Terakura D, Baba A, Ohno T, Tsurumi H, Tanaka T, Moriwaki H (2012) Preventive effects of curcumin on the development of azoxymethane-induced colonic preneoplastic lesions in male C57BL/KsJ-db/db obese mice. *Nutr Cancer* 64(1):72–79
- Kubota M, Shimizu M, Sakai H, Yasuda Y, Ohno T, Kochi T, Tsurumi H, Tanaka T, Moriwaki H (2011) Renin-angiotensin system inhibitors suppress azoxymethane-induced colonic preneoplastic lesions in C57BL/KsJ-db/db obese mice. *Biochem Biophys Res Commun* 410(1):108–113
- Yasuda Y, Shimizu M, Shirakami Y, Sakai H, Kubota M, Hata K, Hirose Y, Tsurumi H, Tanaka T, Moriwaki H (2010) Pitavastatin inhibits azoxymethane-induced colonic preneoplastic lesions in C57BL/KsJ-db/db obese mice. *Cancer Sci* 101(7):1701–1707
- Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D (2011) Global cancer statistics. *CA Cancer J Clin* 61(2):69–90
- Chen CL, Yang HI, Yang WS, Liu CJ, Chen PJ, You SL, Wang LY, Sun CA, Lu SN, Chen DS, Chen CJ (2008) Metabolic factors and risk of hepatocellular carcinoma by chronic hepatitis B/C infection: a follow-up study in Taiwan. *Gastroenterology* 135(1):111–121
- Kaddai V, Negro F (2011) Current understanding of insulin resistance in hepatitis C. *Expert Rev Gastroenterol Hepatol* 5(4):503–516
- Negro F (2010) Abnormalities of lipid metabolism in hepatitis C virus infection. *Gut* 59(9):1279–1287
- Hassan MM, Hwang LY, Hatten CJ, Swaim M, Li D, Abbruzzese JL, Beasley P, Patt YZ (2002) Risk factors for hepatocellular carcinoma: synergism of alcohol with viral hepatitis and diabetes mellitus. *Hepatology* 36(5):1206–1213
- Veldt BJ, Chen W, Heathcote EJ, Wedemeyer H, Reichen J, Hofmann WP, de Knegt RJ, Zeuzem S, Manns MP, Hansen BE, Schalm SW, Janssen HL (2008) Increased risk of hepatocellular carcinoma among patients with hepatitis C cirrhosis and diabetes mellitus. *Hepatology* 47(6):1856–1862
- Wang CS, Yao WJ, Chang TT, Wang ST, Chou P (2009) The impact of type 2 diabetes on the development of hepatocellular carcinoma in different viral hepatitis statuses. *Cancer Epidemiol Biomarkers Prev* 18(7):2054–2060
- Lai SW, Chen PC, Liao KF, Muo CH, Lin CC, Sung FC (2012) Risk of hepatocellular carcinoma in diabetic patients and risk reduction associated with anti-diabetic therapy: a population-based cohort study. *Am J Gastroenterol* 107(1):46–52
- Nkontchou G, Bastard JP, Ziou M, Aout M, Cosson E, Ganne-Carrie N, Grando-Lemaire V, Roulot D, Capeau J, Trinchet JC, Vicaute E, Beaugrand M (2010) Insulin resistance, serum leptin, and adiponectin levels and outcomes of viral hepatitis C cirrhosis. *J Hepatol* 53(5):827–833
- Byrne CD, Olufadi R, Bruce KD, Cagampang FR, Ahmed MH (2009) Metabolic disturbances in non-alcoholic fatty liver disease. *Clin Sci* 116(7):539–564
- Williams CD, Stengel J, Asike MI, Torres DM, Shaw J, Contreras M, Landt CL, Harrison SA (2011) Prevalence of nonalcoholic fatty liver disease and nonalcoholic steatohepatitis among a largely middle-aged population utilizing ultrasound and liver biopsy: a prospective study. *Gastroenterology* 140(1):124–131
- Marchesini G, Brizi M, Morselli-Labate AM, Bianchi G, Bugianesi E, McCullough AJ, Forlani G, Melchionda N (1999) Association of nonalcoholic fatty liver disease with insulin resistance. *Am J Med* 107(5):450–455
- Yang S, Lin HZ, Hwang J, Chacko VP, Diehl AM (2001) Hepatic hyperplasia in noncirrhotic fatty livers: is obesity-related hepatic steatosis a premalignant condition? *Cancer Res* 61(13):5016–5023
- Starley BQ, Calcagno CJ, Harrison SA (2010) Nonalcoholic fatty liver disease and hepatocellular carcinoma: a weighty connection. *Hepatology* 51(5):1820–1832

36. Adams LA, Lymp JF, St Sauver J, Sanderson SO, Lindor KD, Feldstein A, Angulo P (2005) The natural history of nonalcoholic fatty liver disease: a population-based cohort study. *Gastroenterology* 129(1):113–121
37. Ascha MS, Hanouneh IA, Lopez R, Tamimi TA, Feldstein AF, Zein NN (2010) The incidence and risk factors of hepatocellular carcinoma in patients with nonalcoholic steatohepatitis. *Hepatology* 51(6):1972–1978
38. Yatsuji S, Hashimoto E, Tobarai M, Taniai M, Tokushige K, Shiratori K (2009) Clinical features and outcomes of cirrhosis due to non-alcoholic steatohepatitis compared with cirrhosis caused by chronic hepatitis C. *J Gastroenterol Hepatol* 24(2):248–254
39. Day CP, James OF (1998) Steatohepatitis: a tale of two “hits”? *Gastroenterology* 114(4):842–845
40. Sanyal AJ, Campbell-Sargent C, Mirshahi F, Rizzo WB, Contos MJ, Sterling RK, Luketic VA, Shiffman ML, Clore JN (2001) Nonalcoholic steatohepatitis: association of insulin resistance and mitochondrial abnormalities. *Gastroenterology* 120(5):1183–1192
41. Pollak M (2008) Insulin and insulin-like growth factor signalling in neoplasia. *Nat Rev Cancer* 8(12):915–928
42. Clayton PE, Banerjee I, Murray PG, Renehan AG (2011) Growth hormone, the insulin-like growth factor axis, insulin and cancer risk. *Nat Rev Endocrinol* 7(1):11–24
43. Kang S, Song J, Kang H, Kim S, Lee Y, Park D (2003) Insulin can block apoptosis by decreasing oxidative stress via phosphatidylinositol 3-kinase- and extracellular signal-regulated protein kinase-dependent signaling pathways in HepG2 cells. *Eur J Endocrinol* 148(1):147–155
44. Tornkvist A, Parpal S, Gustavsson J, Stralfors P (1994) Inhibition of Raf-1 kinase expression abolishes insulin stimulation of DNA synthesis in H4IIE hepatoma cells. *J Biol Chem* 269(19):13919–13921
45. Sell C, Rubini M, Rubin R, Liu JP, Efstratiadis A, Baserga R (1993) Simian virus 40 large tumor antigen is unable to transform mouse embryonic fibroblasts lacking type 1 insulin-like growth factor receptor. *Proc Natl Acad Sci U S A* 90(23):11217–11221
46. Shimizu M, Shirakami Y, Sakai H, Tatebe H, Nakagawa T, Hara Y, Weinstein IB, Moriwaki H (2008) EGCG inhibits activation of the insulin-like growth factor (IGF)/IGF-1 receptor axis in human hepatocellular carcinoma cells. *Cancer Lett* 262:10–18
47. Alexia C, Fallot G, Lasfer M, Schweizer-Groyer G, Groyer A (2004) An evaluation of the role of insulin-like growth factors (IGF) and of type-I IGF receptor signalling in hepatocarcinogenesis and in the resistance of hepatocarcinoma cells against drug-induced apoptosis. *Biochem Pharmacol* 68(6):1003–1015
48. Tovar V, Alsinet C, Villanueva A, Hoshida Y, Chiang DY, Sole M, Thung S, Moyano S, Toffanin S, Minguez B, Cabellos L, Peix J, Schwartz M, Mazzaferro V, Bruix J, Llovet JM (2010) IGF activation in a molecular subclass of hepatocellular carcinoma and pre-clinical efficacy of IGF-1R blockade. *J Hepatol* 52(4):550–559
49. Iwasa J, Shimizu M, Shiraki M, Shirakami Y, Sakai H, Terakura Y, Takai K, Tsurumi H, Tanaka T, Moriwaki H (2010) Dietary supplementation with branched-chain amino acids suppresses diethylnitrosamine-induced liver tumorigenesis in obese and diabetic C57BL/KsJ-db/db mice. *Cancer Sci* 101(2):460–467
50. Shimizu M, Sakai H, Shirakami Y, Yasuda Y, Kubota M, Terakura D, Baba A, Ohno T, Hara Y, Tanaka T, Moriwaki H (2011) Preventive effects of (–)-epigallocatechin gallate on diethylnitrosamine-induced liver tumorigenesis in obese and diabetic C57BL/KsJ-db/db mice. *Cancer Prev Res* 4(3):396–403
51. Maffei M, Halaas J, Ravussin E, Pratley RE, Lee GH, Zhang Y, Fei H, Kim S, Lallone R, Ranganathan S, Kern PA, Friedman JM (1995) Leptin levels in human and rodent: measurement of plasma leptin and ob RNA in obese and weight-reduced subjects. *Nat Med* 1(11):1155–1161
52. Watanabe N, Takai K, Imai K, Shimizu M, Naiki T, Nagaki M, Moriwaki H (2011) Increased levels of serum leptin are a risk factor for the recurrence of stage I/II hepatocellular carcinoma after curative treatment. *J Clin Biochem Nutr* 49(3):153–158
53. Chen C, Chang YC, Liu CL, Liu TP, Chang KJ, Guo IC (2007) Leptin induces proliferation and anti-apoptosis in human hepatocarcinoma cells by up-regulating cyclin D1 and down-regulating Bax via a Janus kinase 2-linked pathway. *Endocr Relat Cancer* 14(2):513–529
54. Saxena NK, Sharma D, Ding X, Lin S, Marra F, Merlin D, Anania FA (2007) Concomitant activation of the JAK/STAT, PI3K/AKT, and ERK signaling is involved in leptin-mediated promotion of invasion and migration of hepatocellular carcinoma cells. *Cancer Res* 67(6):2497–2507
55. Ikejima K, Takei Y, Honda H, Hirose M, Yoshikawa M, Zhang YJ, Lang T, Fukuda T, Yamashina S, Kitamura T, Sato N (2002) Leptin receptor-mediated signaling regulates hepatic fibrogenesis and remodeling of extracellular matrix in the rat. *Gastroenterology* 122(5):1399–1410
56. Hotta K, Funahashi T, Arita Y, Takahashi M, Matsuda M, Okamoto Y, Iwahashi H, Kuriyama H, Ouchi N, Maeda K, Nishida M, Kihara S, Sakai N, Nakajima T, Hasegawa K, Muraguchi M, Ohmoto Y, Nakamura T, Yamashita S, Hanafusa T, Matsuzawa Y (2000) Plasma concentrations of a novel, adipose-specific protein, adiponectin, in type 2 diabetic patients. *Arterioscler Thromb Vasc Biol* 20(6):1595–1599
57. Kamada Y, Matsumoto H, Tamura S, Fukushima J, Kiso S, Fukui K, Igura T, Maeda N, Kihara S, Funahashi T, Matsuzawa Y, Shimomura I, Hayashi N (2007) Hypoadiponectinemia accelerates hepatic tumor formation in a nonalcoholic steatohepatitis mouse model. *J Hepatol* 47(4):556–564
58. Xu A, Wang Y, Keshaw H, Xu LY, Lam KS, Cooper GJ (2003) The fat-derived hormone adiponectin alleviates alcoholic and nonalcoholic fatty liver diseases in mice. *J Clin Invest* 112(1):91–100
59. Saxena NK, Fu PP, Nagalingam A, Wang J, Handy J, Cohen C, Tighiouart M, Sharma D, Anania FA (2010) Adiponectin modulates C-jun N-terminal kinase and mammalian target of rapamycin and inhibits hepatocellular carcinoma. *Gastroenterology* 139(5):1762–1773, 1773 e1761-1765
60. Shimizu M, Yasuda Y, Sakai H, Kubota M, Terakura D, Baba A, Ohno T, Kochi T, Tsurumi H, Tanaka T, Moriwaki H (2011) Pitavastatin suppresses diethylnitrosamine-induced liver preneoplasms in male C57BL/KsJ-db/db obese mice. *BMC Cancer* 11(1):281
61. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW Jr (2003) Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest* 112(12):1796–1808
62. Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, Sole J, Nichols A, Ross JS, Tartaglia LA, Chen H (2003) Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest* 112(12):1821–1830
63. Hotamisligil GS, Shargill NS, Spiegelman BM (1993) Adipose expression of tumor necrosis factor- α : direct role in obesity-linked insulin resistance. *Science* 259(5091):87–91
64. Hotamisligil GS, Budavari A, Murray D, Spiegelman BM (1994) Reduced tyrosine kinase activity of the insulin receptor in obesity–diabetes. Central role of tumor necrosis factor- α . *J Clin Invest* 94(4):1543–1549
65. Uysal KT, Wiesbrock SM, Marino MW, Hotamisligil GS (1997) Protection from obesity-induced insulin resistance in mice lacking TNF- α function. *Nature* 389(6651):610–614
66. Park EJ, Lee JH, Yu GY, He G, Ali SR, Holzer RG, Osterreicher CH, Takahashi H, Karin M (2010) Dietary and genetic obesity

- promote liver inflammation and tumorigenesis by enhancing IL-6 and TNF expression. *Cell* 140(2):197–208
67. Senn JJ, Klover PJ, Nowak IA, Mooney RA (2002) Interleukin-6 induces cellular insulin resistance in hepatocytes. *Diabetes* 51(12):3391–3399
 68. Pickup JC, Mattock MB, Chusney GD, Burt D (1997) NIDDM as a disease of the innate immune system: association of acute-phase reactants and interleukin-6 with metabolic syndrome X. *Diabetologia* 40(11):1286–1292
 69. Donath MY, Shoelson SE (2011) Type 2 diabetes as an inflammatory disease. *Nat Rev Immunol* 11(2):98–107
 70. Nov O, Kohl A, Lewis EC, Bashan N, Dvir I, Ben-Shlomo S, Fishman S, Wuest S, Konrad D, Rudich A (2010) Interleukin-1beta may mediate insulin resistance in liver-derived cells in response to adipocyte inflammation. *Endocrinology* 151(9):4247–4256
 71. Spranger J, Kroke A, Mohlig M, Hoffmann K, Bergmann MM, Ristow M, Boeing H, Pfeiffer AF (2003) Inflammatory cytokines and the risk to develop type 2 diabetes: results of the prospective population-based European Prospective Investigation into Cancer and Nutrition (EPIC)-Potsdam Study. *Diabetes* 52(3):812–817
 72. Faggioni R, Feingold KR, Grunfeld C (2001) Leptin regulation of the immune response and the immunodeficiency of malnutrition. *FASEB J* 15(14):2565–2571
 73. Molina A, Vendrell J, Gutierrez C, Simon I, Masdevall C, Soler J, Gomez JM (2003) Insulin resistance, leptin and TNF-alpha system in morbidly obese women after gastric bypass. *Obes Surg* 13(4):615–621
 74. Cai D, Yuan M, Frantz DF, Melendez PA, Hansen L, Lee J, Shoelson SE (2005) Local and systemic insulin resistance resulting from hepatic activation of IKK-beta and NF-kappaB. *Nat Med* 11(2):183–190
 75. Solinas G, Karin M (2010) JNK1 and IKKbeta: molecular links between obesity and metabolic dysfunction. *FASEB J* 24(8):2596–2611
 76. Hirosumi J, Tuncman G, Chang L, Gorgun CZ, Uysal KT, Maeda K, Karin M, Hotamisligil GS (2002) A central role for JNK in obesity and insulin resistance. *Nature* 420(6913):333–336
 77. Shoelson SE, Lee J, Goldfine AB (2006) Inflammation and insulin resistance. *J Clin Invest* 116(7):1793–1801
 78. Furukawa S, Fujita T, Shimabukuro M, Iwaki M, Yamada Y, Nakajima Y, Nakayama O, Makishima M, Matsuda M, Shimomura I (2004) Increased oxidative stress in obesity and its impact on metabolic syndrome. *J Clin Invest* 114(12):1752–1761
 79. Solinas G, Naugler W, Galimi F, Lee MS, Karin M (2006) Saturated fatty acids inhibit induction of insulin gene transcription by JNK-mediated phosphorylation of insulin-receptor substrates. *Proc Natl Acad Sci U S A* 103(44):16454–16459
 80. Lee JY, Sohn KH, Rhee SH, Hwang D (2001) Saturated fatty acids, but not unsaturated fatty acids, induce the expression of cyclooxygenase-2 mediated through Toll-like receptor 4. *J Biol Chem* 276(20):16683–16689
 81. Malhi H, Barreiro FJ, Isomoto H, Bronk SF, Gores GJ (2007) Free fatty acids sensitise hepatocytes to TRAIL mediated cytotoxicity. *Gut* 56(8):1124–1131
 82. Feldstein AE, Werneburg NW, Canbay A, Guicciardi ME, Bronk SF, Rydzewski R, Burgart LJ, Gores GJ (2004) Free fatty acids promote hepatic lipotoxicity by stimulating TNF-alpha expression via a lysosomal pathway. *Hepatology* 40(1):185–194
 83. Malhi H, Gores GJ (2008) Molecular mechanisms of lipotoxicity in nonalcoholic fatty liver disease. *Semin Liver Dis* 28(4):360–369
 84. Schattenberg JM, Singh R, Wang Y, Lefkowitz JH, Rigoli RM, Scherer PE, Czaja MJ (2006) JNK1 but not JNK2 promotes the development of steatohepatitis in mice. *Hepatology* 43(1):163–172
 85. Wang Y, Ausman LM, Greenberg AS, Russell RM, Wang XD (2009) Nonalcoholic steatohepatitis induced by a high-fat diet promotes diethylnitrosamine-initiated early hepatocarcinogenesis in rats. *Int J Cancer* 124(3):540–546
 86. Shimizu M, Sakai H, Shirakami Y, Iwasa J, Yasuda Y, Kubota M, Takai K, Tsurumi H, Tanaka T, Moriwaki H (2011) Acyclic retinoid inhibits diethylnitrosamine-induced liver tumorigenesis in obese and diabetic C57BLKS/J-(db)/+Lepr(db) mice. *Cancer Prev Res* 4(1):128–136
 87. He G, Yu GY, Temkin V, Ogata H, Kuntzen C, Sakurai T, Sieghart W, Peck-Radosavljevic M, Leffert HL, Karin M (2010) Hepatocyte IKKbeta/NF-kappaB inhibits tumor promotion and progression by preventing oxidative stress-driven STAT3 activation. *Cancer Cell* 17(3):286–297
 88. Giovannucci E, Michaud D (2007) The role of obesity and related metabolic disturbances in cancers of the colon, prostate, and pancreas. *Gastroenterology* 132(6):2208–2225
 89. Frezza EE, Wachtel MS, Chiriva-Internati M (2006) Influence of obesity on the risk of developing colon cancer. *Gut* 55(2):285–291
 90. Hirose Y, Hata K, Kuno T, Yoshida K, Sakata K, Yamada Y, Tanaka T, Reddy BS, Mori H (2004) Enhancement of development of azoxymethane-induced colonic premalignant lesions in C57BL/KsJ-db/db mice. *Carcinogenesis* 25(5):821–825
 91. Shimizu M, Shirakami Y, Iwasa J, Shiraki M, Yasuda Y, Hata K, Hirose Y, Tsurumi H, Tanaka T, Moriwaki H (2009) Supplementation with branched-chain amino acids inhibits azoxymethane-induced colonic preneoplastic lesions in male C57BL/KsJ-db/db mice. *Clin Cancer Res* 15(9):3068–3075
 92. Shimizu M, Shirakami Y, Sakai H, Adachi S, Hata K, Hirose Y, Tsurumi H, Tanaka T, Moriwaki H (2008) (-)-Epigallocatechin gallate suppresses azoxymethane-induced colonic premalignant lesions in male C57BL/KsJ-db/db mice. *Cancer Prev Res* 1(4):298–304
 93. Moriwaki H, Miwa Y, Tajika M, Kato M, Fukushima H, Shiraki M (2004) Branched-chain amino acids as a protein- and energy-source in liver cirrhosis. *Biochem Biophys Res Commun* 313(2):405–409
 94. Kawaguchi T, Izumi N, Charlton MR, Sata M (2011) Branched-chain amino acids as pharmacological nutrients in chronic liver disease. *Hepatology* 54(3):1063–1070
 95. Muto Y, Sato S, Watanabe A, Moriwaki H, Suzuki K, Kato A, Kato M, Nakamura T, Higuchi K, Nishiguchi S, Kumada H, Long-Term Survival Study Group (2005) Effects of oral branched-chain amino acid granules on event-free survival in patients with liver cirrhosis. *Clin Gastroenterol Hepatol* 3(7):705–713
 96. Marchesini G, Bianchi G, Merli M, Amodio P, Panella C, Loguercio C, Rossi Fanelli F, Abbiati R (2003) Nutritional supplementation with branched-chain amino acids in advanced cirrhosis: a double-blind, randomized trial. *Gastroenterology* 124(7):1792–1801
 97. Kao YH, Chang HH, Lee MJ, Chen CL (2006) Tea, obesity, and diabetes. *Mol Nutr Food Res* 50(2):188–210
 98. Hursel R, Viechtbauer W, Westerterp-Plantenga MS (2009) The effects of green tea on weight loss and weight maintenance: a meta-analysis. *Int J Obes* 33(9):956–961
 99. Yang CS, Wang X, Lu G, Picinich SC (2009) Cancer prevention by tea: animal studies, molecular mechanisms and human relevance. *Nat Rev Cancer* 9(6):429–439
 100. Shimizu M, Adachi S, Masuda M, Kozawa O, Moriwaki H (2011) Cancer chemoprevention with green tea catechins by targeting receptor tyrosine kinases. *Mol Nutr Food Res* 55(6):832–843
 101. Muto Y, Moriwaki H, Ninomiya M, Adachi S, Saito A, Takasaki KT, Tanaka T, Tsurumi K, Okuno M, Tomita E, Nakamura T,

- Kojima T (1996) Prevention of second primary tumors by an acyclic retinoid, polyphenolic acid, in patients with hepatocellular carcinoma. Hepatoma Prevention Study Group. *N Engl J Med* 334(24):1561–1567
102. Muto Y, Moriwaki H, Saito A (1999) Prevention of second primary tumors by an acyclic retinoid in patients with hepatocellular carcinoma. *N Engl J Med* 340(13):1046–1047
 103. Shimizu M, Takai K, Moriwaki H (2009) Strategy and mechanism for the prevention of hepatocellular carcinoma: phosphorylated retinoid X receptor alpha is a critical target for hepatocellular carcinoma chemoprevention. *Cancer Sci* 100(3):369–374
 104. Shimizu M, Sakai H, Moriwaki H (2011) Chemoprevention of hepatocellular carcinoma by acyclic retinoid. *Front Biosci* 16:759–769
 105. Petrides AS, Vogt C, Schulze-Berge D, Matthews D, Strohmeyer G (1994) Pathogenesis of glucose intolerance and diabetes mellitus in cirrhosis. *Hepatology* 19(3):616–627
 106. Mehta SH, Brancati FL, Sulkowski MS, Strathdee SA, Szklo M, Thomas DL (2000) Prevalence of type 2 diabetes mellitus among persons with hepatitis C virus infection in the United States. *Ann Intern Med* 133(8):592–599
 107. She P, Reid TM, Bronson SK, Vary TC, Hajnal A, Lynch CJ, Hutson SM (2007) Disruption of BCATm in mice leads to increased energy expenditure associated with the activation of a futile protein turnover cycle. *Cell Metab* 6(3):181–194
 108. Higuchi N, Kato M, Miyazaki M, Tanaka M, Kohjima M, Ito T, Nakamura M, Enjoji M, Kotoh K, Takayanagi R (2011) Potential role of branched-chain amino acids in glucose metabolism through the accelerated induction of the glucose-sensing apparatus in the liver. *J Cell Biochem* 112(1):30–38
 109. Kawaguchi T, Nagao Y, Matsuoka H, Ide T, Sata M (2008) Branched-chain amino acid-enriched supplementation improves insulin resistance in patients with chronic liver disease. *Int J Mol Med* 22(1):105–112
 110. Urata Y, Okita K, Korenaga K, Uchida K, Yamasaki T, Sakaida I (2007) The effect of supplementation with branched-chain amino acids in patients with liver cirrhosis. *Hepatol Res* 37(7):510–516
 111. Hagiwara A, Nishiyama M, Ishizaki S (2012) Branched-chain amino acids prevent insulin-induced hepatic tumor cell proliferation by inducing apoptosis through mTORC1 and mTORC2-dependent mechanisms. *J Cell Physiol* 227(5):2097–2105
 112. Yoshiji H, Noguchi R, Kitade M, Kaji K, Ikenaka Y, Namisaki T, Yoshii J, Yanase K, Yamazaki M, Tsujimoto T, Akahane T, Kawarata H, Uemura M, Fukui H (2009) Branched-chain amino acids suppress insulin-resistance-based hepatocarcinogenesis in obese diabetic rats. *J Gastroenterol* 44(5):483–491
 113. Qin B, Polansky MM, Harry D, Anderson RA (2010) Green tea polyphenols improve cardiac muscle mRNA and protein levels of signal pathways related to insulin and lipid metabolism and inflammation in insulin-resistant rats. *Mol Nutr Food Res* 54 (Suppl 1):S14–S23
 114. Bose M, Lambert JD, Ju J, Reuhl KR, Shapses SA, Yang CS (2008) The major green tea polyphenol, (–)-epigallocatechin-3-gallate, inhibits obesity, metabolic syndrome, and fatty liver disease in high-fat-fed mice. *J Nutr* 138(9):1677–1683
 115. Ramadan G, El-Beih NM, Abd El-Ghffar E (2009) Modulatory effects of black v. green tea aqueous extract on hyperglycaemia, hyperlipidaemia and liver dysfunction in diabetic and obese rat models. *Br J Nutr* 102(11):1611–1619
 116. Shirakami Y, Shimizu M, Tsurumi H, Hara Y, Tanaka T, Moriwaki H (2008) EGCG and Polyphenon E attenuate inflammation-related mouse colon carcinogenesis induced by AOM plus DDS. *Mol Med Report* 1(3):355–361
 117. Shimizu M, Deguchi A, Hara Y, Moriwaki H, Weinstein IB (2005) EGCG inhibits activation of the insulin-like growth factor-1 receptor in human colon cancer cells. *Biochem Biophys Res Commun* 334(3):947–953
 118. Yasuda Y, Shimizu M, Sakai H, Iwasa J, Kubota M, Adachi S, Osawa Y, Tsurumi H, Hara Y, Moriwaki H (2009) (–)-Epigallocatechin gallate prevents carbon tetrachloride-induced rat hepatic fibrosis by inhibiting the expression of the PDGFRbeta and IGF-1R. *Chem Biol Interact* 182(2–3):159–164
 119. Shimizu M, Fukutomi Y, Ninomiya M, Nagura K, Kato T, Araki H, Suganuma M, Fujiki H, Moriwaki H (2008) Green tea extracts for the prevention of metachronous colorectal adenomas: a pilot study. *Cancer Epidemiol Biomarkers Prev* 17(11):3020–3025
 120. Bettuzzi S, Brausi M, Rizzi F, Castagnetti G, Peracchia G, Corti A (2006) Chemoprevention of human prostate cancer by oral administration of green tea catechins in volunteers with high-grade prostate intraepithelial neoplasia: a preliminary report from a one-year proof-of-principle study. *Cancer Res* 66(2):1234–1240
 121. Li N, Sun Z, Han C, Chen J (1999) The chemopreventive effects of tea on human oral precancerous mucosa lesions. *Proc Soc Exp Biol Med* 220(4):218–224
 122. Germain P, Chambon P, Eichele G, Evans RM, Lazar MA, Leid M, De Lera AR, Lotan R, Mangelsdorf DJ, Gronemeyer H (2006) International Union of Pharmacology. LXIII. Retinoid X receptors. *Pharmacol Rev* 58(4):760–772
 123. Germain P, Chambon P, Eichele G, Evans RM, Lazar MA, Leid M, De Lera AR, Lotan R, Mangelsdorf DJ, Gronemeyer H (2006) International Union of Pharmacology. LX. Retinoic acid receptors. *Pharmacol Rev* 58(4):712–725
 124. Altucci L, Gronemeyer H (2001) The promise of retinoids to fight against cancer. *Nat Rev Cancer* 1(3):181–193
 125. Altucci L, Leibowitz MD, Ogilvie KM, de Lera AR, Gronemeyer H (2007) RAR and RXR modulation in cancer and metabolic disease. *Nat Rev Drug Discov* 6(10):793–810
 126. Ando N, Shimizu M, Okuno M, Matsushima-Nishiwaki R, Tsurumi H, Tanaka T, Moriwaki H (2007) Expression of retinoid X receptor alpha is decreased in 3'-methyl-4-dimethylaminoazobenzene-induced hepatocellular carcinoma in rats. *Oncol Rep* 18(4):879–884
 127. Matsushima-Nishiwaki R, Okuno M, Adachi S, Sano T, Akita K, Moriwaki H, Friedman SL, Kojima S (2001) Phosphorylation of retinoid X receptor alpha at serine 260 impairs its metabolism and function in human hepatocellular carcinoma. *Cancer Res* 61(20):7675–7682
 128. Matsushima-Nishiwaki R, Okuno M, Takano Y, Kojima S, Friedman SL, Moriwaki H (2003) Molecular mechanism for growth suppression of human hepatocellular carcinoma cells by acyclic retinoid. *Carcinogenesis* 24(8):1353–1359
 129. Adachi S, Okuno M, Matsushima-Nishiwaki R, Takano Y, Kojima S, Friedman SL, Moriwaki H, Okano Y (2002) Phosphorylation of retinoid X receptor suppresses its ubiquitination in human hepatocellular carcinoma. *Hepatology* 35(2):332–340
 130. Yoshimura K, Muto Y, Shimizu M, Matsushima-Nishiwaki R, Okuno M, Takano Y, Tsurumi H, Kojima S, Okano Y, Moriwaki H (2007) Phosphorylated retinoid X receptor alpha loses its heterodimeric activity with retinoic acid receptor beta. *Cancer Sci* 98(12):1868–1874
 131. Kanamori T, Shimizu M, Okuno M, Matsushima-Nishiwaki R, Tsurumi H, Kojima S, Moriwaki H (2007) Synergistic growth inhibition by acyclic retinoid and vitamin K2 in human hepatocellular carcinoma cells. *Cancer Sci* 98(3):431–437
 132. Okita K, Matsui O, Kumada H, Tanaka K, Kaneko S, Moriwaki H, Izumi N, Okusaka T, Ohashi Y, Makuuchi M (2010) Effect of peretinoin on recurrence of hepatocellular carcinoma (HCC): results of a phase II/III randomized placebo-controlled trial. *J Clin Oncol* 28(Suppl 7s):4024

Special Report

Nutritional status and quality of life in current patients with liver cirrhosis as assessed in 2007–2011

Makoto Shiraki,¹ Shuhei Nishiguchi,² Masaki Saito,² Yoshitaka Fukuzawa,³ Toshihiko Mizuta,⁴ Masaki Kaibori,⁵ Tatsunori Hanai,¹ Kayoko Nishimura,⁶ Masahito Shimizu,¹ Hisashi Tsurumi¹ and Hisataka Moriwaki¹

¹The First Department of Internal Medicine, Gifu University School of Medicine, Gifu, ²Division of Hepatobiliary and Pancreatic Disease, Department of Internal Medicine, Hyogo College of Medicine, Hyogo, ³Medical Education Center, Aichi Medical University School of Medicine, Aichi, ⁴Department of Internal Medicine, Saga Medical School, Saga University, Saga, ⁵Department of Surgery, Kansai Medical University, Osaka, and ⁶Center for Nutrition Support and Infection Control, Gifu University Hospital, Gifu, Japan

Aim: Current guidelines recommended adequate nutritional support for patients with liver cirrhosis to improve clinical outcome and quality of life (QOL). However, these evidences were obtained more than 10 years ago when malnutrition prevailed. In recent years, the impact of obesity on liver damage and carcinogenesis has grown. We attempted to elucidate the nutritional state and QOL in present cirrhotics.

Methods: A research group supported by the Ministry of Health, Labor and Welfare of Japan recruited 294 cirrhotics between 2007 and 2011. Subjects comprised 171 males and 123 females, 158 of whom had hepatocellular carcinoma (HCC) and Child–Pugh grades A : B : C were 154:91:49. Anthropometry, blood biochemistry and indirect calorimetry were conducted, and QOL was measured using Short Form-8.

Results: The mean body mass index (BMI) of all patients was 23.1 ± 3.4 kg/m², and 31% showed obesity (BMI ≥ 25.0). In subjects without ascites, edema or HCC, mean BMI was

23.6 ± 3.6 , and 34% had obesity. Protein malnutrition defined as serum albumin of less than 3.5 g/dL and energy malnutrition as respiratory quotient of less than 0.85 appeared in 61% and 43%, respectively, and protein-energy malnutrition (PEM) in 27% of all subjects. Among subjects without HCC, each proportion was 67%, 48% and 30%, respectively. QOL was significantly lower on all subscales than Japanese national standard values, but was similar regardless the presence or absence of HCC.

Conclusion: While PEM is still present in liver cirrhosis, an equal proportion has obesity in recent patients. Thus, in addition to guidelines for PEM, establishment of nutrition and exercise guidelines seems essential for obese patients with liver cirrhosis.

Key words: body mass index, energy malnutrition, liver cirrhosis, protein malnutrition, quality of life

INTRODUCTION

BECAUSE THE LIVER plays the central role in nutrient and fuel metabolism, protein-energy malnutrition (PEM) is common in patients with liver cirrhosis.^{1,2} Moreover, such malnutrition leads to poor prognosis and decline in the quality of life (QOL) of cirrhotics.^{3,4}

Branched-chain amino acid (BCAA) administration for protein malnutrition raises the serum albumin level

and improves the QOL and survival of patients with liver cirrhosis.^{5–8} Treatment with late-evening snack (LES) for energy malnutrition improves respiratory quotient (RQ), liver dysfunction and QOL.^{9,10}

Therefore, the guidelines for the treatment of liver cirrhosis by Japanese Society of Gastroenterology,¹¹ American Society for Parenteral and Enteral Nutrition¹² and European Society for Clinical Nutrition and Metabolism¹³ recommend such nutritional therapy.

However, these evidences were obtained in the cirrhotic patients recruited from 1995–2000, where protein or energy malnutrition prevailed in 50–87%.^{1–4} In contrast, in the next 10 years, obesity rate in the cirrhotic patients rose to approximately 30%.¹⁴ More recently, non-alcoholic steatohepatitis (NASH) or the

Correspondence: Dr Makoto Shiraki, The First Department of Internal Medicine, Gifu University School of Medicine, 1-1 Yanagido, Gifu 501-1194, Japan. Email: mshiraki-gif@umin.ac.jp
Received 11 August 2012; revision 7 October 2012; accepted 15 October 2012.

hepatic inflammation, fibrosis and carcinogenesis due to obesity became the topics.^{14–16}

Therefore, it is essential to re-evaluate a nourishment state of the current cirrhotic patients to update the guidelines. In this report, we investigated comprehensive data on the nourishment state and QOL in a large group of patients with liver cirrhosis recruited in the years 2007–2011.

METHODS

Patients

TWO HUNDRED AND ninety-four patients with liver cirrhosis (171 men and 123 women; mean age, 68 ± 10 years) undergoing treatment between 2007 and 2011 were recruited by a Research Group (Gifu University, Hyogo College of Medicine, Aichi Medical University and Saga University) supported by the Ministry of Health, Labor and Welfare of Japan. Liver cirrhosis was diagnosed by clinical and laboratory profiles and by histological examination of liver biopsy specimens. The etiology of cirrhosis was hepatitis B virus in 35 patients, hepatitis C virus in 204, alcohol in 25, NASH in six and others in 24. Child–Pugh classification of the disease severity¹⁷ was A in 154 cases, B in 91 cases and C in 49 cases. One hundred and fifty-eight patients had hepatocellular carcinoma (HCC), and their clinical stage was I

in 41 patients, II in 41, III in 54 and IV in 22. Clinical profiles of the patients are presented in Table 1. The proportion of patients supplemented with BCAA or LES rose in parallel with the increasing grade of Child–Pugh classification. Patients with fever, HIV infection, overt infectious disease (septicemia, pneumonia, urinary tract infection), renal insufficiency or under immunomodulatory therapy were excluded. The study protocol was approved by the Medical Ethics Committee of Gifu University Graduate School of Medicine, and informed consent was obtained from all patients. The study protocol was in agreement with the 1975 Declaration of Helsinki as revised in 1983.

Hematological examinations

Blood was drawn for routine laboratory examinations in the early morning after overnight fasting on the day of metabolic studies. Serum albumin, total bilirubin, alt alanine aminotransferase, prothrombin activity and urinary nitrogen (UN) were measured with a standard clinical analyzer at the central laboratory in each hospital.

Nutritional assessment

Metabolic studies were carried out using an indirect calorimeter (Aeromonitor AE-300S; Minato Medical Science, Osaka, Japan) to estimate non-protein re-

Table 1 Clinical and biochemical profiles of patients with liver cirrhosis

| | Cirrhosis (n = 294) | Child A (n = 154) | Child B (n = 91) | Child C (n = 49) | P |
|--------------------------------------|------------------------|----------------------|---------------------|---------------------|-------|
| Age (years) | 68 ± 10 | 68 ± 10 | 68 ± 10 | 68 ± 12 | n.s. |
| Sex (male/female) | 171/123 | 90/64 | 51/40 | 30/19 | n.s. |
| Height (cm) | 159 ± 9.1 | 159 ± 9.0 | 159 ± 9.1 | 159 ± 9.7 | n.s. |
| Weight (kg) | 59 ± 11 | 58 ± 9.6 | 59 ± 11 | 60 ± 13 | n.s. |
| Body mass index (kg/m ²) | 23.1 ± 3.4 | 22.9 ± 3.0 | 23.4 ± 3.6 | 23.6 ± 4.0 | n.s. |
| Etiology (HBV/HCV/alcohol/others) | 35/204/25/30 | 20/108/11/15 | 11/62/8/10 | 4/34/6/5 | n.s. |
| Hepatocellular carcinoma (+/-)* | 158/136 | 84/69 | 54/38 | 20/29 | n.s. |
| Number of patients | | | | | |
| Treated with BCAA | 97 | 35 | 45 | 17 | <0.01 |
| Supplied with LES | 36 | 8 | 19 | 9 | <0.01 |
| Albumin (g/dL) | 3.3 ± 0.6 | 3.6 ± 0.5 | 3.0 ± 0.4 | 2.6 ± 0.4 | <0.01 |
| Total bilirubin (mg/dL) | 1.4 ± 1.8 | 0.9 ± 0.4 | 1.5 ± 1.2 | 3.2 ± 3.8 | <0.01 |
| Alanine aminotransferase (IU/L) | 44 ± 31 | 43 ± 30 | 44 ± 29 | 45 ± 40 | n.s. |
| Prothrombin time (%) | 81 ± 30 | 91 ± 32 | 75 ± 23 | 66 ± 22 | <0.01 |

HBV, hepatitis B virus; HCV, hepatitis C virus; BCAA, branched-chain amino acids; LES, late-evening snack; n.s., not significant.

Data are presented as number of patients or mean ± standard deviation.

Statistical analysis was performed by one-way ANOVA or contingency table analysis for distribution among Child–Pugh grades A, B and C.

*Clinical stage of hepatocellular carcinoma was I in 41 patients, II in 41, III in 54 and IV in 22.

spiratory quotient (npRQ) from measured oxygen consumption/min (VO_2), carbon dioxide production/min (VCO_2) and total urinary nitrogen using the following equation:^{18–20}

$$\text{npRQ} = (1.44\text{VCO}_2 - 4.890\text{UN}) / (1.44\text{VO}_2 - 6.04\text{UN}).$$

Measurements were performed between 07.00 and 09.00 hours while the patients were still lying in bed. The last meal was served at 18.00 hours on the previous day.

We measured height and bodyweight, and calculated body mass index (BMI).

QOL questionnaire

Health-related QOL was measured using the Short Form-8 (SF-8) questionnaire.^{21–23} The SF-8 contains eight questions that provide a quantitative evaluation on each of eight subscales: (i) physical functioning (PF); (ii) role physical (RP); (iii) bodily pain (BP); (iv) general health perception (GH); (v) vitality (VT); (vi) social functioning (SF); (vii) role emotional (RE); and (viii) mental health (MH).

Statistical analysis

Data were expressed as the mean and standard deviation. Comparisons of measured values among Child–Pugh classification grade A, B and C were performed using one-way ANOVA. Comparisons of sex, etiology and the presence of HCC among Child–Pugh classification grades were performed using contingency table analysis. Measured QOL was analyzed by z-test or Student's *t*-test between each group. Data analysis was performed using JMP ver. 5.1J (SAS Institute Japan, Tokyo, Japan) and $P < 0.05$ was considered statistically significant.

RESULTS

BMI of the patients with liver cirrhosis

THE MEAN BMI of all patients with liver cirrhosis was $23.1 \pm 3.4 \text{ kg/m}^2$.

The ratio of obese subjects with BMI of 25 or higher was 30.6% and that of less than 18.5 kg/m^2 was 5.1%, respectively (Fig. 1).

We then excluded patients with ascites, edema or HCC to match the present cohort with those reported in 2002.⁴ The number of patents in this cohort was 95, and Child–Pugh grades A, B and C were 71:22:2, respectively. Mean BMI was $23.6 \pm 3.6 \text{ kg/m}^2$, and BMI of




Figure 1 Distribution of body mass index (BMI) in patients with liver cirrhosis. Total number of patients = 294. Obese subjects ($\text{BMI} \geq 25$) were present in 30.6%, lean ones ($18.5 \leq \text{BMI} < 25$) were in 64.3% and emaciation ($\text{BMI} < 18.5$) was observed in 5.1%.

less than 18.5 kg/m^2 and 25.0 kg/m^2 or higher were observed in 9.2% and 33.7%, respectively (Fig. 2).

Incidence of protein malnutrition, energy malnutrition and PEM in patients with liver cirrhosis

We examined nutritional status in 181 patients with liver cirrhosis that underwent indirect calorimetry. In these patients, the male : female ratio was 112:69, HCC




Figure 2 Distribution of body mass index (BMI) in cirrhotic patients without ascites, edema or hepatocellular carcinoma. Total number of patients = 95. Obese subjects ($\text{BMI} \geq 25$) were present in 33.7%, lean ones ($18.5 \leq \text{BMI} < 25$) were in 57.1% and emaciation ($\text{BMI} < 18.5$) was observed in 9.2%.

Table 2 Incidence of protein and energy malnutrition in patients with liver cirrhosis

| Energy nutritional state | Protein nutritional state | |
|--------------------------|---------------------------|------------------|
| | Normal (%) | Malnourished (%) |
| Normal (%) | 42 (23%) | 62 (34%) |
| Malnourished (%) | 28 (16%) | 49 (27%) |

Protein malnutrition was defined as serum albumin level of <3.5 g/dL and energy malnutrition as a respiratory quotient of <0.85.

Total number of patients = 181.

Data are presented as number of patients (%).

was present in 94, and Child–Pugh grades A : B : C were 90:58:33. When protein malnutrition was defined as serum albumin level of less than 3.5 g/dL and energy malnutrition as a non-protein respiratory quotient of less than 0.85, protein malnutrition was found in 61%, energy malnutrition in 43% and PEM in 27% (Table 2). Similarly, among 87 patients without HCC (Child–

Table 3 Incidence of protein and energy malnutrition in cirrhotic patients without hepatocellular carcinoma

| Energy nutritional state | Protein nutritional state | |
|--------------------------|---------------------------|------------------|
| | Normal (%) | Malnourished (%) |
| Normal (%) | 13 (15%) | 32 (37%) |
| Malnourished (%) | 16 (18%) | 26 (30%) |

Protein malnutrition was defined as serum albumin level of <3.5 g/dL and energy malnutrition as a respiratory quotient of <0.85.

Total number of patients = 87.

Data are presented as number of patients (%).

Table 4 Comparison of health-related quality of life between the Japanese national standard and the patients with liver cirrhosis

| Subscales | Japanese national standard | Patients with liver cirrhosis | P |
|---------------------------|----------------------------|-------------------------------|-------|
| Physical functioning | 50.1 ± 5.0 | 43.8 ± 5.2 | <0.01 |
| Role physical | 50.2 ± 5.3 | 41.6 ± 6.6 | <0.01 |
| Bodily pain | 51.3 ± 8.3 | 48.3 ± 5.3 | <0.01 |
| General health perception | 50.6 ± 6.6 | 45.2 ± 4.4 | <0.01 |
| Vitality | 52.4 ± 5.5 | 47.5 ± 4.3 | <0.01 |
| Social functioning | 50.2 ± 6.6 | 46.1 ± 5.3 | <0.01 |
| Role emotional | 51.3 ± 4.5 | 45.6 ± 5.7 | <0.01 |
| Mental health | 53.3 ± 5.4 | 47.6 ± 4.0 | <0.01 |

Data are presented as mean ± standard deviation.

Statistical analysis was performed by Student's *t*-test between the Japanese national standard²⁴ and the patients with liver cirrhosis.

Table 5 Comparison of health-related quality of life in cirrhotics by the presence or absence of hepatocellular carcinoma

| Subscales | Absence of hepatocellular carcinoma | Presence of hepatocellular carcinoma | P |
|---------------------------|-------------------------------------|--------------------------------------|------|
| Physical functioning | 43.4 ± 4.9 | 44.2 ± 5.5 | n.s. |
| Role physical | 41.1 ± 6.3 | 42.1 ± 6.8 | n.s. |
| Bodily pain | 47.8 ± 5.3 | 48.7 ± 5.1 | n.s. |
| General health perception | 44.9 ± 4.5 | 45.4 ± 3.9 | n.s. |
| Vitality | 46.5 ± 4.3 | 48.4 ± 4.2 | n.s. |
| Social functioning | 45.3 ± 5.0 | 46.8 ± 5.4 | n.s. |
| Role emotional | 45.3 ± 5.0 | 45.8 ± 6.1 | n.s. |
| Mental health | 46.6 ± 3.9 | 48.5 ± 4.0 | n.s. |

n.s., not significant.

Data are presented as mean ± standard deviation.

Statistical analysis was performed by *z*-test between the presence and absence of hepatocellular carcinoma.

Pugh grades A : B : C, 36:27:24), 67% had protein malnutrition, 48% had energy malnutrition and 30% had PEM (Table 3).

Health-related QOL of the patients with liver cirrhosis

We examined health-related QOL in 114 patients with liver cirrhosis (64 men and 50 women) using the SF-8. Sixty-two patients had HCC, and Child–Pugh grades A : B : C were 63:26:25.

Quality of life of all subjects was significantly lower on all subscales than Japanese national standard values (Table 4),²⁴ but no difference was observed between the presence and the absence of HCC (Table 5).

DISCUSSION

PROTEIN-ENERGY MALNUTRITION is a common manifestation in cirrhotic patients with reported incidences as high as 50–87%.^{1,2} Protein nutrition is usually evaluated by serum albumin level and, for energy nutrition, indirect calorimetry is recommended for precise analysis.¹³ Energy malnutrition typically shows reduced carbohydrate oxidation, increased fat oxidation and decline in nprQ measured by indirect calorimetry. It is reported that PEM worsens prognosis and QOL in patients with liver cirrhosis.^{3,4} Thus, intervention for PEM is an important issue in the clinical management of liver cirrhosis.

For this purpose, BCAA administration for protein malnutrition raises the serum albumin level and improves QOL and survival of patients with liver cirrhosis.^{5–8} LES for energy malnutrition improves nprQ, liver dysfunction and QOL.^{9,10} Thus, many guidelines^{11–13} recommend such nutritional therapy for liver cirrhosis.

However, these evidences were obtained in the cirrhotic patients recruited from 1995 through 2000 where malnutrition prevailed but obesity was apparently less (20%)⁴ than the general cohort (30%).²⁵ In the next 10 years, obesity rose by approximately 1.5 times in the patients with chronic liver disease in Japan.¹⁴ In addition, presence of diabetes mellitus, hyperinsulinemia or obesity is currently regarded as a significant risk factor for liver carcinogenesis.^{14–16} Furthermore, the relationship between obesity and liver inflammation and fibrosis, including NASH has become an important issue in recent years. Therefore, it is necessary to elucidate the nourishment state of the present cirrhotic patients to update guidelines. Thus, we report in this paper a comprehensive survey of the nourishment state and QOL in the present patients with liver cirrhosis.

The etiology of the 294 cirrhotics was hepatitis B virus in 11.9%, hepatitis C virus in 69.4%, alcohol in 8.5%, NASH in 2.0% and others in 8.2% in this study. In the 44th Annual Meeting of Japan Society of Hepatology in 2008 (Matsuyama), the reported etiology of 33 379 cirrhotics was hepatitis B virus in 13.9%, hepatitis C virus in 60.9%, alcohol in 13.6%, NASH in 2.1% and others in 9.5%,²⁶ indicating similar patient composition between two studies.

Obesity is defined by BMI of 25 or higher in Japan but by 30 or higher by World Health Organization. In this study, the mean BMI excluding patients with ascites, edema or HCC was 23.6 ± 3.6 kg/m² and the ratio of obese subjects with BMI of 25 or higher was 33.7% of

these patients (Fig. 2). The proportion of obese people in the general population of Japan at matched age was 30.5% in 2009.²⁵ Thus, an equal or greater proportion of patients with liver cirrhosis has obesity than the general population of Japan at present.

The increase in obesity, or excess energy nutrition status, and subsequent impaired glucose metabolism potentially bring about an unfavorable outcome in cirrhotic patients. Actually, excess energy nutrition contributed to induce carcinogenesis in liver cirrhosis,^{15,27,28} and the number of obese subjects doubled in the candidates for liver transplantation in the previous 10 years in the USA.^{29–31}

As to PEM exactly defined by serum albumin and nprQ, Tajika *et al.* reported that protein malnutrition was identified in 75%, energy malnutrition in 62% and PEM in 50% of 109 patients with liver cirrhosis in 1995.⁴ In our study, 87 patients without HCC composed a group to show comparable backgrounds to those by Tajika *et al.*⁴ Among them, 67% had protein malnutrition, 48% had energy malnutrition and 30% had PEM (Table 3). Taken together, the protein malnutrition remains almost similar in liver cirrhosis, but the patients with energy malnutrition, particularly PEM, substantially decreased.

The above-mentioned results urge that two concerns are addressed. The first is the effect of altered nutritional state of cirrhotics on their QOL, and the second is a question if exercise should be prescribed for obese cirrhotics. Regarding QOL, reduction in bodyweight achieved by chronic liver disease patients with obesity was associated with improved liver dysfunction, histology or QOL.^{32,33}

In this study, basal QOL was estimated by the SF-8, and was significantly lower on all subscales than Japanese national standard values. However, no difference was observed by the presence or absence of HCC. In contrast, QOL of cirrhotic patients significantly correlated with the grade of disease severity as defined by the Child–Pugh classification (data not shown). It was thus suggested that the degree of the hepatic functional reserve contributed to a greater extent than the progression of cancer as for QOL of cirrhotic patients.

In conclusion, while PEM is still present in liver cirrhosis, a greater proportion shows obesity in Japanese patients at present. Because exacerbated inflammation, fibrosis and carcinogenesis has been reported in obese patients with liver cirrhosis, the present findings urge revision of nutritional and, possibly, establishment of exercise guidelines for obese patients with liver cirrhosis, in addition to the current PEM guidelines.

ACKNOWLEDGMENT

THIS WORK WAS supported in part by Grants-in-Aid from the Ministry of Health, Labor, and Welfare of Japan.

REFERENCES

- Morgan AG, McAdam WA, Walmsley GR, Horrocks JC, De Dombal FT. Nutrition in cryptogenic cirrhosis and chronic aggressive hepatitis. *Gut* 1976; 17 (2): 113–8.
- Müller MJ. Malnutrition in cirrhosis. *J Hepatol* 1995; 23 (Suppl 1): 31–5.
- Alberino F, Gatta A, Amodio P *et al.* Nutrition and survival in patients with liver cirrhosis. *Nutrition* 2001; 17: 445–50.
- Tajika M, Kato M, Mohri H *et al.* Prognostic value of energy metabolism in patients with viral liver cirrhosis. *Nutrition* 2002; 18 (3): 229–34.
- Yoshida T, Muto Y, Moriwaki H, Yamato M. Effect of long-term oral supplementation with branched-chain amino acid granules on the prognosis of liver cirrhosis. *J Gastroenterol* 1989; 24: 692–8.
- Marchesini G, Bianci G, Merli M *et al.* Nutritional supplementation with branched-chain amino acids in advanced cirrhosis: a double-blind, randomized trial. *Gastroenterology* 2003; 124: 1792–801.
- Muto Y, Sato S, Watanabe A *et al.* Effects of oral branched-chain amino acid granules on event-free survival in patients with liver cirrhosis. *Clin Gastroenterol Hepatol* 2005; 3 (7): 705–13.
- Fukushima H, Miwa Y, Shiraki M *et al.* Oral branched-chain amino acid supplementation improves the oxidized/reduced albumin ratio in patients with liver cirrhosis. *Hepatol Res* 2007; 37 (9): 765–70.
- Miwa Y, Shiraki M, Kato M *et al.* Improvement of fuel metabolism by nocturnal energy supplementation in patients with liver cirrhosis. *Hepatol Res* 2000; 18 (3): 184–89.
- Nakaya Y, Okita K, Suzuki K *et al.* BCAA-enriched snack improves nutritional state of cirrhosis. *Nutrition* 2007; 23: 113–20.
- The Japanese Society of Gastroenterology. *Guidelines for the Diagnosis and Treatment of Liver Cirrhosis*. Tokyo: Nankodo, 2010. (in Japanese).
- ASPEN Board of Directors and the Clinical Guidelines Task Force. Guidelines for the use of parenteral and enteral nutrition in adult and pediatric patients. *JPEN J Parenter Enteral Nutr* 2002; 26 (Suppl 1): 73–6.
- ESPEN. Guidelines on enteral nutrition: liver disease. *Clin Nutr* 2006; 25: 285–94.
- Moriwaki H, Shiraki M, Fukushima H *et al.* Long-term outcome of branched-chain amino acid treatment in patients with liver cirrhosis. *Hepatol Res* 2008; 38 (s1): S102–6.
- Muto Y, Sato S, Watanabe A *et al.* Over weight and obesity increase the risk for HCC in patients with liver cirrhosis and long-term oral supplementation with branched-chain amino acid granules inhibits liver carcinogenesis in heavier patients with liver cirrhosis. *Hepatol Res* 2006; 35: 204–14.
- Hickman IJ, Powell EE, Prins JB *et al.* In overweight patients with chronic hepatitis C, circulating insulin is associated with hepatic fibrosis: implications for therapy. *J Hepatol* 2003; 39 (6): 1042–8.
- Pugh RN, Murray-Lyon IM, Dawson JL, Pietroni MC, Williams R. Transection of the oesophagus for bleeding oesophageal varices. *Br J Surg* 1973; 60 (8): 646–9.
- Kato M, Miwa Y, Tajika M, Hiraoka T, Muto Y, Moriwaki H. Preferential use of branched-chain amino acids as an energy substrate in patients with liver cirrhosis. *Intern Med* 1998; 37 (5): 429–34.
- Müller MJ, Böttcher J, Selberg O *et al.* Hypermetabolism in clinically stable patients with liver cirrhosis. *Am J Clin Nutr* 1999; 69 (6): 1194–201.
- Süttmann U, Ockenga J, Hoogestraat L *et al.* Resting energy expenditure and weight loss in human immunodeficiency virus-infected patients. *Metabolism* 1993; 42 (9): 1173–9.
- Turner-Bowker DM, Bayliss MS, Ware JE, Jr Kosinski M. Usefulness of the SF-8 Health Survey for comparing the impact of migraine and other conditions. *Qual Life Res* 2003; 12: 1003–12.
- Ware JE, Kosinski M, Dewey JE, Gandek B. *How to Score and Interpret Single-Item Healthstatus Measures. A Manual for Users of the SF-8TM Health Survey*. Lincoln, RI: QualityMetric Inc., 2001.
- Kuroda H, Ushio A, Miyamoto Y *et al.* Effects of branched-chain amino acid-enriched nutrient for patients with hepatocellular carcinoma following radiofrequency ablation: a one-year prospective trial. *J Gastroenterol Hepatol* 2010; 25 (9): 1550–5.
- Fukuhara S, Suzukamo Y. *Manual of the SF-8 Japanese Version*. Kyoto: Institute for Health Outcome and Process Evaluation Research, 2004. (in Japanese).
- Health and Welfare Statistics Association. *Journal of Health and Welfare Statistics 2010/2011*. Tokyo: Kosaido, 2011. (in Japanese).
- Michitaka K, Nishiguchi S, Aoyagi Y *et al.* Etiology of liver cirrhosis in Japan: a nationwide survey. *J Gastroenterol* 2010; 45 (1): 86–94.
- Yoshiji H, Noguchi R, Kitade M *et al.* Branched-chain amino acids suppress insulin-resistance-based hepatocarcinogenesis in obese diabetic rats. *J Gastroenterol* 2009; 44 (5): 483–91.
- Iwasa J, Shimizu M, Shiraki M *et al.* Dietary supplementation with branched-chain amino acids suppresses diethylnitrosamine-induced liver tumorigenesis in obese and diabetic C57BL/KsJ-db/db mice. *Cancer Sci* 2010; 101 (2): 460–7.

- 29 Nair S, Mason A, Eason J *et al.* Is obesity an independent risk factor for hepatocellular carcinoma in cirrhosis? *Hepatology* 2002; 36 (1): 150–5.
- 30 Pelletier SJ, Schaubel DE, Wei G *et al.* Effect of body mass index on the survival benefit of liver transplantation. *Liver Transpl* 2007; 13 (12): 1678–83.
- 31 Thuluvath PJ. Morbid obesity and gross malnutrition are both poor predictors of outcomes after liver transplantation: what can we do about it? *Liver Transpl* 2009; 15 (8): 838–41.
- 32 Oza N, Eguchi Y, Mizuta T *et al.* A pilot trial of body weight reduction for nonalcoholic fatty liver disease with a home-based lifestyle modification intervention delivered in collaboration with interdisciplinary medical staff. *J Gastroenterol* 2009; 44 (12): 1203–8.
- 33 Hickman IJ, Jonsson JR, Prins JB *et al.* Modest weight loss and physical activity in overweight patients with chronic liver disease results in sustained improvements in alanine aminotransferase, fasting insulin, and quality of life. *Gut* 2004; 53 (3): 413–9.