

Fig. 3. Expression of proinflammatory cytokines in response to TLR ligand stimulation was decreased in human monocytes treated with tunicamycin. (A) THP-1 cells pretreated with tunicamycin (5 µg/ml) were cultured in AIM-V serum-free culture medium supplemented with LPS (1 µg/ml). After 3 h incubation with LPS, RNA was isolated from the THP-1 cells, and the expression levels of the TNF-α and IL-1β genes were analyzed by RTD-PCR. The expressions of these cytokines were down-regulated in THP-1 cells treated with tunicamycin. (B) THP-1 cells pretreated with tunicamycin were cultured in AIM-V serum-free culture medium supplemented with LPS (1 µg/ml) for 12 h and the concentration of TNF-α and IL-1β in the culture medium was measured by ELISA. The concentrations of these cytokines were decreased. (C) Primary human monocytes were cultured in AIM-V serum-free culture medium supplemented with LPS (1 µg/ml) and tunicamycin (5 µg/ml). After 4 h incubation, RNA was isolated from the cells, and the expression levels of the TNF-α and IL-1β genes was analyzed by RTD-PCR. The expression of IL-1β was down-regulated in cells treated with tunicamycin and TNF-α tended to be down-regulated. (D) Primary human monocytes were cultured in AIM-V serum-free culture medium supplemented with LPS (1 µg/ml) and tunicamycin (5 µg/ml) for 12 h, and the concentration of TNF-α and IL-1β in the culture medium was measured by ELISA. The concentration of these cytokines was decreased. Data are expressed as means ± SEM of four independent experiments. Filled bars; no treatment. Open bar; treatment with tunicamycin (5 µg/ml). * $P < 0.05$, *** $P < 0.001$.

3.3. ER stress impaired monocyte differentiation into macrophages

Monocytes are progenitor cells that differentiate into macrophages and play an important role for inducing acquired immunity [16]. We tried to differentiate THP-1 cells and primary human monocytes into macrophages by PMA and GM-CSF treatment, respectively. When THP-1 cells were treated with PMA for 3 days, cells appeared morphologically mature, showing increased forward scatter (FSC) and side scatter (SSC) (Fig. 5A and B), and displayed enhanced expression of the macrophage-related markers CD11b and CD68 by flow cytometry (Fig. 5C). However, when THP-1 cells were pretreated with tunicamycin followed by PMA treatment, no alteration of FSC and SSC was observed (Fig. 5A and B) and the expression levels of CD11b and CD68 (Fig. 5C and D) were not enhanced. We confirmed that the ER stress markers

CHOP and BiP were upregulated in the PMA-differentiated THP-1 cells with tunicamycin treatment (Fig. 5E). Furthermore, when primary human monocytes were treated with or without tunicamycin and cultured in media with GM-CSF for 4 days, tunicamycin-treated primary human monocytes showed less mature morphological appearance (Fig. 5F), lower FSC and SSC (Fig. 5G) and lower CD11b expression (Fig. 5H) compared to those of untreated primary human monocytes. In THP-1 cells, the differentiated cells under ER stress showed the same expression levels of TLR and MyD88 regardless of tunicamycin-induced ER stress (Fig. 6A and B). However, TNF-α and IL-1β expression levels were significantly decreased in differentiation-induced THP-1 cells under ER stress (Fig. 6C and D). These results suggest that tunicamycin-induced ER stress disturbed the differentiation capability as well as the signaling pathway of TLR4 in monocytes.




Fig. 4. NF-κB activation in response to LPS stimulation in human monocytes was attenuated under ER stress. (A) THP-1 cells treated with tunicamycin (5 μg/ml) for 12 h were cultured in AIM-V serum-free medium supplemented with LPS. Phosphorylated or total NF-κB quantity in THP-1 cells seeded on the culture plate was assessed by cellular activation of the signaling ELISA kit. Induction of phosphorylated NF-κB by LPS stimulation was not observed in THP-1 cells treated with tunicamycin. (B) Fluorescence microscopy examination of the tunicamycin-treated THP-1 cells followed by LPS stimulation. Cells were permeabilized and were stained with anti- NF-κB p65 and tetramethyl rhodamine isothiocyanate-goat anti-mouse IgG (red). The nucleus is stained with DAPI (4, 6-diamino-2-phenylindole; blue). The sustained localization of NF-κB p65 in the cytoplasm was observed in THP-1 cells treated with tunicamycin followed by LPS stimulation. Original magnification, ×400. Filled bars; no treatment. Open bar; treatment with tunicamycin (5 μg/ml). **P* < 0.01.




Fig. 5. Human monocytes under ER stress demonstrated impaired capability for differentiation into macrophages. (A–E) Conditioning with tunicamycin (5 μg/ml) THP-1 cells were treated with PMA (50 ng/ml) for 72 h. Cells were examined under microscopy for morphology and flow cytometry for surface molecules of macrophage markers CD11b and CD68. Gene expression of cells was also analyzed by RTD-PCR for CD11b, CD68, CHOP and BiP. Morphological analysis (A) and scatter and forward scatter by flow cytometry (B) PMA-differentiated THP-1 cells conditioned with tunicamycin were observed to be smaller and less granular compared to cells not treated with tunicamycin. Flow cytometry analysis (C) and gene expression analysis (D) for markers of macrophages, CD11b and CD68 showed decreased expression of these markers in THP-1 cells treated with tunicamycin and PMA. (E) Gene expressions of CHOP and BiP were increased in THP-1 cells conditioned with tunicamycin. (F–H) Primary human monocytes treated with tunicamycin (0.1 μg/ml) were cultured with GM-CSF (100 ng/ml) for 4 days for differentiation into macrophages. Cultured cells were examined by microscopy and flow cytometry. Tunicamycin-treated primary human monocytes were smaller (F) and less granular (G), and CD11b expression was repressed (H) after 4 days culture in media with GM-CSF. Data are expressed as means ± SEM of four independent experiments. Filled bars; no treatment. Open bar; treatment with tunicamycin (5 μg/ml). ***P* < 0.01, ****P* < 0.001.




Fig. 5. (continued)




Fig. 6. The expression of proinflammatory cytokines in response to LPS stimulation was decreased in PMA-driven differentiated macrophage-like cells under ER stress without alteration of the expression of TLR4 and accessory molecules. THP-1 cells were treated with tunicamycin (5 μ g/ml) followed by PMA treatment for differentiation induction into macrophages. Expression analysis by RTD-PCR (A) and flow cytometry (B) was performed for TLR4 and MyD88. Expressions of both molecules were not affected. (C–D) THP-1 cells were treated with tunicamycin (5 μ g/ml) followed by PMA treatment for 72 h followed by LPS stimulation. Gene expression analysis (C) of cells stimulated with LPS for 3 h as well as ELISA quantification (D) of concentrations of cells stimulated with LPS for 12 h in culture medium were assessed. The expressions of TNF- α and IL-1 β in response to LPS stimulation were significantly decreased in differentiated macrophage-like cells when they were under ER stress by tunicamycin. Filled bars; no treatment. Open bar; treatment with tunicamycin (5 μ g/ml).

4. Discussion

We observed that ER stress induced susceptibility to apoptosis in THP-1 cells, a human monocytic cell line. Functionally, the response of LPS-stimulated human monocytes to express proinflammatory cytokines under ER stress was attenuated prior to the obvious appearance of apoptosis, with decreased activation of NF- κ B. Furthermore, the differentiation capability of monocytes into macrophages was inhibited with a hypo-responsiveness to LPS under ER stress. These observations suggest that ER stress is an important pathological condition affecting a variety of monocyte functions.

The ER system is indispensable for synthesizing properly functioning proteins [1]. Under ER stress, cells malfunction due to the improper production or folding of proteins. Therefore, ER stress can result in the retention of harmful unfolded proteins as a consequence of impaired function of ER-related molecules such as protein kinase R-like ER kinase (PERK), inositol requiring enzyme 1 (IRE1), and activating transcription factor 6 (ATF6), all of which are important molecules of the ER system [17,18]. In the immune system, an unfolded protein response (UPR) related to ER stress was observed in myeloma cells, malignant cells of plasma cells [19]. The affected immune response due to an UPR was also observed in the development of nonmalignant immunological disorders, such as rheumatoid arthritis and neurodegenerative disease [2]. UPRs affect the survival and function of DCs because of their pivotal capacity to process peptides for presentation and secrete cytokines [20]. These findings, as well as our previous observation that diabetic monocytes were under ER stress and were impaired [11], demonstrate that monocytes are an important subpopulation in immune cells in ER stress-related diseases.

Prolonged ER stress leads to apoptotic cell death, which is mediated by CHOP [21], a crucial specific molecule for ER stress-induced apoptosis. Apoptosis is also accompanied by alterations of the transcriptional expression of the BCL-2 gene family [22]. We observed that THP-1 monocytic cells under ER stress underwent apoptosis with decreased expression of the anti-apoptotic molecule BCL-2, indicating that ER stress induced THP-1 cell death using conventional apoptotic pathways [23,24].

The TLR pattern recognition by the innate immune system induces proinflammatory cytokines through the activation of NF- κ B [25,26]. In the quiescent state, NF- κ B remains inactive in the cytoplasm through binding to the inhibitory protein I κ B. The phosphorylated I κ B releases NF- κ B to translocate into the nucleus where it mediates the transcription of its target genes [14]. We observed that the tunicamycin-induced ER stress diminished TLR4 signaling without altering the expressions of TLR4 and MyD88. With regard to the transcriptional factor, we observed a lack of the activated form of NF- κ B and sustained cytoplasmic localization of NF- κ B in LPS-stimulated THP-1 cells under ER stress. The activation of NF- κ B is transient and cyclic upon continuous stimulation, which is due to specific negative feedback control systems such as the NF- κ B inducible synthesis of I κ B and A20 [27]. A previous report indicated that ER stress-induced A20, a deubiquitinating protease [28], acts as an inhibitor of NF- κ B [29,30]. We observed that ER stress by tunicamycin treatment induced the expression of BiP, an important chaperone involved in quality control. BiP was previously reported to decrease the activation of NF- κ B [31], and is a potent extracellular anti-inflammatory molecule [32]. Thus, these molecules may be related to the underlining mechanism of the attenuated TLR4 signaling by ER stress involving decreased activation of NF- κ B. Although some molecules related to ER stress, such as A20 and BiP, were previously reported to be involved in decreased NF- κ B activation, we were unable to identify a specific mechanistic ER stress pathway directly related to impaired TLR4-induced expression of TNF- α and IL-1 β . Impaired TLR signaling in

the presence of tunicamycin was not observed following treatment with specific inhibitors of IRE-1, ATF6, or PERK (data not shown), suggesting that multiple pathways regulate this effect. Further investigation of how ER stress-related molecules are involved in inhibition of NF- κ B under ER stress in terms of cellular machinery is, therefore, warranted.

Diabetes is associated with chronic inflammation [33,34]. Although smoldering inflammation is a fundamental pathological condition of diabetic patients, a characteristic of their immunity is the hypo-responsiveness to pathogenic stimulation [35]. We observed that NF- κ B was, to some extent, activated in THP-1 cells under ER stress induced by tunicamycin before TLR4 ligand stimulation, despite the attenuated signal transduction to TLR ligand stimulation in monocytes (Fig. 4A). ER stress slightly increased the activated form of NF- κ B in the quiescent condition before external ligand stimulation. This finding may correspond to the smoldering inflammation of diabetes, whereas the hypo-responsiveness to impaired TLR4 signaling under ER stress is consistent with the impaired immunity of diabetes. Further investigations are needed to elucidate the details of the chronic inflammation status and the systemic immunity condition.

Monocytes are progenitor cells that can differentiate into mature resident macrophages in various human tissues and become important immune regulators that control the acquired immune response [36]. Monocyte differentiation to macrophages is characterized as a reduction in the nucleocytoplasmic ratio [37] and enhanced granularity [38], which was observed in THP-1 cells (Fig. 5A) and primary human monocytes (Fig. 5C). The morphological maturity of macrophage differentiation was not observed in cells under tunicamycin-induced ER stress. Furthermore, the characteristic change of surface markers in the monocyte/macrophage lineage was also affected, showing decreased expressions of macrophage differentiation markers. A previous report suggested that the inhibition of NF- κ B activation is involved in the impairment of monocyte differentiation into macrophages in THP-1 cells [39]. Therefore, the affected NF- κ B activation and subcellular localization under ER stress may explain, in part, the affected differentiation capability of monocytes into macrophages.

We observed that TLR4 signaling was impaired in differentiated macrophages when they were under tunicamycin-induced ER stress (Fig. 6). Therefore, ER stress may broadly affect the monocytic lineage cells that express the pattern recognition molecules related to innate immunity, implying that ER stress is also a significant condition that broadly impairs host immunity.

In conclusion, our findings demonstrate that ER stress affects the pathogenic ligand-induced TLR signaling as a consequence of attenuated activation of NF- κ B, as well as the differentiation capability into macrophages, which are the important functional regulators of the immune reaction. Further investigations are needed to elucidate the mechanisms of ER stress-related perturbations of the immune reaction in association with various diseases.

Conflict of interest

The authors declare no competing financial interests.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.cellimm.2013.04.006>.

References

- [1] B. Kleizen, I. Braakman, Protein folding and quality control in the endoplasmic reticulum, *Curr. Opin. Cell Biol.* 16 (2004) 343–349.

- [2] M. Aridor, W.E. Balch, Integration of endoplasmic reticulum signaling in health and disease, *Nat. Med.* 5 (1999) 745–751.
- [3] C. Xu, B. Bailly-Maitre, J.C. Reed, Endoplasmic reticulum stress: cell life and death decisions, *J. Clin. Invest.* 115 (2005) 2656–2664.
- [4] Y. Ma, L.M. Hendershot, The role of the unfolded protein response in tumor development: friend or foe?, *Nat. Rev. Cancer* 4 (2004) 966–977.
- [5] U. Ozcan, Q. Cao, E. Yilmaz, A.H. Lee, N.N. Iwakoshi, E. Ozdelen, G. Tuncman, C. Görgün, L.H. Glimcher, G.S. Hotamisligil, Endoplasmic reticulum stress links obesity, insulin action, and type 2 diabetes, *Science* 306 (2004) 457–461.
- [6] G.K. Hansson, Inflammation, atherosclerosis, and coronary artery disease, *N. Engl. J. Med.* 352 (2005) 1685–1695.
- [7] J.P. Taylor, J. Hardy, K.H. Fischbeck, Toxic proteins in neurodegenerative disease, *Science* 296 (2002) 1991–1995.
- [8] S. Akira, S. Uematsu, O. Takeuchi, Pathogen recognition and innate immunity, *Cell* 124 (2006) 783–801.
- [9] C. Pasare, R. Medzhitov, Toll-like receptors: linking innate and adaptive immunity, *Microbes. Infect.* 6 (2004) 1382–1387.
- [10] S. Gordon, P.R. Taylor, Monocytes and macrophage heterogeneity, *Nat. Rev. Immunol.* 5 (2005) 953–964.
- [11] T. Komura, Y. Sakai, M. Honda, T. Takamura, K. Matsushima, S. Kaneko, CD14⁺ monocytes are vulnerable and functionally impaired under ER stress in patients with type 2 diabetes, *Diabetes* 59 (2010) 634–643.
- [12] M. Tateno, M. Honda, T. Kawamura, H. Honda, S. Kaneko, Expression profiling of peripheral-blood mononuclear cells from patients with chronic hepatitis C undergoing interferon therapy, *J. Infect. Dis.* 195 (2007) 255–267.
- [13] S. Thoma-Urszynski, S. Stenger, O. Takeuchi, M.T. Ochoa, M. Engele, P.A. Sieling, P.F. Barnes, M. Rollinghoff, P.L. Bolcskei, M. Wagner, S. Akira, M.V. Norgard, J.T. Belisle, P.J. Godowski, B.R. Bloom, R.L. Modlin, Induction of direct antimicrobial activity through mammalian toll-like receptors, *Science* 291 (2001) 1544–1547.
- [14] M.S. Hayden, S. Ghosh, Shared principles in NF- κ B signaling, *Cell* 132 (2008) 344–362.
- [15] F. Wong, C. Hull, R. Zhande, J. Law, A. Karsan, Lipopolysaccharide initiates a TRAF6-mediated endothelial survival signal, *Blood* 103 (2004) 4520–4526.
- [16] N.V. Serbina, T. Jia, T.M. Hohl, E.G. Pamer, Monocyte-mediated defense against microbial pathogens, *Annu. Rev. Immunol.* 26 (2008) 421–452.
- [17] D. Ron, P. Walter, Signal integration in the endoplasmic reticulum unfolded protein response, *Nat. Rev. Mol. Cell Biol.* 8 (2007) 519–529.
- [18] B. Bukau, J. Weissman, A. Horwich, Molecular chaperones and protein quality control, *Cell* 125 (2006) 443–451.
- [19] D.T. Rutkowski, R.J. Kaufman, A trip to the ER: coping with stress, *Trends Cell Biol.* 14 (2004) 20–28.
- [20] N.N. Iwakoshi, M. Pypaert, L.H. Glimcher, The transcription factor XBP-1 is essential for the development and survival of dendritic cells, *J. Exp. Med.* 204 (2007) 2267–2275.
- [21] X.Z. Wang, D. Ron, Stress-induced phosphorylation and activation of the transcription factor CHOP (GADD153) by p38 MAP kinase, *Science* 272 (1996) 1347–1349.
- [22] K.D. McCullough, J.L. Martindale, L.O. Klotz, T.Y. Aw, N.J. Holbrook, Gadd153 sensitizes cells to endoplasmic reticulum stress by down-regulating Bcl2 and perturbing the cellular redox state, *Mol. Cell Biol.* 21 (2001) 1249–1259.
- [23] T. Gotoh, K. Terada, S. Oyadomari, M. Mori, Hsp70-DnaJ chaperone pair prevents nitric oxide- and CHOP-induced apoptosis by inhibiting translocation of Bax to mitochondria, *Cell Death Differ.* 11 (2004) 390–402.
- [24] I. Tabas, D. Ron, Integrating the mechanisms of apoptosis induced by endoplasmic reticulum stress, *Nat. Cell Biol.* 13 (2011) 184–190.
- [25] A. Iwasaki, R. Medzhitov, Toll-like receptor control of the adaptive immune responses, *Nat. Immunol.* 10 (2004) 987–995.
- [26] I. Sabroe, L.C. Parker, S.K. Dower, M.K. Whyte, The role of TLR activation in inflammation, *J. Pathol.* 214 (2008) 126–135.
- [27] S.L. Werner, J.D. Kearns, V. Zadorozhnaya, C. Lynch, C. O'Lynch, M.P. Boldin, A. Ma, D. Baltimore, A. Hoffmann, Encoding NF- κ B temporal control in response to TNF: distinct roles for the negative regulators I κ B α and A20, *Gene Dev.* 22 (2008) 2093–2101.
- [28] K. Hayakawa, N. Hiramatsu, M. Okamura, H. Yamazaki, S. Nakajima, J. Yao, A.W. Paton, J.C. Paton, M. Kitamura, Acquisition of anergy to proinflammatory cytokines in nonimmune cells through endoplasmic reticulum stress response: a mechanism for subsidence of inflammation, *J. Immunol.* 182 (2009) 1182–1191.
- [29] K. Heyninck, R. Beyaert, A20 inhibits NF- κ B activation by dual ubiquitin-editing functions, *Trends Biochem. Sci.* 30 (2005) 1–4.
- [30] A. Krikos, C.D. Laherty, V.M. Dixit, Transcriptional activation of the tumor necrosis factor alpha-inducible zinc finger protein, A20, is mediated by kappa B elements, *J. Biol. Chem.* 267 (1992) 17971–17976.
- [31] S. Nakajima, N. Hiramatsu, K. Hayakawa, Y. Saito, H. Kato, T. Huang, J. Yao, A.W. Paton, J.C. Paton, M. Kitamura, Selective Abrogation of BiP/GRP78 Blunts Activation of NF- κ B through the ATF6 Branch of the UPR: Involvement of C/EBP and mTOR-Dependent Dephosphorylation of Akt, *Mol. Cell Biol.* 31 (2011) 1710–1718.
- [32] V.M. Corrigan, M.D. Bodman-Smith, M. Brunst, H. Cornell, G.S. Panayi, Inhibition of Antigen-Presenting Cell Function and Stimulation of Human Peripheral Blood Mononuclear Cells to Express an Antiinflammatory Cytokine Profile by the Stress Protein BiP, *Arthritis. Rheum.* 50 (2004) 1164–1171.
- [33] K.E. Wellen, G.S. Hotamisligil, Inflammation, stress and diabetes, *J. Clin. Invest.* 115 (2005) 1111–1119.
- [34] M. Stumvoll, B.J. Goldstein, T.W. van Haefen, Type 2 diabetes: principles of pathogenesis and therapy, *Lancet* 365 (2005) 1333–1346.
- [35] C.P. Liang, S. Han, T. Senokuchi, A.R. Tall, The macrophage at the crossroads of insulin resistance and atherosclerosis, *Circ. Res.* 100 (2007) 1546–1555.
- [36] G. Siamon, R.T. Philip, Monocyte and macrophage heterogeneity, *Nat. Rev. Immunol.* 5 (2005) 953–964.
- [37] R.J. Sokol, G. Hudson, N.T. James, I.J. Frost, J. Wales, Human macrophage development: a morphometric study, *J. Anat.* 151 (1987) 27–35.
- [38] K.C. McCullough, S. Basta, S. Knötig, H. Gerber, R. Schaffner, Y.B. Kim, A. Saalmüller, A. Summerfield, Intermediate stages in monocyte-macrophage differentiation modulate phenotype and susceptibility to virus infection, *Immunology* 98 (1999) 203–212.
- [39] Y.S. Chuang, S.H. Yang, T.Y. Chen, J.H. Pang, Cilostazol inhibits matrix invasion and modulates the gene expressions of MMP-9 and TIMP-1 in PMA-differentiated THP-1 cells, *Eur. J. Pharmacol.* 670 (2011) 419–426.

RESEARCH ARTICLE

Open Access

Peretinoin, an acyclic retinoid, improves the hepatic gene signature of chronic hepatitis C following curative therapy of hepatocellular carcinoma

Masao Honda^{1,2*}, Taro Yamashita¹, Tatsuya Yamashita¹, Kuniaki Arai¹, Yoshio Sakai¹, Akito Sakai¹, Mikiko Nakamura¹, Eishiro Mizukoshi¹ and Shuichi Kaneko¹

Abstract

Background: The acyclic retinoid, peretinoin, has been shown to be effective for suppressing hepatocellular carcinoma (HCC) recurrence after definitive treatment in a small-scale randomized clinical trial. However, little has been documented about the mechanism by which peretinoin exerts its inhibitory effects against recurrent HCC in humans *in vivo*.

Methods: Twelve hepatitis C virus-positive patients whose HCC had been eradicated through curative resection or ablation underwent liver biopsy at baseline and week 8 of treatment with either a daily dose of 300 or 600 mg peretinoin. RNA isolated from biopsy samples was subjected to gene expression profile analysis.

Results: Peretinoin treatment elevated the expression levels of *IGFBP6*, *RBP1*, *PRB4*, *CEBPA*, *GOS2*, *TGM2*, *GPRC5A*, *CYP26B1*, and many other retinoid target genes. Elevated expression was also observed for interferon-, Wnt-, and tumor suppressor-related genes. By contrast, decreased expression levels were found for mTOR- and tumor progression-related genes. Interestingly, gene expression profiles for week 8 of peretinoin treatment could be classified into two groups of recurrence and non-recurrence with a prediction accuracy rate of 79.6% ($P < 0.05$). In the liver of patients with non-recurrence, expression of *PDGFC* and other angiogenesis genes, cancer stem cell marker genes, and genes related to tumor progression was down-regulated, while expression of genes related to hepatocyte differentiation, tumor suppression genes, and other genes related to apoptosis induction was up-regulated.

Conclusions: Gene expression profiling at week 8 of peretinoin treatment could successfully predict HCC recurrence within 2 years. This study is the first to show the effect of peretinoin in suppressing HCC recurrence *in vivo* based on gene expression profiles and provides a molecular basis for understanding the efficacy of peretinoin.

Keywords: Acyclic retinoid, Gene expression, Hepatocellular carcinoma

Background

Hepatocellular carcinoma (HCC) is the sixth most common form of cancer worldwide, and it is estimated that there are more than 740,000 new cases each year [1]. Early-stage HCC is indicated for definitive treatment by surgical resection or local therapy [2-4]; however, the

prognosis of HCC is typically poor, and around 50% of patients experience recurrence within 3 years of definitive therapy [5-7]. Indeed, some researchers estimate that the 3-year recurrence rate is higher than 70% for hepatitis C virus (HCV)-positive patients [8], and past clinical experience with interferon-based therapy, systemic chemotherapy, and other treatment modalities has shown the lack of effective standard therapy for suppressing tumor recurrence after definitive treatment for HCC [9-11].

Peretinoin (NIK-333) has only been reported to suppress HCC recurrence in a small-scale randomized controlled

* Correspondence: mhonda@m-kanazawa.jp

¹Department of Gastroenterology, Graduate School of Medicine, Kanazawa University, 13-1 Takara-machi, Kanazawa 920-0934, Japan

²Department of Advanced Medical Technology, Graduate School of Health Medicine, Kanazawa University, 13-1 Takara-machi, Kanazawa 920-8641, Japan

trial [12] in which patients who were disease-free after definitive treatment received oral administration of 600 mg peretinoin daily for one year. The results showed that peretinoin significantly reduced the incidence of recurrent or new HCC [12] and improved patient survival rates [13]. Based on the results of rat pharmacological studies [14,15] and a phase I clinical study of peretinoin [16], a phase II/III clinical study of peretinoin was conducted in which the doses were set at 300 and 600 mg daily. The study demonstrated that, in the Child-Pugh A subgroup, 600 mg/day peretinoin (n=100) reduced the risk of HCC recurrence or death by approximately 40% compared to placebo (n=106) [hazard ratio (HR)=0.60; 95% confidence interval (CI): 0.40–0.89] [17]. On the other hand, 300 mg daily doses of peretinoin were insufficient for tumor control and showed no substantial difference from the placebo [17]. A large-scale clinical study including several countries is now planned to confirm the clinical efficacy of peretinoin.

Little is known about the mechanism by which peretinoin exerts its inhibitory effects against recurrent HCC in humans *in vivo*. In order to investigate this mechanism, we conducted here a comparative study recruiting HCV-positive patients who successfully completed definitive treatment for HCC (similar to the phase II/III clinical study mentioned above). Patients underwent liver biopsy before and after 8 weeks of treatment with repeated doses of peretinoin, and the collected samples were analyzed for gene expression profiling using the remnant liver after eradication of HCC. We found that changes in the gene expression signature observed in this study help us to understand the means by which peretinoin suppresses HCC, in particular its inhibition against *de novo* carcinogenesis.

Methods

Patients

We enrolled 12 HCV-positive patients who were cured of their primary and first recurrent HCC by surgical hepatectomy or radiofrequency ablation therapy and other non-surgical local treatments (Table 1). Complete tumor removal was confirmed by dynamic computed tomography (CT) scans. Inclusion criteria were as follows: positive presence of HCV-RNA in the serum; Child-Pugh class A or B liver function; platelet counts $\geq 50,000/\mu\text{L}$; and age ≥ 20 years. Exclusion criteria included the following: positive for hepatitis B surface antigen; tumor infiltration into the portal vein; use of transarterial embolization or transarterial chemoembolization (TAE/TACE) for definitive therapy; postoperative use of investigational medicinal products, antitumor agents, interferon, or vitamin K2 formulations; blood pressure unmanageable even with medication (systolic pressure ≥ 160 mmHg or diastolic pressure ≥ 100 mmHg); complication with renal impairment, cardiovascular disease, diabetes mellitus,

autoimmune disease, asthma, or other severe disease; presence of neoplasm; allergy to CT contrast media; allergy to retinoids; history of total gastrectomy; possible pregnancy during study; and lactating mothers.

Study design

This trial was a randomized, parallel-group, open-label study. Twelve eligible patients signed the informed consent form for registration. They were randomized to receive one of the two peretinoin doses: 600 or 300 mg per day. Each dose group consisted of 6 patients. After randomization, patients underwent liver biopsy before the start of peretinoin treatment, then orally received peretinoin twice daily for 8 weeks. At the end of the 8-week therapy, they underwent a second liver biopsy (Figure 1A). The collected biopsy samples were kept in RNeasy[®] solution (Ambion Inc., Austin, TX) at 4°C overnight or longer. Within 3 days, the biopsy samples were removed from the RNeasy solution and partially subjected to RNA extraction and purification. The purified RNA samples were stored at -80°C until required for gene expression profiling. The remaining part of the biopsy samples was used to determine the intrahepatic peretinoin concentration. Samples were placed in polypropylene bottles containing 99.5% ethanol, and the air in the bottle was purged with argon. The bottles were tightly closed and stored at -80°C protected from light. Peripheral blood samples were also collected for the analysis of gene expression signatures and to determine plasma peretinoin levels.

After the second biopsy, patients were orally administered peretinoin twice daily for 88 weeks. During the treatment period, patients visited the hospital every 4 weeks for check-ups, drug compliance, and protocol-specified medical examinations. Drug compliance was assessed by pill counts. During the study, use of anticancer agents, interferon, vitamins K and A, and antiviral drugs (e.g., rivabirin) was prohibited. The study was registered at the Japan Pharmaceutical Information Center (JapicCTI-121757). This protocol was approved by the Institutional Review Board of Kanazawa University for clinical investigation following the provisions of Helsinki, Good Clinical Practice guidelines, local laws, and regulations. Written informed consent was obtained from all patients involved in this study. The detail protocol of this study is presented in Additional file 1: Study protocol.

Plasma peretinoin concentration

A 5-mL blood sample was drawn into an EDTA-2Na tube, immediately mixed, and centrifuged to obtain a plasma sample. The air in the sample tubes was replaced with argon, and the tubes were stored at -80°C protected

Table 1 Patient characteristics and prognosis

Patient no.	Dose	Age	Sex	P/R	Curative treatment	MTD	Tumor no.	Tumor histology	Background liver		CP	ALT	PLT	Prognosis	
									F	A				2 yrs	4.5 yrs
1	300	70	F	P	RFA	15	2	m-p	4	2	A	112	7.9	Rec	Rec
2	600	72	F	R	RFA	20	2	w	4	2	A	40	7.9	Rec	Rec λ
3	300	58	M	P	resection	25	1	m-p	2	1	A	16	19.2	nonRec	nonRec
4	600	54	M	P	resection	25	1	m-p	3	2	A	57	16.4	nonRec	Rec
5	600	60	F	P	RFA	23	1	m-p	4	2	B	23	6.4	nonRec	nonRec
6	300	73	F	P	RFA	20	2	m-p	3	2	A	31	14.2	Rec	Rec λ
7	300	69	F	P	RFA	11	3	w-m	4	1	A	38	11.5	Rec	Rec λ
8	600	74	F	P	RFA	16	2	m-p	4	1	A	45	5.1	nonRec	Rec
9	600	65	M	R	RFA	10	1	m-p	2	1	A	29	16.5	nonRec	nonRec
10	600	59	M	P	resection	34	1	m-p	4	2	B	60	9.4	nonRec	nonRec
11	300	70	F	R	RFA	15	1	w-m	4	2	B	98	7	nonRec	nonRec
12	300	66	M	P	RFA	15	1	m-p	4	1	A	90	10.6	nonRec	nonRec

Dose (mg/day), ALT(U/L), PLT($\times 10^4/\mu\text{L}$), MTD (mm).
 F; female, M; male, P; primary HCC, R; (first) recurrent HCC, MTD; maximum tumor diameter, w; well-differentiated, m; moderately differentiated, p; poorly differentiated, F; fibrosis stage, A; activity grade, CP; Child-Pugh classification, ALT; alanine aminotransferase, PLT; platelet.
 Rec; recurrence, nonRec; non-recurrence, λ; death.




Figure 1 Peretinoin pharmacokinetics study design and change of gene expression profiling. **A:** Peretinoin pharmacokinetics study design. Twelve patients were enrolled in the study and two groups of 6 patients were randomly administered one of two doses of peretinoin (600 or 300 mg per day) for 8 weeks. A liver biopsy was obtained before the start of peretinoin administration and 8 weeks into the treatment. After the second liver biopsy, oral administration of peretinoin twice daily was resumed for 88 weeks. **B:** Hierarchical clustering analysis of patients using all expressed genes. Patient numbers (Table 1) and peretinoin dose are listed. Patients with HCC recurrence are shown in red and boxed. There was no significant association between hepatic gene expression and HCC recurrence before starting peretinoin treatment, while distinct clusters of patients were observed after week 8 depending on HCC recurrence.

from light. The plasma concentrations of the unchanged form of peretinoin and its lipid-bound form were determined as follows: first, the peretinoin-containing fractions were extracted from the plasma samples, then subjected to derivatization of peretinoin, and the concentration of the derivative was measured by liquid chromatography-atmospheric pressure chemical ionization-tandem mass spectrometry.

Liver peretinoin concentration

Collected liver tissue samples were immersed in 99.5% ethanol in containers, and the internal air was replaced with argon. The samples were stored at -80°C protected from light. The liver concentrations of the unchanged form of peretinoin and its lipid-bound form were determined as for the plasma concentrations above.

Microarray analysis

For gene expression profiling of the liver, in-house cDNA microarrays containing a representative panel of 10,000 liver-specific genes (Kanazawa liver chip 10K ver. 2.0) were used. RNA isolation, amplification of antisense RNA, labeling, and hybridization were conducted as previously described [18].

To identify genetic variants, paired *t*-tests were performed using BRB-Array Tools software (<http://linus.nci.nih.gov/BRB-ArrayTools.html>) to define *P*-values <0.05 as gene variants. Hierarchical cluster analysis, exploration of significantly expressed genes, and class prediction were also performed using the BRB-Array Tools.

Hierarchical clustering was carried out using centered correlation and average linkage. The class comparison tool in the BRB-Array Tools was used to extract significantly expressed genes. Genes whose expression levels were significantly different between two groups were located by the *t*-test at the $P<0.002$ significance level. Univariate permutation tests were repeated 1,000–2,000 times to control for errors. Class prediction was performed using the above-mentioned significantly differentiated genes as discriminators, and the results were cross-validated using seven algorithms: compound-covariate predictor, diagonal linear discriminant analysis, 1-nearest neighbor, 3-nearest neighbors, nearest centroid, support vector machine, and Bayesian compound covariate. The mean value of the seven success rates for class prediction was defined as the prediction accuracy rate [18].

Pathway analysis was performed using MetaCore™ (Thomson Reuters, New York, NY) and functional ontology enrichment analysis was performed to find differentially expressed pathway using differentially expressed genes [18,19].

The microarray data have been submitted to the Gene Expression Omnibus (GEO) public database at NCBI (Accession No. GSE29302).

Quantitative real-time detection polymerase chain reaction

Quantitative real-time detection polymerase chain reaction (RTD-PCR) was performed using the TaqMan Universal Master Mix (PE Applied Biosystems, Foster City, CA). Primer pairs and probes were purchased from the TaqMan assay reagents library. Standard curves were generated for each assay using RNA derived from normal human liver tissue. Expression data were normalized by GAPDH, and the results are shown as the relative fold expression to the normal liver.

Statistical analysis

Results are expressed as means \pm S.D. Significance was tested by one-way ANOVA with Bonferroni's method, and differences were considered statistically significant at $P<0.05$.

Results

Safety

In this study, 88 adverse events were recorded in 12 patients (100%). Major adverse events included rhinopharyngitis ($n=7$), blood pressure elevation ($n=5$), peripheral edema ($n=3$), and enteritis ($n=3$). Most of these adverse events were mild or moderate, and were adequately controlled. Nine serious adverse events were documented in 5 patients, including hyperglycemia ($n=2$) and coronary stenosis ($n=1$). However, all reported serious adverse events were alleviated with appropriate treatment, and there was no substantial concern identified regarding the safety of peretinoin.

Plasma peretinoin concentration

Plasma peretinoin concentrations were determined at week 8 of treatment. The mean (\pm SD) plasma concentrations of the unchanged form of peretinoin were 82.3 (\pm 90.0) and 201.2 (\pm 111.4) ng/mL at 4 h post-dose and 35.8 (\pm 49.2) and 29.0 (\pm 17.9) ng/mL at 8 h post-dose for the 300 and 600 mg per day groups, respectively. The plasma concentrations of the unchanged peretinoin measured at 4 h post-dose ($\approx t_{\text{max}}$) were dose-dependent. The mean (\pm SD) plasma concentrations of the lipid-bound form of peretinoin were 1478.8 (\pm 853.7) and 2789.8 (\pm 1630.0) ng/mL at 4 h post-dose and 1227.8 (\pm 942.7) and 2213.2 (\pm 1156.1) ng/mL at 8 h post-dose for the 300 and 600 mg per day groups, respectively. The plasma concentrations of the lipid-bound form of peretinoin were dose-dependent at 4 and 8 h post-dose.

Liver peretinoin concentration

Liver peretinoin concentrations were determined at week 8 of treatment. The measurements of the liver concentration of the unchanged form of peretinoin were all below the lower limit of quantitation at 4 h post-dose for all 6 patients in the 300 mg per day group. For the

600 mg per day group, 2 patients yielded measurements of 0.052 and 0.059 $\mu\text{g/g}$, while the remaining 4 patients produced results under the lower limit of quantitation (0.050 $\mu\text{g/g}$). The mean (\pm SD) concentrations of the lipid-bound form of peretinoin were 13.7508 (\pm 11.1097) and 12.8345 (\pm 8.7048) $\mu\text{g/g}$ for the 300 and 600 mg per day groups, respectively.

Gene expression analysis

To analyze the gene expression signature of the liver tissue, we identified genes whose expression levels were significantly different before and after the start of the peretinoin treatment (Figure 1A). The identified genes were candidates for peretinoin-responsive genes. The phase II/III clinical study showed that a daily dose of 600 mg peretinoin reduced the risk of HCC recurrence, while a 300 mg dose was not significantly different from the placebo [17]. Therefore, gene expression patterns were compared before and after the start of the 600 mg peretinoin therapy ($n=6$). Consequently, 424 hepatic genes showed significantly different expression levels from baseline at week 8 (enhancement and suppression seen for 190 and 234 genes, respectively). Typical examples of these genes are represented in Table 2 where fold changes of gene expression for the 300 mg and 600 mg doses are shown respectively. In addition to the retinoid-induced genes, genes related to interferon, tumor suppressors, negative regulators of Wnt signaling, insulin-like growth factor (IGF) signaling, and hepatocyte differentiation were significantly up-regulated by peretinoin. By contrast, genes related to the mammalian target of rapamycin (mTOR), tumor progression, cell cycle, and metastasis/angiogenesis were down-regulated. Serial changes in peretinoin-responsive gene expression are shown in Additional file 2: Figure S1. Significant changes in expression were observed in response to 600 mg of peretinoin, while changes in expression were minimal with 300 mg of peretinoin.

Hierarchical clustering of patients using hepatic gene expression prior to administering peretinoin revealed no significant association with clinical outcome, but a significant association became clearly apparent 8 weeks after peretinoin treatment (Figure 1B). The patients were clustered into two groups: one containing patients with HCC recurrence (4 of 5 patients had recurrence) and the other containing those without recurrence (all 6 patients were recurrence free) within 2 years. Supervised learning methods using seven different algorithms showed that the patients receiving treatment could be differentiated into two groups with or without recurrence by 224 gene predictors ($P<0.002$) at 79.6% accuracy ($P<0.05$) (Table 3). Interestingly, 44 of 224 (20%) genes were peretinoin induced.

Although peretinoin-responsive genes were more induced in patients treated with the 600 mg dosage, gene

expression profiling 8 weeks after peretinoin treatment could not be classified according to the dosage (Table 3). This might be because two patients treated with the 300 mg dosage (No. 11 and No. 12) had already expressed high levels of peretinoin-response genes before starting peretinoin treatment (Additional file 2: Figure S1). Interestingly, patients with high levels of peretinoin-response genes before treatment (No. 9–12) did not show HCC recurrence during the entire observation period (4.5 years; Table 1).

Hierarchical clustering of all 12 patients using 224 gene predictors is shown in Figure 2A. Clear gene clusters were observed according to patients with recurrence and those without, with the exception of one patient (No. 3, Table 1). Interestingly, in the liver of patients with non-recurrence, genes related to angiogenesis, cancer stem cells, Wnt signaling, and tumor progression were repressed, while genes inducing differentiation, tumor suppression, and apoptosis were up-regulated (Figure 2B, Table 4). Interestingly, PDGF-C was the most significant predictor to differentiate patients who will experience recurrence within 2 years (Table 4).

Consistent with these results, hierarchical clustering using pre-defined curated gene sets based on the NCBI's Cancer Genome Anatomy Project similarly differentiated patients into two groups with or without HCC recurrence (Figure 3). Among angiogenesis-related genes, PDGF-C, PDGF-B, vascular endothelial growth factor (VEGF)-B, VEGF-D, and fibroblast growth factor-basic (FGF-2) were repressed in patients without recurrence. As for cell signaling-related genes, MYC, SRC, and RAS-related genes were also repressed; retinoid X receptor alpha (RXRA) and CCAAT/enhancer binding protein (C/EBP), alpha were up-regulated in patients without recurrence. Some cytokines (IL-7, IL-13, and IL-18) and chemokines (e.g. CXCL7) were repressed, while major histocompatibility complex molecules and interferon-related molecules (e.g. IFNAR2) were up-regulated in patients without recurrence (Figure 3).

cDNA microarray analysis revealed that among these predictors, the mRNA level of PDGF-C was the most significant predictor for differentiating patients who will experience recurrence within 2 years (Table 4). This observation was also assessed by RTD-PCR (Figure 4). The expression of the catalytic enzyme of retinoic acid, CYP26B1, was significantly up-regulated at around 200 fold by peretinoin treatment, but its expression was equally induced in patients with or without recurrence. However, the expression of RAR- β , a retinoid receptor, was significantly up-regulated by peretinoin in patients without HCC recurrence (Figure 4).

Patients were followed up for a further 3 years (mean: 2.5 ± 0.5 years) after the cessation of peretinoin treatment. Other two patients experienced recurrence during

Table 2 Representative genes significantly up-regulated or down-regulated in response to peretinoin treatment

Parametric p-value	Ratios (Under/Pre)		Description	Symbol	GB acc
	600 mg	300 mg			
Up-regulated genes in response to peretinoin treatment					
Retinoid target genes					
0.0002	1.85	1.25	Cytochrome P450, family 26, subfamily B, polypeptide 1	CYP26B1	NM_019885
0.004	1.75	1.33	Insulin-like growth factor binding protein 6	IGFBP6	NM_002178
0.005	1.42	1.16	Regulatory factor X-associated ankyrin-containing protein	RFXANK	NM_134440
0.006	1.33	1.30	Putative lymphocyte G0/G1 switch gene	G0S2	NM_015714
0.013	1.54	0.90	Retinol binding protein 1	RBP1	NM_002899
0.014	1.56	0.87	Retinol binding protein 4	RBP4	NM_006744
0.034	1.27	1.07	Retinoic acid induced 3	GPRC5A	AI923823
0.040	1.22	1.19	Transglutaminase 2	TGM2	AI962033
0.044	1.23	1.14	CCAAT/enhancer binding protein (C/EBP), alpha	CEBPA	NM_004364
Interferon-related genes					
0.029	1.45	0.93	Guanylate binding protein 1, interferon-inducible, 67kDa	GBP1	NM_002053
0.047	1.39	0.94	Interferon-induced protein 44	IFI44	NM_006417
0.048	1.28	1.05	Chemokine (C-X-C motif) ligand 9	CXCL9	NM_002416
Negative regulator of Wnt and TGF-β signaling					
0.004	1.54	1.06	BMP and activin membrane-bound inhibitor homolog	BAMBI	NM_012342
0.008	1.45	1.11	Secreted frizzled-related protein 5	SFRP5	NM_003015
Anti-angiogenesis					
0.021	1.37	0.98	Thrombomodulin	THBD	NM_000361
0.038	1.28	0.99	Protein C receptor, endothelial (EPCR)	PROCR	NM_006404
Tumor suppressor related					
0.029	1.35	0.96	Jumonji domain containing 3	JMJD3	XM_043272
0.029	1.39	0.91	Jumping translocation breakpoint	JTB	NM_006694
0.034	1.39	1.32	Protein kinase, AMP-activated, alpha 2 catalytic subunit	PRKAA2	NM_006252
Down-regulated genes in response to peretinoin treatment					
mTOR-related-gene					
0.045	0.78	0.94	FK506 binding protein 12-rapamycin associated protein 1	FRAP1	NM_004958
Cytokine and growth factor					
0.019	0.77	1.25	Interleukin 13	IL13	NM_002188
0.031	0.74	1.00	Hepatocyte growth factor	HGF	NM_000601
Tumor progression related					
0.011	0.73	0.94	Junctional adhesion molecule 3	JAM3	NM_032801
0.013	0.70	1.00	V-myc myelocytomatosis viral oncogene homolog	Myc	NM_002467
0.017	0.73	1.12	Src-like-adaptor	SLA	NM_006748
0.028	0.78	1.10	Cell division cycle 2, G1 to S and G2 to M	CDC2	NM_001786
0.030	0.66	0.95	BCL2-associated athanogene	BAG1	NM_004323
0.039	0.64	0.93	Chemokine (C-C motif) receptor 9	CCR9	NM_031200
0.043	0.76	1.13	Pre-B-cell leukemia transcription factor 1	PBX1	H08835

The peretinoin-response genes were identified by comparing hepatic gene expression in the pre and under treatment of 6 patients who were treated with 600 mg dose of peretinoin. The fold changes of gene expression are shown in 300 mg and 600 mg dosage respectively.

Table 3 Supervised learning methods

Class		No. of predictors	Prediction (%)	p-value
		($p < 0.002$)		
Pre-treatment	Recurrence vs non-recurrence	6	47.1	N.S.
On-treatment	Recurrence vs non-recurrence	224	79.6	< 0.05
On-treatment	300 mg vs 600 mg	38	72.7	N.S.

Seven algorithms of Compound-Covariate Predictor, Diagonal Linear Discriminant Analysis 1-Nearest Neighbor, 3-Nearest Neighbors, Nearest Centroid, Support Vector Machine, and Bayesian Compound Covariate were used for class prediction. Prediction % was calculated as the average of these seven algorithms.

further follow up period (No. 4 and No. 8 in Figure 2A, Table 1). Three patients with recurrence died at 0.3, 1.9, and 2.5 years after the cessation of peretinoin treatment. The Kaplan-Meier estimation of the recurrence-free ratio deduced from 224 gene predictors showed significant differences in HCC recurrence between patients with the recurrence expression pattern and those with non-recurrence expression ($P=0.04$). Moreover, Kaplan-Meier estimation of the survival ratio deduced from the same gene predictors showed a trend for improved survival of patients with non-recurrence expression patterns compared with those with the recurrence expression pattern ($P=0.12$) (Figure 2C, D).

With the exception of the number of tumors at the time of curative therapy, none of the other clinical parameters (e.g. peretinoin dose, tumor, background liver histology, or background liver function) were associated with the recurrence-free or survival ratio. Thus, the peretinoin response during the early period of administration deduced from the hepatic gene expression pattern can successfully predict HCC recurrence and, potentially, patient survival.

Discussion

Peretinoin [(2E,4E,6E,10E)-3,7,11,15-tetramethylhexadeca-2,4,6,10,14-pentaenoic acid] is expected to be a powerful agent against HCC recurrence. This synthetic retinoid induces the transcriptional activation of the retinoic acid receptor (RAR) and retinoid X receptor (RXR), which are both members of the retinoid receptor family. One primary pathway of HCC development involves sustained hepatitis virus infection, which causes repeated cycles of hepatocellular necrosis and proliferation. During increased cell proliferation, mutations occur that lead to the development of HCC unless the dedifferentiated tumor cells are eliminated by apoptosis. The anti-HCC mechanism of action of peretinoin has previously been suggested to be a result of induction of cell apoptosis [20,21], enhancement of cell differentiation [21,22], suppression of cell proliferation by elevation of P21 protein expression and suppression of cyclin D1 expression [23,24]. The first route of action is

independent of retinoid receptors, while the others are retinoid receptor-dependent, although all mechanisms remain largely speculative.

Peretinoin was previously shown to suppress *in vivo* hepatocarcinogenesis in 3'-methyl-4-dimethylaminoazobenzene- and *N*-diethylnitrosamine-induced rats [14,15,25], and in hepatoma-bearing mice and transgenic mice expressing a dominant-negative retinoic acid receptor [25,26]. Recently, we revealed that peretinoin effectively inhibits hepatic fibrosis and HCC development in *Pdgf-c Tg* mice. This demonstrated that PDGF signaling is a target of peretinoin in preventing the development of hepatic fibrosis and HCC [27]. The purpose of this study was to investigate how peretinoin exerts its therapeutic potential by analyzing its effects on the gene expression patterns in clinical samples.

Gene expression profiling in patients without HCC recurrence demonstrated the promotion of *RAR-β* expression, the most common retinoid target gene identified by basic research. Moreover, the expression of other retinoid target genes such as *C/EBP-α*, *IGFBP6*, *TGM2*, *GOS2*, *RBPI*, *RBP4*, and *GPRC5A* was also enhanced. Of these, *C/EBP-α*, *IGFBP6*, and *TGM2* have been shown to inhibit HCC proliferation when co-expressed with *RAR-β* by all-trans-retinoic acid [28,29]. In addition, the RXR-selective agonist (rexinoid)-induced expression of *IGFBP6*, which occurs following *RAR-β*-mediated transcriptional activation of *RAR/RXR*, has been shown to suppress tumor growth [30]. Moreover, *GOS2* and *GPRC5A* have been reported to possess tumor suppressive or apoptosis-inducing effects [31,32]. These primary response retinoid target genes are presumably retinoid-responsive genes. In addition to enhancing retinoid target gene expression, peretinoin induced changes in the expression levels of a variety of genes involved in hepatocarcinogenesis, such as those related to Wnt signaling, IGF signaling, interferon, mTOR, and cell cycle regulation. These results suggest that peretinoin modulates multiple signaling cascades involved in carcinogenesis, either directly or indirectly. Abnormalities in the genes regulating Wnt signaling, IGF signaling, interferon, mTOR, and the cell cycle have been indicated to play a crucial role in the development of HCC [33,34]. We argue that peretinoin suppresses HCC cell proliferation by improving the expression of these genes, thereby preventing HCC recurrence.

The cluster analysis performed in this study successfully differentiated patients with recurrence within 2 years and those without it. Supervised learning methods identified 224 genes as predictors for HCC recurrence ($p < 0.002$). Importantly, 44 (20%) of these were peretinoin-responsive genes, suggesting that recurrence-related genes might be regulated by peretinoin-responsive genes.

A comparison of these groups of patients revealed that the non-recurrence group was associated with the




Figure 2 Expression profiling of 224 gene predictors and the prognosis of patients. **A:** Hierarchical clustering using 224 gene predictors of patients with or without HCC recurrence. Patients with HCC recurrence within 2 years are shown in red and patients with HCC recurrence after the cessation of peretinoin are boxed in red. **B:** Pathway analysis of differentially expressed genes using MetaCore (GeneGo). Functional ontology enrichment analysis was performed to find differentially expressed pathway maps or map folders using 224 differentially expressed genes ($p < 0.002$) between patients with and without HCC recurrence. **C, D:** Kaplan-Meier estimation of recurrence-free ratio (**C**) and survival ratio (**D**) of patients with recurrence expression patterns (red) and those with non-recurrence expression (blue).

enhanced expression of genes related to hepatocellular differentiation and tumor suppression. The non-recurrence group also showed reduced expression of the genes promoting liver fibrosis and steatosis and the liver cancer stem cell marker genes. The genes related to hepatocellular differentiation, *MT1H*, *MT2A*, *FOXA1* (*HNF3 α*), and *FOXA3* (*HNF3 γ*), may be secondary response genes regulated by *C/EBP- α* [35,36]. Indeed, *C/EBP- α* manifested a significant shift in expression level before and during treatment with peretinoin, and could also differentiate

between recurrence and non-recurrence within 2 years. Even after the cessation of peretinoin treatment, the expression of these genes was still significantly related to HCC recurrence (Figure 2C, D). Thus, we speculate that the differences in expression levels of peretinoin-response genes would determine the expression of recurrence-related genes (Additional file 3: Figure S2).

Interestingly, PDGF-C was the most significant predictor to differentiate those patients who will experience recurrence. Using a mouse model of PDGF-C over-

Table 4 Representative genes differentially expressed between HCC recurrence and non-recurrence groups

Parametric p-value	t-values	Description	Symbol	GB acc
Up-regulated genes in the recurrence group				
Angiogenesis related				
0.0001	-5.19	Platelet derived growth factor C	PDGFC	A1446155
0.0006	-4.37	Sperm equatorial segment protein 1	NOX5	NM_145658
0.0010	-4.13	Interleukin 18	IL18	A1800476
Cancer stem cell related				
0.0004	-4.63	Prominin 1	PROM1	NM_006017
0.0018	-3.83	Pyruvate kinase, muscle	PKM2	NM_002654
Positive regulator of Wnt				
0.0018	-3.84	TGFB-induced factor (TALE family homeobox)	TGIF1	A1866302
0.0018	-3.84	NK2 transcription factor related, locus 5	NKX2-5	NM_004387
Tumor progression related				
0.0005	-4.47	Transcriptional co-activator with PDZ-binding motif	WWTR1	AK025216
0.0017	-3.87	Roundabout, axon guidance receptor, homolog 1	ROBO1	NM_133631
0.0018	-3.84	Hepatitis B virus x associated protein	RSF1	NM_016578
0.0019	-3.79	Heat shock 27kDa protein 2	HSPB2	NM_001541
Up-regulated genes in the non-recurrence group				
Liver function and hepatocytes differenti related				
0.0002	4.88	Metallothionein 2A	MT2A	NM_005953
0.0002	4.08	CCAAT/enhancer binding protein (C/EBP), alpha	CEBPA	NM_004364
0.0003	4.72	Forkhead box A3	FOXA3	NM_004497
0.0006	4.42	Hemopexin	HPX	NM_000613
0.0006	4.35	Metallothionein 1H	MT1H	NM_005951
0.0013	4.01	Forkhead box A1	FOXA1	NM_004496
0.0014	3.97	FK506 binding protein 8, 38kDa	FKBP8	NM_012181
Tumor suppressor related				
0.0005	4.51	Deleted in colorectal carcinoma	DCC	X76132
0.0018	3.84	NK3 transcription factor related, locus 1	NKX3-1	NM_006167
Apoptosis inducing				
0.0015	3.93	BH3 interacting domain death agonist	BID	NM_197967
0.0019	3.82	Programmed cell death 8	AIFM1	NM_145813

expression resulting in hepatic fibrosis, steatosis, and eventually HCC development, peretinoin was previously shown to significantly repress the development of hepatic fibrosis and tumors [27].

Although gene expression profiling analysis was conducted using the remnant liver after definitive treatment in the present study, past similar research has demonstrated the possibility of predicting recurrent metachronous and multicentric HCC [37,38]. The exact mechanisms of how the expression profile of non-tumor tissues might determine the recurrence risk are not known. However, the degree of differentiation of hepatocytes and microenvironments such as angiogenesis and

fibrogenesis in non-tumor lesions of the liver is likely to be closely associated with hepatocarcinogenesis. Interestingly, patients with pre-activated peretinoin-response genes were resistant to HCC recurrence for the entire observation period (4.5 years).

This study demonstrated that the patient response to peretinoin during the early period of administration could predict HCC recurrence and, potentially, patient survival. However, it should be noted that the current study protocol consisted of 600 mg peretinoin as the subsequent maintenance treatment for all patients after the 8-week start phase (Figure 1A). In addition, we did not conduct a placebo control to observe serial changes



Figure 3 Hierarchical clustering using pre-defined curated gene sets based on NCBI's Cancer Genome Anatomy Project. Presented genes were differentially expressed at P -values <0.05 between patients with and without HCC recurrence.

of hepatic gene expression without peretinoin administration. Therefore, there might be some limitations in drawing concrete conclusions from this study.

Although we attempted to analyze the liver peretinoin concentration in the present study to investigate its possible relationship with gene expression, peretinoin levels

were too low to yield a meaningful result. However, considering that gene expression profiling identified significant changes in the expression levels of retinoid-related and other genes before and during peretinoin treatment, we believe that sufficient levels of peretinoin reached the liver.




Figure 4 RTD-PCR evaluation of PDGF-C, PROM1, MDK, CYP26B1, and RAR β in the liver of patients with or without HCC recurrence.

The previous peretinoin phase II/III clinical study demonstrated that daily doses of 600 mg peretinoin significantly reduced the incidence of HCC recurrence in HCV-positive patients who underwent definitive treatment. The findings of the present study are complementary to this as we successfully identified candidates for the peretinoin-responsive and recurrence-related genes. These genes are probably involved in the inhibition of HCC recurrence and could be beneficial as future candidate biomarkers of the effectiveness of peretinoin.

Conclusions

In this study, patients underwent liver biopsy before and after 8 weeks of treatment with repeated doses of peretinoin. Gene expression profiling at week 8 of peretinoin treatment could successfully predict HCC recurrence within 2 years. This study is the first to show the effect of peretinoin in suppressing HCC recurrence *in vivo* based on gene expression profiles and provides a molecular basis for understanding the efficacy of peretinoin.

Additional files

Additional file 1: Study protocol.

Additional file 2: Figure S1. One-way hierarchical clustering of up-regulated or down-regulated genes in the liver by the administration of peretinoin (300 mg and 600 mg). Changes in gene expression in the liver before the start of peretinoin administration and 8 weeks into the treatment are shown. Patients with HCC recurrence within 2 years are shown in red and those with HCC recurrence after the cessation of peretinoin are boxed in red.

Additional file 3: Figure S2. Schematic representation of peretinoin action in the liver.

Abbreviations

ACR: Acyclic Retinoid; CH-C: Chronic Hepatitis C; HCC: Hepatocellular Carcinoma; HCV: Hepatitis C Virus.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MH: study concept and design, manuscript preparation. TY: gene expression analysis. TY: acquisition of data of clinical data. KA: acquisition of data of clinical data. YS: gene expression analysis. AS: acquisition of data of clinical data. MN: gene expression analysis. EM: acquisition of data of clinical. SK: study concept and design. All authors read and approved the final manuscript.

Acknowledgements

The authors thank Nami Nishiyama for excellent technical assistance. This work was supported in part by KOWA Co. Ltd., Tokyo, Japan.

Received: 13 September 2012 Accepted: 8 March 2013

Published: 15 April 2013

References

1. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM: Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 2010, **127**(12):2893–2917.

2. Altekruse SF, McGlynn KA, Reichman ME: Hepatocellular carcinoma incidence, mortality, and survival trends in the United States from 1975 to 2005. *J Clin Oncol* 2009, **27**(9):1485–1491.
3. Llovet JM, Burroughs A, Bruix J: Hepatocellular carcinoma. *Lancet* 2003, **362**(9399):1907–1917.
4. Llovet JM, Bru C, Bruix J: Prognosis of hepatocellular carcinoma: the BCLC staging classification. *Semin Liver Dis* 1999, **19**(3):329–338.
5. Mazzaferro V, Regalia E, Docì R, Andreola S, Pulvirenti A, Bozzetti F, Montalto F, Ammatuna M, Morabito A, Gennari L: Liver transplantation for the treatment of small hepatocellular carcinomas in patients with cirrhosis. *N Engl J Med* 1996, **334**(11):693–699.
6. Llovet JM, Schwartz M, Mazzaferro V: Resection and liver transplantation for hepatocellular carcinoma. *Semin Liver Dis* 2005, **25**(2):181–200.
7. Castells A, Bruix J, Bru C, Fuster J, Vilana R, Navasa M, Ayuso C, Boix L, Visa J, Rodes J: Treatment of small hepatocellular carcinoma in cirrhotic patients: a cohort study comparing surgical resection and percutaneous ethanol injection. *Hepatology* 1993, **18**(5):1121–1126.
8. Shiratori Y, Shiina S, Teratani T, Imamura M, Obi S, Sato S, Koike Y, Yoshida H, Omata M: Interferon therapy after tumor ablation improves prognosis in patients with hepatocellular carcinoma associated with hepatitis C virus. *Ann Intern Med* 2003, **138**(4):299–306.
9. Breitenstein S, Dimitroulis D, Petrowsky H, Puhon MA, Mülhaupt B, Clavien PA: Systematic review and meta-analysis of interferon after curative treatment of hepatocellular carcinoma in patients with viral hepatitis. *Br J Surg* 2009, **96**(9):975–981.
10. Samuel M, Chow PK, Chan Shih-Yen E, Machin D, Soo KC: Neoadjuvant and adjuvant therapy for surgical resection of hepatocellular carcinoma. *Cochrane Database Syst Rev* 2009, **1**:CD001199.
11. Mathurin P, Raynard B, Dharancy S, Kirzin S, Fallik D, Pruvot FR, Roumilhac D, Canva V, Paris JC, Chaput JC, et al: Meta-analysis: evaluation of adjuvant therapy after curative liver resection for hepatocellular carcinoma. *Aliment Pharmacol Ther* 2003, **17**(10):1247–1261.
12. Muto Y, Moriwaki H, Ninomiya M, Adachi S, Saito A, Takasaki KI, Tanaka T, Tsurumi K, Okuno M, Tomita E, et al: Prevention of second primary tumors by an acyclic retinoid, polyphenolic acid, in patients with hepatocellular carcinoma. Hepatoma Prevention Study Group. *N Engl J Med* 1996, **334**(24):1561–1567.
13. Muto Y, Moriwaki H, Saito A: Prevention of second primary tumors by an acyclic retinoid in patients with hepatocellular carcinoma. *N Engl J Med* 1999, **340**(13):1046–1047.
14. Sano T, Kagawa M, Okuno M, Ishibashi N, Hashimoto M, Yamamoto M, Suzuki R, Kohno H, Matsushima-Nishiwaki R, Takano Y, et al: Prevention of rat hepatocarcinogenesis by acyclic retinoid is accompanied by reduction in emergence of both TGF- α -expressing oval-like cells and activated hepatic stellate cells. *Nutr Cancer* 2005, **51**(2):197–206.
15. Kagawa M, Sano T, Ishibashi N, Hashimoto M, Okuno M, Moriwaki H, Suzuki R, Kohno H, Tanaka T: An acyclic retinoid, NIK-333, inhibits N-diethylnitrosamine-induced rat hepatocarcinogenesis through suppression of TGF- α expression and cell proliferation. *Carcinogenesis* 2004, **25**(6):979–985.
16. Okusaka T, Ueno H, Ikeda M, Morizane C: Phase I and pharmacokinetic clinical trial of oral administration of the acyclic retinoid NIK-333. *Hepatol Res* 2011, **41**(6):542–552.
17. Okita K, Matsui O, Kumada H, Tanaka K, Kaneko S, Moriwaki H, Izumi N, Okusaka T, Ohashi Y, Makuuchi M: Effect of peretinoin on recurrence of hepatocellular carcinoma (HCC): Results of a phase II/III randomized placebo-controlled trial. *J Clin Oncol* 2010, **15**(28Suppl):4024.
18. Honda M, Yamashita T, Ueda T, Takatori H, Nishino R, Kaneko S: Different signaling pathways in the livers of patients with chronic hepatitis B or chronic hepatitis C. *Hepatology* 2006, **44**(5):1122–1138.
19. Honda M, Sakai A, Yamashita T, Nakamoto Y, Mizukoshi E, Sakai Y, Yamashita T, Nakamura M, Shirasaki T, Horimoto K, et al: Hepatic ISG expression is associated with genetic variation in interleukin 28B and the outcome of IFN therapy for chronic hepatitis C. *Gastroenterology* 2010, **139**(2):499–509.
20. Nakamura N, Shidoji Y, Yamada Y, Hatakeyama H, Moriwaki H, Muto Y: Induction of apoptosis by acyclic retinoid in the human hepatoma-derived cell line, HuH-7. *Biochem Biophys Res Commun* 1995, **207**(1):382–388.
21. Yasuda I, Shiratori Y, Adachi S, Obora A, Takemura M, Okuno M, Shidoji Y, Seishima M, Muto Y, Moriwaki H: Acyclic retinoid induces partial differentiation, down-regulates telomerase reverse transcriptase mRNA

- expression and telomerase activity, and induces apoptosis in human hepatoma-derived cell lines. *J Hepatol* 2002, **36**(5):660–671.
22. Yamada Y, Shidoji Y, Fukutomi Y, Ishikawa T, Kaneko T, Nakagama H, Imawari M, Moriwaki H, Muto Y: Positive and negative regulations of albumin gene expression by retinoids in human hepatoma cell lines. *Mol Carcinog* 1994, **10**(3):151–158.
 23. Suzui M, Masuda M, Lim JT, Albanese C, Pestell RG, Weinstein IB: Growth inhibition of human hepatoma cells by acyclic retinoid is associated with induction of p21(CIP1) and inhibition of expression of cyclin D1. *Cancer Res* 2002, **62**(14):3997–4006.
 24. Suzui M, Shimizu M, Masuda M, Lim JT, Yoshimi N, Weinstein IB: Acyclic retinoid activates retinoic acid receptor beta and induces transcriptional activation of p21(CIP1) in HepG2 human hepatoma cells. *Mol Cancer Ther* 2004, **3**(3):309–316.
 25. Muto Y, Moriwaki H: Antitumor activity of vitamin A and its derivatives. *J Natl Cancer Inst* 1984, **73**(6):1389–1393.
 26. Sakabe T, Tsuchiya H, Endo M, Tomita A, Ishii K, Gonda K, Murai R, Takubo K, Hoshikawa Y, Kurimasa A, et al: An antioxidant effect by acyclic retinoid suppresses liver tumor in mice. *Biochem Pharmacol* 2007, **73**(9):1405–1411.
 27. Okada H, Honda M, Campbell JS, Sakai Y, Yamashita T, Takebuchi Y, Hada K, Shirasaki T, Takabatake R, Nakamura M, et al: Acyclic retinoid targets platelet-derived growth factor signaling in the prevention of hepatic fibrosis and hepatocellular carcinoma development. *Cancer Res* 2012, **72**(17):4459–4471.
 28. Tomaru Y, Nakanishi M, Miura H, Kimura Y, Ohkawa H, Ohta Y, Hayashizaki Y, Suzuki M: Identification of an inter-transcription factor regulatory network in human hepatoma cells by Matrix RNAi. *Nucleic Acids Res* 2009, **37**(4):1049–1060.
 29. Nakanishi M, Tomaru Y, Miura H, Hayashizaki Y, Suzuki M: Identification of transcriptional regulatory cascades in retinoic acid-induced growth arrest of HepG2 cells. *Nucleic Acids Res* 2008, **36**(10):3443–3454.
 30. Uray IP, Shen Q, Seo HS, Kim H, Lamph WW, Bissonnette RP, Brown PH: Retinoid-induced expression of IGFBP-6 requires RARbeta-dependent permissive cooperation of retinoid receptors and AP-1. *J Biol Chem* 2009, **284**(1):345–353.
 31. Ma Y, Koza-Taylor PH, DiMattia DA, Hames L, Fu H, Dragnev KH, Turi T, Beebe JS, Freemantle SJ, Dmitrovsky E: Microarray analysis uncovers retinoid targets in human bronchial epithelial cells. *Oncogene* 2003, **22**(31):4924–4932.
 32. Ye X, Tao Q, Wang Y, Cheng Y, Lotan R: Mechanisms underlying the induction of the putative human tumor suppressor GPRC5A by retinoic acid. *Cancer Biol Ther* 2009, **8**(10):951–962.
 33. Llovet JM, Bruix J: Molecular targeted therapies in hepatocellular carcinoma. *Hepatology* 2008, **48**(4):1312–1327.
 34. Zender L, Villanueva A, Tovar V, Sia D, Chiang DY, Llovet JM: Cancer gene discovery in hepatocellular carcinoma. *J Hepatol* 2010, **52**(6):921–929.
 35. Datta J, Majumder S, Kutay H, Motiwala T, Frankel W, Costa R, Cha HC, MacDougald OA, Jacob ST, Ghoshal K: Metallothionein expression is suppressed in primary human hepatocellular carcinomas and is mediated through inactivation of CCAAT/enhancer binding protein alpha by phosphatidylinositol 3-kinase signaling cascade. *Cancer Res* 2007, **67**(6):2736–2746.
 36. Fields AL, Soprano DR, Soprano KJ: Retinoids in biological control and cancer. *J Cell Biochem* 2007, **102**(4):886–898.
 37. Hoshida Y, Villanueva A, Kobayashi M, Peix J, Chiang DY, Camargo A, Gupta S, Moore J, Wrobel MJ, Lerner J, et al: Gene expression in fixed tissues and outcome in hepatocellular carcinoma. *N Engl J Med* 2008, **359**(19):1995–2004.
 38. Utsunomiya T, Shimada M, Imura S, Morine Y, Ikemoto T, Mori M: Molecular signatures of noncancerous liver tissue can predict the risk for late recurrence of hepatocellular carcinoma. *J Gastroenterol* 2010, **45**(2):146–152.

doi:10.1186/1471-2407-13-191


Cite this article as: Honda et al.: Peretinoin, an acyclic retinoid, improves the hepatic gene signature of chronic hepatitis C following curative therapy of hepatocellular carcinoma. *BMC Cancer* 2013 **13**:191.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit





Gene expression profiling of hepatitis B- and hepatitis C-related hepatocellular carcinoma using graphical Gaussian modeling[☆]

Teruyuki Ueda^a, Masao Honda^{a,b,*}, Katsuhisa Horimoto^c, Sachiyo Aburatani^c, Shigeru Saito^c, Taro Yamashita^a, Yoshio Sakai^a, Mikiko Nakamura^a, Hajime Takatori^a, Hajime Sunagozaka^a, Shuichi Kaneko^a

^a Department of Gastroenterology, Graduate School of Medicine, Kanazawa University, Kanazawa, Takara-Machi 13-1, Kanazawa 920-8641, Japan

^b Department of Advanced Medical Technology, Kanazawa University Graduate School of Health Medicine, Kanazawa, Takara-Machi 13-1, Kanazawa 920-8641, Japan

^c Biological Network Team, Computational Biology Research Center, National Institute of Advanced Industrial Science and Technology, Aomi 2-4-7, Koto-ku, Tokyo 135-0064, Japan

ARTICLE INFO

Article history:

Received 1 October 2012

Accepted 11 February 2013

Available online 26 February 2013

Keywords:

Hepatitis B virus

Hepatitis C virus

Hepatocellular carcinoma

Gene expression

ABSTRACT

Background & aims: Gene expression profiling of hepatocellular carcinoma (HCC) and background liver has been studied extensively; however, the relationship between the gene expression profiles of different lesions has not been assessed.

Methods: We examined the expression profiles of 34 HCC specimens (17 hepatitis B virus [HBV]-related and 17 hepatitis C virus [HCV]-related) and 71 non-tumor liver specimens (36 chronic hepatitis B [CH-B] and 35 chronic hepatitis C [CH-C]) using an in-house cDNA microarray consisting of liver-predominant genes. Graphical Gaussian modeling (GGM) was applied to elucidate the interactions of gene clusters among the HCC and non-tumor lesions.

Results: In CH-B-related HCC, the expression of vascular endothelial growth factor-family signaling and regulation of T cell differentiation, apoptosis, and survival, as well as development-related genes was up-regulated. In CH-C-related HCC, the expression of ectodermal development and cell proliferation, wnt receptor signaling, cell adhesion, and defense response genes was also up-regulated. Many of the metabolism-related genes were down-regulated in both CH-B- and CH-C-related HCC. GGM analysis of the HCC and non-tumor lesions revealed that DNA damage response genes were associated with AP1 signaling in non-tumor lesions, which mediates the expression of many genes in CH-B-related HCC. In contrast, signal transducer and activator of transcription 1 and phosphatase and tensin homolog were associated with early growth response protein 1 signaling in non-tumor lesions, which potentially promotes angiogenesis, fibrogenesis, and tumorigenesis in CH-C-related HCC.

Conclusions: Gene expression profiling of HCC and non-tumor lesions revealed the predisposing changes of gene expression in HCC. This approach has potential for the early diagnosis and possible prevention of HCC.

© 2013 Elsevier Inc. All rights reserved.

1. Introduction

Hepatocellular carcinoma (HCC) is one of the most common malignancies worldwide with a particularly poor patient outcome [1]. It often develops as a result of chronic liver disease associated with hepatitis B

Abbreviations: CH-B, chronic hepatitis B; CH-C, chronic hepatitis C; CLL, cells in liver lobules; CPA, cells in the portal area; EF, early fibrosis; EGR1, early growth response protein 1; ESR1, estrogen receptor 1; GGM, graphical Gaussian modeling; HBV, hepatitis B virus; HCC, hepatocellular carcinoma; HCV, hepatitis C virus; hTERT, human telomerase reverse transcriptase; LCM, laser capture microdissection; LF, late fibrosis; PCCM, partial correlation coefficient matrix; PTEN, phosphatase and tensin homolog; SD, standard deviation; SHC, src homology 2 domain containing; STAT1, signal transducer and activator of transcription 1; TCA, tricarboxylic acid cycle; VEGF, vascular endothelial growth factor.

[☆] Conflict of interest: The authors declare no conflict of interest.

* Corresponding author at: Department of Gastroenterology, Graduate School of Medicine, Kanazawa University, Takara-Machi 13-1, Kanazawa 920-8641, Japan. Fax: +81 76 234 4250.

E-mail address: mhonda@m-kanazawa.jp (M. Honda).

(HBV) or hepatitis C virus (HCV) infection or with other etiologies such as long-term alcohol abuse, autoimmunity, and hemochromatosis [2]. HBV and HCV infections are the leading cause of HCC in the world [3]. In Japan, approximately 85% of patients with HCC are positive for the HBV surface antigen or anti-HCV antibody. Approximately 7% of patients with HCV-related liver cirrhosis develop HCC [4] and 3% of patients with HBV-related liver cirrhosis develop HCC [5].

Gene expression analysis of HCC has revealed from previous work, the activation of the wnt/ β -catenin, pRb, p53, transforming growth factor- β , mitogen-activated protein kinase, and Janus kinase/signal transducer and activator of transcription pathways, stress response signaling, and epidermal growth factor receptor [6–8]. In addition, we have previously reported that the gene expression profiles in the livers of patients with chronic hepatitis B (CH-B) and chronic hepatitis C (CH-C) were different. Pro-apoptotic and DNA repair responses were predominant in CH-B, while inflammatory and anti-apoptotic phenotypes were predominant in CH-C [9,10]. Furthermore, we optimized

the laser capture microdissection (LCM) method to isolate cells in liver lobules (CLL) and cells in the portal area (CPA) for detailed gene expression analysis [10,11]. However, it is still unknown how cancer signaling pathways are activated in HCC. As HCC frequently develops from a cirrhotic liver, analyzing the relationship of signaling pathways between HCC and non-cancerous liver tissue might be a useful approach for revealing the mechanism that ultimately leads to the development of HCC.

Graphical Gaussian modeling (GGM) is utilized widely to infer or test the relationships between multiple variables [12–14]. Previously, we developed a method that combines cluster analysis with GGM to infer a genetic network on the basis of expression profile data. Analysis of the expression profile of *Saccharomyces cerevisiae* revealed a model of its genetic network, and the accuracy of the inferred network was confirmed by its agreement with the cumulative results of experimental studies [15]. Therefore, GGM has the potential to be a useful analytical tool to identify the relationship between the gene expression profiles of HCC and non-cancerous liver tissue.

In the present study, we extended the analysis of gene expression in HCC and applied GGM analysis [15,16]. Indeed, our procedure inferred the relationships between gene groups defined by clustering, and its application enabled us to elucidate the framework of the gene clusters in relation to the hepatocellular carcinogenesis of CH-B and CH-C.

2. Results

2.1. Expressed genes in CH-B-related HCC

The gene expression profiles of whole liver biopsy specimens and surgically resected liver were obtained from 36 patients with CH-B, 17 with CH-B-related HCC, 35 with CH-C, and 17 with CH-C-related HCC. The clinical characteristics of the patients are shown in Supplemental Tables A and B. We categorized the F1 and F2 fibrosis stages as early fibrosis (EF; $n = 13$ for CH-B and $n = 12$ for CH-C) and the F3 and F4 fibrosis stages as late fibrosis (LF; $n = 22$ for CH-B and $n = 23$ for CH-C).

The 783 differentially expressed genes in CH-B-related HCC were identified across 20 clusters, of which 4 (Nos. 8, 9, 11, and 20) were up-regulated and 12 (No. 1–7, 12–14, 16, and 17) were down-regulated (Fig. 1 and Supplemental Table C). The up-regulated clusters were comprised angiogenesis, cell cycle, apoptosis, and survival-related genes. Placental growth factor, vascular endothelial growth factor (VEGF)-related protein, SUMO-activating enzyme subunit 2, cyclin E1, and baculoviral IAP repeat-containing 5 were up-regulated (Nos. 8, 9, and 11). In addition, oncogene-related proteins, such as *v-myc myelocytomatosis viral-related oncogene* (No. 9), telomerase-associated protein 1, and stathmin 1/oncoprotein 18 (No. 8), tumor marker genes, such as glypican 3, and growth factors, such as midkine (No. 9), were also up-regulated. In cluster No. 20, the proliferation and invasiveness-related gene and protein tyrosine kinase 2 were up-regulated.

Down-regulation was prominent in many metabolism-related genes including ornithine aminotransferase, insulin receptor substrate 1, glutamate dehydrogenase 2, acyl-coenzyme A oxidase 2, and acetyl-coenzyme A acyltransferase 2, as well as many cytochrome P450 family genes, suggesting impaired xenobiotic, amino acid, and lipid metabolism (Nos. 6, 7, 12, 13, 16, and 17). The characteristic genes expressed in CH-B-related HCC are shown in Table 1.

2.2. Expressed genes in CH-C-related HCC

The 668 differentially expressed genes in CH-C-related HCC were identified across 18 genetic clusters, of which 5 (Nos. 10, 12, 14, 15, and 18) were up-regulated and 11 (Nos. 1–7, 11, 13, 16, and 17) were down-regulated (Fig. 2 and Supplemental Table D). Cluster No. 12 comprised immune defense response genes, such as chemokine (the C-C motif) ligand 19, natural killer cell transcript 4, major

histocompatibility complex class I B, major histocompatibility complex class II DQ beta 1, and ubiquitin-specific protease 8. Cluster No. 14 comprised cytoskeleton-associated, cell cycle, mitosis-related, and MAPKKK cascade-related genes, such as tubulin, src homology 2 domain containing (SHC) transforming protein 1, sterile alpha motif domain containing 9, S100 calcium binding protein A10, annexin A2, cyclin B1, platelet-activating factor acetylhydrolase, isoform 1b, and vimentin. In cluster No. 15, glypican 3, aldo-keto reductase family 1, member B10, ATP citrate lyase, farnesyl diphosphate synthase, serine protease inhibitor, and Kazal type 1 were up-regulated. Cluster No. 15 included many candidate tumor markers of HCC. Interestingly, LCM analysis revealed that many of the up-regulated genes in clusters Nos. 12, 14, and 15 were preferentially expressed in CPA. Cluster No. 18 comprised regulation of G1/S checkpoint, signal transduction, and ectoderm development-related genes, such as bone morphogenetic protein 4, cyclin-dependent kinase inhibitor 2A, fibroblast growth factor 9, and ornithine decarboxylase 1. Similar to CH-B-related HCC, many of the metabolism-related genes, including glucose, lipid, and amino acid genes, were down-regulated. The unique feature of lipid metabolism in CH-C-related HCC was the up-regulation of cholesterol and fatty acid synthesis genes and down-regulation of cholesterol metabolism and β oxidation genes. It was characterized by the up-regulation of stearyl-CoA desaturase, farnesyl diphosphate synthase (No. 14), and ATP citrate lyase (No. 15), and down-regulation of acetyl-coenzyme A acetyltransferase 1. The characteristic genes expressed in CH-C-related HCC are shown in Table 2. Representative gene expression levels confirmed by TaqMan PCR are shown in Supplemental Fig. C1.

Pathway analysis of the combined up- and down-regulated clusters is shown in Supplemental Fig. D and Supplemental Table E. In CH-C-related HCC, immune response- and cytoskeleton-related genes, such as actin, tubulin, and vimentin, were up-regulated, while in CH-B-related HCC, cell matrix interaction genes, such as collagen IV and matrix metalloproteinase, were up-regulated. Immune-related genes were shown to be down-regulated in both CH-C- and CH-B-related HCC by MetaCore™ database analysis (Thomson Reuters, New York, NY) (Supplemental Fig. D). Gene annotation by DAVID Bioinformatics Resources 6.7 (<http://david.abcc.ncifcrf.gov/>) [17] showed that oxidative phosphorylation and ATP synthesis coupled electron transport were up-regulated more in CH-C-related HCC than in CH-B-related HCC (Supplemental Table E).

2.3. Expressed genes in CH-B and CH-C

Differentially expressed genes in CH-B or CH-C were identified by backward selection, which did not include genes that were differentially expressed in CH-B- or CH-C-related HCC. As HCC frequently develops in the LF stage of liver disease, gene expression was evaluated in this stage. A total of 352 genes were differentially expressed in the LF stage of CH-B and classified into 21 clusters, of which 7 (Nos. 2, 3, 9, 10, 15, 16, and 18) were up-regulated and 11 (No. 5–7, 8, 11–14, 17, 20, and 21) were down-regulated (Supplemental Fig. B and Supplemental Table F).

In the CH-B fibrotic liver, genes involved in apoptosis, survival, and response to stress, as well as chemokine- and cytokine-related genes and *wnt* beta-catenin and angiogenesis-related genes, were up-regulated. Interestingly, these genes were already up-regulated in the EF stage of CH-B. In contrast, metabolism-related genes, such as those for pyruvate, cholesterol, and retinol metabolism and the mitochondrial tricarboxylic acid (TCA) cycle, were down-regulated.

In total, 214 genes were differentially expressed in the LF stage of CH-C and classified into 7 gene clusters, of which 1 was up-regulated (No. 1) and 3 were down-regulated (Nos. 3, 5, and 6) (Supplemental Fig. B and Supplemental Table G). In CH-C, genes involved in the interferon signaling pathway, leukocyte chemotaxis, and immune response were preferentially up-regulated. These genes were expressed at a significantly higher level in CPA than in CLL in the liver (No. 1). Conversely, many metabolism and liver function-related genes were down-regulated (Nos. 3, 5, and 6). These genes were expressed at significantly higher levels in CLL compared to CPA in the liver.