

- 41 **Mennan C**, Wright K, Bhattacharjee A, Balain B, Richardson J, Roberts S. Isolation and characterisation of mesenchymal stem cells from different regions of the human umbilical cord. *Biomed Res Int* 2013; **2013**: 916136 [PMID: 23984420 DOI: 10.1155/2013/916136]
- 42 **Dominici M**, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini F, Krause D, Deans R, Keating A, Prockop Dj, Horwitz E. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy* 2006; **8**: 315-317 [PMID: 16923606 DOI: 10.1080/14653240600855905]
- 43 **Seo KW**, Lee SR, Bhandari DR, Roh KH, Park SB, So AY, Jung JW, Seo MS, Kang SK, Lee YS, Kang KS. OCT4A contributes to the stemness and multi-potency of human umbilical cord blood-derived multipotent stem cells (hUCB-MSCs). *Biochem Biophys Res Commun* 2009; **384**: 120-125 [PMID: 19394308]
- 44 **Karahuseyinoglu S**, Cinar O, Kilic E, Kara F, Akay GG, Demiralp DO, Tukun A, Uckan D, Can A. Biology of stem cells in human umbilical cord stroma: in situ and in vitro surveys. *Stem Cells* 2007; **25**: 319-331 [PMID: 17053211 DOI: 10.1634/stemcells.2006-0286]
- 45 **Schugar RC**, Chirieleison SM, Wescoe KE, Schmidt BT, Askew Y, Nance JJ, Evron JM, Peault B, Deasy BM. High harvest yield, high expansion, and phenotype stability of CD146 mesenchymal stromal cells from whole primitive human umbilical cord tissue. *J Biomed Biotechnol* 2009; **2009**: 789526 [PMID: 20037738 DOI: 10.1155/2009/789526]
- 46 **Conconi M**, Di Liddo R, Tommasini M, Calore C, Parnigotto P. Phenotype and differentiation potential of stromal populations obtained from various zones of human umbilical cord: An overview. *Open Tissue Eng Regen Med J* 2011; **4**: 6-20
- 47 **Greco SJ**, Liu K, Rameshwar P. Functional similarities among genes regulated by OCT4 in human mesenchymal and embryonic stem cells. *Stem Cells* 2007; **25**: 3143-3154 [PMID: 17761754 DOI: 10.1634/stemcells.2007-0351]
- 48 **Gang EJ**, Bosnakovski D, Figueiredo CA, Visser JW, Perlingeiro RC. SSEA-4 identifies mesenchymal stem cells from bone marrow. *Blood* 2007; **109**: 1743-1751 [PMID: 17062733 DOI: 10.1182/blood-2005-11-010504]
- 49 **Schrobbach K**, Wrobel J, Hutmacher DW, Woodfield TB, Klein TJ. Stage-specific embryonic antigen-4 is not a marker for chondrogenic and osteogenic potential in cultured chondrocytes and mesenchymal progenitor cells. *Tissue Eng Part A* 2013; **19**: 1316-1326 [PMID: 23301556 DOI: 10.1089/ten.TEA.2012.0496]
- 50 **Huang P**, Lin LM, Wu XY, Tang QL, Feng XY, Lin GY, Lin X, Wang HW, Huang TH, Ma L. Differentiation of human umbilical cord Wharton's jelly-derived mesenchymal stem cells into germ-like cells in vitro. *J Cell Biochem* 2010; **109**: 747-754 [PMID: 20052672 DOI: 10.1002/jcb.22453]
- 51 **Fong CY**, Gauthaman K, Cheyyatraivendran S, Lin HD, Biswas A, Bongso A. Human umbilical cord Wharton's jelly stem cells and its conditioned medium support hematopoietic stem cell expansion ex vivo. *J Cell Biochem* 2012; **113**: 658-668 [PMID: 21976004 DOI: 10.1002/jcb.23395]
- 52 **Fong CY**, Subramanian A, Gauthaman K, Venugopal J, Biswas A, Ramakrishna S, Bongso A. Human umbilical cord Wharton's jelly stem cells undergo enhanced chondrogenic differentiation when grown on nanofibrous scaffolds and in a sequential two-stage culture medium environment. *Stem Cell Rev* 2012; **8**: 195-209 [PMID: 21671058 DOI: 10.1007/s12015-011-9289-8]
- 53 **Kim JW**, Kim SY, Park SY, Kim YM, Kim JM, Lee MH, Ryu HM. Mesenchymal progenitor cells in the human umbilical cord. *Ann Hematol* 2004; **83**: 733-738 [PMID: 15372203]
- 54 **Anzalone R**, Lo Iacono M, Corrao S, Magno F, Loria T, Capello F, Zummo G, Farina F, La Rocca G. New emerging potentials for human Wharton's jelly mesenchymal stem cells: immunological features and hepatocyte-like differentiative capacity. *Stem Cells Dev* 2010; **19**: 423-438 [PMID: 19958166]
- 55 **Conconi MT**, Burra P, Di Liddo R, Calore C, Turetta M, Bellini S, Bo P, Nussdorfer GG, Parnigotto PP. CD105(+) cells from Wharton's jelly show in vitro and in vivo myogenic differentiative potential. *Int J Mol Med* 2006; **18**: 1089-1096 [PMID: 17089012]
- 56 **Zhang HT**, Fan J, Cai YQ, Zhao SJ, Xue S, Lin JH, Jiang XD, Xu RX. Human Wharton's jelly cells can be induced to differentiate into growth factor-secreting oligodendrocyte progenitor-like cells. *Differentiation* 2010; **79**: 15-20 [PMID: 19800163]
- 57 **Campard D**, Lysy PA, Najimi M, Sokal EM. Native umbilical cord matrix stem cells express hepatic markers and differentiate into hepatocyte-like cells. *Gastroenterology* 2008; **134**: 833-848 [PMID: 18243183 DOI: 10.1053/j.gastro.2007.12.024]
- 58 **Dongmei H**, Jing L, Mei X, Ling Z, Hongmin Y, Zhidong W, Li D, Zikuan G, Hengxiang W. Clinical analysis of the treatment of spinocerebellar ataxia and multiple system atrophy-cerebellar type with umbilical cord mesenchymal stromal cells. *Cytotherapy* 2011; **13**: 913-917 [PMID: 21545234]
- 59 **Wang L**, Li J, Liu H, Li Y, Fu J, Sun Y, Xu R, Lin H, Wang S, Lv S, Chen L, Zou Z, Li B, Shi M, Zhang Z, Wang FS. Pilot study of umbilical cord-derived mesenchymal stem cell transfusion in patients with primary biliary cirrhosis. *J Gastroenterol Hepatol* 2013; **28 Suppl 1**: 85-92 [PMID: 23855301 DOI: 10.1111/jgh.12029]
- 60 **Shi M**, Zhang Z, Xu R, Lin H, Fu J, Zou Z, Zhang A, Shi J, Chen L, Lv S, He W, Geng H, Jin L, Liu Z, Wang FS. Human mesenchymal stem cell transfusion is safe and improves liver function in acute-on-chronic liver failure patients. *Stem Cells Transl Med* 2012; **1**: 725-731 [PMID: 23197664 DOI: 10.5966/sctm.2012-0034]
- 61 **Rasmusson I**, Ringdén O, Sundberg B, Le Blanc K. Mesenchymal stem cells inhibit lymphocyte proliferation by mitogens and alloantigens by different mechanisms. *Exp Cell Res* 2005; **305**: 33-41 [PMID: 15777785]
- 62 **Prasanna SJ**, Gopalakrishnan D, Shankar SR, Vasandan AB. Pro-inflammatory cytokines, IFNgamma and TNFalpha, influence immune properties of human bone marrow and Wharton jelly mesenchymal stem cells differentially. *PLoS One* 2010; **5**: e9016 [PMID: 20126406]
- 63 **Girdlestone J**, Limbani VA, Cutler AJ, Navarrete CV. Efficient expansion of mesenchymal stromal cells from umbilical cord under low serum conditions. *Cytotherapy* 2009; **11**: 738-748 [PMID: 19878060]
- 64 **Stubbendorff M**, Deuse T, Hua X, Phan TT, Bieback K, Atkinson K, Eiermann TH, Velden J, Schröder C, Reichenspurner H, Robbins RC, Volk HD, Schrepfer S. Immunological properties of extraembryonic human mesenchymal stromal cells derived from gestational tissue. *Stem Cells Dev* 2013; **22**: 2619-2629 [PMID: 23711207 DOI: 10.1089/scd.2013.0043]
- 65 **Ennis J**, Götherström C, Le Blanc K, Davies JE. In vitro immunologic properties of human umbilical cord perivascular cells. *Cytotherapy* 2008; **10**: 174-181 [PMID: 18368596]
- 66 **Wang D**, Chen K, Du WT, Han ZB, Ren H, Chi Y, Yang SG, Bayard F, Zhu D, Han ZC. CD14+ monocytes promote the immunosuppressive effect of human umbilical cord matrix stem cells. *Exp Cell Res* 2010; **316**: 2414-2423 [PMID: 20420825]
- 67 **Weiss ML**, Anderson C, Medicetty S, Seshareddy KB, Weiss RJ, VanderWerff I, Troyer D, McIntosh KR. Immune properties of human umbilical cord Wharton's jelly-derived cells. *Stem Cells* 2008; **26**: 2865-2874 [PMID: 18703664 DOI: 10.1634/stemcells.2007-1028]

- 68 **Polchert D**, Sobinsky J, Douglas G, Kidd M, Moabsiri A, Reina E, Genrich K, Mehrotra S, Setty S, Smith B, Bartholomew A. IFN-gamma activation of mesenchymal stem

cells for treatment and prevention of graft versus host disease. *Eur J Immunol* 2008; 38: 1745-1755 [PMID: 18493986 DOI: 10.1002/eji.200738129]

P- Reviewers: Forte A, Kim SJ **S- Editor:** Wen LL
L- Editor: Roemmele A **E- Editor:** Zhang DN



