

Table 1. Laboratory data

<i>Hematology</i>	
White blood cells	3,180/ μ l
Hemoglobin	14.0 mg/dl
Platelets	6.7 \times 10 4 / μ l
<i>Blood chemistry</i>	
Total bilirubin	1.4 mg/dl
Direct bilirubin	0.5 mg/dl
AST	19 U/l
ALT	17 U/l
Alkaline phosphatase	213 U/l
γ -GTP	22 U/l
Total protein	7.6 g/dl
Albumin	4.8 g/dl
Immunoglobulin G	1,580 mg/dl
Immunoglobulin A	272 mg/dl
Immunoglobulin M	146 mg/dl
<i>Coagulation</i>	
PT%	77.0%
<i>Serology test</i>	
HBs-Ag	negative
HCV-Ab	negative
Anti-nuclear antibody	negative
Anti-mitochondrial M2 antibody	negative
EBV-VCA IgM	negative
EBV-VCA IgG	80+
EBNA	80+
Anti-platelet antibody	negative
PAIgG	166 ng/10 7 cells
Anti-DNA antibody	40+
Anti- <i>H. pylori</i> antibody IgG	negative
<i>Others</i>	
Acid phosphatase (8.3–21.4)	26.3 U/l
ACE (<14.4)	42.5 IU/ μ l
Leukocyte β -glucuronidase (56.76–74.85)	11.67 nmol/h/mg

ALT = Alanine aminotransferase; AST = aspartate aminotransferase; EBNA = Epstein-Barr virus nuclear antigen; EBV-VCA IgG = Epstein-Barr virus-viral capsid antigen antibody immunoglobulin G; EBV-VCA IgM = Epstein-Barr virus-viral capsid antigen antibody immunoglobulin M; γ -GTP = gamma-glutamyl transpeptidase; HBs-Ag = hepatitis B surface antigen; HCV-Ab = hepatitis C virus antibody; PT% = prothrombin time activity.

Ito et al.: A Case of Adult Type 1 Gaucher Disease Complicated by Temporal Intestinal Hemorrhage




Fig. 3. Changes in platelets count and ACE concentration during the course of ERT.

