

University Hospital, University of Miyazaki Hospital, Yokohama City University Hospital, Yokohama Minami Kyousai Hospital, and Wakayama Medical University Hospital.

This study was supported in part by a grant from the Ministry of Education, Culture, Sports, Science and Technology of Japan (21591384) and a Health and Labor Science Research Grant on intractable diseases from the Ministry of Health, Labor and Welfare of Japan (H22-Nanchi-080 to H.K.).

Authorship

Contribution: H.K. designed the study, followed the patients, analyzed the data, and wrote the manuscript; Y.I. contributed to the

study design, followed the patients, and helped to edit the manuscript; S. Kawabe, K.G., and S.E. performed the experiments; Y.T., S. Kojima, and T.N. followed the patients, collected the clinical data, and helped to edit the manuscript; A.K., A.S., and K.K. followed the patients and collected the clinical data; K.O. performed the experiments and helped to edit the manuscript; and S.N. contributed to the study design, performed the experiments, and wrote the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Hiroshi Kimura, MD, PhD, Department of Virology, Nagoya University Graduate School of Medicine, 65 Tsurumai-cho, Showa-ku, Nagoya 466-8550, Japan; e-mail: hkimura@med.nagoya-u.ac.jp.

References

- Cohen JI. Epstein-Barr virus infection. *N Engl J Med*. 2000;343(7):481-492.
- Williams H, Crawford DH. Epstein-Barr virus: the impact of scientific advances on clinical practice. *Blood*. 2006;107(3):862-869.
- Rickinson AB, Kieff E. Epstein-Barr virus. In: Knipe DM, Howley PM, eds. *Virology*. Vol 2. Philadelphia, PA: Wolters Kluwer/Lippincott Williams & Wilkins; 2006:2655-2700.
- Nakamura S, Jaffe ES, Swerdlow SH. EBV positive diffuse large B-cell lymphoma of the elderly. In: Swerdlow S, Campo E, Harris NL, eds; International Agency for Research on Cancer. *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissue*. Geneva, Switzerland: World Health Organization; 2006:243-244.
- Chan JKC, Quintanilla-Martinez L, Ferry JA, Peh S-C. Extranodal NK/T-cell lymphoma, nasal type. In: Swerdlow S, Campo E, Harris NL, eds; International Agency for Research on Cancer. *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissue*. Geneva, Switzerland: World Health Organization; 2008:285-288.
- Song SY, Kim WS, Ko YH, Kim K, Lee MH, Park K. Aggressive natural killer cell leukemia: clinical features and treatment outcome. *Haematologica*. 2002; 87(12):1343-1345.
- Quintanilla-Martinez L, Kimura H, Jaffe ES. EBV+ T-cell lymphoma of childhood. In: Swerdlow S, Campo E, Harris NL, eds; International Agency for Research on Cancer. *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissue*. Geneva, Switzerland: World Health Organization; 2008: 278-280.
- Jaffe ES. The 2008 WHO classification of lymphomas: implications for clinical practice and translational research. *Hematology Am Soc Hematol Educ Program*. 2009;523-531.
- Ishihara S, Okada S, Wakiguchi H, Kurashige T, Hirai K, Kawa-Ha K. Clonal lymphoproliferation following chronic active Epstein-Barr virus infection and hypersensitivity to mosquito bites. *Am J Hematol*. 1997;54(4):276-281.
- Iwatsuki K, Xu Z, Takata M, et al. The association of latent Epstein-Barr virus infection with hydroa vacciniforme. *Br J Dermatol*. 1999;140(4):715-721.
- Henter JI, Horne A, Arico M, et al. HLH-2004: Diagnostic and therapeutic guidelines for hemophagocytic lymphohistiocytosis. *Pediatr Blood Cancer*. 2007;48(2):124-131.
- Kimura H, Hoshino Y, Kanegane H, et al. Clinical and virologic characteristics of chronic active Epstein-Barr virus infection. *Blood*. 2001;98(2):280-286.
- Quintanilla-Martinez L, Kumar S, Fend F, et al. Fulminant EBV(+) T-cell lymphoproliferative disorder following acute/chronic EBV infection: a distinct clinicopathologic syndrome. *Blood*. 2000; 96(2):443-451.
- Straus SE. The chronic mononucleosis syndrome. *J Infect Dis*. 1988;157(3):405-412.
- Jones J, Shurin S, Abramowsky C, et al. T-cell lymphomas containing Epstein-Barr viral DNA in patients with chronic Epstein-Barr virus infections. *N Engl J Med*. 1988;318(12):733-741.
- Okano M, Kawa H, Kimura H, et al. Proposed guidelines for diagnosing chronic active Epstein-Barr virus infection. *Am J Hematol*. 2005;80(1): 64-69.
- Kikuta H, Taguchi Y, Tomizawa K, et al. Epstein-Barr virus genome-positive T lymphocytes in a boy with chronic active EBV infection associated with Kawasaki-like disease. *Nature*. 1988; 333(6172):455-457.
- Kawa-Ha K, Ishihara S, Ninomiya T, et al. CD3-negative lymphoproliferative disease of granular lymphocytes containing Epstein-Barr viral DNA. *J Clin Invest*. 1989;84(1):51-55.
- Savoldo B, Huls MH, Liu Z, et al. Autologous Epstein-Barr virus (EBV)-specific cytotoxic T cells for the treatment of persistent active EBV infection. *Blood*. 2002;100(12):4059-4066.
- Cohen JI, Jaffe ES, Dale JK, et al. Characterization and treatment of chronic active Epstein-Barr virus disease: a 28-year experience in the United States. *Blood*. 2011;117(22):5835-5849.
- Lay JD, Tsao CJ, Chen JY, Kadin ME, Su IJ. Up-regulation of tumor necrosis factor-alpha gene by Epstein-Barr virus and activation of macrophages in Epstein-Barr virus-infected T cells in the pathogenesis of hemophagocytic syndrome. *J Clin Invest*. 1997;100(8):1969-1979.
- Kasahara Y, Yachie A, Takei K, et al. Differential cellular targets of Epstein-Barr virus (EBV) infection between acute EBV-associated hemophagocytic lymphohistiocytosis and chronic active EBV infection. *Blood*. 2001;98(6):1882-1888.
- Fox CP, Shannon-Lowe C, Gothard P, et al. Epstein-Barr virus-associated hemophagocytic lymphohistiocytosis in adults characterized by high viral genome load within circulating natural killer cells. *Clin Infect Dis*. 2010;51(1):66-69.
- Banionuevo C, Anderson VM, Zevallos-Giampietri E, et al. Hydroa-like cutaneous T-cell lymphoma: a clinicopathologic and molecular genetic study of 16 pediatric cases from Peru. *Appl Immunohistochem Mol Morphol*. 2002;10(1):7-14.
- Cho KH, Lee SH, Kim CW, et al. Epstein-Barr virus-associated lymphoproliferative lesions presenting as a hydroa vacciniforme-like eruption: an analysis of six cases. *Br J Dermatol*. 2004;151(2): 372-380.
- Iwatsuki K, Satoh M, Yamamoto T, et al. Pathogenic link between hydroa vacciniforme and Epstein-Barr virus-associated hematologic disorders. *Arch Dermatol*. 2006;142(5):587-595.
- Kawa K, Okamura T, Yagi K, Takeuchi M, Nakayama M, Inoue M. Mosquito allergy and Epstein-Barr virus-associated T/natural killer-cell lymphoproliferative disease. *Blood*. 2001;98(10): 3173-3174.
- Tokura Y, Ishihara S, Ohshima K, et al. Severe mosquito bite hypersensitivity, natural killer cell leukaemia, latent or chronic active Epstein-Barr virus infection and hydroa vacciniforme-like eruption. *Br J Dermatol*. 1998;138(5):905-906.
- Kimura H, Morishima T, Kanegane H, et al. Prognostic factors for chronic active Epstein-Barr virus infection. *J Infect Dis*. 2003;187(4):527-533.
- Nitta Y, Iwatsuki K, Kimura H, et al. Fatal natural killer cell lymphoma arising in a patient with a crop of Epstein-Barr virus-associated disorders. *Eur J Dermatol*. 2005;15(6):503-506.
- Ishii E, Ohga S, Imashuku S, et al. Nationwide survey of hemophagocytic lymphohistiocytosis in Japan. *Int J Hematol*. 2007;86(1):58-65.
- Kimura H, Morita M, Yabuta Y, et al. Quantitative analysis of Epstein-Barr virus load by using a real-time PCR assay. *J Clin Microbiol*. 1999;37(1): 132-136.
- Kimura H, Ito Y, Suzuki R, Nishiyama Y. Measuring Epstein-Barr virus (EBV) load: the significance and application for each EBV-associated disease. *Rev Med Virol*. 2008;18(5):305-319.
- Kimura H, Hoshino Y, Hara S, et al. Differences between T-cell-type and natural killer cell-type chronic active Epstein-Barr virus infection. *J Infect Dis*. 2005;191(4):531-539.
- Gotoh K, Ito Y, Ohta R, et al. Immunologic and virologic analyses in pediatric liver transplant recipients with chronic high Epstein-Barr virus loads. *J Infect Dis*. 2010;202(3):461-469.
- Kimura H, Miyake K, Yamauchi Y, et al. Identification of Epstein-Barr virus (EBV)-infected lymphocyte subtypes by flow cytometric *in situ* hybridization in EBV-associated lymphoproliferative diseases. *J Infect Dis*. 2009;200(7):1078-1087.
- Chan JKC, Jaffe ES, Ralfkjaer E, Ko YH. Aggressive NK-cell leukaemia. In: Swerdlow S, Campo E, Harris NL, eds; International Agency for Research on Cancer. *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissue*. Geneva, Switzerland: World Health Organization; 2008:276-277.
- Pileri SA, Weisenburger DD, Sng I, Jaffe ES. Peripheral T-cell lymphoma, not otherwise specified. In: Swerdlow S, Campo E, Harris NL, eds; International Agency for Research on Cancer. *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissue*. Geneva, Switzerland: World Health Organization; 2008:306-308.
- Cohen JI, Kimura H, Nakamura S, Ko YH, Jaffe ES. Epstein-Barr virus-associated lymphoproliferative disease in non-immunocompromised hosts: a status report and summary of an international meeting, 8-9 September 2008. *Ann Oncol*. 2009;20(9):1472-1482.

40. Kawa K, Sawada A, Sato M, et al. Excellent outcome of allogeneic hematopoietic SCT with reduced-intensity conditioning for the treatment of chronic active EBV infection. *Bone Marrow Transplant*. 2011;46(1):77-83.
41. Gotoh K, Ito Y, Shibata-Watanabe Y, et al. Clinical and virological characteristics of 15 patients with chronic active Epstein-Barr virus infection treated with hematopoietic stem cell transplantation. *Clin Infect Dis*. 2008;46(10):1525-1534.
42. van Dongen JJ, Langerak AW, Bruggemann M, et al. Design and standardization of PCR primers and protocols for detection of clonal immunoglobulin and T-cell receptor gene recombinations in suspect lymphoproliferations: report of the BIOMED-2 Concerted Action BMH4-CT98-3936. *Leukemia*. 2003;17(12):2257-2317.
43. Takahashi E, Asano N, Li C, et al. Nodal T/NK-cell lymphoma of nasal type: a clinicopathological study of six cases. *Histopathology*. 2008;52(5):585-596.
44. Ohshima K, Kimura H, Yoshino T, et al. Proposed categorization of pathological states of EBV-associated T/natural killer-cell lymphoproliferative disorder (LPD) in children and young adults: overlap with chronic active EBV infection and infantile fulminant EBV T-LPD. *Pathol Int*. 2008;58(4):209-217.
45. Tabiasco J, Vercellone A, Meggetto F, Hudrisier D, Brousset P, Fournie JJ. Acquisition of viral receptor by NK cells through immunological synapse. *J Immunol*. 2003;170(12):5993-5998.
46. Arai A, Imadome K, Watanabe Y, et al. Clinical features of adult-onset chronic active Epstein-Barr virus infection: a retrospective analysis. *Int J Hematol*. 2011;93(5):602-609.
47. Imashuku S, Teramura T, Tauchi H, et al. Longitudinal follow-up of patients with Epstein-Barr virus-associated hemophagocytic lymphohistiocytosis. *Haematologica*. 2004;89(2):183-188.
48. Sato E, Ohga S, Kuroda H, et al. Allogeneic hematopoietic stem cell transplantation for Epstein-Barr virus-associated T/natural killer-cell lymphoproliferative disease in Japan. *Am J Hematol*. 2008;83(9):721-727.
49. Gottschalk S. Reduced-intensity SCT for chronic active EBV infection: excellent outcomes should trigger future investigations on how EBV-positive recipient cells are eradicated. *Bone Marrow Transplant*. 2011;46(1):18-19.
50. Takahashi E, Ohshima K, Kimura H, et al. Clinicopathological analysis of the age-related differences in patients with Epstein-Barr virus (EBV)-associated extranasal natural killer (NK)/T-cell lymphoma with reference to the relationship with aggressive NK cell leukaemia and chronic active EBV infection-associated lymphoproliferative disorders. *Histopathology*. 2011;59(4):660-671.

