

- 275 : 29643.
- 32) Williams JK, Armstrong ML, Heistad DD. Vasa vasorum in atherosclerotic coronary arteries : responses to vasoactive stimuli and regression of atherosclerosis. *Circ Res* 1988 ; 62 : 515.
- 33) Kumamoto M, Nakashima Y, Sueishi K. Intimal neovascularization in human coronary atherosclerosis : its origin and pathophysiological significance. *Hum Pathol* 1995 ; 26 : 450.
- 34) Kaneko H, Anzai T, Takahashi T, et al. Role of vascular endothelial growth factor-A in development of abdominal aortic aneurysm. *Cardiovasc Res* 2011 ; 91 : 358.
- 35) Brown RD, Mitchell MD, Long CS. Pro-inflammatory cytokines and cardiac extracellular matrix : regulation of fibroblast phenotype. In : Villarreal FJ, editor. *Interstitial Fibrosis in Heart Failure*. New York : Springer Science ; 2005. p. 57.
- 36) Hao Q, Wang L, Tang H. Vascular endothelial growth factor induces protein kinase D-dependent production of pro-inflammatory cytokines in endothelial cells. *Am J Physiol Cell Physiol* 2009 ; 296 : C821.
- 37) Sluimer JC, Gasc JM, van Wanroij JL, et al. Hypoxia, hypoxia-inducible transcription factor, and macrophages in human atherosclerotic plaques are correlated with intraplaque angiogenesis. *J Am Coll Cardiol* 2008 ; 51 : 1258.
- 38) Higgins DF, Biju MP, Akai Y, et al. Hypoxic induction of Ctgf is directly mediated by Hif-1. *Am J Physiol Renal Physiol* 2004 ; 287 : F1223.
- 39) Spinale FG. Matrix metalloproteinases : regulation and dysregulation in the failing heart. *Circ Res* 2002 ; 90 : 520.
- 40) Robert V, Besse S, Sabri A, et al. Differential regulation of matrix metalloproteinases associated with aging and hypertension in the rat heart. *Lab Invest* 1997 ; 76 : 729.
- 41) Galis ZS, Khatri JJ. Matrix metalloproteinases in vascular remodeling and atherogenesis : the good, the bad, and the ugly. *Circ Res* 2002 ; 90 : 251.
- 42) Spinale FG, Janicki JS, Zile MR. Membrane-associated matrix proteolysis and heart failure. *Circ Res* 2013 ; 112 : 195.
- 43) Tsuruda T, Kato J, Hatakeyama K, et al. Adventitial mast cells contribute to pathogenesis in the progression of abdominal aortic aneurysm. *Circ Res* 2008 ; 102 : 1368.
- 44) Mizoguchi M, Tahara N, Tahara A, et al. Pioglitazone attenuates atherosclerotic plaque inflammation in patients with impaired glucose tolerance or diabetes a prospective, randomized, comparator-controlled study using serial FDG PET/CT imaging study of carotid artery and ascending aorta. *JACC Cardiovasc Imaging* 2011 ; 4 : 1110.
- 45) Tsuruda T, Hatakeyama K, Nagamachi S, et al. Inhibition of development of abdominal aortic aneurysm by glycolysis restriction. *Arterioscler Thromb Vasc Biol* 2012 ; 32 : 1410.

\* \* \*

