

1994;8(10):1184-97.

20. Fujiwara Y, Browne CP, Cunniff K, Goff SC, Orkin SH. Arrested development of embryonic red cell precursors in mouse embryos lacking transcription factor GATA-1. *Proc Natl Acad Sci U S A*. 1996;93(22):12355-8.

21. Surinya KH, Cox TC, May BK. Identification and characterization of a conserved erythroid-specific enhancer located in intron 8 of the human 5-aminolevulinate synthase 2 gene. *J Biol Chem*. 1998;273(27):16798-809.

22. Fujiwara T, O'Geen H, Keles S, Blahnik K, Linnemann AK, Kang YA, et al. Discovering hematopoietic mechanisms through genome-wide analysis of GATA factor chromatin occupancy. *Mol Cell*. 2009;36(4):667-81.

23. Vargas PD, Furuyama K, Sassa S, Shibahara S. Hypoxia decreases the expression of the two enzymes responsible for producing linear and cyclic tetrapyrroles in the heme biosynthetic pathway. *FEBS J*. 2008;275(23):5947-59.

24. Bekri S, May A, Cotter PD, Al-Sabah AI, Guo X, Masters GS, et al. A promoter mutation in the erythroid-specific 5-aminolevulinate synthase (ALAS2) gene causes X-linked sideroblastic anemia. *Blood*. 2003;102(2):698-704.

25. Muto A, Hoshino H, Madisen L, Yanai N, Obinata M, Karasuyama H, et al. Identification of Bach2 as a B-cell-specific partner for small maf proteins that negatively regulate the immunoglobulin heavy chain gene 3' enhancer. *EMBO J*. 1998;17(19):5734-43.

26. Kadirvel S, Furuyama K, Harigae H, Kaneko K, Tamai Y, Ishida Y, et al. The carboxyl-terminal region of erythroid-specific 5-aminolevulinate synthase acts as an intrinsic modifier for its catalytic activity and protein stability. *Exp Hematol.* 2012;40(6):477-86 e1.
27. Szpurka H, Tiu R, Murugesan G, Aboudola S, Hsi ED, Theil KS, et al. Refractory anemia with ringed sideroblasts associated with marked thrombocytosis (RARS-T), another myeloproliferative condition characterized by JAK2 V617F mutation. *Blood.* 2006;108(7):2173-81.
28. Miller IJ, Bieker JJ. A novel, erythroid cell-specific murine transcription factor that binds to the CACCC element and is related to the Kruppel family of nuclear proteins. *Mol Cell Biol.* 1993;13(5):2776-86.
29. Ney PA, Sorrentino BP, McDonagh KT, Nienhuis AW. Tandem AP-1-binding sites within the human beta-globin dominant control region function as an inducible enhancer in erythroid cells. *Genes Dev.* 1990;4(6):993-1006.
30. Harigae H, Furuyama K. Hereditary sideroblastic anemia: pathophysiology and gene mutations. *Int J Hematol.* 2010;92(3):425-31.
31. Ducamp S, Kannengiesser C, Touati M, Garcon L, Guerci-Bresler A, Guichard JF, et al. Sideroblastic anemia: molecular analysis of the ALAS2 gene in a series of 29 probands and functional studies of 10 missense mutations. *Hum Mutat.* 2011;32(6):590-7.
32. May A, Barton C, Masters G, Kingston J, Lawless S, Jenner M. Severe

Sideroblastic Anaemia in an ALAS2 Compound Heterozygote for -206G, a Common Polymorphism, and a Novel Mutation in Exon 11 (Lys535del) Linked to Lack of Haemoglobinisation In Vitro and Ineffective Erythropoiesis In Vivo. *Blood* (ASH Annual Meeting Abstracts). 2005;106(11):3541.

33. Dou QP, Fridovich-Keil JL, Pardee AB. Inducible proteins binding to the murine thymidine kinase promoter in late G1/S phase. *Proc Natl Acad Sci U S A*. 1991;88(4):1157-61.

34. Lagrange T, Kapanidis AN, Tang H, Reinberg D, Ebright RH. New core promoter element in RNA polymerase II-dependent transcription: sequence-specific DNA binding by transcription factor IIB. *Genes Dev*. 1998;12(1):34-44.

35. Gallagher PG, Sabatino DE, Romana M, Cline AP, Garrett LJ, Bodine DM, et al. A human beta-spectrin gene promoter directs high level expression in erythroid but not muscle or neural cells. *J Biol Chem*. 1999;274(10):6062-73.

36. Faisst S, Meyer S. Compilation of vertebrate-encoded transcription factors. *Nucleic Acids Res*. 1992;20(1):3-26.

37. Rikitake Y, Moran E. DNA-binding properties of the E1A-associated 300-kilodalton protein. *Mol Cell Biol*. 1992;12(6):2826-36.

38. Moser M, Imhof A, Pscherer A, Bauer R, Amselgruber W, Sinowatz F, et al. Cloning and characterization of a second AP-2 transcription factor: AP-2 beta. *Development*. 1995;121(9):2779-88.

39. Gilthorpe J, Vandromme M, Brend T, Gutman A, Summerbell D, Totty N, et al. Spatially specific expression of Hoxb4 is dependent on the ubiquitous transcription factor NFY. *Development*. 2002;129(16):3887-99.
40. Nallur GN, Prakash K, Weissman SM. Multiplex selection technique (MuST): an approach to clone transcription factor binding sites. *Proc Natl Acad Sci U S A*. 1996;93(3):1184-9.
41. Sorensen KD, Quintanilla-Martinez L, Kunder S, Schmidt J, Pedersen FS. Mutation of all Runx (AML1/core) sites in the enhancer of T-lymphomagenic SL3-3 murine leukemia virus unmasks a significant potential for myeloid leukemia induction and favors enhancer evolution toward induction of other disease patterns. *J Virol*. 2004;78(23):13216-31.
42. Spandidos DA, Yiagnisis M, Pintzas A. Human immunodeficiency virus long terminal repeat responds to transformation by the mutant T24 H-ras1 oncogene and it contains multiple AP-1 binding TPA-inducible consensus sequence elements. *Anticancer Res*. 1989;9(2):383-6.
43. Mullhaupt B, Feren A, Jones A, Fodor E. DNA sequence and functional characterization of the human and rat epidermal growth factor promoter: regulation by cell growth. *Gene*. 2000;250(1-2):191-200.

Table 1. Hematologic status of each proband on diagnosis of the disease

	onset of the anemia	Age on diagnosis of SA	family history for XLSA	Hb (g/L)	MCV (fl)	MCH (pg)	platelet (x10E9/L)	serum iron (μmol/L)	ferritin (pmol/L)
Proband 1	0 y.o. (4 months)	0 y.o. (4 months)	yes	39 [136-183]	65 [83-101]	18.7 [28-35]	246 [140-379]	63.9 [10.7-37.6]	399.7 [49.4-270]
Proband 2	4 y.o.	4 y.o.	no	84 [126-165]	73.4 [87-104]	22 [29-35]	610 [138-309]	49.1 [12.5-25.0]	670.1 [67.4-725]
Proband 3	2 y.o.	19 y.o.	no	78 [120-165]	73.9 [80-100]	22.2 [28-34]	373 [160-420]	39.6 [14.3-21.5]	2489.7 [40.4-288]

SI unit and normal value of each clinical examination was in parentheses and brackets, respectively.

Figure legends

Figure 1. Identification of a functional GATA1 element in the first intron of ALAS2 gene.

A: Chromatin immunoprecipitation assay. Fragmented genomic DNA segments were immunoprecipitated with anti-GATA1 antibody or control IgG, and then precipitated fragments were quantified using real-time PCR as described in “supplemental methods”. PC or NC indicates positive control or negative control, respectively, for ChIP assay using anti-GATA1 in K562 cells.²² One GATA element is present in the proximal promoter region and 17 GATA elements in the first intron (black symbols). Shaded double arrow indicates the region corresponding to ChIP-peak.

B: Nucleotide sequence of ChIPmini. The GATA binding site, ALAS2int1GATA, is located in the center of ChIPmini (boxed). Box also indicates the consensus for E-box that is bound by Scl/TAL1²². The sequence of ChIPmini was further analyzed for putative transcription factor binding sites using GeneQuest software (DNASTAR Inc., Madison, WI), and the results are indicated with the horizontal bar. Yi-consensus, Yi transcription

factor consensus site³³; BRE, transcription factor IIB binding site³⁴; EKLF, erythroid/kruppel like factor consensus site³⁵; Sp1, stimulatory protein 1 binding site³⁶; P300, P300 transcriptional coactivator consensus site³⁷; AP-2, AP-2 beta consensus site³⁸; Hox-TF, C1 element binding factor binding site³⁹; OctT3, OctT3 binding site⁴⁰; Runx, Runx proteins binding site⁴¹; AP-1, Activator Protein 1 binding site⁴²; and AP4/E1247, AP4/E1247 binding site⁴³. The sequence for the wild-type probe used in EMSA is indicated with the dashed line. A double arrow indicates the deleted region of delGATA mutation.

Figure 2. Functional analyses of CHIPmini present in the first intron of ALAS2

gene.

A: Electrophoretic mobility shift assay (EMSA). Wild-type (wt) probe was incubated with nuclear extracts prepared from K562 cells (lanes 2-4) or HEK293 cells expressing GATA1-FLAG (lanes 6-9). HEK293 cells were transfected with mock vector (lane 5) or FLAG-fused GATA1 expression vector before preparation of nuclear extracts. The protein-probe complex was detected as a retarded band (arrow). An excess amount of

unlabeled probe (lanes 3, 7), anti-GATA1 antibody (G1) (lanes 4, 8) or anti-FLAG antibody (F) (lane 9) was included in the reaction mixture. Lane 1 shows the control without nuclear extracts. Asterisk indicates the super-shifted band (lane 9).

B. Functional analysis of ChIPmini as an enhancer for ALAS2 gene. Details for the fragments consisting each plasmid, such as intron1, ChIP-peak and ChIPmini, was described in Methods. Each DNA fragment was inserted at the upstream of ALAS2 proximal promoter or at the downstream of luciferase cDNA, indicated as (U) or (D), respectively. Results are expressed as a relative activity compared to that of pGL3-AEpro(-267), and are presented as the mean \pm standard deviation (S.D.) of three independent experiments.

C: Functional analysis of ChIPmini as an enhancer for non-erythroid gene promoter. The enhancer activity of the first intron was examined using the herpes simplex virus TK promoter as a non-erythroid promoter. ChIP-peak or ChIPmini was inserted at the downstream of luciferase gene of pGL3-TKpro, yielding pGL3-TKpro+ChIP-peak(D) or pGL3-TKpro+ChIPmini(D). Each of these reporter vectors was introduced into K562 cells or HEK293 cells for measuring enhancer activity. Results are expressed as a relative

activity compared to that of pGL3-TKpro, and are presented as the mean \pm S.D. of three independent experiments.

Figure 3. Family trees of three unrelated probands

Shown are the family trees of proband 1 with XLSA (A), proband 2 with nfCSA (B), and Proband 3 with nfCSA (C). Shaded boxes indicate affected individuals in each pedigree. Asterisks indicate the individuals in whom mutation in the first intron of ALAS2 gene was detected.

Figure 4. Identification of mutations in the first intron of ALAS2 gene in a patient with XLSA and two patients with nfCSA.

A. ALAS2 mutations of three probands. Upper, middle and lower panels show the sequences of the flanking regions of ALAS2int1GATA (boxed in the wild-type sequence) in ALAS2 gene of probands 1, 2 and 3, respectively. Asterisk indicates the T to C transition in the sense strand identified in ALAS2 gene of proband 1 and proband 2 with CSA. Broken line between middle and lower panels indicates the deleted region

identified in proband 3 with CSA. Solid horizontal bar in each panel indicates the sequence of the sense strand of each probe used for EMSA (see Fig. 3A and 5B).

B: ALAS2 mRNA expression in erythroblasts of proband 3. ALAS2 mRNA levels were determined in purified erythroblasts isolated from proband 3 and three independent normal individuals using real-time PCR. Results are expressed as the mean \pm S.D. of independent three experiments.

Figure 5. Effects of the mutations of ALAS2int1GATA on the GATA1-binding activity.

A: DNA probes used in EMSA. Shown are the nucleotide sequences in the antisense strand of the probes. The position of each probe is also indicated in the Figure 1B as the solid horizontal bar. ALAS2int1GATA is boxed in the sequence of WT probe, and the single nucleotide transition (GGTA mutation) is underlined in the sequence of GGTA probe. The delGATA probe represents the 5'- and 3'-flanking sequences of the deleted 35-bp segment (see Fig. 3B).

B: Effect of each mutation of ALAS2int1GATA on the GATA1-binding activity.

Wild-type probe (lanes 3-7) or each mutant probe (lanes 8, 9) was incubated with the nuclear extracts prepared from HEK293 cells transfected with GATA1-FLAG expression vector. An excess amount of unlabeled wild-type probe (lane 5), each of unlabeled mutant probes (lanes 6, 7), or anti-FLAG antibody (lane 4) was included in the reaction mixture. Lane 2 shows the negative control with nuclear extracts from HEK293 cells transfected with mock vector.

Figure 6. Identification of cis-elements essential for erythroid-specific enhancer activity of ChIPmini.

A: Effect of each mutation of ALAS2int1GATA on the enhancer activity of ALAS2 ChIPmini. The region corresponding to +intron1, ChIP-peak or ChIPmini, derived from proband 1 or proband 3, was subcloned into pGL3-AEpro(-267) for constructing the reporter vector containing the GGTA mutation or the deletion of ALAS2int1GATA, respectively.

B: Effect of the deletion at the 5'- or 3'-flanking region of ALAS2int1GATA on the enhancer activity of ChIPmini.

The 5'- and 3'-flanking regions of ALAS2int1GATA contain potential transcription factor-binding sites (cis-elements), and a portion of each flanking region was deleted, as schematically shown. The enhancer activity of each deletion mutant was determined in K562 erythroleukemia cells.

C: The nucleotide sequence of the 3'-flanking region of ALAS2int1GATA. Note that the Sp1 site is overlapping with AP-1 site and E-box. Each number, *1, *2 or *3, indicates the nucleotide at the 3' end of the deletion mutant, delAP1, delSP1 or delE-box, respectively.

Thus, delSP1 also lacks the 3' portion of the AP-1 site.

D: Effect of deletion of the 5'- and 3'-flanking regions of ALAS2int1GATA on the enhancer activity of ChIPmini.

The construct, delEKL2F2/delSP1, lacks two EKL2F sites in the 5'-flanking region and both Sp1 element and E-box in the 3'-flanking region. The AP-1 element at the 3'-flanking region was deleted from delEKL2F2/delSP1, yielding delEKL2F2/delAP1.

Results are expressed as a relative activity compared to that of pGL3-AEpro(-267), and are presented as the mean \pm S.D. of at least three independent experiments.

Figure 1A




Figure 1B




Figure 2A

antibody	-	-	-	G1	-	-	-	G1	F
competitor	-	-	wt	-	-	-	wt	-	-
nuclear extracts	-	<u>K562</u>		293	<u>293(GATA1-FLAG)</u>				





Figure 2

B



C







Figure 3

A



B



C





Figure 4A

wild type


5' GTGGTCAGACCCCAATGGCGATAAACTCTGGCAACTTTATCTGTGGTCTGCAGGCTCAGCCCCAAGTGCTTTAGCTT
3' CACCAGTCTGGGGTTACCGCTATTTGAGACCGTTGAATAGACACCAGACGTCCGAGTCCGGGTTTCACGAAATCGAA

Proband 1



“GGTA” mutation

Proband 2



Proband 3
“delGATA”
mutation




Figure 4B




Figure 5A

ALAS2int1GATA

wt probe: GAGCCTGCAGACCACAGATAAAGTTGCCAGAGTTTA

GGTA probe: GAGCCTGCAGACCACAGGTAAAGTTGCCAGAGTTTA

delGATA probe: TTGGGGCTGAGCCTGCAGGGGTCTGACCACTCCCCA