

Fig. 3. Immunohistochemical analysis of pERK in the trigeminal ganglion following dural nociceptive stimulation. Immunohistochemical analysis of the time course of phosphorylation of ERK1/2 after the treatment of trigeminal ganglion neurons with capsaicin. (A–D) In the controls, few pERK-IR-exhibiting neurons were observed (A). At 1 min after the application of capsaicin, only a few pERK-IR-exhibiting neurons were observed (B). At 3 min, several neurons exhibited pERK-IR, after which pERK-IR labeling was subsequently decreased at 5 min (C and D; scale bar = 50 μm). (E–G): Double-labeling for pERK (red; E) and NeuN (green; F) immunoreactivity in the trigeminal ganglion 3 min after capsaicin stimulation. The arrow indicates a pERK-IR neuron co-localized with NeuN. (G) Merged image of (E) and (F). Scale bar = 50 μm. (H) Histogram summarizing the quantitative data for pERK-IR in the trigeminal ganglion (A–D). As shown in the histogram, the phosphorylation of ERK after capsaicin stimulation of the dura mater shows a temporal profile similar to that obtained by Western blot analysis (* $p < 0.05$; $n = 4$ in each group; error bars indicate SD).

shown in Fig. 3A, the vehicle-administered animals (controls) showed only a small number of pERK-immunoreactive (pERK-IR) cells. We observed only a few pERK-IR neurons at 1 min after capsaicin application (Fig. 3B), but the number increased 3 min after capsaicin application and was reduced at 5 min (Fig. 3C and D). Quantitatively, the proportion of pERK-IR cells was 0.6% among the control animals (5220 neurons from 4 animals). Following capsaicin administration, pERK immunoreactivity was detectable in 1.5% of all of the cells examined at 1 min after application (5927 neurons),

4.0% at 3 min (4054 neurons), and 2.3% at 5 min (8375 neurons). The ratios of the pERK-IR neurons at 3 min after capsaicin application differed significantly from those of the control animals (Fig. 3H). To demonstrate that ERK was phosphorylated in the neurons, we performed double immunostaining using anti-pERK and anti-NeuN antibodies. As shown in Fig. 3E and F, the immunoreactivity of pERK was restricted to NeuN-IR neurons. Control tissue sections lacking the primary antisera showed no specific staining patterns (data not shown).




Fig. 4. The size of p-ERK-IR neurons with retrograde labeling. Sections of the trigeminal ganglion containing neurons retrogradely labeled with true blue (A), TRPV1-IR (B), and pERK-IR (C). Two weeks after administering the tracer, the skull bone was again removed, and capsaicin was applied to the dura mater. Neurons that retrogradely labeled with true blue (A) and co-localized with TRPV1-IR (B) and pERK-IR (C) are indicated by arrows. Scale bar, 50 µm for all. (D) The size distribution of retrogradely labeled neurons (open bars), retrogradely labeled neurons with TRPV1-IR (gray bars), and retrogradely labeled neurons with TRPV1 and pERK-IR (solid bars).

3.4. Combined use of a neuronal retrograde tracer technique and immunohistochemical analysis of pERK

Following the administration of true blue to the dura mater, retrogradely labeled neurons were observed in the bilateral trigeminal ganglion (Fig. 4A). Additionally, some of these retrogradely labeled neurons showed TRPV1- and pERK-IR (Fig. 4B and C). The mean cross-sectional area of the retrogradely labeled neurons was $478 \pm 250 \mu\text{m}^2$ (mean \pm SD; 4 animals, 480 neurons). The mean cross-sectional area of soma labeling for pERK and TRPV1 containing the retrogradely labeled neurons was $414 \pm 171 \mu\text{m}^2$ (mean \pm SD; 4 animals, 68 neurons) and was significantly smaller compared to that of the retrogradely labeled neurons ($p < 0.05$). A total of 20% of the neurons that accumulated the tracer showed TRPV1-IR, and 83% of those neurons exhibited pERK-IR. The size distributions of all retrogradely labeled neurons, retrogradely labeled neurons with TRPV1-IR and retrogradely labeled neurons with TRPV1 and pERK-IR were calculated. As shown in Fig. 4D and Table 1, the area of most of the neurons retrogradely labeled with TRPV1 and pERK-IR was between 200 and 600 μm^2 (arrows in Fig. 4A–C). Although some large neurons with tracer accumulation were observed, no neurons showing TRPV1-IR had an area of over 1000 μm^2 .

3.5. Effects of CSD on the phosphorylation of ERK in the trigeminal ganglion of rat

Representative blots for pERK and ERK in the trigeminal ganglion are shown in Fig. 5A and D. The ratios of pERK/total-ERK

were 1.82 ± 0.48 for ERK1 and 2.97 ± 0.76 for ERK2 in group-3; 1.12 ± 0.45 for ERK1 and 1.24 ± 0.63 for ERK2 in group-2. The ratios of pERK/total-ERK showed significantly higher levels in group-3 compared to group-1 ($p < 0.05$); however, there was no significant difference between group-1 and group-2 (Fig. 5B and C).

The ratios of pERK/total-ERK were 0.23 ± 0.35 for ERK1 and 0.35 ± 0.43 for ERK2 in group-5. The ratios of pERK/total-ERK showed significantly attenuated levels in group-5 compared to group-4 ($p < 0.05$; Fig. 5E and F). The frequencies of CSD occurrence during the 30 min were 4.7 ± 0.5 times in group-4 and 4.7 ± 1.4 times in group-5, and there was no significant difference between group-4 and group-5.

4. Discussion

4.1. Temporal profile of ERK phosphorylation resulting from dural nociceptive stimulation

Since Ji and colleagues first reported the involvement of ERK following the nociceptive stimulation of spinal neurons (Ji et al., 1999), many studies involving the stimulation (e.g., chemical, electrical, or mechanical) of peripheral C-fibers have shown ERK phosphorylation in the spinal cord, trigeminal nucleus, and dorsal root ganglion (Dai et al., 2002; Shimizu et al., 2006; Noma et al., 2008). Among these reports was a study by Dai and co-workers, who investigated the phosphorylation status of ERK during the very early stages immediately following TRPV1 stimulation. They observed a transient upregulation of ERK phosphorylation only minutes after TRPV1 stimulation in the dorsal root ganglion, which returned to

Table 1
Size distribution of all retrogradely labeled neurons and those retrogradely labeled neurons showing TRPV1- and p-ERK-IR.

Cross-sectional area of neurons (μm^2)	Number of retrogradely labeled cells	Number of cells retrogradely labeled with TRPV1-IR	Number of cells retrogradely labeled with TRPV1 and pERK-IR	Ratio of retrogradely labeled cells with TRPV1 and pERK-IR (%)
100–199	36	7	6	16.7
200–399	181	33	28	15.5
400–599	143	31	26	18.2
600–799	70	9	6	8.6
800–999	29	2	2	6.9
1000–1199	12	0	0	0
1200–1399	50	0	0	0
1400–1599	3	0	0	0
1600 \leq	1	0	0	0

baseline by 120 min (Dai et al., 2002). Our study shows a similar time course in terms of the upregulation of ERK phosphorylation after dural stimulation. We also observed ERK phosphorylation 60 min after dural stimulation and found that the upregulation of ERK phosphorylation gradually decreased to baseline levels. This time course of ERK phosphorylation is consistent with our *in vitro* experiment, which also demonstrated the translocation of pERK to the nucleus (supplementary data). These results imply a role for ERK, which translocates to the nucleus from the cytosol, in the regulation of gene expression.

Our study shows that the dural application of MEK inhibitors attenuated ERK phosphorylation at 3 min following the application of capsaicin. A previous study by Dai and colleagues demonstrated that intradermal administration of a MEK inhibitor prevented the increase of pERK-IR in peripheral nerve terminals and fibers of the skin after capsaicin stimulation, indicating the possibility that the activation of the ERK signaling pathway occurred within the nerve terminals and fibers in addition to the cell bodies in response to peripheral receptor stimulation (Dai et al., 2002). Furthermore, an increase in the expression of pERK in dural nerve fibers was reported after dural mast cell degradation (Levy et al., 2007). Based on these findings, it is likely that the phosphorylation of ERK primarily occurs at the nerve terminals with subsequent propagation to the soma.

We used the right and left side of the trigeminal ganglia for analysis. This methodology is based on our preliminary observations, which have indicated a similar degree of ERK phosphorylation in the bilateral trigeminal ganglia following the stimulation of dural TRPV1 receptors. These findings are also consistent with our tracer experiment, which shows the accumulation of the retrograde tracer in the bilateral trigeminal ganglion after its application to the dura mater around the confluent sinus.

4.2. Size of the neurons in the trigeminal ganglion showing ERK phosphorylation following dural nociceptive stimulation

In line with the findings of our Western blot analysis, immunohistochemical studies also demonstrated peak ERK phosphorylation 3 min after the dural application of capsaicin. To clarify the size of the trigeminal ganglionic neurons responding to dural nociceptive stimulation, we applied a retrograde tracer to the dura mater prior to administering the stimulus and then collected the immunohistochemical data 3 min later. Although the accumulation of the tracer was observed in neurons of varying sizes, most of the labeled neurons that showed TRPV1-IR were small-sized (area <600 μm^2), which includes neurons known to be related to thermal sensation and nociception (Gardner et al., 2000). This result is consistent with our previous results showing that TRPV1-IR neurons that are distributed in the dura mater are also small-sized neurons in the trigeminal ganglion (Shimizu et al., 2007).

In addition, our immunohistochemical analysis, which examined the co-localization of pERK and TRPV1 revealed that most of these TRPV1-IR neurons that innervate the dura mater exhibit pERK-IR. Taken together, this study verifies that the nociceptive stimulation of the dura mater leads to the phosphorylation of ERK in small-sized neurons of the trigeminal ganglion.

4.3. Effects of CSD on the phosphorylation of ERK in the trigeminal ganglion

This study revealed that CSD affects the nociceptive stimulation to the trigeminal ganglion. It has been thought that CSD and the activation of the trigeminovascular system play important roles in the mechanism of migraines. Although CSD has been reported to be related to the pathophysiology of the migraine aura, it is still controversial whether CSD activates the trigeminovascular system. Zhang et al. reported that the trigeminal ganglion was activated at 14–25 min after CSD, and these temporal profiles in terms of the activation of the trigeminal ganglion after CSD were consistent with the results of this study (Zhang et al., 2010). Although the reason for the delay between the induction of CSD and the activation of the trigeminal ganglion system was unclear, this delay could explain why migraine headaches occur after the aura. Karatas et al. reported that CSD activates neuronal pannexin 1 channels, initiates inflammatory responses and promotes the sustained release of inflammatory mediators (Karatas et al., 2013). These inflammatory mediators are continuously released into the subarachnoid space via astrocytes forming the glia limitans and stimulate trigeminal nerve endings around pial vessels. Kosaras et al. revealed that these trigeminal nerve fibers innervating the pial vessels were connected to the dura mater by traversing the arachnoid space (Kosaras et al., 2009). These anatomical findings make it possible to explain the activation of the trigeminal ganglion following CSD.

More importantly, our pharmacological study has raised the possibility that TRPV1 mediates the CSD-induced nociceptive signaling to the trigeminal system (Fig. 6). In animal experiments, CSD is known to evoke a transient release of excitatory amino acids and to increase the extracellular level of proton, which is one of the agonists of TRPV1 (Levy, 2012). Furthermore, CSD is reported to cause degranulation of mast cells in the dura mater (Karatas et al., 2013). The mast cells contain serotonin, histamine and prostaglandin E_2 (Bergerot et al., 2000; Zimmermann et al., 2003), and these substances are known to induce TRPV1 sensitization, which decreases the threshold of TRPV1 activation in terms of temperature and acidity (Sugiura et al., 2002; Dux et al., 2012). Taken together, CSD causes the sensitization and activation of TRPV1, and increases the level of the phosphorylation of ERK.

The ERK signal transduction pathways were originally identified as the primary effectors of growth factor receptor such as TrkA.




Fig. 5. Effects of CSD on ERK activation. The level of phosphorylation of ERK1/2 in trigeminal ganglion neurons following CSD. In group-2, bilateral trigeminal ganglia were dissected immediately after the first wave of CSD. In group-3, they were removed 30 min after the first wave of CSD. In group-1 (control), saline was applied to the open cranial window instead of the 1 M KCl solution used to evoke CSD. A representative image is shown in (A), and the ratios of the intensities of pERK1 and pERK2 are shown in (B) and (C), respectively. A marked increase in the phosphorylation of ERK1/2 was observed in group-3 compared to control ($*p < 0.05$; $n = 5$ in each group; error bars indicate SD). In another experiment, capsazepine (50 nM) was injected into the cisterna magna (group-5). In group-4, vehicle (20% DMSO in saline) was injected instead of capsazepine. In both groups, 1 M KCl was applied to the cranial window to induce CSD, and 30 min later, the bilateral trigeminal ganglia were dissected for Western blot analysis. A representative image is shown in (D), and the ratios of the intensities of pERK1 and pERK2 are shown in (E) and (F), respectively. A marked decrease in the phosphorylation of ERK1/2 was observed in group-5 compared to group-4 ($*p < 0.05$; $n = 5$ in each group; error bars indicate SD).

In this pathway, after GTP loading and activation of the small G-protein Ras were provoked by activated receptors, Ras-GTP recruits a three-tiered enzyme cascade in which a MAPK kinase (Raf) phosphorylates MEK, which phosphorylates and activates ERK (English et al., 1999). Although the precise pathway for the ERK signal

transduction responsible for nociceptive stimulation is still obscure, the activation of the NMDA receptor or voltage-gated calcium channels such as TRPV1 is considered to cause calcium influx or release from intracellular stores, and activates several calcium-dependent kinases in the ERK signal transduction pathway,




Fig. 6. Schematic representation for the possible relationship between CSD and the phosphorylation of ERK via TRPV1 activation. CSD causes a transient release of excitatory amino acids and to increase the extracellular level of proton. Furthermore, CSD facilitates degranulation of mast cells in the dura mater. The mast cells contain serotonin, histamine and prostagrandin E₂, and these substances are known to induce TRPV1 sensitization, which decreases the threshold of TRPV1 activation by temperature and proton. Influx of Ca²⁺ via TRPV1 increases the level of the phosphorylation of ERK, and it may cause peripheral hypersensitivity via transcriptional regulation.

including protein kinase C, tyrosine kinase, calcium/calmodulin-dependent kinase, adenylyl cyclase and tyrosine kinase Pyk2, leading to ERK phosphorylation (Ji and Woolf, 2001; Obata and Noguchi, 2004).

As previously described, recent studies revealed the relationship between ERK phosphorylation and sensitization of the trigeminal ganglion (Yan et al., 2012; Zhang et al., 2013). Since the results of this experiment exhibited the potentiation of the level of the phosphorylation of ERK in the trigeminal ganglion after CSD, our data indicate the possibility that CSD may contribute to migraine-related sensitization.

In clinical studies, Diamond et al. have been reported the efficaciousness of civamide, which is a TRPV1 agonist, for the acute treatment of migraine headache (Diamond et al., 2000). They presumed that the anti-migraine action of civamide was due to desensitization of TRPV1 function. Their finding conferred clinical evidence for the involvement of TRPV1 in the development of migraine headache. Furthermore, Evans et al. have reported that sumatriptan, which is one of the effective migraine abortive drugs, inhibits TRPV1 channels in trigeminal neurons, implying that TRPV1 plays an important role in the migraine-aborting action of sumatriptan (Evans et al., 2012). In addition to these findings, our results imply a pivotal role of TRPV1 for the pathophysiology of migraine, and provide an important clue to novel therapeutic strategies against migraine.

5. Conclusions

The present findings demonstrate that the nociceptive stimulation of TRPV1 receptors in the dura mater causes the phosphorylation of ERK in small-sized neurons of the trigeminal ganglion. CSD also induces the phosphorylation of ERK via TRPV1 stimulation in the trigeminal ganglion. These results indicate the possibility that CSD may act to provide nociceptive stimulation to the trigeminal ganglion and provide important clues regarding the

relationship between CSD and the activation of the trigeminovascular system.

Conflict of interest statement

The authors have no conflicts of interest to declare.

Acknowledgements

This research was supported in part by a Grant-in-Aid for Scientific Research (grant number 22390182 to N. Suzuki) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.neures.2013.08.001>.

References

- Andres, K.H., von Düring, M., Muszynski, K., Schmidt, R.F., 1987. Nerve fibres and their terminals of the dura mater encephali of the rat. *Anat. Embryol. (Berl.)* 175, 289–301.
- Bergerot, A., Reynier-Rebuffel, A.M., Callebert, J., Aubineau, P., 2000. Long-term superior cervical sympathectomy induces mast cell hyperplasia and increases histamine and serotonin content in the rat dura mater. *Neuroscience* 96, 205–213.
- Dai, Y., Iwata, K., Fukuoka, T., Kondo, E., Tokunaga, A., Yamanaka, H., Tachibana, T., Liu, Y., Noguchi, K., 2002. Phosphorylation of extracellular signal-regulated kinase in primary afferent neurons by noxious stimuli and its involvement in peripheral sensitization. *J. Neurosci.* 22, 7737–7745.
- Diamond, S., Freitag, F., Phillips, S.B., Bernstein, J.E., Saper, J.R., 2000. Intranasal civamide for the acute treatment of migraine headache. *Cephalalgia* 20, 597–602.
- Dux, M., Santha, P., Jancso, G., 2012. The role of chemosensitive afferent nerves and TRP ion channels in the pathomechanism of headaches. *Pflugers Arch.* 464, 239–248.

- English, J., Pearson, G., Wilsbacher, J., Swantek, J., Karandikar, M., Xu, S., Cobb, M.H., 1999. New insights into the control of MAP kinase pathways. *Exp. Cell Res.* 253, 255–270.
- Evans, M.S., Cheng, X., Jeffry, J.A., Disney, K.E., Premkumar, L.S., 2012. Sumatriptan inhibits TRPV1 channels in trigeminal neurons. *Headache* 52, 773–784.
- Gardner, E.P., Martin, J.H., Jessell, T.M., 2000. The bodily senses. In: Kandel, E.R., Schwartz, J.H., Jessell, T.M. (Eds.), *Principles of neural science*. , Fourth edition. McGraw-Hill, New York, pp. 430–450.
- Hadjikhani, N., Sanchez Del Rio, M., Wu, O., Schwartz, D., Bakker, D., Fischl, B., Kwong, K.K., Cutrer, F.M., Rosen, B.R., Tootell, R.B., Sorensen, A.G., Moskowitz, M.A., 2001. Mechanisms of migraine aura revealed by functional MRI in human visual cortex. *Proc. Natl. Acad. Sci. U.S.A.* 98, 4687–4692.
- Ji, R.R., Woolf, C.J., 2001. Neuronal plasticity and signal transduction in nociceptive neurons: implications for the initiation and maintenance of pathological pain. *Neurobiol. Dis.* 8, 1–10.
- Ji, R.R., Baba, H., Brenner, G.J., Woolf, C.J., 1999. Nociceptive-specific activation of ERK in spinal neurons contributes to pain hypersensitivity. *Nat. Neurosci.* 2, 1114–1119.
- Ji, R.R., Gereau, R.W.t., Malcangio, M., Strichartz, G.R., 2009. MAP kinase and pain. *Brain Res. Rev.* 60, 135–148.
- Karatas, H., Erdener, S.E., Gursoy-Ozdemir, Y., Lule, S., Eren-Kocak, E., Sen, Z.D., Dalkara, T., 2013. Spreading depression triggers headache by activating neuronal Panx1 channels. *Science* 339, 1092–1095.
- Kosaras, B., Jakubowski, M., Kainz, V., Burstein, R., 2009. Sensory innervation of the calvarial bones of the mouse. *J. Comp. Neurol.* 515, 331–348.
- Leão, A.A.P., 1944. Spreading depression of activity in the cerebral cortex. *J. Neurophysiol.* 78, 359–390.
- Levy, D., 2012. Endogenous mechanisms underlying the activation and sensitization of meningeal nociceptors: the role of immuno-vascular interactions and cortical spreading depression. *Curr. Pain Headache Rep.* 16, 270–277.
- Levy, D., Burstein, R., Kainz, V., Jakubowski, M., Strassman, A.M., 2007. Mast cell degranulation activates a pain pathway underlying migraine headache. *Pain* 130, 166–176.
- Messlinger, K., Hanesch, U., Baumgartel, M., Trost, B., Schmidt, R.F., 1993. Innervation of the dura mater encephali of cat and rat: ultrastructure and calcitonin gene-related peptide-like and substance P-like immunoreactivity. *Anat. Embryol. (Berl.)* 188, 219–237.
- Noma, N., Tsuboi, Y., Kondo, M., Matsumoto, M., Sessle, B.J., Kitagawa, J., Saito, K., Iwata, K., 2008. Organization of pERK-immunoreactive cells in trigeminal spinal nucleus caudalis and upper cervical cord following capsaicin injection into oral and craniofacial regions in rats. *J. Comp. Neurol.* 507, 1428–1440.
- Obata, K., Noguchi, K., 2004. MAPK activation in nociceptive neurons and pain hypersensitivity. *Life Sci.* 74, 2643–2653.
- Olesen, J., Burstein, R., Ashina, M., Tfelt-Hansen, P., 2009. Origin of pain in migraine: evidence for peripheral sensitisation. *Lancet Neurol.* 8, 679–690.
- Olesen, J., Friberg, L., Olsen, T.S., Iversen, H.K., Lassen, N.A., Andersen, A.R., Karle, A., 1990. Timing and topography of cerebral blood flow, aura, and headache during migraine attacks. *Ann Neurol.* 28, 791–798.
- Reuss, S., Riemann, R., Vollrath, L., 1992. Substance P- and calcitonin gene-related peptide-like immunoreactive neurons in the rat trigeminal ganglion – With special reference to meningeal and pineal innervation. *Acta Histochem.* 92, 104–109.
- Roux, P.P., Blenis, J., 2004. ERK and p38 MAPK-activated protein kinases: a family of protein kinases with diverse biological functions. *Microbiol. Mol. Biol. Rev.* 68, 320–344.
- Shimizu, K., Asano, M., Kitagawa, J., Ogiso, B., Ren, K., Oki, H., Matsumoto, M., Iwata, K., 2006. Phosphorylation of extracellular signal-regulated kinase in medullary and upper cervical cord neurons following noxious tooth pulp stimulation. *Brain Res.* 1072, 99–109.
- Shimizu, T., Toriumi, H., Sato, H., Shibata, M., Nagata, E., Gotoh, K., Suzuki, N., 2007. Distribution and origin of TRPV1 receptor-containing nerve fibers in the dura mater of rat. *Brain Res.* 1173, 84–91.
- Sugiura, T., Tominaga, M., Katsuya, H., Mizumura, K., 2002. Bradykinin lowers the threshold temperature for heat activation of vanilloid receptor 1. *J. Neurophysiol.* 88, 544–548.
- Uddman, R., Hara, H., Edvinsson, L., 1989. Neuronal pathways to the rat middle meningeal artery revealed by retrograde tracing and immunocytochemistry. *J. Auton. Nerv. Syst.* 26, 69–75.
- Yan, J., Melemedjian, O.K., Price, T.J., Dussor, G., 2012. Sensitization of dural afferents underlies migraine-related behavior following meningeal application of interleukin-6 (IL-6). *Mol. Pain* 8, 6.
- Zhang, X., Kainz, V., Zhao, J., Strassman, A.M., Levy, D., 2013. Vascular extracellular signal-regulated kinase mediates migraine-related sensitization of meningeal nociceptors. *Ann. Neurol.* 73, 741–750.
- Zhang, X., Levy, D., Nosedá, R., Kainz, V., Jakubowski, M., Burstein, R., 2010. Activation of meningeal nociceptors by cortical spreading depression: implications for migraine with aura. *J. Neurosci.* 30, 8807–8814.
- Zimmermann, K., Reeh, P.W., Averbek, B., 2003. S⁺-flurbiprofen but not 5-HT₁ agonists suppress basal and stimulated CGRP and PGE₂ release from isolated rat dura mater. *Pain* 103, 313–320.

DIFFERENTIAL CELLULAR LOCALIZATION OF ANTIOXIDANT ENZYMES IN THE TRIGEMINAL GANGLION

H. SATO,^{a,b,e} M. SHIBATA,^{a*} T. SHIMIZU,^a S. SHIBATA,^c
H. TORIUMI,^a T. EBINE,^a T. KUROI,^a T. IWASHITA,^a
M. FUNAKUBO,^a Y. KAYAMA,^a C. AKAZAWA,^d
K. WAJIMA,^b T. NAKAGAWA,^b H. OKANO^c AND
N. SUZUKI^a

^a Department of Neurology, School of Medicine, Keio University, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan

^b Department of Dentistry and Oral Surgery, School of Medicine, Keio University, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan

^c Department of Physiology, School of Medicine, Keio University, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan

^d Department of Biochemistry and Biophysics, Graduate School of Health and Sciences, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8510, Japan

^e Japan Society for the Promotion of Science, 8 Ichiban-cho, Chiyoda-ku, Tokyo 102-8472, Japan

Abstract—Because of its high oxygen demands, neural tissue is predisposed to oxidative stress. Here, our aim was to clarify the cellular localization of antioxidant enzymes in the trigeminal ganglion. We found that the transcriptional factor Sox10 is localized exclusively in satellite glial cells (SGCs) in the adult trigeminal ganglion. The use of transgenic mice that express the fluorescent protein Venus under the Sox10 promoter enabled us to distinguish between neurons and SGCs. Although both superoxide dismutases 1 and 2 were present in the neurons, only superoxide dismutase 1 was identified in SGCs. The enzymes relevant to hydrogen peroxide degradation displayed differential cellular localization, such that neurons were endowed with glutathione peroxidase 1 and thioredoxin 2, and catalase and thioredoxin 2 were present in SGCs. Our immunohistochemical finding showed that only SGCs were labeled by the oxidative damage marker 8-hydroxy-2'-deoxyguanosine, which indicates that the antioxidant systems of SGCs were less

potent. The transient receptor potential vanilloid subfamily member 1 (TRPV1), the capsaicin receptor, is implicated in inflammatory hyperalgesia, and we demonstrated that topical capsaicin application causes short-lasting mechanical hyperalgesia in the face. Our cell-based assay revealed that TRPV1 agonist stimulation in the presence of TRPV1 overexpression caused reactive oxygen species-mediated caspase-3 activation. Moreover, capsaicin induced the cellular demise of primary TRPV1-positive trigeminal ganglion neurons in a dose-dependent manner, and this effect was inhibited by a free radical scavenger and a pancaspase inhibitor. This study delineates the localization of antioxidative stress-related enzymes in the trigeminal ganglion and reveals the importance of the pivotal role of reactive oxygen species in the TRPV1-mediated caspase-dependent cell death of trigeminal ganglion neurons. Therapeutic measures for antioxidative stress should be taken to prevent damage to trigeminal primary sensory neurons in inflammatory pain disorders. © 2013 IBRO. Published by Elsevier Ltd. All rights reserved.

Key words: trigeminal ganglion, glutathione peroxidase, catalase, transient receptor potential vanilloid subfamily member 1, caspase, inflammatory pain.

INTRODUCTION

Reactive oxygen species (ROS) are defined as oxygen-containing chemicals that are damaging to cellular components, such as lipids, proteins and nucleic acids. Oxidative stress stems from an imbalance in the generation and disposal of ROS, which leads to an elevated intracellular concentration of ROS and also leads to oxidative cell damage (Dringen et al., 2005). A superoxide anion ($O_2^{\cdot-}$) is generated, primarily by the mitochondrial respiratory chain and by cellular oxidases. The superoxide anion is quickly converted to hydrogen peroxide (H_2O_2) and oxygen by mitochondrial superoxide dismutase 1 (SOD1, Mn-SOD) and cytoplasmic SOD2 (Cu/Zn-SOD). H_2O_2 is degraded by several peroxide-decomposing enzyme systems, which include catalase, glutathione peroxidase (GPx) and the thioredoxin (Trx)/peroxiredoxin system (Salvemini et al., 2011). Of these systems, GPx1, a member of the GPx family, is known to play a major role in peroxide degradation in neural cells (Mitoyo et al., 2011). Recent studies indicate that significant analgesic effects in both neuropathic pain (Khalil et al., 1999; Kim et al., 2004, 2010; Mao et al., 2009; Yowtak et al., 2011) and inflammatory pain (Thiemermann, 2003; Mika et al., 2007; Kuhad et al., 2008) are produced by the removal

*Corresponding author. Tel: +81-3-5363-3788; fax: +81-3-3353-1272.

E-mail address: mshibata@a7.keio.jp (M. Shibata).

Abbreviations: 8-OHdG, 8-hydroxy-2'-deoxyguanosine; ANOVA, analysis of variance; BAC, bacterial artificial chromosome; DMEM, Dulbecco's modified Eagle medium; EGFP, enhanced green fluorescent protein; flTRPV1, full-length TRPV1; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; GLAST, glutamate/aspartate transporter; GPx, glutathione peroxidase; GST, glutathione S-transferase; HIV, human immunodeficiency virus; H_2O_2 , hydrogen peroxide; HRP, horse radish peroxidase; JNK, c-Jun N-terminal kinase; $O_2^{\cdot-}$, superoxide anion; PBS, phosphate-buffered saline; Prx, peroxiredoxin; ROS, reactive oxygen species; RT-PCR, reverse transcription-polymerase chain reaction; SGCs, satellite glial cells; SOD, superoxide dismutase; STS, staurosporine; TBS-T, Tris-buffered saline/0.1% Tween-20; TEMPOL, 4-hydroxy-2,2,6,6-tetramethylpiperidine; TG, trigeminal ganglion; TRPV1, transient receptor potential vanilloid subfamily member 1; Trx, thioredoxin.

of excess ROS by free radical scavengers. Either systemic or intrathecal administration of ROS scavengers assuaged hyperalgesia that was induced by a formalin injection (Hacimuftuoglu et al., 2006) and mechanical allodynia in neuropathic pain models (Khalil et al., 1999; Kim et al., 2004), implying that the site of algogenic ROS production was within the peripheral nervous system (Kim et al., 2004). The trigeminal ganglia (TG) harbor trigeminal primary neurons, and intraganglionic modifications of nociceptive signals have been described (Rappaport and Devor, 1994; Ceruti et al., 2011). In addition to primary sensory neurons, many specialized glial cells, termed satellite glial cells (SGCs), are present in TG (Hanani, 2010). Similar to TG neurons and Schwann cells, SGCs are derived from the neural crest (Maro et al., 2004). Morphologically, SGCs surround the neurons and form an envelope-like structure that ensheaths the neurons. Because of their small size and the relative paucity of specific markers, not much attention has been paid to these glial cells until recently. Nevertheless, evidence is now accumulating that shows that SGCs play important modulatory roles in the development of pain disorders (Takeda et al., 2009; Katagiri et al., 2012).

The capsaicin receptor, TRPV1 (transient receptor vanilloid subfamily, member 1), is a non-selective cation channel and serves as a transducer of various stimuli into pain signals in the nociceptors (Tominaga et al., 1998; Caterina et al., 2000; Khairatkar-Joshi and Szallasi, 2009). TRPV1 function is upregulated by inflammatory mediators using posttranslational mechanisms, which are critically implicated in the development of inflammatory hyperalgesia (Davis et al., 2000; Khairatkar-Joshi and Szallasi, 2009; Campribi-Robles et al., 2009). In addition, excess TRPV1 activation has been shown to result in the activation of apoptosis-related cysteine proteases (called caspases) and cell death (Shin et al., 2003; Jin et al., 2005a; Amantini et al., 2007; Czaja et al., 2008).

In the present study, we show that Sox10, a transcription factor that is relevant to the development of neural crest-derived cells (Britsch et al., 2001), is a specific marker of SGCs in the adult TG and that antioxidant enzymes exhibit differential cellular localization in TG neurons and SGCs. Moreover, our cell-based studies demonstrate that excess TRPV1 activation is toxic to TRPV1-positive TG neurons because of ROS production and caspase-3 activation.

EXPERIMENTAL PROCEDURES

Animals

The *Sox10-Venus* bacterial artificial chromosome (BAC) transgenic mouse is described elsewhere (Shibata et al., 2010). Sox10 is a marker protein that is closely related to neural crest-lineage cells (Britsch et al., 2001). In the transgenic mouse, the fluorescent protein, Venus (Nagai et al., 2002), is expressed under the control of the *Sox10* promoter. As such, the generation of this transgenic mouse rendered the identification of Sox10-expressing cells feasible without immunostaining.

Male *Sox10-Venus* transgenic mice ($n = 8$; body weight, 20–25 g) were used for immunohistochemistry. Newborn rats ($n = 25$, postnatal day 2–3) were used for the preparation of primary TG cultures. All animal procedures were approved by the Laboratory Animal Care and Use Committee of our institution and were performed in accordance with the guidelines for the Care and Use of Laboratory Animals of the Keio University School of Medicine (No. 08076). All surgeries and animal care were undertaken with the utmost caution to minimize the suffering of the animals.

Immunohistochemistry

Under deep anesthesia by excess pentobarbital sodium (Somnopentyl, Schering-Plough Animal Health Corp., Summit, NJ, USA), the animals were transcardially perfused with 4% paraformaldehyde in 0.1 M phosphate buffer, pH 7.0. Immediately after the perfusion fixation, TG were dissected out and immersed in the same fixative for 4 h at 4 °C and were then kept in 0.01 M phosphate-buffered saline (PBS) solution containing 20% sucrose (w/v) for 12 h for cryoprotection. Subsequently, the TGs were embedded in Tissue TEK (Sakura Finetek, Torrance, CA, USA) and were frozen in liquid nitrogen. Serial sections of 8- μ m thickness were prepared on a cryostat (Reichert-Jung Cryocut 1800; Leica Microsystems, Mannheim, Germany) in the horizontal plane along the long axis. Every 15th section was thaw-mounted on MAS-GP micro slide glass (Matsunami, Osaka, Japan) and air-dried overnight at room temperature.

The sections were pre-incubated with 10% normal donkey serum/0.1 M phosphate buffer for 30 min for blocking. They were incubated with specific primary antibodies for 48 h at room temperature. After they were washed with 0.01 M PBS, the sections were incubated with species-specific fluorophore-labeled secondary antibodies for 2 h at room temperature. After rinsing with 0.01 M PBS, the sections were coverslipped in mounting medium (buffered glycerol: pH 8.6). The primary antibodies and dilutions used in the procedure were as follows: anti-glutamate/aspartate transporter (GLAST) polyclonal antibody (raised in guinea pig; code, AB1782; Millipore, Billerica, MA, USA; 1:200) (Benediktsson et al., 2012), anti-glutamine synthase monoclonal antibody (raised in mouse; code, MAB302; Millipore; 1:500) (Marin-Valencia et al., 2012), anti-glutathione S-transferase-pi (GST- π) polyclonal antibody (raised in rabbit; code, 311; Medical & Biological Laboratories Co., Ltd., Nagoya, Japan; 1:200), anti-tubulin, β III isoform (raised in mouse; code MAB1637; Millipore), anti-catalase polyclonal antibody (raised in rabbit; code, ab52477; Abcam, Cambridge, MA, USA; 1:200), anti-Trx2 polyclonal antibody (raised in rabbit; sc-50336; Santa Cruz Biotechnology, Santa Cruz, CA, USA; 1:50) (McCommis et al., 2011), anti-SOD1 (Cu/Zn Enzyme) polyclonal antibody (raised in sheep; code, 574597; Calbiochem, Darmstadt, Germany; 1:200), anti-SOD2 (Mn Enzyme) polyclonal antibody (raised in sheep; code, 574596; Calbiochem, 1:200), and anti-GPx1 polyclonal antibody (raised in goat; code, AF3798; R&D

Systems, Minneapolis, MN, USA, 1:100). Then, 8-hydroxy-2'-deoxyguanosine (8-OHdG) immunolabeling was accomplished using the anti-8-OHdG monoclonal antibody (raised in mouse; code, MOG-020P; Japan institute for the Control of Aging, Tokyo, Japan; 1:200) (Toyokuni et al., 1997), with the aid of a staining kit (MOM; Vector Laboratories, Burlingame, CA, USA), the latter of which was specifically designed to localize mouse primary antibodies in mouse tissues and was used according to the manufacturer's instructions. The species-specific secondary antibodies were raised in donkeys and were conjugated to Cy3, Dylight 649, or Alexa Fluor 647; all of them were obtained from Jackson ImmunoResearch Laboratories (West Grove, PA, USA). The immunolabeled specimens were examined under a Leica TCS-SP5 confocal laser scanning microscopy (Leica Microsystems). In order to quantify SOD1 and SOD2 immunoreactivity, we counted SOD1- and SOD2-positive cells in randomly selected 300 neurons or SGCs. For the determination of GST- π -positive TG neurons, we measured neuronal soma diameters using image analysis software (Keyence, Osaka, Japan). The proportion of GST- π -positive cells in total TG neurons ($n = 47$) by diameter size was evaluated.

Behavioral testing for assessment of head withdrawal responses

A group of male Sprague–Dawley (SD) rats ($n = 6$) was intraperitoneally injected with 4-hydroxy-2,2,6,6-tetramethyl piperidine (TEMPOL) (Sigma, St. Louis, MO) dissolved in saline (60 mg/ml, 300 mg/kg). Another set of male SD rats ($n = 6$) was intraperitoneally injected with the same volume of saline (Schwartz et al., 2008). After 10 min, mechanical sensitivity of the whisker pad was evaluated by the amount of positive head withdrawals in response to von Frey stimuli. A von Frey monofilament (3.5 g = 34.3 mN) was applied perpendicular to the stimulation site with sufficient force to bend the filament for 1 s and then slowly removed. In each session of trials, von Frey stimulation was applied 10 times, and the number of head withdrawals was counted. To assess the role of ROS in the induction and maintenance of capsaicin-induced facial hyperalgesia, we applied a piece of square filter paper soaked with capsaicin (10 mM in DMSO) (Sigma) to the left whisker pad for 30 min under light anesthesia with sodium pentobarbital (40 mg/kg, i.p.) 30 min after the pretreatment with either TEMPOL or saline (Honda et al., 2008). For reference, DMSO-soaked square filter paper was applied to the right whisker pad. After 30 min, the filter paper was removed, and the development of flare in the skin treated with capsaicin was recognized. During the procedure, body temperature was maintained at 37 °C using a rectal thermistor-controlled heating pad. Two hours after the removal of filter paper, we confirmed that anesthesia had worn off in all the rats, and mechanical sensitivity was assessed by applying a von Frey monofilament to the site of the flared skin on the side of capsaicin application and the corresponding site on the

contralateral side. Subsequently, the assessment of mechanical sensitivity was repeated at 6 and 24 h after the capsaicin treatment. All the animals had been habituated to the monofilament stimulation for the last 7 days before the behavioral testing. The statistical analysis was performed using a two-way ANOVA with one repeated factor (time) and *t*-test for between-group difference. $p < 0.05$ was considered statistically significant.

Stable transformants expressing an enhanced GFP-full-length rat TRPV1 fusion protein

Total RNA was prepared from the TG of an adult male Sprague–Dawley rat using TRIZOL LS Reagent (Invitrogen, Carlsbad, CA, USA). From an aliquot of 1 μ g of total RNA, cDNA was synthesized employing the SuperScript III RT First-Strand Synthesis System (Invitrogen) according to the manufacturer's instructions. Full-length *TRPV1* cDNA was amplified by PCR using a set of sequence-specific primers (forward; 5'-ggaattctg gaacaacgggctagcttagac-3', reverse; 5'-gggttacctatttct cccctgggacat-3'). The full-length *TRPV1* cDNA PCR fragment was subcloned into the pEGFP-C3 plasmid (Clontech, Mountain View, CA, USA), which had a G418 resistance gene expression cassette, and the resultant vector was designed to express an enhanced green fluorescent protein (EGFP)-full length rat TRPV1 fusion protein in mammalian cells. The integrity of the vector was confirmed by DNA sequencing. The vector was transfected into the rat pheochromocytoma-derived cell line, PC12, using the Lipofectamine 2000 Transfection Reagent. PC12 cells were grown in Dulbecco's modified Eagle's medium (DMEM, Invitrogen) supplemented with 15% horse serum, 2.5% fetal bovine serum, 50 units penicillin and 50- μ g streptomycin. Successfully transfected cells were identified by GFP signals under fluorescence microscopy. Stable transformants were selected by continuously adding G418 (final concentration, 700 μ g/ml) to the medium and were grown until clonal colonies could be isolated.

Capsaicin treatment and Western blot analysis

Capsaicin (Sigma), the TRPV1 antagonist, capsazepine (Sigma), and the cell permeant ROS scavenger, TEMPOL (Sigma), were dissolved in dimethylsulfoxide (DMSO) at 100 mM, 50 mM, and 1 M, respectively. PC12 cells were then divided into several groups. The control cell group was treated with vehicle (0.1% DMSO) for 30 min, and the capsaicin-treated cell groups were exposed to capsaicin at concentrations of 1, 10 or 100 μ M for 30 min. In separate groups, cells were treated with 100 μ M capsaicin for 30 min, which was accompanied by capsazepine (50 μ M) or the free radical scavenger, TEMPOL (0.5 M or 1 M), with treatment starting 30 min prior to and continuing throughout the capsaicin treatment. Following the pharmacological treatment, the cells were harvested, and cell lysates were prepared using Laemmli sample buffer (Biorad, Hercules, CA, USA). The samples were subjected to SDS-PAGE, and proteins were transferred

to PVDF membranes (Immobilon-P, Millipore). The membranes were incubated with primary antibodies overnight at 4 °C. After rinsing with Tris-buffered saline/0.1% Tween-20 (TBS-T), the blots were then incubated with species-specific horse radish peroxidase (HRP)-conjugated secondary antibodies for two hours at room temperature. Immunoreactive bands were visualized by the enhanced chemiluminescence reaction with Western Lightning-ECL Plus (PerkinElmer, Waltham, MA, USA), and images were captured with a CCD camera connected to the ImageQuant LAS 4000 system (Fuji Film, Tokyo, Japan). The primary antibodies and dilutions used in this study were as follows: anti-TRPV1 (TransGenic, Kumamoto, Japan, 1:1000), anti-GFP (MBL, Nagoya, Japan, 1:1000), anti-phospho-c-Jun N-terminal kinase (phospho-JNK) (Cell Signaling Technology, Danvers, MA, 1:1000), anti-total c-Jun N-terminal kinase (total JNK) (Santa Cruz Biotechnology, Santa Cruz, CA, 1:1000), anti-active caspase-3 (Cell Signaling, 1:1000), anti- α -tubulin (Sigma, 1:100,000) and anti-GAPDH (Cell Signaling, 1:1000). All of the secondary antibodies were purchased from Jackson ImmunoResearch (West Grove, PA, USA). Densitometric analysis of immunoreactive bands for phospho-JNK and total JNK was performed using Multigauge software ver. 3.3 (Fuji Film, Tokyo, Japan). The ratio of phospho-JNK to total JNK (phospho-JNK/total JNK) was calculated in each sample. Data shown represent the mean \pm SD. The fold change of the ratio of phospho-JNK/total JNK from the corresponding ratio in the vehicle sample was evaluated by employing a one-way analysis of variance (ANOVA) and Bonferroni's post hoc test. We used IBM SPSS, ver. 19 (Chicago, IL) for statistical analysis. Statistical significance was set at $p < 0.05$.

Reverse transcription-polymerase chain reaction for TRPV1 and GAPDH

For a rat TG tissue sample, RNA extraction and reverse transcription-polymerase chain reaction (RT-PCR) for *TRPV1* were performed as described in the Section 4. For PC12 cells, approximately 1×10^7 cells grown on a culture dish were collected as a pellet by centrifugation. RNA extraction and RT-PCR for *TRPV1* were conducted as described above. The primers for the *TRPV1* transcript were designed to amplify both endogenous and recombinant rat full-length *TRPV1* cDNA. For the internal control of the RT-PCR, we amplified a rat *GAPDH* cDNA fragment using a set of sequence-specific primers: forward, 5'-atggtgaaggtcgggtgaacgga-3', and reverse, 5'-acgatgccaagttgcatggatg-3'. PCR products were electrophoresed in 1% agarose gels.

TRPV1-positive neuron survival assay using rat primary TG cultures

Rat primary TG cultures were prepared from neonatal Sprague–Dawley rats (2–3 days old). Briefly, after dissection, TG tissue samples were minced in ice-cold HBSS (Hank's balanced saline solution; Invitrogen).

After centrifugation, the pellet was resuspended in Leibovitz L15 (Invitrogen) containing 250 U/ml collagenase (Worthington, Lakewood, NJ, USA), 1 U/ml elastase (Worthington), and 5 U/ml papain (Worthington), and the suspension was incubated for 1.5 h at 37 °C. Cells were grown in DMEM/F12 (Invitrogen) supplemented with 10% FBS, 50 ng/ml NGF (Sigma), and 1% penicillin/streptomycin (Invitrogen). At Day 2 of culture, the cells were treated with different concentrations (100 nM, 1 μ M and 10 μ M) of capsaicin for 30 min. In parallel, for 10 μ M capsaicin, treatment with either TEMPOL or the pancapsase inhibitor, zVAD, 30 min prior to and continuing capsaicin treatment was performed. After the pharmacological interventions, the cultured cells were fixed with 2% paraformaldehyde/PBS and immunostained with anti-TRPV1 antibody (raised in a rabbit; code KM 018; TransGenic, Kumamoto, Japan; 1:200). Immunofluorescent images were randomly captured using a fluorescence microscope. The number of TRPV1-positive neurons in each treatment group was counted and expressed as the mean \pm SD% of that in the vehicle-treated control culture. Statistical analysis was performed by a one-way ANOVA and Bonferroni's post hoc test using IBM SPSS, ver. 19. Statistical significance was set at $p < 0.05$.

RESULTS

Sox10-Venus expression in TG

We examined the distribution of Venus expression in TG of adult *Sox10-Venus* BAC transgenic mice. In the TG cell architecture, the Venus-derived fluorescence signal was localized exclusively to small cells that ensheathed TG neurons. Meanwhile, TG neurons were devoid of Venus signals. Judging from the morphology and distribution of the Venus-positive cells, we surmised that they were SGCs. To confirm this observation, we conducted immunostaining for two established markers for SGCs in TG: glutamine synthase and GLAST. There was clear co-localization of Venus with both markers (Fig. 1), indicating that Sox10 expression is restricted to SGCs in adult mouse TG. There was no difference in the distribution pattern of glutamine synthase in TG between the wild-type and *Sox10-Venus* BAC transgenic mouse (Fig. 1E).

Expression of proteins related to reactive oxygen species catabolism in TG

As described above, the *Sox10-Venus* BAC transgenic mouse proved to be a useful tool for identifying SGCs in TG. We performed an immunohistochemical screening for protein expression in SGCs by employing this transgenic mouse, and we found that immunoreactivity for the oxidative stress marker, 8-OHdG, was prominent in SGCs (Fig. 2). Hence, this observation prompted us to study the localization of antioxidant enzymes in TG. Because 8-OHdG immunoreactivity was not observed in neurons, we reasoned that SGCs were subjected to more oxidative stress compared to neurons. We then




Fig. 1. Co-localization of the Venus signal with the SGC markers, glutamine synthase and GLAST, in *Sox10-Venus* BAC transgenic mouse TG. (A) Sox10 (Venus), (B) glutamine synthase, (C) merged image, (D) DAPI (nuclear staining), Scale bars = 20 μm , (E) glutamine synthase in wild-type mouse TG, Scale bar = 20 μm , (F) Sox10, (G) GLAST, (H) DAPI, (I) merged image, Scale bar = 25 μm .




Fig. 2. The oxidative marker, 8-OHdG, was recognized in Sox10-expressing SGCs in TG. (A) 8-OHdG, (B) Sox10 (Venus), (C) merged image. Scale bars = 25 μm .

conducted single immunohistochemical labeling experiments using antibodies against SOD1 and 2 (Fig. 3), GPx1 (Fig. 4A–C), catalase (Fig. 4D–F), Trx2 (Fig. 4G–I) and GST- π (Fig. 5). SOD 1 and 2 displayed distinct cellular localizations, such that SOD1 was present in both neurons and SGCs, whereas SOD2

was localized exclusively in neurons (Fig. 3). Quantitatively, SOD1 immunoreactivity was recognized in 96.3% of TG neurons ($n = 300$) and 41.0% of SGCs ($n = 300$), whereas SOD2 immunoreactivity was identified in 94% of TG neurons ($n = 300$). GPx1 was identified only in neurons (Fig. 4A–C). In contrast, catalase expression was restricted to SGCs (Fig. 4D–F). Meanwhile, Trx 2 was present in both neurons and SGCs (Fig. 4G–I). Immunoreactivity for GST- π was discernible primarily in SGCs (Fig. 5A–C, arrows). However, weak immunoreactivity was also present in a subset of small- to medium-sized (diameter < 35 μm) neurons (Fig. 5A–C, arrowheads; Table 1 for quantification), which were positive for the neuronal marker, β III-tubulin (Fig. 5D–H).

ROS is involved in capsaicin-induced mechanical hyperalgesia in the trigeminal territory

Mechanical sensitivity of the rat whisker pad was assessed by the amount of positive head withdrawals in




Fig. 3. Localization of SOD1 and 2 in TG. (A) SOD1 (left), Sox10 (middle), merged image (right). Arrowheads indicate SOD1-positive SGCs. (B) SOD2 (left), Sox10 (middle), merged image (right). Scale bars = 25 μ m. The images were obtained from a TG section of the *Sox10-Venus* BAC transgenic mouse.




Fig. 4. Cellular localization of GPx1, catalase and Trx2 in TG. (A) GPx1, (B) Sox10, (C) merged image, (D) catalase, (E) Sox10, (F) merged image, (G) Trx2, (H) Sox10, (I) merged image. Scale bars = 25 μ m. The images were obtained from a TG section of the *Sox10-Venus* transgenic mouse.




Fig. 5. Localization of GST- π in TG. (A) GST- π , (B) Sox10, (C) merged image. Immunoreactivity for GST- π is recognized in Sox10-positive SGCs (arrows). Small- to middle-sized neurons exhibit weak immunoreactivity for GST- π (asterisk). Scale bars = 25 μ m. (D) β III tubulin, (E) GST- π , (F) Sox10, (G) merged image, (H) DAPI. Scale bar = 15 μ m. The two sets of images (A–C and D–H) were obtained from two distinct TG sections of the Sox10-Venus transgenic mouse.

Table 1. Proportion of GST- π -positive cells in TG neurons by diameter size. A total of 47 TG neurons were analyzed

Diameter (μ m)	Number	GST- π -positive neurons (%)
10–15	5	60.0
16–20	12	33.3
21–25	13	69.2
26–30	9	33.3
31–35	4	25.0
36–40	3	0
41–45	1	0

response to von Frey monofilament stimulation. Topical capsaicin application to the whisker pad induced mechanical hyperalgesia, which was evidenced by increased frequency of head withdrawals. The mechanical hyperalgesia was most prominent at 2 h after capsaicin treatment (1.8 ± 1.8 at pretreatment to 5.7 ± 4.1 at 2 h, $p = 0.026$, paired t -test, $n = 6$ in each group, Fig. 6). With treatment with TEMPOL, a well-established ROS scavenger, the mechanical hyperalgesia was attenuated (5.7 ± 4.1 in the TEMPOL-treated group vs. 1.6 ± 2.4 in the saline-treated group, $p = 0.044$, unpaired t -test, $n = 6$ in each group).

Capsaicin treatment induces ROS-mediated caspase-3 activation in high TRPV1-expressing PC12 cells

Several PC12 cell lines that stably expressed an EGFP-full-length TRPV1 fusion protein (EGFP-fTRPV1) were obtained. In most of these cell lines, Western blot

analysis identified a distinct band that was immunoreactive for both TRPV1 and GFP at the molecular weight of 115 kDa, which was consistent with the expected molecular weight of the EGFP-fTRPV1 fusion protein (Fig. 7A, lanes 2 and 3). Incidentally, we obtained a PC12 cell line that failed to express the fusion protein in spite of acquiring resistance to G418 after the transfection procedure (Fig. 7A, lane 1). This cell line was designated as the EGFP-fTRPV1-null cell line. In this cell line, there was no discernible endogenous TRPV1 protein expression by Western blot. Because RT-PCR analysis revealed the presence of *TRPV1* mRNA, it seemed likely that there was translational down-regulation of the *TRPV1* transcript (Fig. 7B, lane 2). JNK phosphorylation is known to be an important step in the signal transduction pathway activated by ROS exposure (McCubrey et al., 2006). Hence, we assayed JNK phosphorylation in cells exposed to capsaicin treatment. There was a robust increase in JNK phosphorylation in the high EGFP-fTRPV1-expressing cell line after the capsaicin treatment in a dose-dependent manner (for p54, 26.81 ± 0.61 -fold at 10 μ M and 52.09 ± 2.97 -fold at 100 μ M compared to the vehicle-treated cells, $p < 0.001$; for p46, 35.50 ± 1.07 -fold at 10 μ M and 64.90 ± 3.17 -fold at 100 μ M compared to the vehicle-treated cells, $p < 0.001$) (Fig. 8A, B). Capsaicin treatment at 100 μ M beyond 30 min was found to cause JNK protein degradation (data not shown). Therefore, we reasoned that capsaicin treatment at 100 μ M for 30 min was the




Fig. 6. Capsaicin-induced nocifensive head withdrawals. The numbers of head withdrawals in response to von Frey stimulation were counted on the capsaicin (10 mM)-treated side (A) and vehicle (DMSO)-treated side. To explore the involvement of ROS production, TEMPOL (300 mg/kg) was intraperitoneally injected 30 min before the capsaicin application in six animals (dotted lines). For controls, the same volume of saline was intraperitoneally injected in six animals (solid lines). The data are expressed as the mean \pm SD. The interaction between the time course and head withdrawal numbers was assessed by a two-way ANOVA. Comparison within the same group between two time points was made by paired *t*-test. Between group comparisons were performed by unpaired *t*-test. #*p* < 0.05 vs. the pretreatment group (Pre), **p* < 0.05 between the saline- and TEMPOL-treated groups.




Fig. 7. Characterization of EGFP-fTRPV1-expressing PC12 cell lines by Western blot and RT-PCR. (A) Western blot data for TRPV1 (left) and GFP (right) expression. Lane 1: EGFP-fTRPV1-null cell line, Lane 2: moderate EGFP-fTRPV1-expressing cell line, Lane 3: high EGFP-fTRPV1-expressing cell line. The arrow indicates immunoreactive bands that represent the EGFP-fTRPV1 fusion protein. (B) RT-PCR for rat *TRPV1* and *GAPDH* transcripts. Lane 1: negative control, Lane 2: EGFP-fTRPV1-null cell line, Lane 3: recombinant rat full-length *TRPV1* cDNA. Lane 4: rat TG. M, markers.

optimal condition for examining capsaicin-induced JNK phosphorylation in our experimental conditions. JNK phosphorylation was lessened by 100 μ M capsazepine treatment (for p54, 55.64 ± 3.98 -fold at 100 μ M CAPS vs. 24.58 ± 1.78 -fold at 100 μ M CAPS + 100 μ M capsazepine compared to the vehicle-treated cells, *p* < 0.001; for p46, 75.54 ± 11.19 -fold at 100 μ M CAPS vs. 20.52 ± 2.14 -fold at 100 μ M CAPS + 100 μ M capsazepine compared to the vehicle-treated cells, *p* < 0.001) (Fig. 8C), indicating that the capsaicin-induced JNK phosphorylation was a TRPV1 ligand-dependent process. TEMPOL attenuated capsaicin-induced JNK phosphorylation in a dose-dependent manner (Fig. 8C). Meanwhile, the EGFP-fTRPV1-null cell line did not exhibit any significant alteration in JNK phosphorylation (Fig. 8A, B). Several lines of evidence show that ROS overproduction induces caspase-3-mediated cell death (Franklin, 2011). Exposure of the high EGFP-fTRPV1-expressing cell line to 100 μ M capsaicin induced cell rounding and a detachment of cells from the culture dish, starting at as early as 30 min; these changes are morphological changes that are indicative of cell demise (Fig. 9A). These morphological alterations were prevented by capsazepine. Next, we examined caspase-3 activation in the high EGFP-fTRPV1-expressing cell line. Our preliminary experiments showed that 3 h of capsaicin treatment is the most appropriate time point for examining the activation of caspase-3 (data not shown). Caspase-3 activation became obvious at this time point in a dose-dependent manner (Fig. 9B). Such capsaicin-induced caspase-3 activation was abolished by TEMPOL, indicating that ROS production was implicated in this process (Fig. 9C).

Capsaicin treatment induces caspase- and ROS-mediated cell demise of TRPV1-positive TG neurons

To assess the neurotoxic effects of capsaicin on TG neurons, we applied capsaicin to primary TG cultures and measured the number of remaining TRPV1-positive TG neurons. Capsaicin induced a dose-dependent loss of TRPV1-positive neurons (Fig. 10). The capsaicin-induced neuronal loss was prevented by TEMPOL and z-VAD pretreatment.

DISCUSSION

Sox10 expression in adult TG

Sox10 is a transcriptional factor that plays a pivotal role in the development of neural crest-derived cells. In DRG, Sox10 expression is primarily recognized in neurons at embryonic day 11.5, but neurons have ceased to express Sox10 by embryonic day 15.5 (Shibata et al., 2010). However, little is known about Sox10 expression in TG, especially in adulthood. To our knowledge, the present study provides the first evidence that Sox10 is expressed exclusively in SGCs in adult TG. In addition to the previous observation that spontaneous or targeted disruption of Sox10 resulted in a loss of




Fig. 8. Capsaicin enhances JNK phosphorylation via TRPV1 activation in the high EGFP-fTRPV1-expressing cell line. (A) Western blot data for phosphorylated (phospho-JNK) and total JNK. JNK is identified as two distinct bands that represent p54 and p46. High: high EGFP-fTRPV1-expressing cell line, Null: EGFP-fTRPV1-null cell line, Veh: vehicle (0.1% DMSO), CAPS: capsaicin. The duration of capsaicin exposure was 30 min. (B) Densitometric data for phospho-JNK/total JNK (separately for p54 and p46) ratio against the vehicle sample (Veh). The data are expressed as the mean \pm SD ($n = 4$ in each group). * $p < 0.001$, versus Veh, a one-way ANOVA, Bonferroni's post hoc test. (C) Effects of capsazepine and TEMPOL on capsaicin-induced JNK phosphorylation in high EGFP-fTRPV1-expressing cell line. Treatment with either capsazepine (50 μ M) or the free radical scavenger, TEMPOL (0.5 M or 1 M), was started 30 min prior to capsaicin treatment and was continued throughout the capsaicin treatment. CAPS, capsaicin; +CZP, capsaicin + capsazepine.

Schwann cells and satellite cells in DRG (Britsch et al., 2001), our finding suggests that Sox10 is important in the development and maintenance of the SGCs in neural crest-derived ganglia.

Cellular distribution of antioxidant enzymes in TG

Our immunohistochemical analysis revealed that antioxidant enzymes are differentially distributed in TG. Both SOD1 and 2 are localized in neurons, whereas




Fig. 9. Capsaicin treatment leads to caspase-3 activation via ROS production in the high EGFP-TRPV1-expressing cell line. (A) Micrographs showing cell morphology treated with the vehicle (0.1% DMSO, Veh), capsaicin (100 μM, CAPS) or capsaicin (100 μM) + capsazepine (100 μM) (+CZP). The duration of capsaicin treatment was 30 min, with capsazepine treatment starting 30 min prior to and continuing throughout capsaicin treatment. (B) Western blot data for active (cleaved) caspase-3. Staurosporine (STS) was used as a positive control agent for activation caspase-3. The blot is representative of 3 independent studies. Veh, vehicle (0.1% DMSO); CAPS, capsaicin; STS, staurosporine (2.5 μM). The duration of staurosporine treatment was 16 h. (C) Effect of TEMPOL pretreatment on capsaicin-induced caspase-3 activation. The duration of capsaicin treatment was 3 h, and TEMPOL administration was started 30 min prior to capsaicin treatment, and was continued throughout capsaicin treatment. Veh, vehicle (0.1% DMSO); CAPS, capsaicin (100 μM); +TEMP, capsaicin (100 μM) + TEMPOL (1 M). The blot is representative of three independent studies.




Fig. 10. Capsaicin-induced cell demise of TRPV1-positive TG neurons. The number of TRPV1-positive TG neurons in each group was shown as mean ± SD% of the vehicle-treated cells. * $p < 0.001$, ** $p < 0.01$, # $p < 0.05$, a one-way ANOVA, Bonferroni's post hoc test.

SGCs are provided with SOD1 alone. The enzymes with H_2O_2 -decomposing activity, GPx1 and catalase, displayed differential cellular localization patterns, such that GPx1 and catalase are exclusively distributed in neurons and SGCs, respectively. The predominant localization of GPx1 in neurons over glial cells has also been reported in human brain tissue (Power and Blumbergs, 2009). Recently, the Trx/peroxiredoxin system has been recognized as another important peroxide-removing system in a variety of cellular systems, including neural cells (Patenaude et al., 2005; Bell and Hardingham, 2011). Trx2 is a mitochondrial enzyme that contains two redox-active cysteine residues (Cys-Gly-Pro-Cys) within its catalytic site. These cysteines exist in their -SH form in the reduced state and form an intramolecular disulfide bond in the oxidized state. Trx2 activates the peroxide scavengers, peroxiredoxins, by regulating their redox state (Bell and Hardingham, 2011). The present study showed that Trx2 is localized to both neurons and SGCs. Superoxide anion ($O_2^{\cdot-}$) reacts with nitric oxide to form peroxynitrite, a molecule that is exquisitely detrimental to cells. SOD1 and 2 protect cells from the toxicity of the superoxide anion by converting it to H_2O_2 . H_2O_2 per se is not readily damaging to biological molecules in physiological micromolar concentrations. However, unless it is efficiently eliminated from tissue, hydroxyl radical ($\cdot OH$) formation by the iron-catalyzed Fenton reaction follows. Hydroxyl radical can cause lipid peroxidation, which oxidizes amino acid side chains in proteins and induces aberrant DNA modifications, thus causing cell damage (Dringen et al., 2005). To maintain the integrity of H_2O_2 -removing systems, it is therefore imperative to bypass such oxidative damage, especially in neural tissue with high oxygen demands. Because the cell viability of neurons must be maintained over a lifetime, it appears reasonable that TG neurons are endowed with more thorough antioxidant systems than SGCs. The relative importance of GPx, catalase and peroxiredoxins in neural cells has been a matter of debate (Mitozo et al., 2011). Generally, GPx1 is considered to be the primary peroxide-degrading enzyme, and this view has been corroborated by several *in vivo* experiments. The GPx1 knockout mouse brain exhibited increased apoptosis following ischemia/reperfusion injury and cold injury (Crack et al., 2001; Flentjar et al., 2002) and exacerbated dopaminergic cell damage induced by *N*-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) (Zhang et al., 2000) compared to the wild-type mouse brain. The contribution of the Trx/peroxiredoxin system to the clearance of H_2O_2 in neural tissue is poorly understood. The inactivation of peroxiredoxins by the Trx reductase inhibitor, auranofin, caused DNA damage in astrocytoma-derived C6 cells but not in N2a neuroblastoma cells or in hippocampal tissue slices (Mitozo et al., 2011). Catalase deficiency was found to compromise the NAD-linked electron transfer activities and energy coupling capacities of brain mitochondria (Ho et al., 2004). Moreover, pharmacological catalase inhibition retarded H_2O_2 decomposition in hippocampal slices and N2a neuroblastoma cells (Mitozo et al.,

2011). These findings imply that catalase plays an important role in the clearance of H₂O₂ in brain tissue. H₂O₂ removal by the catalytic activity of catalase follows the pattern of a first-order reaction, and its rate relies on the H₂O₂ concentration in the environment. Given such enzymatic kinetics, catalase appears to be fully functional at removing H₂O₂ at higher concentrations. Furthermore, the contribution of catalase to the control of endogenous H₂O₂ in erythrocytes is negligible (Johnson et al., 2010). In addition, the *ex vivo* analysis using lung slices from wild-type (*Cat*^{+/+}) and *catalase* knockout mice (*Cat*^{+/-} and *Cat*^{-/-}) revealed that the rates of H₂O₂ decomposition were proportional to the levels of catalase that were expressed, when examined at higher H₂O₂ concentrations, whereas the rates did not significantly differ among the genotypes at lower concentrations (Ho et al., 2004). Our study demonstrated that only SGCs were labeled by the oxidative damage marker, 8-OHdG, in TG. This finding is likely to reflect the inefficient H₂O₂-scavenging activity of catalase relative to GPx1 at physiologically low concentrations in SGCs. Alternatively, ROS may be actively involved in physiological signaling functions in SGCs under the normal state. Recent evidence shows that H₂O₂ functions as a second messenger by protein thiol oxidation of key proteins such as protein tyrosine phosphatases (Forman et al., 2010).

GST family members exert cytoprotective actions by catalyzing the conjugation of reactive chemicals with glutathione. Specific isoforms are known to be expressed in distinct cell populations, and GST- π is expressed primarily in oligodendrocytes in the central nervous system. The present study provides the first histological evidence that GST- π is localized in SGCs and in small- to medium-sized neurons (diameter < 35 μ m) in TG. The significance of this finding is unclear. However, because GST family members are critically involved in the maintenance of the cellular antioxidant glutathione in different cellular compartments (Raza, 2011), GST- π might play a compensatory role against the absence of GPx1 in SGCs, which are equipped with less potent antioxidant machinery compared to neurons. Intriguingly, the monomeric form of GST- π is known to bind to the C-terminal of JNK and suppress its kinase activity (Wang et al., 2001). JNK activation is known to lead to the development of cell death (Shen and Liu, 2006). Although we performed the TUNEL method on TG tissue specimens, there was no evidence of apoptotic cell demise (data not shown). In spite of the evidence of oxidative damage in SGCs, they appear healthy. This enigma can be explained by the suppressive effect of GST- π on JNK activation.

TRPV1-mediated ROS production and cell death

TRPV1 activation is known to be sensitized by inflammatory mediators, such as NGF and bradykinin, via its increased expression and/or its phosphorylation, which is mediated by several kinases (Khairatkar-Joshi and Szallasi, 2009). Inflammation-mediated potentiation

of the TRPV1 function is regarded as a mechanism that underlies inflammatory hyperalgesia. TRPV1 agonist stimulation of the high EGFP-flTRPV1-expressing cell line appears to mimic this situation. Under the circumstance, capsaicin treatment led to ROS-mediated caspase-3 activation. Such situations appear to be commonly encountered in meningitis and pulpitis. In addition, recent studies show that ROS are capable of augmenting TRPV1 activity by different mechanisms. NADPH oxidase-derived ROS were shown to promote protein kinase C ϵ (PKC ϵ) translocation from the cytosolic to the membrane fractions, which led to TRPV1 activation by phosphorylation (Ibi et al., 2008). Moreover, covalent modification of conserved cysteines by oxidative stress results in TRPV1 sensitization (Chuang and Lin, 2009). Therefore, it is likely that inflammatory conditions favor the long-lasting enhancement of TRPV1 activation by ROS-mediated posttranslational modifications. In the present study, capsaicin treatment caused cellular demise of TRPV1-positive primary TG neurons in a dose-dependent manner, which indicates that the enhancement of TRPV1 function renders TRPV1-positive TG neurons more predisposed to cell death. Moreover, our pharmacological experiments demonstrated that the TRPV1-induced TG neuron death is mediated by ROS production and caspase activation. Although a previous study implicated ROS production in TRPV1-mediated apoptotic cell death of a nasopharyngeal carcinoma-derived cell line (Ip et al., 2012), our results provide the first evidence for the involvement of ROS in TRPV1-mediated TG neuron death. TRPV1-induced death of TG neurons does not seem to be a phenomenon restricted to an artificial *in vitro* situation because systemic capsaicin administration has been shown to induce the apoptotic cell death of primary sensory neurons in TG, DRG and nodose ganglia (Jin et al., 2005a,b; Czaja et al., 2008). Caspases constitute a family of cysteine proteases that execute apoptosis and that are implicated in a variety of neurological disease conditions, such as stroke and neurodegenerative diseases (Namura et al., 1998; Shibata et al., 2000; Yuan and Yankner, 2000; Friedlander, 2003). More specifically, caspase-3 is known to be a principal executioner caspase. Judging from the pharmacological data obtained with the high EGFP-flTRPV1-expressing cell line, ROS generation can be responsible for the caspase-3 activation. Our finding implies that the efficient elimination of ROS is likely to be a key action for rescuing primary sensory neurons from TRPV1-mediated apoptosis. What is the role of sensory neuronal death in the pathophysiology of pain disorders? Intuitively, the demise of sensory neurons appears to result in the development of sensory loss. However, there is ample evidence that loss of sensory neurons and fibers occurs in pain disorders. Diabetic neuropathic pain is recognized in approximately 10–20% of the entire diabetic population, and in 40–60% of patients with documented neuropathy (Daousi et al., 2004; Abbott et al., 2011). Diabetic neuropathy is pathologically characterized by considerable loss of small-fibers, including C- and A δ -fibers, and oxidative stress is

postulated to play an important role in the nerve damage (Callaghan et al., 2012). Human immunodeficiency virus (HIV)-associated sensory neuropathy presents a big clinical challenge because of its increasing prevalence and its association with debilitating pain. Its viral proteins, Tat and viral protein R, are known to cause hyperexcitability of DRG neurons via excess calcium influx and impaired cyclin-dependent kinase 5 activity, which culminates in apoptosis of DRG neurons (Acharjee et al., 2010; Chi et al., 2011). Burdo et al. (2012) reported pathological features of DRG damage in simian immunodeficiency virus-infected rhesus macaques, including satellitosis, presence of Nageotte nodules, and neuronophagia as well as macrophage infiltration and abundant viral replication. Collectively, it is likely that neurodegenerative processes, which eventually lead to sensory neuronal death, can be responsible for the development of painful conditions. We surmise that the capsaicin-induced demise of TRPV1-expressing sensory neurons, which was found to be mediated by ROS production, have significant implications in the development of neuropathic pain.

Clinical implications

Our behavioral testing revealed that ROS is involved in the development of mechanical hyperalgesia in the trigeminal territory. Although mechanical hyperalgesia driven by ROS has been demonstrated in several models in rats, there seems to be controversy about the exact site of ROS production relevant to the development of mechanical hyperalgesia. Systemic and intrathecal administration of ROS scavengers showed similar efficacy against hyperalgesia induced by formalin injection and xanthine oxidase activity was elevated in the sciatic nerve in a chronic nerve constriction model, which supports peripheral ROS production (Khalil et al., 1999; Hacimuftuoglu et al., 2006). Meanwhile, some authors claim the importance of spinal involvement mainly because secondary, but not primary, capsaicin-induced hyperalgesia can be attenuated by ROS scavengers (Lee et al., 2007; Schwartz et al., 2008). As we administered the ROS scavenger, TEMPOL, intraperitoneally, we cannot determine where the agent exerted its action. Nevertheless, our cell experiments using TRPV1-expressing PC12 cell lines demonstrated the cell autonomous ROS production in response to capsaicin application and, more importantly, ROS was found to be involved in the capsaicin-induced demise of TRPV1-expressing neurons in the primary TG culture. These findings seem to lend support to the peripheral action of TEMPOL with respect to its antihyperalgesic effect.

The capsaicin-induced mechanical hyperalgesia was recognized at 2 h after capsaicin application, and face sensitivity then returned to normal levels. Such transient nature of mechanical hyperalgesia implies the presence of tight regulation of ROS levels by scavenging enzymes in the trigeminal system. Our primary culture data show that TRPV1-expressing neurons are vulnerable to capsaicin treatment in the culture condition and that capsaicin-induced TRPV1-expressing neuronal death is

mediated by ROS production. Although TG neurons possess the potent antioxidant system, the culture conditions are quite different from *in vivo* situations. The cultured TG neurons are not in close contact with SGCs, and, because of the lack of elongated axons, TRPV1 stimulation occurs in the proximity to the neuronal soma. In this sense, the capsaicin stimulation in our primary culture experiments may mimic the situation in which endogenous TRPV1 agonists, such as protons and anandamide, stimulate TRPV1 in the activated state as seen in the inflammatory conditions. The trigeminal nociceptors are not infrequently involved in inflammatory disease conditions, such as meningitis, pulpitis and migraine (Moskowitz, 1984). Under these circumstances, TRPV1-mediated ROS overproduction could surpass the antioxidative capacity of trigeminal primary neurons and could contribute to compromising their viability, possibly by caspase activation. Concomitantly, the diffusion of H₂O₂ from the initially affected neurons is expected to be detrimental to the surrounding neurons. Such propagation of H₂O₂-induced neurotoxicity can be prevented by catalase, which is known to be effective at degrading H₂O₂ at higher concentrations and is located in neuron-ensheathing cells (Ho et al., 2004). In this sense, the localization of ROS-scavenging enzymes in TG is strategically important in protecting TG neurons from ROS-mediated cellular injury under the pathological conditions.

CONCLUSION

The present study has determined the cellular localization of antioxidant enzymes in TG. In terms of the decomposition of O₂⁻ and peroxide, TG neurons are endowed with more potent antioxidant systems compared to SGCs. Catalase, predominantly located in SGCs, may play an important role in degrading large quantities of H₂O₂ generated in pathological conditions. We also have found that excess TRPV1 stimulation can cause toxicity to TG neurons via ROS production, which implies that therapeutic measures for antioxidative stress should be taken to prevent damage to trigeminal primary sensory neurons in inflammatory pain disorders.

Acknowledgements—This research was supported in part by a Grant-in-Aid for Scientific Research (Grant No. 22390182 to N.S.) from the Ministry of Education, Culture, Sports, Science and Technology of Japan, and a grant from the Takeda Science Foundation to M.S. There are no conflicts of interest concerning this paper.

REFERENCES

- Abbott CA, Malik RA, van Ross ER, Kulkarni J, Boulton AJ (2011) Prevalence and characteristics of painful diabetic neuropathy in a large community-based diabetic population in the U.K.. *Diabetes Care* 34:2220–2224.
- Acharjee S, Noorbakhsh F, Stemkowski PL, Olechowski C, Cohen EA, Ballanyi K, Kerr B, Pardo C, Smith PA, Power C (2010) HIV-1 viral protein R causes peripheral nervous system injury associated with *in vivo* neuropathic pain. *FASEB J* 24:4343–4353.
- Amantini C, Mosca M, Nabissi M, Lucciarini R, Caprodossi S, Arcella A, Giangaspero F, Santoni G (2007) Capsaicin-induced apoptosis

- of glioma cells is mediated by TRPV1 vanilloid receptor and requires p38 MAPK activation. *J Neurochem* 102:977–990.
- Bell KF, Hardingham GE (2011) CNS peroxiredoxins and their regulation in health and disease. *Antioxid Redox Signal* 14:1467–1477.
- Benediktsson AM, Marrs GS, Tu JC, Worley PF, Rothstein JD, Bergles DE, Dailey ME (2012) Neuronal activity regulates glutamate transporter dynamics in developing astrocytes. *Glia* 60:175–188.
- Britsch S, Goerich DE, Riethmacher D, Peirano RI, Rossner M, Nave KA, Birchmeier C, Wegner M (2001) The transcription factor Sox10 is a key regulator of peripheral glial development. *Genes Dev* 15:66–78.
- Burdo TH, Orzechowski K, Knight HL, Miller AD, Williams K (2012) Dorsal root ganglia damage in SIV-infected rhesus macaques: an animal model of HIV-induced sensory neuropathy. *Am J Pathol* 180:1362–1369.
- Callaghan BC, Cheng HT, Stables CL, Smith AL, Feldman EL (2012) Diabetic neuropathy: clinical manifestations and current treatments. *Lancet Neurol* 11:521–534.
- Camprubi-Robles M, Planells-Cases R, Ferrer-Montiel A (2009) Differential contribution of SNARE-dependent exocytosis to inflammatory potentiation of TRPV1 in nociceptors. *FASEB J* 23:3722–3733.
- Caterina MJ, Leffler A, Malmberg AB, Martin WJ, Trafton J, Petersen-Zeitz KR, Koltzenburg M, Basbaum AI, Julius D (2000) Impaired nociception and pain sensation in mice lacking the capsaicin receptor. *Science* 288:306–313.
- Ceruti S, Villa G, Fumagalli M, Colombo L, Magni G, Zanardelli M, Fabbretti E, Verderio C, van den Maagdenberg AM, Nistri A, Abbracchio MP (2011) Calcitonin gene-related peptide-mediated enhancement of purinergic neuron/glia communication by the algogenic factor bradykinin in mouse trigeminal ganglia from wild-type and R192Q Cav2.1 Knock-in mice: implications for basic mechanisms of migraine pain. *J Neurosci* 31:3638–3649.
- Chi X, Amet T, Byrd D, Chang KH, Shah K, Hu N, Grantham A, Hu S, Duan J, Tao F, Nicol G, Yu Q (2011) Direct effects of HIV-1 Tat on excitability and survival of primary dorsal root ganglion neurons: possible contribution to HIV-1-associated pain. *PLoS One* 6:e24412.
- Chuang HH, Lin S (2009) Oxidative challenges sensitize the capsaicin receptor by covalent cysteine modification. *Proc Natl Acad Sci U S A* 106:20097–20102.
- Crack PJ, Taylor JM, Flentjar NJ, de Haan J, Hertzog P, Iannello RC, Kola I (2001) Increased infarct size and exacerbated apoptosis in the glutathione peroxidase-1 (Gpx-1) knockout mouse brain in response to ischemia/reperfusion injury. *J Neurochem* 78:1389–1399.
- Czaja K, Burns GA, Ritter RC (2008) Capsaicin-induced neuronal death and proliferation of the primary sensory neurons located in the nodose ganglia of adult rats. *Neuroscience* 154:621–630.
- Daousi C, MacFarlane IA, Woodward A, Nurmikko TJ, Bundred PE, Benbow SJ (2004) Chronic painful peripheral neuropathy in an urban community: a controlled comparison of people with and without diabetes. *Diabet Med* 21:976–982.
- Davis JB, Gray J, Gunthorpe MJ, Hatcher JP, Davey PT, Overend P, Harries MH, Latcham J, Clapham C, Atkinson K, Hughes SA, Rance K, Grau E, Harper AJ, Pugh PL, Rogers DC, Bingham S, Randall A, Sheardown SA (2000) Vanilloid receptor-1 is essential for inflammatory thermal hyperalgesia. *Nature* 405:183–187.
- Dringen R, Pawlowski PG, Hirrlinger J (2005) Peroxide detoxification by brain cells. *J Neurosci Res* 79:157–165.
- Flentjar NJ, Crack PJ, Boyd R, Malin M, de Haan JB, Hertzog P, Kola I, Iannello R (2002) Mice lacking glutathione peroxidase-1 activity show increased TUNEL staining and an accelerated inflammatory response in brain following a cold-induced injury. *Exp Neurol* 177:9–20.
- Forman HJ, Maiorino M, Ursini F (2010) Signaling functions of reactive oxygen species. *Biochemistry* 49:835–842.
- Franklin JL (2011) Redox regulation of the intrinsic pathway in neuronal apoptosis. *Antioxid Redox Signal* 14:1437–1448.
- Friedlander RM (2003) Apoptosis and caspases in neurodegenerative diseases. *N Engl J Med* 348:1365–1375.
- Hacimuftuoglu A, Handy CR, Goettl VM, Lin CG, Dane S, Stephens Jr RL (2006) Antioxidants attenuate multiple phases of formalin-induced nociceptive response in mice. *Behav Brain Res* 173:211–216.
- Hanani M (2010) Satellite glial cells: more than just 'rings around the neuron'. *Neuron Glia Biol* 6:1–2.
- Ho YS, Xiong Y, Ma W, Spector A, Ho DS (2004) Mice lacking catalase develop normally but show differential sensitivity to oxidant tissue injury. *J Biol Chem* 279:32804–32812.
- Honda K, Kitagawa J, Sessle BJ, Kondo M, Tsuboi Y, Yonehara Y, Iwata K (2008) Mechanisms involved in an increment of multimodal excitability of medullary and upper cervical dorsal horn neurons following cutaneous capsaicin treatment. *Mol Pain* 4:59.
- Ibi M, Matsuno K, Shiba D, Katsuyama M, Iwata K, Takechi T, Nakagawa T, Sango K, Shirai Y, Yokoyama T, Kaneko S, Saito N, Yabe-Nishimura C (2008) Reactive oxygen species derived from NOX1/NADPH oxidase enhance inflammatory pain. *J Neurosci* 28:9486–9494.
- Ip SW, Lan SH, Lu HF, Huang AC, Yang JS, Lin JP, Huang HY, Lien JC, Ho CC, Chiu CF, Wood W, Chung JG (2012) Capsaicin mediates apoptosis in human nasopharyngeal carcinoma NPC-TW 039 cells through mitochondrial depolarization and endoplasmic reticulum stress. *Hum Exp Toxicol* 31:539–549.
- Jin HW, Ichikawa H, Fujita M, Yamaai T, Mukae K, Nomura K, Sugimoto T (2005a) Involvement of caspase cascade in capsaicin-induced apoptosis of dorsal root ganglion neurons. *Brain Res* 1056:139–144.
- Jin HW, Ichikawa H, Nomura K, Mukae K, Terayama R, Yamaai T, Sugimoto T (2005b) Activation of the caspase cascade underlies the rat trigeminal primary neuronal apoptosis induced by neonatal capsaicin administration. *Arch Histol Cytol* 68:301–310.
- Johnson RM, Ho YS, Yu DY, Kuypers FA, Ravindranath Y, Goyette GW (2010) The effects of disruption of genes for peroxiredoxin-2, glutathione peroxidase-1, and catalase on erythrocyte oxidative metabolism. *Free Radic Biol Med* 48:519–525.
- Katagiri A, Shinoda M, Honda K, Toyofuku A, Sessle BJ, Iwata K (2012) Satellite glial cell P2Y12 receptor in the trigeminal ganglion is involved in lingual neuropathic pain mechanisms in rats. *Mol Pain* 8:23.
- Khairatkar-Joshi N, Szallasi A (2009) TRPV1 antagonists: the challenges for therapeutic targeting. *Trends Mol Med* 15:14–22.
- Khalil Z, Liu T, Helme RD (1999) Free radicals contribute to the reduction in peripheral vascular responses and the maintenance of thermal hyperalgesia in rats with chronic constriction injury. *Pain* 79:31–37.
- Kim HK, Park SK, Zhou JL, Tagliatalata G, Chung K, Coggeshall RE, Chung JM (2004) Reactive oxygen species (ROS) play an important role in a rat model of neuropathic pain. *Pain* 111:116–124.
- Kim HK, Zhang YP, Gwak YS, Abdi S (2010) Phenyl N-tert-butyl nitron, a free radical scavenger, reduces mechanical allodynia in chemotherapy-induced neuropathic pain in rats. *Anesthesiology* 112:432–439.
- Kuhad A, Sharma S, Chopra K (2008) Lycopene attenuates thermal hyperalgesia in a diabetic mouse model of neuropathic pain. *Eur J Pain* 12:624–632.
- Lee I, Kim HK, Kim JH, Chung K, Chung JM (2007) The role of reactive oxygen species in capsaicin-induced mechanical hyperalgesia and in the activities of dorsal horn neurons. *Pain* 133:9–17.
- Mao YF, Yan N, Xu H, Sun JH, Xiong YC, Deng XM (2009) Edaravone, a free radical scavenger, is effective on neuropathic pain in rats. *Brain Res* 1248:68–75.
- Marin-Valencia I, Good LB, Ma Q, Duarte J, Bottiglieri T, Sinton CM, Heilig CW, Pascual JM (2012) Glut1 deficiency (G1D): epilepsy and metabolic dysfunction in a mouse model of the most common human phenotype. *Neurobiol Dis* 48:92–101.

- Maro GS, Vermeren M, Voiculescu O, Melton L, Cohen J, Charnay P, Topilko P (2004) Neural crest boundary cap cells constitute a source of neuronal and glial cells of the PNS. *Nat Neurosci* 7:930–938.
- McCommis KS, McGee AM, Laughlin MH, Bowles DK, Baines CP (2011) Hypercholesterolemia increases mitochondrial oxidative stress and enhances the MPT response in the porcine myocardium: beneficial effects of chronic exercise. *Am J Physiol Regul Integr Comp Physiol* 301: R1250–R1258.
- McCubrey JA, Lahair MM, Franklin RA (2006) Reactive oxygen species-induced activation of the MAP kinase signaling pathways. *Antioxid Redox Signal* 8:1775–1789.
- Mika J, Osikowicz M, Makuch W, Przewlocka B (2007) Minocycline and pentoxifylline attenuate allodynia and hyperalgesia and potentiate the effects of morphine in rat and mouse models of neuropathic pain. *Eur J Pharmacol* 560:142–149.
- Mitozo PA, de Souza LF, Loch-Neckel G, Flesch S, Maris AF, Figueiredo CP, Dos Santos AR, Farina M, Dafre AL (2011) A study of the relative importance of the peroxiredoxin-, catalase-, and glutathione-dependent systems in neural peroxide metabolism. *Free Radic Biol Med* 51:69–77.
- Moskowitz MA (1984) The neurobiology of vascular head pain. *Ann Neurol* 16:157–168.
- Nagai T, Iyata K, Park ES, Kubota M, Mikoshiba K, Miyawaki A (2002) A variant of yellow fluorescent protein with fast and efficient maturation for cell-biological applications. *Nat Biotechnol* 20:87–90.
- Namura S, Zhu J, Fink K, Endres M, Srinivasan A, Tomaselli KJ, Yuan J, Moskowitz MA (1998) Activation and cleavage of caspase-3 in apoptosis induced by experimental cerebral ischemia. *J Neurosci* 18:3659–3668.
- Patenaude A, Murthy MR, Mirault ME (2005) Emerging roles of thioredoxin cycle enzymes in the central nervous system. *Cell Mol Life Sci* 62:1063–1080.
- Power JH, Blumbergs PC (2009) Cellular glutathione peroxidase in human brain: cellular distribution, and its potential role in the degradation of Lewy bodies in Parkinson's disease and dementia with Lewy bodies. *Acta Neuropathol* 117:63–73.
- Rappaport ZH, Devor M (1994) Trigeminal neuralgia: the role of self-sustaining discharge in the trigeminal ganglion. *Pain* 56:127–138.
- Raza H (2011) Dual localization of glutathione S-transferase in the cytosol and mitochondria: implications in oxidative stress, toxicity and disease. *FEBS J* 278:4243–4251.
- Salvemini D, Little JW, Doyle T, Neumann WL (2011) Roles of reactive oxygen and nitrogen species in pain. *Free Radic Biol Med* 51:951–966.
- Schwartz ES, Lee I, Chung K, Chung JM (2008) Oxidative stress in the spinal cord is an important contributor in capsaicin-induced mechanical secondary hyperalgesia in mice. *Pain* 138:514–524.
- Shen HM, Liu ZG (2006) JNK signaling pathway is a key modulator in cell death mediated by reactive oxygen and nitrogen species. *Free Radic Biol Med* 40:928–939.
- Shibata M, Hisahara S, Hara H, Yamawaki T, Fukuuchi Y, Yuan J, Okano H, Miura M (2000) Caspases determine the vulnerability of oligodendrocytes in the ischemic brain. *J Clin Invest* 106:643–653.
- Shibata S, Yasuda A, Renault-Mihara F, Suyama S, Katoh H, Inoue T, Inoue YU, Nagoshi N, Sato M, Nakamura M, Akazawa C, Okano H (2010) Sox10-Venus mice: a new tool for real-time labeling of neural crest lineage cells and oligodendrocytes. *Mol Brain* 3:31.
- Shin CY, Shin J, Kim BM, Wang MH, Jang JH, Surh YJ, Oh U (2003) Essential role of mitochondrial permeability transition in vanilloid receptor 1-dependent cell death of sensory neurons. *Mol Cell Neurosci* 24:57–68.
- Takeda M, Takahashi M, Matsumoto S (2009) Contribution of the activation of satellite glia in sensory ganglia to pathological pain. *Neurosci Biobehav Rev* 33:784–792.
- Thiemermann C (2003) Membrane-permeable radical scavengers (tempol) for shock, ischemia–reperfusion injury, and inflammation. *Crit Care Med* 31:S76–84.
- Tominaga M, Caterina MJ, Malmberg AB, Rosen TA, Gilbert H, Skinner K, Raumann BE, Basbaum AI, Julius D (1998) The cloned capsaicin receptor integrates multiple pain-producing stimuli. *Neuron* 21:531–543.
- Toyokuni S, Tanaka T, Hattori Y, Nishiyama Y, Yoshida A, Uchida K, Hiai H, Ochi H, Osawa T (1997) Quantitative immunohistochemical determination of 8-hydroxy-2'-deoxyguanosine by a monoclonal antibody N45.1: its application to ferric nitrilotriacetate-induced renal carcinogenesis model. *Lab Invest* 76:365–374.
- Wang T, Arifoglu P, Ronai Z, Tew KD (2001) Glutathione S-transferase P1-1 (GSTP1-1) inhibits c-Jun N-terminal kinase (JNK1) signaling through interaction with the C terminus. *J Biol Chem* 276:20999–21003.
- Yowtak J, Lee KY, Kim HY, Wang J, Kim HK, Chung K, Chung JM (2011) Reactive oxygen species contribute to neuropathic pain by reducing spinal GABA release. *Pain* 152:844–852.
- Yuan J, Yankner BA (2000) Apoptosis in the nervous system. *Nature* 407:802–809.
- Zhang J, Graham DG, Montine TJ, Ho YS (2000) Enhanced N-methyl-4-phenyl-1,2,3,6-tetrahydropyridine toxicity in mice deficient in CuZn-superoxide dismutase or glutathione peroxidase. *J Neuropathol Exp Neurol* 59:53–61.

(Accepted 1 June 2013)
(Available online 15 June 2013)