研究成果の刊行に関する一覧表

書籍(日本語)

	T	I	T	I		1	-
著者氏名	論文タイトル名	書籍全体 の 編集者名	書籍名	出版社名	出版地	出版年	ページ
的場元弘	疼痛マネジメント	日本緩和 医療薬学 会	緩和医療薬学	株式会社南江堂	東京	2013	12-14
的場元弘鳥 越一宏	第3章 薬による痛 み治療の基本戦略	World Health Organizat ion	WHOガイドラ イン 病態に 起因した小児 の持続性の痛 みの薬による 治療	金原出版株式会社	東京	2013	41-58
的場元弘鳥 越一宏	第4章 保健医療機 関網における痛み治 療へのアクセス改善 を目指して	World Health Organizat ion	WHOガイドラ イン 病態に 起因した小児 の持続性の痛 みの薬による 治療	金原出版株式会社	東京	2013	59-66
小川朝生	癌患者の心理的反 応・サイコオンコロ ジー	小川修、 田裕作、 井陽郎、 田 新一、 松 田 一、 、 、 、 、 、 、 、 、 、 、 、 、 、 、 、 、 、	ベッドサイド 泌尿器科学改 定第4版	南江堂	東京	2013	617-20
小川朝生	意識障害(せん妄)	日本緩和 医療薬学 会	緩和医療薬学	南江堂	東京	2013	80-1
小川朝生	がん領域における抑 うつの現状と対応	村松公美 子、伊藤弘 人	身体疾患患者 精神的支援ス トラテジー	NOVA 出版	東京	2013	23-7
小川朝生	入院患者の不眠に注 意	小川修、谷 口充孝	内科医のため の不眠診療は じめの一歩	羊土社	東京	2013	27-32
小川朝生	せん妄を発症する疑 いがある場合	小川修、谷 口充孝	内科医のため の不眠診療は じめの一歩	羊土社	東京	2013	156-7

小川朝生	せん妄になってしま った場合	小川修、谷 口充孝	内科医のため の不眠診療は	羊土社	東京	2013	158-60
			じめの一歩				
明智龍男	がん患者の抑うつの		NAGOYA		名古屋	2013	51-55
	評価と治療.		MEDICAL				
			JOURNAL				
明智龍男	一般身体疾患による	山口物 北	今日の治療指	医学書院	東京	2013	868
<u>마이터 RE 기</u>	気分障害	原光夫,福		区于自机	木小	2013	000
	メルド古	井次矢	亚				
	w+++	(編)	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	> <u>></u>			
明智龍男	精神症状マネジメン	日本緩和	緩和医療薬学	南江堂	東京	2013	79
	ト概論	医療薬学					
		会 (編)					
宮下光令	ナーシング・グラフ	宮下光令	緩和ケア	メディカ	大阪	2013	
(編集), 森	ィカ成人看護学	(編集),		出版			
田達也(医		森田達也					
学監修),他		(医学監					
,		修),他					
日本緩和医		日本緩和	終末期がん患	金原出版	東京	2013	
療学会緩		医療学会	者の輸液療法	株式会社	>/>,	20.0	
和医療ガイ		緩和医療	に関するガイ	1/1.2VZ IZ			
ドライン作		ガイドラ	ドライン				
成委員会		イン作成	2013年版				
<u> 双安貝云</u>			2013 11 71X				
本田法山	短切 ケフ並ひのも は	委員会	ナラルタラルダギロ	/B+\ [] +	+-	0040	00.07
森田達也	緩和ケア普及のため	(財)日本	ホスピス緩和	(財)日本	東京	2013	28-37
	の地域プロジェクト	ホスピ	ケア白書2013	ホスピ			
	(OPTIM-study.)か			ス・緩和ケ			
	ら得られたものをど	ア研究振		ア研究振			
	う生かすか	興財団「ホ		興財団			
		スピス緩					
		和ケア白					
		書」編集委					
		員会					
木澤義之,		木澤義之.	3ステップ実	青海社	東京	2013	
森田達也,			践緩和ケア				
他		他					
日本アプラ			2ページで理	南山堂	東京	2013	
イド・セラ			解する標準薬		214731	•	
ピューティ		ラピュー	物治療ファイ				
<u>しューティ</u> クス学会		ティクス	がたまったり				
ノヘチム			10				
		学会					

雑誌 (外国語)

翌 ≢老爪々	論文タイトル名	発表誌名	巻号	ページ	出版年
発表者氏名 Ogawa Y <u>, Hotta T</u> , et al.	bortezomib-melphalan-prednis olone for previously untreated Japanese patients with	Cancer Sci	104(7)	912-919	2013
Ogura M, Itoh K, Ishizawa K, Kobayashi Y, Tobinai K, Kinoshita T, Hirano M, Ueda R, Shibata T, Nakamura S, Tsukasaki K, Hotta T, Shimoyama M, Morishima Y	Hodgkin lymphoma: Japan Clinical Oncology Group study,	Leukemia & Lymphoma	54(1)	46-52 DOI: 10.3109/1 0428194.2 012.70500 0	2013
Sasahara T,_ <u>Kinoshita H</u> , <u>Morita T</u> , et al	Assessment of reasons for referral and activities of hospital palliative care teams using a standard format: A multicenter 1000 case description.	J Pain Symptom Manage	Aug 21	[epub ahead of print]	2013
Imura C, <u>Morita T</u> , <u>Kinoshita H</u> , et al	How and why did a regional palliative care program lead to changes in region? A qualitative analysis of the Japan OPTIM Study.	J Pain Symptom Manage	Aug 24	[epub ahead of print]	2013
Morita T, Sato K, Miyashita M, Akiyama M, Kato M, Kawagoe S, Kinoshita H, Shirahige Y, Yamakawa S, Yamada M, Eguchi K.	Exploring the perceived changes and the reasons why expected outcomes were not obtained in individual levels in a successful regional palliative care intervention	Support Care Cancer.	Dec;21 (12)	3393-402	2013
Asai M, Akechi T, Kinoshita H, Uchitomi Y, et al	Impaired mental health among the bereaved spouses of cancer patients	Psychooncology	22(5)	995-1001	2013
Inagaki M, <u>Akechi</u> <u>T, Kinoshita H,</u> <u>Uchitomi Y</u> , et al	Associations of interleukin-6 with vegetative but not affective depressive symptoms	Cancer	21(8)	2097-2106	2013

Description			T		, ,	
Yamaguchi T, Shima Clinical Guideline for Y, Morita T, Hosoya Clinical Guideline for Pharmacological Management of Cancer Pain: The Japanese Society of Palliative Medicine Recommendations. James Pain The Japanese Society of Palliative Medicine Recommendations. Palliat Med 27(2) 179-184 2013 Komura K, Kawagoe S, Morita T, et al S, Morita T, et al Standard A, Morita T, Miyashita M, et al Patient-perceived usefulness of patient-held records for cancer patients in Japan: OPTIM Study. J Palliat Med 27(2) 179-184 2013 Shirado A, Morita T, Miyashita M, et al Usefulness of the left-based intervention for family members of terminally cancer patients with delirium. J Pain Symptom Manage 45(5) 848-858 2013 Shirado A, Morita T, Miyashita M, et al Palliative care in Japan: such with cancer: a mixed-methods study. J Pain Symptom Manage 45(5) 848-858 2013 Morita T, Miyashita M, et al Palliative care in Japan: a mixed-methods study. Lancet Oncol 14(7) 638-646 2013 Kunieda K, Morita T, et al Specialized palliative care services in Japan: a nationwide survey of resources and utilization by patients with cancer: a mixed-methods study. Am J Hosp Palliat Med 30(6) 552-555 2013 Kizawa Y, Morita T, et al Pspecialized palliative care services in Japa		in terminally ill cancer				
V, Morita T, Hosoya M, Matoba M. Pharmacological Management of Cancer Pain: The Japanese Society of Palliative Medicine Recommendations. Patient Perceived usefulness Society of Palliative Medicine Recommendations. Komura K, Kawagoo S, Morita T, et al Patient -perceived usefulness of patient-held records for cancer patients in Japan: OPTIM study. Palliat Med 27(2) 179-184 2013 Otani H, Morita T, et al Usefulness of the leaflet-based intervention for family members of terminally cancer patients with delirium. J Palliat Med 16(4) 419-22 2013 Shirado A, Morita T, Miyashita M, et al Both maintaining hope and preparing for death: Effects of physicians' and nurses behaviors from bereaved family members' perspectives. J Pain Symptom Manage 45(5) 848-858 2013 Morita T, et al Palliative care in Japan: a review focusing on care delivery system. Curr Opin Support Palliat Care Curr Opin Support Palliat Care Care Lancet Oncol 14(7) 638-646 2013 M, et al mixed-methods study. Reliability and validity of a J Pain Symptom to patients with cancer: a mixed-methods study. Am J Hosp Palliat 30(6) 552-555 2013 Kunieda K, Morita T, Spicalized palliative care services in Japan: a nationwide survey of resources and utilization by patients with cancer. Am J Hosp Palliat 30(6) 552-555 2013 <td></td> <td>patients</td> <td></td> <td></td> <td></td> <td></td>		patients				
M, Matoba M. Cancer Pain: The Japanese Society of Palliative Medicine Recommendations. Komura K, Kawagoe S, Morita T, et al patient-perceived usefulness and practical obstacles of patient-led records for cancer patients in Japan: OPTIM study. Otani H, Morita T, et al leaflet-based intervention for family members of terminally cancer patients with delirium. Shirado A, Morita Both maintaining hope and preparing for death: Effects of physicians' and nurses' behaviors from bereaved family members' perspectives. Morita T, et al Palliative care in Japan: a review focusing on care delivery system. Morita T, Miyashita Effects of a programme of interventions on regional comprehensive palliative care for patients with cancer: a mixed-methods study. Kunieda K, Morita T, et al Colt measure the severity of dysphagia: The food intake LEVEL scale. Kizawa Y, Morita T, Specialized palliative care and utilization by patients with cancer. Kizawa Y, Morita T, Miyashita M, et al Services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Morita T, et al agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Palliat twedions Palliat Support 11(5) 383-388 2013	Yamaguchi T, Shima	Clinical Guideline for	Jpn J Clin Oncol.	43(9)	896-909	2013
Society of Palliative Medicine Recommendations. Recommendations. Palliat Med P	Y, <u>Morita T</u> , Hosoya	Pharmacological Management of				
Recommendations. Patient-perceived usefulness Palliat Med 27(2) 179-184 2013	M, <u>Matoba M.</u>	Cancer Pain: The Japanese				
Romura K, Kawagoe S, Morita T, et al and practical obstacles of patient-held records for cancer patients in Japan: OPTIM study.		Society of Palliative Medicine				
S, Morita T, et al and practical obstacles of patient-held records for cancer patients in Japan: OPTIM study. Otani H, Morita T, et al leaflet-based intervention for family members of terminally cancer patients with delirium. Shirado A, Morita Both maintaining hope and T, Miyashita M, et al preparing for death: Effects of physicians' and nurses' behaviors from bereaved family members' perspectives. Morita T, et al Palliative care in Japan: a review focusing on care delivery system. Morita T, Miyashita Effects of a programme of interventions on regional comprehensive palliative care for patients with cancer: a mixed-methods study. Kunieda K, Morita T, Specialized palliative care for patients with cancer: a nationwide survey of resources and utilization by patients with cancer. Kizawa Y, Morita T, Specialized palliative care antionwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Poshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013		Recommendations.				
patient-held records for cancer patients in Japan: OPTIM study. Otani H, Morita T, et al	Komura K, <u>Kawagoe</u>	Patient-perceived usefulness	Palliat Med	27(2)	179-184	2013
cancer patients in Japan: OPTIM study. Otani H, Morita T, et al Morita T, et al Morita T, Miyashita M, et al Morita T, Miyashita M, et al Morita T, et al Morita T, et al Morita T, et al Morita T, Miyashita M, et al Morita T, et al Morita T, et al Morita T, et al Morita T, Miyashita M, et al Morita T, et al Morita T, Miyashita Effects of a programme of interventions on regional comprehensive palliative care for patients with cancer: a mixed-methods study. Kunieda K, Morita T, et al Kizawa Y, Morita T, Specialized palliative care for patients with cancer: a nationwide survey of resources and utilization by patients with cancer. Miyashita M, et al Kizawa Y, Morita T, Specialized palliative care anationwide survey of resources and utilization by patients with cancer. Kizawa Y, Morita T, Specialized palliative care anationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Morita T, et al Morita T, et al Morita T, et al Morita T, Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013	S, Morita T, et al	and practical obstacles of				
OPTIM study. Otani H, Morita T, et al		patient-held records for				
Otani H, Morita T, et al Shirado A, Morita T, et al Shirado A, Morita T, Miyashita M, et al Shirado A, Morita T, Miyashita M, et al Shirado A, Morita T, et al Shirado A, Morita T, Miyashita M, et al Shirado A, Morita T, et al Shirado A, Morita T, Miyashita M, et al Shirado A, Morita T, et al Shirado A, Morita T, Miyashita M, et al Shirado A, Morita T, et al Shirado A, Morita T, Morita T, Morita T, et al Shirado A, Morita T, et al Shirado A, Morita T, Morita T, Morita T, et al Shirado A, Morita T, Specialized palliative care and utilization by patients with cancer. Kanbayashi Y, Morita T, Morita T, et al Shirado A, Morita T, et al Shirado		cancer patients in Japan:				
Part		OPTIM study.				
for family members of terminally cancer patients with delirium. Shirado A, Morita T, Miyashita M, et al morita T, et al Palliative care in Japan: a review focusing on care delivery system. Morita T, Miyashita Effects of a programme of interventions on regional comprehensive palliative care for patients with cancer: a mixed-methods study. Kunieda K, Morita T, et al Palliative care the severity of dysphagia: The food intake LEVEL scale. Kizawa Y, Morita T, Morita T, Miyashita M, et al Practices and utilization by patients with cancer. Kanbayashi Y, Morita T, et al Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. For family members of terminally cancer patients with adiractions with adiractions of Palliat Symptom 45(5) 848-858 2013 J Pain Symptom 45(5) 848-858 2013 Curr Opin Support Palliat Care Out Opin Support Palliat Care Oncol 14(7) 638-646 2013 T, et al Symptom 46(2) 207-215 2013 Lancet Oncol 14(7) 638-646 2013 The food intake Level and validity of a tool to measure the severity of dysphagia: The food intake Level and the severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013	Otani H, <u>Morita T</u> ,	Usefulness of the	J Palliat Med	16(4)	419-22	2013
terminally cancer patients with delirium. Shirado A, Morita T, Miyashita M, et al Morita T, et al Morita T, Miyashita M, et al Morita T, Miyashita M, et al Morita T, et al Morita T, Miyashita M, et al Morita T, et al Morita T, Miyashita M, et al Morita T, Miyashita M, et al Morita T, et al Morita T, Miyashita M, et al Morita T, Morita T, Miyashita M, et al Morita T, Morita T, Miyashita M, et al Morita T, et al Miyashita M, et al Morita T, Morita T, Miyashita M, et al Morita T, Morita T, Miyashita M, et al Morita T, et al Morita T, et al Morita T, et al Morita T, Predictive factors for an antionwide survey of resources and utilization by patients with cancer. Morita T, et al Morita T, Predictive factors for antionwide survey of resources and utilization by patients with cancer. Morita T, et al	et al	leaflet-based intervention				
Shirado A, Morita Both maintaining hope and T, Miyashita M, et al Sof physicians ' and nurses' behaviors from bereaved family members' perspectives.		for family members of				
Shirado A, Morita T, Miyashita M, et al Palliative care in Japan: a review focusing on care delivery system. Morita T, Miyashita M, et al Palliative care in Japan: a review focusing on care delivery system. Morita T, Miyashita M, et al Palliative care in Japan: a review focusing on care delivery system. Morita T, Miyashita Effects of a programme of interventions on regional comprehensive palliative care for patients with cancer: a mixed-methods study. Kunieda K, Morita T, et al tool to measure the severity of dysphagia: The food intake LEVEL scale. Kizawa Y, Morita T, Miyashita M, et al services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Morita T, et al services in Japan: a nationwide survey of resources and utilization severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Practices and evaluations of Palliat Support 11(5) 383-388 2013		terminally cancer patients				
T, Miyashita M, et al preparing for death: Effects of physicians ' and nurses' behaviors from bereaved family members' perspectives. Morita T, et al Palliative care in Japan: a review focusing on care delivery system. Morita T, Miyashita M, et al Effects of a programme of interventions on regional comprehensive palliative care for patients with cancer: a mixed-methods study. Kunieda K, Morita T, et al Eliability and validity of a tool to measure the severity of dysphagia: The food intake LEVEL scale. Kizawa Y, Morita T, Specialized palliative care services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita P, Practices and evaluations of Palliat Support 11(5) 383-388 2013		with delirium.				
T, Miyashita M, et al preparing for death: Effects of physicians ' and nurses' behaviors from bereaved family members' perspectives. Morita T, et al Palliative care in Japan: a review focusing on care delivery system. Morita T, Miyashita M, et al Effects of a programme of interventions on regional comprehensive palliative care for patients with cancer: a mixed-methods study. Kunieda K, Morita T, et al Eliability and validity of a tool to measure the severity of dysphagia: The food intake LEVEL scale. Kizawa Y, Morita T, Specialized palliative care services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita P, Practices and evaluations of Palliat Support 11(5) 383-388 2013	Shirado A, Morita	Both maintaining hope and	J Pain Symptom	45(5)	848-858	2013
al of physicians ' and nurses ' behaviors from bereaved family members' perspectives. Morita T, et al Palliative care in Japan: a review focusing on care delivery system. Morita T, Miyashita Effects of a programme of interventions on regional comprehensive palliative care for patients with cancer: a mixed-methods study. Kunieda K, Morita Reliability and validity of a tool to measure the severity of dysphagia: The food intake LEVEL scale. Kizawa Y, Morita T, Miyashita M, et al Services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013		preparing for death: Effects		, ,		
behaviors from bereaved family members ' perspectives. Morita T, et al Palliative care in Japan: a review focusing on care delivery system. Morita T, Miyashita Effects of a programme of interventions on regional comprehensive palliative care for patients with cancer: a mixed-methods study. Kunieda K, Morita T, et al Reliability and validity of a tool to measure the severity of dysphagia: The food intake LEVEL scale. Kizawa Y, Morita T, Specialized palliative care services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013						
Morita T, et al						
Morita T, et al						
review focusing on care delivery system. Morita T, Miyashita Effects of a programme of interventions on regional comprehensive palliative care for patients with cancer: a mixed-methods study. Kunieda K, Morita Reliability and validity of a tool to measure the severity of dysphagia: The food intake LEVEL scale. Kizawa Y, Morita T, Specialized palliative care services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013	Morita T, et al		Curr Opin	7(2)	207-215	2013
delivery system. Morita T, Miyashita Effects of a programme of interventions on regional comprehensive palliative care for patients with cancer: a mixed-methods study. Kunieda K, Morita Reliability and validity of a tool to measure the severity of dysphagia: The food intake LEVEL scale. Kizawa Y, Morita T, Specialized palliative care services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Morita T, et al Practices and evaluations of Palliat Support 11(5) 383-388 2013			•	()		
Morita T, Miyashita M, et al Effects of a programme of interventions on regional comprehensive palliative care for patients with cancer: a mixed-methods study. Kunieda K, Morita T, et al Effects of a programme of interventions on regional comprehensive palliative care for patients with cancer: a mixed-methods study. Kunieda K, Morita T, et al Eliability and validity of a tool to measure the severity of dysphagia: The food intake LEVEL scale. Kizawa Y, Morita T, Specialized palliative care services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Predictive factors for a gitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013		_				
M, et al interventions on regional comprehensive palliative care for patients with cancer: a mixed-methods study. Kunieda K, Morita T, et al Reliability and validity of a tool to measure the severity of dysphagia: The food intake LEVEL scale. Kizawa Y, Morita T, Specialized palliative care services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013	Morita T, Miyashita	• •	Lancet Oncol	14(7)	638-646	2013
comprehensive palliative care for patients with cancer: a mixed-methods study. Kunieda K, Morita T, et al tool to measure the severity of dysphagia: The food intake LEVEL scale. Kizawa Y, Morita T, Specialized palliative care services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013		I		,		
for patients with cancer: a mixed-methods study. Kunieda K, Morita T, et al		_				
Mixed-methods study. Kunieda K, Morita T, et al Reliability and validity of a tool to measure the severity of dysphagia: The food intake LEVEL scale. Specialized palliative care services in Japan: a nationwide survey of resources and utilization by patients with cancer. Am J Hosp Palliat		-				
Kunieda K, Morita T, et al Reliability and validity of a tool to measure the severity of dysphagia: The food intake LEVEL scale. Kizawa Y, Morita T, Specialized palliative care services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Morita T, et al Services for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013		· ·				
T, et al tool to measure the severity of dysphagia: The food intake LEVEL scale. Kizawa Y, Morita T, Specialized palliative care services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita T tool to measure the severity of dysphagia: The food intake LEVEL scale. Am J Hosp Palliat 30(6) 552-555 2013 Care J Palliat Med 16(9) 1020-1025 2013	Kunieda K, Morita	Reliability and validity of a	J Pain Symptom	46(2)	201-206	2013
dysphagia: The food intake LEVEL scale. Kizawa Y, Morita T, Specialized palliative care services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013	· · · · · · · · · · · · · · · · · · ·	1		,		
LEVEL scale. Kizawa Y, Morita T, Miyashita M, et al services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013		-				
Miyashita M, et al services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013						
Miyashita M, et al services in Japan: a nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013	Kizawa Y, Morita T,	Specialized palliative care	Am J Hosp Palliat	30(6)	552-555	2013
nationwide survey of resources and utilization by patients with cancer. Kanbayashi Y, Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013		·	•	,		
With cancer. Kanbayashi Y, Morita T, et al Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita With cancer. J Palliat Med 16(9) 1020-1025 2013 Palliat Med 16(9) 1020-1025 Palliat Support 11(5) 383-388		•				
With cancer. Kanbayashi Y, Morita T, et al Predictive factors for agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita With cancer. J Palliat Med 16(9) 1020-1025 2013 Palliat Med 16(9) 1020-1025 Palliat Support 11(5) 383-388		and utilization by patients				
Morita T, et al agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013		1				
Morita T, et al agitation severity of hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013	Kanbayashi Y,	Predictive factors for	J Palliat Med	16(9)	1020-1025	2013
hyperactive delirium in terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013	-	agitation severity of		,		
terminally cancer patients in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013	,	,				
in a general hospital using ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013						
ordered logistic regression analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013		,				
analysis. Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013						
Yoshida S, Morita Practices and evaluations of Palliat Support 11(5) 383-388 2013						
· · · · · · · · · · · · · · · · · · ·	Yoshida S, Morita		Palliat Support	11(5)	383-388	2013
	<u>T</u> , et al	prognostic disclosure for		` ,		

		T	T		
	Japanese cancer patients and				
	their families from the				
	family's point of view.				
Imai K, <u>Morita T</u> , et	Sublingually administered	Support Care	21(10)	2777-2781	2013
al	scopolamine for nausea in	Cancer			
	terminally cancer				
	patients.				
Yamamoto R, <u>Morita</u>	The palliative care knowledge	J Palliat Med	16(11)	1423-1428	2013
<u>T</u> , et al	questionnaire for PEACE:				
	Reliability and validity of an				
	instrument to measure				
	palliative care knowledge				
	among physicians.				
Amano K, Morita T,	Effect of nutritional support	Am J Hosp Palliat	30(7)	730-733	2013
et al	on terminally ill patients	Care			
	with cancer in a palliative				
	care unit.				
Igarashi A,	A population-based survey on	Am J Hosp Palliat	Mar 15	[epub	2013
Miyashita M,	perceptions of opioid	Care		ahead of	
Morita T, et al	treatment and palliative care			print]	
	units: OPTIM Study.				
Amano K, <u>Morita T</u> ,	The determinants of patients	Am J Hosp Palliat	Apr 2	[epub	2013
et al	in a palliative care unit being	· ·		ahead of	
	discharged home in Japan.			print]	
Otani H, Morita T,	Effect of leaflet-based	Am J Hosp Palliat	Apr 23	[epub	2013
et al	intervention on family members	·		ahead of	
	of terminally ill patients			print]	
	with cancer having delirium:				
	Historical control study.				
Muta R, <u>Miyashita</u>	What bereavement follow-up	Am J Hosp Palliat	May 16	[epub	2013
	does family members request in	-		ahead of	
	Japanese palliative care			print]	
	units?: A qualitative study.			F	
Ando M, <u>Morita T</u> ,	A pilot study of adaptation of	Am J Hosp Palliat	Mav 20	[epub	2013
Miyashita M, et al	the transtheoretical model to	•	,	ahead of	- · •
,	narratives of bereaved family			print]	
	members in the bereavement			p	
	life review.				
lse Y. Morita T. et	The activity of palliative	J Pain Symptom	Sep 6	[epub	2013
al	care team pharmacists in	Manage	COP C	ahead of	_0.0
	designated cancer hospitals: A	•		print]	
	nationwide survey in Japan.			h 1	
Yamaguchi T. Morita	Pneumocystic pneumonia in	Am J Hos Palliat	Sep 30	[epub	2013
T, et al	patients treated with ong-term		202 00	ahead of	_0.0
<u>-</u> , or a.	steroid therapy for symptom			print]	
	palliation: A neglected			h	
	infection in palliative care.				
	Innoction in partialive cale.				

		T	T	1	
Shimizu Y,	Care strategy for death rattle		0ct 22		2013
<u>Miyashita M</u> ,	in terminally ill cancer	Manage		ahead of	
<u>Morita T</u> , et al	patients and their family			print]	
	members: Recommendations from				
	a cross-sectional nationwide				
	survey of bereaved family				
	members' perceptions.				
Nakaya N, Uchitomi	All-cause mortality among men	Epidemiology	24(1)	96-99	2013
<u>Y</u> , et al	whose cohabiting partner has		, ,		
_,	been diagnosed with cancer				
Terada S, Uchitomi	Person-centered care and	Psychiatry Res	205(1-	103-108	2013
\underline{Y} , et al	quality of life of patients	l sysmatry kos	2)	100 100	2010
	with dementia in long-term				
	care facilities				
Kishimoto Y,	Abuse of people with cognitive	Develiatry Poe	209(3)	699-704	2013
1	1	rsychiatry Nes	209(3)	099-704	2013
<u>Uchitomi Y</u> , et al	impairment by family				
	caregivers in Japan (a				
Namas C. Habitani	cross-sectional study)	I Name I Cai	200/4	70.74	0040
Nagao S, <u>Uchitomi</u>	Progressive supranuclear	J Neurol Sci	329(1-	70-71	2013
\underline{Y} , et al	palsy presenting as primary		2)		
	lateral sclerosis		4->		
Oshima E, <u>Uchitomi</u>	,	Brain Pathol	23(6)	633-644	2013
\underline{Y} , et al	Apoptosis and Neurological				
	Dysfunction Caused by Chronic				
	Oral Administration of				
	Aluminum in a Mouse Model of				
	Tauopathies				
Hayashi S, <u>Uchitomi</u>	Burden of caregivers for	Int	25(8)	1357-1363	2013
<u>Y</u> , et al	patients with mild cognitive	Psychogeriatr			
	impairment in Japan				
Shindo A, <u>Uchitomi</u>	Trail making test part a and	Dement Geriatr	3(1)	E39-40	2013
<u>Y</u> , et al	brain perfusion imaging in	Cogn Dis Extra			
	mild Alzheimer's disease				
Kondo K, Ogawa A,	Characteristics associated	Patient Educ	93(2)	350-353	2013
Uchitomi Y, et al	with empathic behavior in	Couns	, ,		
,	Japanese oncologists				
Terada S, <u>Uchitomi</u>		Psychiatry Res	213(3)	249-255	2013
\underline{Y} , et al	perfusion imaging in mild			- 10 - 200	_0.0
	cognitive impairment and mild				
	Alzheimer's disease				
Fujimori M,	Development and preliminary	Palliat Support			2013
Uchitomi Y, et al	evaluation of communication	Care			2010
oon tom i, et al	skills training program for				
	oncologists based on patient				
	preferences for communicating				
Nogoo C Halaiteani	bad news	Fur Arab			2042
Nagao S, <u>Uchitomi</u>	Argyrophilic grain disease as				2013
Y, et al	a neurodegenerative substrate	rsydmatry Clin			

	in late-onset schizophrenia	Neurosci			
	and delusional disorders				
Terada S, <u>Uchitomi</u> <u>Y</u> , et al	Depressive symptoms and regional cerebral blood flow in Alzheimer's disease	Psychiatry Res	221(1)	86-91	2013
Akechi T, et al.	Assessing medical decision making capacity among cancer patients: Preliminary clinical experience of using a competency assessment instrument.	Palliat Support Care	21	1-5	2013
Fielding R, Akechi T, et al.	Attributing Variance in Supportive Care Needs during Cancer: Culture-Service, and Individual Differences, before Clinical Factors.	PLOS ONE	8 (5)	e65099	2013
Furukawa TA, <u>Akechi</u> <u>T</u> , et al	Cognitive-behavioral therapy modifies the naturalistic course of social anxiety disorder: Findings from an ABA design study in routine clinical practices.	Psychiatry Clin Neurosci	67 (3)	139-47	2013
Kawaguchi A, <u>Akechi</u> <u>T</u> , et al.	Group cognitive behavioral therapy for patients with generalized social anxiety disorder in Japan: outcomes at 1-year follow up and outcome predictors.	Neuropsychiatr Dis Treat	9	267-75	2013
Nakaguchi T, Akechi T, et al.	Oncology nurses' recognition of supportive care needs and symptoms of their patients undergoing chemotherapy.	Jpn J Clin Oncol	43 (4)	369-76	2013
Nakano Y, Akechi T, et al.	Cognitive behavior therapy for psychological distress in patients with recurrent miscarriage.	Psychol Res Behav Manag	6	37-43	2013
Igawa S, <u>Sasaki J</u> , et al.	Evaluation of amrubicin as a Third or Later Line of Chemotherapy for Advanced Non-Small Cell Lung Cancer.	Chemotherapy	59	99-105	2013
Otani S, Hamada A, <u>Sasaki J</u> , et al.	Phase I and Pharmacokinetic Study of Erlotinib Administered in Combination With Amrubicin in Patients With Previously Treated, Advanced Non-Small Cell Lung Cancer.	Am J. Clin Oncol	In press		In press

<u>Fukui S</u> , Yoshiuchi K, Fujita J, Ikezaki S.	Determinants of financial performance of home-visit nursing agencies in Japan.	BMC Health Services Research.	DOI: 10.118 6/1472 -6963- 14-11	オンライ ンジャー ナル	2014
<u>Fukui S</u> , Fujita J, Yoshiuchi K.	Associations between Japanese People's Concern about Family Caregiver Burden and Preference for End-of-Life care Location		29(1)	22-28	2013
Ishikawa Y, <u>Fukui</u> <u>S</u> , Saito T, et al.	Family preference for place of death mediates the relationship between patient preference and actual place of death: A nationwide retrospective cross-sectional study.	PLOS ONE	8 e56848	オンライ ンジャー ナル	2013
Kinoshita S. Miyashita M	Evaluation of end-of-life cancer care in the ICU: perceptions of the bereaved family in Japan	Am J Hosp Palliat Med	30(3)	225-30	2013

雑誌(日本語)

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
	在宅医療におけるがん疼痛治療 の評価と治療の実態について.	Palliat Care Res.	8(1)	158-167	2013
森田達也	せん妄マネジメントの実際とケアの具体策がんによる「せん妄」の原因と出現するメカニズム	がん患者ケア	6(2)	62-66	2013
森田達也	せん妄マネジメントの実際とケアの具体策 「せん妄」の薬物治療とケアの注意点	がん患者ケア	6(3)	67-72	2013
山内敏宏, 森田達也	EAPC (European Association of Palliative Care) 疼痛ガイドラインを読む. 第5回1代替全身投与経路 2突出痛に対するオピオイド	緩和ケア	23(1)	60-63	2013
森田達也	社会の力を最大化する「顔の見える関係」緩和ケアプログラムの地域介入研究(OPTIM-study)	週刊医学界新聞	第 3019 号	4	2013

	を終えて				
	FADO (F	/⊠10 /_ →	00 (0)	101 100	0040
厨芽衣子, 森田達也	EAPC (European Association of	綾杣ケ <i>ソ</i> 	23(2)	161-162	2013
	Palliative Care) 疼痛ガイドラ				
	インを読む. 最終回 1オピオイ				
	ドスイッチング, 2オピオイド力				
// ** * -	価	/= /= - <.1 \\			
佐藤一樹,宮下光		保健の科学	55(4)	230-235	2013
<u>令</u> , <u>森田達也</u> ,他	和ケア) 緩和ケア普及のための				
	地域プロジェクト(1)緩和ケ				
	ア普及のための地域プロジェク				
	トで使用した評価尺度				
森田達也	地域における緩和ケア(在宅緩	保健の科学	55(4)	236-241	2013
	和ケア) 緩和ケア普及のための				
	地域プロジェクト(2)地域プ				
	ロジェクト(OPTIM-study)の効				
	果				
<u>森田達也</u> ,他	「緩和ケアに関する地域連携評	Palliat Care Res	8(1)	116-126	2013
	価尺度」の開発				
本田海山 仙	がん患者のこころのケアと地域	 精神科	22(2)	307-314	2013
<u>森田達也</u> ,他		个月个中个子 	23(3)	307-314	2013
	ネットワーク OPTIM-studyの				
本田法山	知見から	and the trans	50/44	507 504	0040
森田達也	苦痛緩和のための鎮静	medicina	50(11	527-531	2013
			増刊		
******	中央、中央の短行とつの時間	/⊠το - →	号)	407.504	0040
森田達也, 宮下光	患者・遺族の緩和ケアの質評	緩和ケア	23(6)	497-501	2013
<u>令,</u> 他	価・quality of life,医師・看				
	護師の困難感と施設要因との関				
	連				
内富庸介	がん患者の抑うつ対策 医療者	Clinic magazine	524	18-21	2013
	が積極的に抑うつの症状を聞く				
	ことが重要				
井上真一郎, 内富庸	せん妄の要因と予防	精神科治療学	42(3)	289-297	2013
<u>介</u>					
井上真一郎,内富庸	がん診断早期から行うべき緩和	Mebio	30(7)	23-29	2013
	薬物療法の実際・精神的ストレ	INICUTO	30(1)	23-29	2013
<u>介</u>	スの観点から -				
井上真一郎,内富庸	せん妄を見逃さないための注意	 精神科治療学	28(8)	1011-1017	2012
	点	▗▗▗ ▗ ▗ ▗ ▗ ▗ ▗ ▎ ▎ ▎ ▎ ▎ ▎ ▎ ▎ ▎ ▎ ▎ ▎	20(0)	1011-1017	2013
<u>介,</u> 他					
浅井真理子,内富庸	配偶者をがんで亡くした遺族の	心理学研究	84(5)	498-507	2013
<u>介</u> ,他	対処行動パターン				
	大ウダエロケラレン・日子学	口/本のひど	55 (A \	054 057	0040
川越博美	在宅緩和ケアと訪問看護	保健の科学	55(4)	254-257	2013
L	1	1			

川越博美	践力 "とは	訪問看護と介護	18(7)	534-537	2013
	その育成に向けて				
小川朝生	がん領域における精神疾患と緩 和ケアチームの役割	PSYCHIATRIST	18	54-61	2013
小川朝生	一般病棟における精神的ケアの 現状	看護技術	59(5)	422-6	2013
小川朝生	せん妄の予防-エビデンスに基づいた薬物療法と非薬物療法・	緩和ケア	23(3)	196-9	2013
小川朝生	高齢がん患者のこころのケア	精神科	23(3)	283-7	2013
小川朝生	がん患者の終末期のせん妄	精神科治療学	28(9)	1157-62	2013
小川朝生	がん領域における精神心理的ケ アの連携	日本社会精神医 学会雑誌	22(2)	123-30	2013
伊藤嘉規, <u>明智龍男、</u> 他	小児がん患者とその家族のこころのケア.	精神科	23	288-292	2013
明智龍男	がんとこころのケア-サイコオ ンコロジー.	精神科	23	271-275	2013
明智龍男	せん妄の向精神薬による対症療 法と処方計画.	精神科治療学	28	1041-1047	2013
明智龍男	緩和医療とせん妄.	臨床精神医学	42	307-312	2013
明智龍男	術後せん妄.	消化器外科	36	1643-1646	2013
明智龍男	抑うつとがん.	レジデントノー ト	15	2440-2443	2013
明智龍男,森田達也	臨床で役立つサイコオンコロジーの最新エビデンス-特集にあたって.	緩和ケア	23	191	2013
明智龍男	がん患者の自殺に関する最新データ.	緩和ケア	23	195	2013
明智龍男	希死念慮を有する患者のアセス メントとケア.	緩和ケア	23	200	2013
佐々木治一郎	喫煙と健康障害-禁煙支援の理解・普及から「脱タバコ社会」を目指して- タバコ煙の成分と疾患発症メカニズム.	日本臨床	71	383-89	2013
福井 朋也,佐々木治一郎	がんの治療薬を知る 薬剤選択 のための知識 ノギテカン	臨床腫瘍プラク ティス	9	180-83	2013

井川 聡, 大谷 咲 子, 佐々木 治一郎	非小細胞肺癌の維持療法	呼吸	32	332-38	2013
福井小紀子、乙黒千鶴、石川孝子、藤田淳子、秋山正子	都市部公営団地に在住する健康 相談未利用者における健康相談 の必要性に関する認識とその関 連要因の検討.	日本公衆衛生学会誌	60(12	745-753	2013
福井小紀子	訪問看護事業所の黒字化のため の経営指標の提案	社会保険旬報	2545	22-29	2013
藤田淳子、渡辺美奈 子、 <u>福井小紀子</u>	介護支援専門員・介護職に対す る訪問看護師の連携行動とその 関連要因-死亡前1か月の高齢 者終末期ケアに関して-	日本地域看護学会誌	16(1)	40-47	2013
	訪問看護事業所の質管理の取り 組みと関連要因の検討	コミュニティケア	15(11	67-71	2013
子、清水準一、高砂	平成24年度の介護報酬・診療報 酬の改定内容が訪問看護事業所 の経営へ与えた影響	コミュニティケア	15(10	67-74	2013
子、清水準一、高砂裕子、上野 桂子、齋		コミュニティケア	15(9)	60-66	2013
福井小紀子	がん対策推進基本計画見直しと 診療報酬・介護報酬改定の全体 像	緩和ケア	23(2)	104-107	2013